



# Simulation of Biomolecules

## Basic Simulation Analysis



Dr Matteo Degiacomi  
Durham University

[matteo.t.degiacomini@durham.ac.uk](mailto:matteo.t.degiacomini@durham.ac.uk)





Dr Antonia Mey  
University of Edinburgh

[antonia.mey@ed.ac.uk](mailto:antonia.mey@ed.ac.uk)


# Volume and pressure equilibration

Steps until production:

Production ensemble



# Sampling the conformational space





Probability of sampling a conformation is inversely proportional to its energy:  $p_i \propto e^{-\epsilon_i/kT}$

# Timescales in biology

Different regions, different timescales:

- Side chains faster than backbone
- Loops faster than helices and sheets
- backbone faster than side chains
- Protein surface faster than core





# Root Mean Square Deviation (RMSD)

Given a system with  $N$  atoms, and a reference arrangement  $x_0$ :


$$RMSD = \sqrt{\frac{1}{N} \sum_{i=0}^N (X_i - x_0)^2}$$

In MD,  $x_0$  is often the first conformation in the simulation.







# Convergence?







*Refrain from using RMSD as a single indicator of simulation convergence.*




HIV-1 capsomer simulation from: *Degiacomi & Dal Peraro, Structure, 2013*

# RMDS is alignment- and selection-dependent



# Pairwise RMSD

- Two structures with same RMSD from a reference are not forcefully similar to each other.
- Pairwise RMSD helps seeing if protein re-visits conformations throughout the simulation.







# Root Mean Square Fluctuation (RMSF)

The RMSF  $\sigma_i$  of atom  $i$  calculates how much it fluctuates around its mean position  $\langle X_i \rangle$ .

$$\sigma_i = \sqrt{\langle (X_i - \langle X_i \rangle)^2 \rangle}$$




Typically done on  $C_\alpha$  atoms.  
Result depends on alignment!


# end-to-end distance and Radius of Gyration (Rg)

$$d(p, q) = \|p - q\|$$

$$R_g = \sqrt{\frac{1}{N} \sum |r_k - r_{mean}|^2}$$




Internal properties do *not* depend on alignment



# Some quantities are not *directly* measurable


Submit MD conformers to external software simulating experimental data, e.g.:

## Chemical cross-linking




DynamXL, Xwalk, ...

## SAXS



CRY SOL, FoXS, ...


## Collision cross-section



IMPACT, MobCal,  
CollisionCode, ...

# Warning: time averaging may hide processes


**Thought experiment:** typically hydrogen bond is considered established if donor-acceptor distance  $< 2.5 \text{ \AA}$ , and donor-acceptor-hydrogen angle  $< 20^\circ$ .



Reporting % time a bond is established in simulation can be misleading!




# Large ecosystem of software and packages for data analysis



NumPy



**SciPy**



Open-Source Cheminformatics  
and Machine Learning

VMD

Visual Molecular Dynamics



**MDTRAJ**



**PyEMMA**

And many more...

# MDAnalysis



The **Universe** contains everything about a MD system

- *Static information*: atoms and their connectivities
- *Dynamic information*: The trajectory

Data accessible via a hierarchy of containers

