



# Simulation of Biomolecules



## Setting up a protein simulation



Dr Matteo Degiacomi  
University of Edinburgh  
[matteo.degiacomini@ed.ac.uk](mailto:matteo.degiacomini@ed.ac.uk)

Dr Antonia Mey  
University of Edinburgh  
[antonia.mey@ed.ac.uk](mailto:antonia.mey@ed.ac.uk)



# Useful resources to learn running simulations

## Best Practices for Foundations in Molecular Simulations [Article v1.0]

Efrem Braun<sup>1</sup>, Justin Gilmer<sup>2</sup>, Heather B. Mayes<sup>3</sup>, David L. Mobley<sup>4</sup>, Jacob I. Monroe<sup>5</sup>, Samarjeet Prasad<sup>6</sup>, Daniel M. Zuckerman<sup>7</sup>

## A suite of tutorials for the BioSimSpace framework for interoperable biomolecular simulation [Article v1.0]

Lester O. Hedges<sup>1,2\*</sup>, Sofia Bariami<sup>3†</sup>, Matthew Burman<sup>2</sup>, Finlay Clark<sup>3</sup>, Benjamin P. Cossins<sup>4</sup>, Adele Hardie<sup>3</sup>, Anna M. Herz<sup>3</sup>, Dominykas Lukauskis<sup>5</sup>, Antonia S.J.S. Mey<sup>3</sup>, Julien Michel<sup>2,3\*</sup>, Jenke Scheen<sup>3‡</sup>, Miroslav Suruzhon<sup>4</sup>, Christopher J. Woods<sup>1</sup>, Zhiyi Wu<sup>4</sup>



## From Proteins to Perturbed Hamiltonians: A Suite of Tutorials for the GROMACS-2018 Molecular Simulation Package [Article v1.0]


Justin A. Lemkul

Department of Biochemistry, Virginia Polytechnic Institute and State University

<https://orcid.org/0000-0001-6661-8653>

DOI: <https://doi.org/10.33011/livecoms.1.1.5068>

Keywords: tutorials, gromacs, molecular dynamics simulation, computational chemistry



 PDF


 ARTICLE CODE REPOSITORY

GROMACS: [tutorials.gromacs.org](https://tutorials.gromacs.org)

Amber: [ambermd.org/tutorials](https://ambermd.org/tutorials)

OpenMM: [docs.openmm.org/latest/userguide/library/03\\_tutorials.html](https://docs.openmm.org/latest/userguide/library/03_tutorials.html)

# A typical workflow for molecular dynamics



Getting your protein structure

AlphaFold Protein Structure Database




Getting ligands/co-factors




ZINC20



# A typical workflow for molecular dynamics




# A typical workflow for molecular dynamics



*mostly C++  
command line* *Python API*


# Disclaimer!



Running biomolecular MD can take days on specialised hardware.


Today we will *not* run any of them, and instead will focus on fundamental principles using small molecules.

# A Molecular Dynamics timestep





# Molecular dynamics require multiple steps for the setup of simulations






# There are many choices for force fields to be made

$$\vec{F}_i = -\frac{dU}{d\vec{x}_i}$$

$$U = \sum_{\text{All Bonds}} \frac{1}{2} K_b (b - b_0)^2 + \sum_{\text{All Angles}} \frac{1}{2} K_\theta (\theta - \theta_0)^2$$

$$+ \sum K_\phi [1 - \cos(n\phi + \delta)]$$

$$+ \sum \epsilon \left[ \left( \frac{r_0}{r} \right)^{12} - 2 \left( \frac{r_0}{r} \right)^6 \right]$$

$$+ \sum \frac{332 q_i q_j}{r}$$



The diagrams show potential energy curves for different molecular interactions:
 

- Bonds:** A parabolic potential energy curve  $U(b)$  versus bond length  $b$ .
- Angles:** A parabolic potential energy curve  $U(\theta)$  versus bond angle  $\theta$ .
- Torsion Angles:** A periodic potential energy curve  $U(\phi)$  versus torsion angle  $\phi$ .
- Nonbonded Pairs:** A potential energy curve  $U(r)$  versus distance  $r$ , showing a steep repulsive wall and a shallower attractive well.
- Partial Charges:** A diagram showing two spheres with opposite charges (+ and -) and a distance  $r$ .

- **Amber** (glycam params cover most sugars)
- **CHARMM** (incl. POPC, POPE, DPPC lipids)
- **OPLS**
- **GROMOS**
- ...

*There is no “best force field”!*

## Small molecule force fields



$$E_{\text{pair}} = \sum_{\text{bonds}} K_r (r - r_{eq})^2 + \sum_{\text{angles}} K_\theta (\theta - \theta_{eq})^2 +$$

$$\sum_{\text{dihedrals}} \frac{V_n}{2} [1 + \cos(n\phi - \gamma)] + \sum_{i < j} \left[ \frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6} + \frac{q_i q_j}{\epsilon R_{ij}} \right]$$

**GAFF**

## Machine learned force fields




**Espaloma**

**Ani-2x**

**SchNet**

**PhysNet**

**BandNN**





*Coarse-grained force fields...*

# Choosing your thermodynamic ensemble

Simulations replicate a specific *thermodynamic ensemble* (typically NVT or NPT), or even grand canonical ( $\mu$ VT)


You will have different options to include *thermostats* (scaling atom velocities) and *barostats* (scaling positions) in your calculations:

- Nose-Hoover
- Berendsen
- Parrinello-Rahman
- Langevin piston
- ...




# Periodic boundary conditions (PBC) and pressure coupling

Useful to reduce finite-size effect and simulate bulk



Typically, PBC applied in x, y and z direction



For membrane systems, use semi-isotropic pressure coupling ( $x, y \neq z$ , lipids compressibility is direction-dependent)

# Sampling timescales for protein systems

The steepest gradient determines the smallest timestep.

Timestep size is imposed by the fastest phenomenon we want to observe.


In atomistic simulations:

- Covalent bond hydrogen-heavy atom ( $10^{14}$  Hz): 0.5 fs
- Covalent bond heavy atom-heavy atom: 1 fs
- Angles fluctuations: 2 fs

Restraining covalent bond distances allows to use 1-2 fs timesteps (restraining methods: SHAKE, RATTLE, LINCS,...)

Hydrogen Mass repartitioning: 4 fs

Other integrators (e.g., Langevin): 4 fs - 6 fs.



# “equilibration” and “convergence”: what do they mean?

Equilibration phase:  
is the system in a  
“relaxed” state?


Production phase: do  
we have good sampling  
and convergence?



Thinking about the problem holistically: an integrated framework for the analysis of equilibration, sampling, and convergence.

# Example equilibration protocols

**YOU WANT:** constant volume, pressure, and temperature, healthy Ramachandran plot, no exotic chemistry, bulk water (if used)





# An *example* simulation protocol

**YOU WANT:** constant volume, pressure, and temperature, healthy Ramachandran plot, no exotic chemistry, bulk water (if used)

## Equilibration:

1. Minimize energy, 1000 steepest descent
2. Heat system from 0 to 300 K in 500 ps, NPT, Berendsen barostat 1 atm.  $\alpha$ -carbon restrained with 10 kcal/mol harmonic potential. 2 fs timestep, LINCS all bonds
3. 1 ns nVT equilibration with Langevin dynamics, no atom constrained.

## Production:

4. 1  $\mu$ s NPT, Nose-Hoover barostat, PME for electrostatics

### DETERMINE HANDLING OF CUTOFFS

- ☐ As a general rule, electrostatics are long-range enough that either the cutoff needs to be larger than the system size (for finite systems) or periodicity is needed along with full treatment of long-range electrostatics (Section 3.4)
- ☐ Nonpolar interactions can often be safely treated with cutoffs of 1-1.5 nm as long as the system size is at least twice that, but long-range dispersion corrections may be needed (Section 4.1)

### CHOOSE APPROPRIATE SETTINGS FOR THE DESIRED ENSEMBLE

- ☐ Pick a thermostat that gives the correct distribution of temperatures, not just the correct average temperature; if you have a small system or a system with weakly interacting component choose one which works well even in the small-system limit.
- ☐ Pick a barostat that gives the correct distribution of pressures
- ☐ Consider the known shortcomings and limitations of certain integrators and thermostats/barostats and whether your choices will impact the properties you are calculating

### CHOOSE AN APPROPRIATE TIMESTEP FOR STABILITY AND AVOIDING ENERGY DRIFT

- ☐ Determine the highest-frequency motion in the system (typically bond vibrations unless bond lengths are constrained)
- ☐ As a first guess, set the timestep to approximately one tenth of the highest-frequency motion's characteristic period
- ☐ Test this choice by running a simulation in the microcanonical ensemble, and ensure that energy is conserved



# Which MD engine should I use?


Consider:

- Support for force field of choice
  - Enables running desired simulation protocol
  - Performance for available hardware
  - Ease of use
- depends on number of atoms, hardware, simulation protocol, MD engine*


**Graphical Processing Units (GPUs) are especially effective for MD**

From: [developer.nvidia.com/hpc-application-performance](https://developer.nvidia.com/hpc-application-performance)


**NVIDIA H200**



**NVIDIA A100**



# Calculating runtimes: example on UK Tier 2 systems



https://www.hecbiosim.ac.uk/access-hpc/hpc-calculator

**HECBioSim**

Home Contact Us Covid-19 Events Access HPC Highlights Publications Downloads About Us

Search ... Log in

## The HECtime Calculator

Enter the following information about your simulation:

JADE2

1000


100000

GROMACS 2020.4


Reset Calculate

<https://www.hecbiosim.ac.uk/access-hpc/hpc-calculator>

# Calculating runtimes: example on UK Tier 2 systems




[as of 2024]



<https://www.hecbiosim.ac.uk/access-hpc/hpc-calculator>

← ↻ 🔒 <https://docs.openforcefield.org/en/latest/examples.html> 🔊 ☆ 📄 ⌵ 📌 📌 📌 ⌵ ⋮

 **open**  
forcefield

OpenFF Documentation documentation / Examples

Search 🔍

v: latest

Overview

OpenFF Standards

GETTING STARTED

Installation

Modelling with OpenFF

Examples

PROJECTS

OpenFF Toolkit

Interchange

Units


BespokeFit


QCSubmit


Fragmenter


Tutorials

Tutorials describing key workflows from start to finish, with detailed explanations of what's going on.

  
**open**  
forcefield  
Protein-ligand-water systems with Interchange  
openff-interchange

  
**open**  
forcefield  
Solvate and equilibrate a ligand in a box of water  
openff-interchange

  
**open**  
forcefield  
Toolkit Showcase: Prepare and run a protein-ligand simulation  
openff-toolkit

  
**open**  
forcefield  
Visualizations in the OpenFF Toolkit  
openff-toolkit

PARAMETRIZATION AND EVALUATION

CONTENTS

Examples

Tutorials

Parametrization and Evaluation

Combining Force Fields

Force Field Development

Uncategorized

Running examples locally

