orbital relaxation pt.1

TCC:

TCC:

Ψ_{cc}

TCC:

$$\Psi_{CC} = \exp(T_1 + T_2 + T_3 + \ldots)\,\Phi$$

TCC:

$$\Psi_{CC} = \exp(T_1 + T_2 + T_3 + \ldots)\,\Phi$$

$$= \exp(T_2 + T_3 + \ldots)\,\tilde{\Phi}$$

TCC:

$$\Psi_{CC} = \exp(T_1 + T_2 + T_3 + \ldots)\,\Phi$$

$$= \exp(T_2 + T_3 + \ldots)\,\tilde{\Phi}$$

$\tilde{\psi}_i$

TCC:

$$\Psi_{cc} = \exp(T_1 + T_2 + T_3 + \dots)\,\Phi$$

$$= \exp(T_2 + T_3 + \dots)\,\tilde{\Phi}$$

$$\tilde{\psi}_i = \psi_i + \sum_a t_a^i \psi_a$$

TCC:

$$\Psi_{CC} = \exp(T_1 + T_2 + T_3 + \ldots)\,\Phi$$

$$= \exp(T_2 + T_3 + \ldots)\,\tilde{\Phi}$$

$$\tilde{\psi}_i = \psi_i + \sum_a t_a^i \psi_a$$

T_1 is big when Φ is bad

$$\tilde{\Phi} = \exp(T_1)\,\Phi$$ is intermediately normalized

$$\tilde{\Phi} = \exp(T_1)\,\Phi$$ is intermediately normalized

$$\tilde{\Phi} = \exp(T_1 - T_1^{\dagger})\,\Phi$$ is normalized

Alternative treatments of orb. relaxation:

Alternative treatments of orb. relaxation:

1. best overlap: maximize $\|\langle\Phi|\Psi\rangle\|^2$

Alternative treatments of orb. relaxation:

1. best overlap: maximize $\| \langle \Phi | \Psi \rangle \|^2$

2. best energy: minimize $\langle \Psi | H | \Psi \rangle$

Alternative treatments of orb. relaxation:

1. best overlap: maximize $\|\langle\Phi|\Psi\rangle\|^2$

"Brueckner orbitals"

2. best energy: minimize $\langle\Psi|H|\Psi\rangle$

Alternative treatments of orb. relaxation:

1. best overlap: maximize $\|\langle\Phi|\Psi\rangle\|^2$

"Brueckner orbitals"

2. best energy: minimize $\langle\Psi|H|\Psi\rangle$

"optimized orbitals"