

• SECOND • QUANTIZATION •

1-particle Hilbert space:

1-particle Hilbert space:


\mathcal{H}

1-particle Hilbert space:

$$\mathcal{H} = \text{span}\{\psi_p\}$$

1-particle Hilbert space:

$$\mathcal{H} = \text{span}\{\psi_p\}$$



n -particle Hilbert space:

n -particle Hilbert space:

H^n

n -particle Hilbert space:

$$\mathcal{H}^{\otimes n} = \text{span} \{ \psi_{p_1} \otimes \cdots \otimes \psi_{p_n} \}$$

n -particle Hilbert space:

$$\mathcal{H}^{\otimes n} = \text{span} \{ \psi_{p_1} \otimes \dots \otimes \psi_{p_n} \}$$



$$\langle 1 \otimes \dots \otimes n | \psi_{p_1} \otimes \dots \otimes \psi_{p_n} \rangle = \psi_{p_1}(1) \dots \psi_{p_n}(n)$$

n -particle Hilbert space:

$$\mathcal{H}^n = \text{span} \{ \psi_{p_1} \otimes \dots \otimes \psi_{p_n} \}$$



$$\langle 1 \otimes \dots \otimes n | \psi_{p_1} \otimes \dots \otimes \psi_{p_n} \rangle = \psi_{p_1}(1) \dots \psi_{p_n}(n)$$

"Hartree products"

$$\psi_{p_1}(1) \cdots \psi_{p_n}(n)$$

$$\psi_{p_1}(1) \cdots \psi_{p_n}(n)$$

- Hartree products are not antisymmetric

$$\psi_{p_1}(1) \cdots \psi_{p_n}(n)$$

- Hartree products are not antisymmetric
 \Rightarrow combine into Slater determinants

$$\psi_{p_1}(1) \cdots \psi_{p_n}(n)$$

- Hartree products are not antisymmetric

\Rightarrow combine into Slater determinants

$$\Phi_{(p_1, \dots, p_n)}(1, \dots, n) = \frac{1}{\sqrt{n!}} \sum_{\pi} \psi_{p_{\pi(1)}}(1) \cdots \psi_{p_{\pi(n)}}(n)$$

$$\psi_{p_1}(1) \cdots \psi_{p_n}(n)$$

- Hartree products are not antisymmetric

\Rightarrow combine into Slater determinants

$$\Phi_{(p_1, \dots, p_n)}(1, \dots, n) = \frac{1}{\sqrt{n!}} \sum_{\pi} \psi_{p_{\pi(1)}}(1) \cdots \psi_{p_{\pi(n)}}(n)$$

upshot: Hilbert space is too big!

$$\text{use } F_n(\mathcal{H}) = \text{span} \{ \Phi_{(p_1, \dots, p_n)} \}$$

use $F_n(\mathcal{H}) = \text{span} \{ \Phi_{(p_1, \dots, p_n)} \}$

↪ space of physically realistic (a.s.)
n-particle wavefunctions

consider the operator \hat{a}_p def'd by

consider the operator \hat{a}_p def'd by

$$\hat{a}_p: F_n(H) \rightarrow F_{n-1}(H)$$

consider the operator \hat{a}_p def'd by

$$\hat{a}_p: \mathcal{F}_n(H) \rightarrow \mathcal{F}_{n-1}(H)$$

$$\hat{a}_p \equiv \sqrt{n} \int d(l) \psi_p^*(l)$$

consider the operator \hat{a}_p def'd by

$$\hat{a}_p: \mathcal{F}_n(H) \rightarrow \mathcal{F}_{n-1}(H)$$

$$\hat{a}_p \equiv \sqrt{n} \int d(1) \psi_p^*(1)$$

$$\hat{a}_{p_1} \Phi_{(p_1, p_2, \dots, p_n)}(1, 2, \dots, n)$$

consider the operator \hat{a}_p def'd by

$$\hat{a}_p: \mathcal{F}_n(H) \rightarrow \mathcal{F}_{n-1}(H)$$

$$\hat{a}_p \equiv \sqrt{n} \int d(1) \psi_p^*(1)$$

$$\hat{a}_{p_1} \Phi_{(p_1, p_2 \dots p_n)}(1, 2, \dots, n) = \Phi_{(\cancel{p_1}, p_2 \dots p_n)}(2, \dots, n)$$

consider the operator \hat{a}_p def'd by

$$\hat{a}_p: \mathcal{F}_n(H) \rightarrow \mathcal{F}_{n-1}(H)$$

$$\hat{a}_p \equiv \sqrt{n} \int d(l) \psi_p^*(l)$$

$$\hat{a}_{p_1} \Phi_{(p_1, p_2, \dots, p_n)}(1, 2, \dots, n) = \Phi_{(\cancel{p_1}, p_2, \dots, p_n)}(2, \dots, n)$$

$$\Psi(1, \dots, n) = \frac{1}{\sqrt{n}} \sum_{p=1}^{\infty} \psi_p(1) (\hat{a}_p \Psi)(2, \dots, n)$$

$$\langle \psi | \hat{H}_e | \psi' \rangle$$

$$\langle \Psi | \hat{H}_e \Psi' \rangle$$

$$= \sum_{i=1}^n \langle \Psi | \hat{h}(i) \Psi' \rangle + \sum_{i < j}^n \langle \Psi | \hat{g}(i, j) \Psi' \rangle$$

$$\langle \psi | \hat{H}_e \psi' \rangle$$

$$= n \langle \psi | \hat{h}(1) \psi' \rangle + \frac{n(n-1)}{2} \langle \psi | \hat{g}(1,2) \psi' \rangle$$

$$\langle \Psi | \hat{H}_e \Psi' \rangle$$

$$= n \langle \Psi | \hat{h}(1) \Psi' \rangle + \frac{n(n-1)}{2} \langle \Psi | \hat{g}(1,2) \Psi' \rangle$$

$$= \sum_{pq} \langle \psi_p(1) | \hat{h}(1) \psi_q(1) \rangle \langle \hat{a}_p \Psi | \hat{a}_q \Psi' \rangle$$

$$\langle \Psi | \hat{H}_e | \Psi' \rangle$$

$$= n \langle \Psi | \hat{h}(1) | \Psi' \rangle + \frac{n(n-1)}{2} \langle \Psi | \hat{g}(1,2) | \Psi' \rangle$$

$$= \sum_{pq} \langle \psi_p(1) | \hat{h}(1) | \psi_q(1) \rangle \langle \hat{a}_p \Psi | \hat{a}_q \Psi' \rangle$$

$$+ \frac{1}{2} \sum_{pqrs} \langle \psi_p(1) \psi_q(2) | \hat{g}(1,2) | \psi_r(1) \psi_s(2) \rangle \cdot$$

$$\langle \hat{a}_q \hat{a}_p \Psi | \hat{a}_s \hat{a}_r \Psi' \rangle$$

$$\langle \Psi | \hat{H}_e | \Psi' \rangle$$

$$= n \langle \Psi | \hat{h}(1) | \Psi' \rangle + \frac{n(n-1)}{2} \langle \Psi | \hat{g}(1,2) | \Psi' \rangle$$

$$= \sum_{pq} \langle \psi_p(1) | \hat{h}(1) | \psi_q(1) \rangle \langle \hat{a}_p \Psi | \hat{a}_q \Psi' \rangle$$

$$+ \frac{1}{2} \sum_{pqrs} \langle \psi_p(1) \psi_q(2) | \hat{g}(1,2) | \psi_r(1) \psi_s(2) \rangle \cdot$$

$$\langle \hat{a}_q \hat{a}_p \Psi | \hat{a}_s \hat{a}_r \Psi' \rangle$$

$$\Rightarrow \hat{H}_e |_{\mathcal{F}_n} = \sum_{pq} h_{pq} \hat{a}_p^\dagger \hat{a}_q + \frac{1}{2} \sum_{pqrs} \langle pq | rs \rangle \hat{a}_p^\dagger \hat{a}_q^\dagger \hat{a}_s \hat{a}_r$$

$$\langle \Psi | \hat{H}_e \Psi' \rangle$$

$$= n \langle \Psi | \hat{h}(1) \Psi' \rangle + \frac{n(n-1)}{2} \langle \Psi | \hat{g}(1,2) \Psi' \rangle$$

$$= \sum_{pq} \langle \psi_p(1) | \hat{h}(1) \psi_q(1) \rangle \langle \hat{a}_p \Psi | \hat{a}_q \Psi' \rangle$$

$$+ \frac{1}{2} \sum_{pqrs} \langle \psi_p(1) \psi_q(2) | \hat{g}(1,2) \psi_r(1) \psi_s(2) \rangle \cdot$$

$$\langle \hat{a}_q \hat{a}_p \Psi | \hat{a}_s \hat{a}_r \Psi' \rangle$$

$$\Rightarrow \hat{H}_e |_{F_n} = \sum_{pq} h_{pq} \hat{a}_p^\dagger \hat{a}_q + \frac{1}{2} \sum_{pqrs} \langle pq | rs \rangle \hat{a}_p^\dagger \hat{a}_q^\dagger \hat{a}_s \hat{a}_r$$

$\underbrace{\hspace{10em}}_{\text{same for all } F_n!}$

Fock space:

Fock space:

$$\mathcal{F}(\mathcal{H}) = \mathcal{F}_0(\mathcal{H}) \oplus \mathcal{F}_1(\mathcal{H}) \oplus \mathcal{F}_2(\mathcal{H}) \oplus \dots$$

Fock space:

$$\mathcal{F}(\mathcal{H}) = \mathcal{F}_0(\mathcal{H}) \oplus \mathcal{F}_1(\mathcal{H}) \oplus \mathcal{F}_2(\mathcal{H}) \oplus \dots$$

Fock space Hamiltonian:

Fock space:

$$\mathcal{F}(\mathcal{H}) = \mathcal{F}_0(\mathcal{H}) \oplus \mathcal{F}_1(\mathcal{H}) \oplus \mathcal{F}_2(\mathcal{H}) \oplus \dots$$

Fock space Hamiltonian:

$$H_c = \sum_{pq} h_{pq} a_p^\dagger a_q + \frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle a_p^\dagger a_q^\dagger a_s a_r$$

a_p is an annihilation operator

a_p is an annihilation operator 😡

a_p is an annihilation operator 😡

$$a_{p_1} \Phi_{(p_1, p_2, \dots, p_n)} = \Phi_{(p_2, \dots, p_n)}$$

a_p is an annihilation operator 😡

$$a_{p_1} \Phi_{(p_1, p_2, \dots, p_n)} = \Phi_{(p_2, \dots, p_n)}$$

$$a_p \Phi_{(p_1, \dots, p_n)} = 0 \quad \text{if } p \notin (p_1, \dots, p_n)$$

a_p is an annihilation operator 😡

$$a_{p_1} \Phi_{(p_1, p_2, \dots, p_n)} = \Phi_{(p_2, \dots, p_n)}$$

$$a_p \Phi_{(p_1, \dots, p_n)} = 0 \quad \text{if } p \notin (p_1, \dots, p_n)$$

can we define c_p , creation operator?

a_p is an annihilation operator 😡

$$a_p \Phi_{(p_1 p_2 \dots p_n)} = \Phi_{(p_2 \dots p_n)}$$

$$a_p \Phi_{(p_1 \dots p_n)} = 0 \quad \text{if } p \notin (p_1 \dots p_n)$$

can we define c_p , creation operator? 🌸

a_p is an annihilation operator 😡

$$a_{p_1} \Phi_{(p_1, p_2, \dots, p_n)} = \Phi_{(p_2, \dots, p_n)}$$

$$a_p \Phi_{(p_1, \dots, p_n)} = 0 \quad \text{if } p \notin (p_1, \dots, p_n)$$

can we define c_p , creation operator? 🌸

$$c_p \Phi_{(p_1, \dots, p_n)} = \Phi_{(p, p_1, \dots, p_n)} \quad \text{if } p \notin (p_1, \dots, p_n)$$

a_p is an annihilation operator 😡

$$a_{p_1} \Phi_{(p_1, p_2, \dots, p_n)} = \Phi_{(p_2, \dots, p_n)}$$

$$a_p \Phi_{(p_1, \dots, p_n)} = 0 \quad \text{if } p \notin (p_1, \dots, p_n)$$

can we define c_p , creation operator? 🌸

$$c_p \Phi_{(p_1, \dots, p_n)} = \begin{cases} \Phi_{(p, p_1, \dots, p_n)} & \text{if } p \notin (p_1, \dots, p_n) \\ 0 & \text{otw.} \end{cases}$$

a little bit of proofing shows
that

a little bit of proofing shows
that

$$c_p = a_p^+$$

a little bit of proofing shows
that

$$c_p = a_p^\dagger$$

$$|\Phi_{(p_1 \dots p_n)}\rangle = a_{p_1}^\dagger \dots a_{p_n}^\dagger |\Phi_{(1)}\rangle$$

a little bit of proofing shows
that

$$c_p = a_p^\dagger$$

$$|\Phi_{(p_1 \dots p_n)}\rangle = a_{p_1}^\dagger \dots a_{p_n}^\dagger |\text{vac}\rangle$$

commutation relations:

commutation relations:

$$[a_p, a_q]_+ = 0$$

commutation relations:

$$[a_p, a_q]_+ = 0$$

$$[a_p^\dagger, a_q^\dagger]_+ = 0$$

commutation relations:

$$[a_p, a_q]_+ = 0$$

$$[a_p^+, a_q^+]_+ = 0$$

$$[a_p, a_q^+]_+ = 0 \quad \text{if } p \neq q$$

commutation relations:

$$[a_p, a_q]_+ = 0$$

$$[a_p^+, a_q^+]_+ = 0$$

$$[a_p, a_q^+]_+ = 0 \quad \text{if } p \neq q$$

$$[a_p, a_p^+]_+ = 1$$

Let's work through some examples:

let's work through some examples:

1. $\langle \text{vac} | a_p a_q^\dagger | \text{vac} \rangle$

2. $\langle \text{vac} | a_p a_q a_s^\dagger a_r^\dagger | \text{vac} \rangle$

strategy: move the daggers to the left

brief interlude: occupation number formalism

brief interlude: occupation number formalism

$$|n_1, n_2, n_3, \dots\rangle$$

brief interlude: occupation number formalism

$$|n_1, n_2, n_3, \dots\rangle \quad n_p = \begin{cases} 1 & \psi_p \text{ occupied} \\ 0 & \psi_p \text{ unoccupied} \end{cases}$$

brief interlude: occupation number formalism

$$|n_1, n_2, n_3, \dots\rangle \quad n_p = \begin{cases} 1 & \psi_p \text{ occupied} \\ 0 & \psi_p \text{ unoccupied} \end{cases}$$

$$|vac\rangle = |0, \dots, 0, 0, \dots\rangle$$

brief interlude: occupation number formalism

$$|n_1, n_2, n_3, \dots\rangle \quad n_p = \begin{cases} 1 & \psi_p \text{ occupied} \\ 0 & \psi_p \text{ unoccupied} \end{cases}$$

$$|vac\rangle = |0, \dots, 0, 0, \dots\rangle$$

$$|\Phi\rangle = |1, \dots, 1, 0, \dots\rangle$$

p.h. isomorphism:

p.h. isomorphism:

$|n_1, \dots, n_m, n_{m+1}, \dots\rangle$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$$|vac\rangle$$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$|\text{vac}\rangle$

\downarrow

$|\Phi\rangle$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$$|vac\rangle \quad |\Phi\rangle$$



$$|\Phi\rangle$$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$
\downarrow	\downarrow
$ \Phi\rangle$	$ vac\rangle$

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$	a_i	a_i^\dagger
\downarrow	\downarrow		
$ \Phi\rangle$	$ vac\rangle$		

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$	a_i	a_i^+
\downarrow	\downarrow	\downarrow	\downarrow
$ \Phi\rangle$	$ vac\rangle$	a_i^+	a_i

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$	a_i	a_i^\dagger	a_a	a_a^\dagger
\downarrow	\downarrow	\downarrow	\downarrow		
$ \Phi\rangle$	$ vac\rangle$	a_i^\dagger	a_i		

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$	a_i	a_i^+	a_a	a_a^+
\downarrow	\downarrow	\downarrow	\downarrow	\downarrow	\downarrow
$ \Phi\rangle$	$ vac\rangle$	a_i^+	a_i	a_a	a_a^+

p.h. isomorphism:

$$|n_1, \dots, n_m, n_{m+1}, \dots\rangle \mapsto |\bar{n}_1, \dots, \bar{n}_m, n_{m+1}, \dots\rangle$$

$$\bar{n}_i = 1 - n_i$$

$ vac\rangle$	$ \Phi\rangle$	a_i	a_i^+	a_a	a_a^+
\downarrow	\downarrow	\downarrow	\downarrow	\downarrow	\downarrow
$ \widetilde{\Phi}\rangle$	$ \widetilde{vac}\rangle$	b_i^+	b_i	b_a	b_a^+

$$\sum_{pq} h_{pq} a_p^\dagger a_q$$

$$\sum_{pq} h_{pq} a_p^\dagger a_q$$

$$= \sum_{ab} h_{ab} a_a^\dagger a_b + \sum_{ai} h_{ai} a_a^\dagger a_i$$

$$+ \sum_{ia} h_{ia} a_i^\dagger a_a + \sum_{ij} h_{ij} a_i^\dagger a_j$$

$$\sum_{pq} h_{pq} a_p^\dagger a_q$$

$$= \sum_{ab} h_{ab} b_a^\dagger b_b + \sum_{ai} h_{ai} b_a^\dagger b_i^\dagger$$

$$+ \sum_{ia} h_{ia} b_i b_a + \sum_{ij} h_{ij} b_i b_j^\dagger$$

$$\sum_{pq} h_{pq} a_p^\dagger a_q$$

$$= \sum_{ab} h_{ab} b_a^\dagger b_b + \sum_{ai} h_{ai} b_a^\dagger b_i^\dagger$$

$$+ \sum_{ia} h_{ia} b_i b_a + \sum_{ij} h_{ij} b_i b_j^\dagger$$

examples to work through:

$$\sum_{pq} h_{pq} a_p^\dagger a_q$$

$$= \sum_{ab} h_{ab} b_a^\dagger b_b + \sum_{ai} h_{ai} b_a^\dagger b_i^\dagger$$

$$+ \sum_{ia} h_{ia} b_i b_a + \sum_{ij} h_{ij} b_i b_j^\dagger$$

examples to work through:

$$1. \langle \Phi | H_{\text{core}} | \Phi \rangle$$

$$2. \langle \Phi | H_{\text{core}} | \Phi_i^a \rangle$$

$$3. \langle \Phi | H_{\text{core}} | \Phi_{ij}^{ab} \rangle$$

if there's time:

if there's time:

expand $\frac{1}{4} \sum_{pqrs} \langle pq || rs \rangle a_p^\dagger a_q^\dagger a_s a_r$

* work out matrix elements

the end.