

the quasi-energy formalism

the quasi-energy formalism
pt. 1

$$V(t)$$

$$V(t) = \sum_{\beta} V_{\beta} f_{\beta}(t)$$

$$V(t) = \sum_{\beta} V_{\beta} f_{\beta}(t)$$

$$f_{\beta}(t)$$

$$V(t) = \sum_{\beta} V_{\beta} f_{\beta}(t)$$

$$f_{\beta}(t) = \int_{-\infty}^{\infty} d\omega \, f_{\beta}(\omega_{\epsilon}) e^{-i\omega_{\epsilon}t}$$

$$V(t) = \sum_{\beta} V_{\beta} f_{\beta}(t)$$

$$f_{\beta}(t) = \int_{-\infty}^{\infty} d\omega \, f_{\beta}(\omega_{\epsilon}) e^{-i\omega_{\epsilon}t}$$

$$f_{\beta}(t) \in \mathbb{R}$$

$$V(t) = \sum_{\beta} V_{\beta} f_{\beta}(t)$$

$$f_{\beta}(t) = \int_{-\infty}^{\infty} d\omega \, f_{\beta}(\omega_{\epsilon}) e^{-i\omega_{\epsilon}t}$$

$$f_{\beta}(t) \in \mathbb{R} \quad \Rightarrow \quad f_{\beta}^*(\omega_{\epsilon}) = f_{\beta}(-\omega_{-\epsilon})$$

$$\langle \psi(t) | W | \psi(t) \rangle - \langle \psi_0 | W | \psi_0 \rangle$$

$$\langle \psi(t) | W | \psi(t) \rangle - \langle \psi_0 | W | \psi_0 \rangle$$

$$\approx \sum_{\beta} \int_{-\infty}^{\infty} dt' f_{\beta}(t') \ll \tilde{W}(t); \tilde{V}_{\beta}(t') \gg$$

$$\langle \psi(t) | W | \psi(t) \rangle - \langle \psi_0 | W | \psi_0 \rangle$$

$$\approx \sum_{\beta} \int_{-\infty}^{\infty} dt' f_{\beta}(t') \ll \tilde{W}(t); \tilde{V}_{\beta}(t') \gg$$

$$= \sum_{\beta} \int_{-\infty}^{\infty} d\omega f_{\beta}(\omega) \ll W; V_{\beta} \gg_{\omega} e^{-i\omega t}$$

$$\langle \psi(t) | W | \psi(t) \rangle - \langle \psi_0 | W | \psi_0 \rangle$$

$$\approx \sum_{\beta} \int_{-\infty}^{\infty} dt' f_{\beta}(t') \ll \tilde{W}(t); \tilde{V}_{\beta}(t') \gg$$

$$= \sum_{\beta} \int_{-\infty}^{\infty} d\omega f_{\beta}(\omega) \ll W; V_{\beta} \gg_{\omega} e^{-i\omega t}$$

($\epsilon \rightarrow 0$ limit)

Store this for later:

Store this for later:

$$\frac{\partial H(t)}{\partial f_{\alpha}(\omega')}$$

Store this for later:

$$\frac{\partial H(t)}{\partial f_{\alpha}(w')} = V_{\alpha} \frac{\partial f_{\alpha}(t)}{\partial f_{\alpha}(w')}$$

Store this for later:

$$\frac{\partial H(t)}{\partial f_{\alpha}(\omega')} = V_{\alpha} \frac{\partial f_{\alpha}(t)}{\partial f_{\alpha}(\omega')} = V_{\alpha} e^{-i\omega' t}$$

Store this for later:

$$\frac{\partial H(t)}{\partial f_{\alpha}(\omega')} = V_{\alpha} \frac{\partial f_{\alpha}(t)}{\partial f_{\alpha}(\omega')} = V_{\alpha} e^{-i\omega' t}$$

($\epsilon \rightarrow 0$ limit)

phase-isolated wfn:

phase-isolated wfn:

$$\Psi(t)$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t dt' Q(t')} \bar{\Psi}(t)$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t dt' Q(t')} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t dt' Q(t')} \bar{\Psi}(t)$$

$$Q(t) = Q^\dagger(t)$$

Q.E. Schrödinger eq.:

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t Q(t') dt'} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

Q.E. Schrödinger eq.:

$$(H(t) - i \frac{\partial}{\partial t}) \bar{\Psi}(t) = Q(t) \bar{\Psi}(t)$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t Q(t') dt'} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

Q.E. Schrödinger eq.:

$$(H(t) - i \frac{\partial}{\partial t}) \bar{\Psi}(t) = Q(t) \bar{\Psi}(t)$$

↑
"quasi-energy"

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t Q(t') dt'} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

Q.E. Schrödinger eq.:

$$(H(t) - i \frac{\partial}{\partial t}) \bar{\Psi}(t) = Q(t) \bar{\Psi}(t)$$

$$Q(t) = \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle \quad \begin{matrix} \uparrow \\ \text{"quasi-energy"} \end{matrix}$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t Q(t') dt'} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

Q.E. Schrödinger eq.:

$$(H(t) - i \frac{\partial}{\partial t}) \bar{\Psi}(t) = Q(t) \bar{\Psi}(t)$$

$$Q(t) = \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle \quad \begin{matrix} \uparrow \\ \text{"quasi-energy"} \end{matrix}$$

$$\bar{\Psi}^{(0)} = \Psi_0$$

phase-isolated wfn:

$$\Psi(t) = e^{-i \int_0^t Q(t') dt'} \bar{\Psi}(t)$$

$$Q(t) = Q^*(t)$$

Q.E. Schrödinger eq.:

$$(H(t) - i \frac{\partial}{\partial t}) \bar{\Psi}(t) = Q(t) \bar{\Psi}(t)$$

$$Q(t) = \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle \quad \begin{matrix} \uparrow \\ \text{"quasi-energy"} \end{matrix}$$

$$\bar{\Psi}^{(0)} = \Psi_0$$

$$Q^{(0)} = E_0$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re} [\langle \delta \bar{\Psi}(t) | H(t) | \Psi(t) \rangle]$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re} [\langle \delta \bar{\Psi}(t) | H(t) | \Psi(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \Psi(t) \rangle$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle]$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle \rightarrow 0$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle \xrightarrow{0}$$

$$2 \operatorname{Re}\left[i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle \right]$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle \xrightarrow{0}$$

$$2 \operatorname{Re}\left[i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle\right] = i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle - i \left\langle \frac{\partial \bar{\Psi}(t)}{\partial t} \left| \delta \bar{\Psi}(t) \right. \right\rangle$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle \xrightarrow{0}$$

$\frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \bar{\Psi}(t) \rangle - \langle \bar{\Psi}(t) | \frac{\partial}{\partial t} \delta \bar{\Psi}(t) \rangle$

$$2 \operatorname{Re}\left[i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle\right] = i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle - i \left\langle \frac{\partial \bar{\Psi}(t)}{\partial t} \left| \delta \bar{\Psi}(t) \right. \right\rangle$$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

project by $\delta \bar{\Psi}(t)$ and determine real part

$$2 \operatorname{Re}[\langle \delta \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle] = \delta \langle \bar{\Psi}(t) | H(t) | \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}[Q(t) \langle \delta \bar{\Psi}(t) | \bar{\Psi}(t) \rangle] = Q(t) \delta \langle \bar{\Psi}(t) | \bar{\Psi}(t) \rangle \xrightarrow{0} \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \bar{\Psi}(t) \rangle - \langle \bar{\Psi}(t) | \frac{\partial}{\partial t} \delta \bar{\Psi}(t) \rangle$$

$$2 \operatorname{Re}\left[i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle \right] = i \left\langle \delta \bar{\Psi}(t) \left| \frac{\partial \bar{\Psi}(t)}{\partial t} \right. \right\rangle - i \left\langle \frac{\partial \bar{\Psi}(t)}{\partial t} \left| \delta \bar{\Psi}(t) \right. \right\rangle$$

$$= \delta \langle \bar{\Psi}(t) | i \frac{\partial}{\partial t} \bar{\Psi}(t) \rangle - i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \bar{\Psi}(t) \rangle$$

Combine results:

Combine results:

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \bar{\Psi}(t) \rangle = 0$$

Combine results:

$$\underbrace{\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle}_{Q(t)} + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

Combine results:

$$\delta \underbrace{\langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle}_{Q(t)} + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \bar{\Psi}(t) \rangle = 0$$

almost a variational condition

Assume periodic

Assume periodic $H(t) = H(t+T)$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

Assume periodic $H(t) = H(t+T)$

$$\begin{array}{c} (H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t) \\ \parallel \\ H(t+T) \end{array}$$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

\parallel

$$H(t+T)$$

\parallel

$$U(T) H(t) U^\dagger(T)$$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

\parallel

$$H(t+T)$$

\parallel

$$U(T) H(t) U^\dagger(T)$$

$$(H(t) - i \frac{\partial}{\partial t}) U^\dagger(T) \Psi(t) = Q(t) U^\dagger(T) \bar{\Psi}(t)$$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

\parallel

$$H(t+T)$$

\parallel

$$U(T) H(t) U^\dagger(T)$$

$$(H(t) - i \frac{\partial}{\partial t}) U^\dagger(T) \Psi(t) = Q(t) U^\dagger(T) \bar{\Psi}(t)$$

$$\Rightarrow \bar{\Psi}(t)$$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

\parallel

$$H(t+T)$$

\parallel

$$U(T) H(t) U^\dagger(T)$$

$$(H(t) - i \frac{\partial}{\partial t}) U^\dagger(T) \Psi(t) = Q(t) U^\dagger(T) \bar{\Psi}(t)$$

$$\Rightarrow \bar{\Psi}(t) = U^\dagger(T) \bar{\Psi}(t)$$

Assume periodic $H(t) = H(t+T)$

$$(H(t) - i \frac{\partial}{\partial t}) \Psi(t) = Q(t) \bar{\Psi}(t)$$

\parallel

$$H(t+T)$$

\parallel

$$U(T) H(t) U^\dagger(T)$$

$$(H(t) - i \frac{\partial}{\partial t}) U^\dagger(T) \Psi(t) = Q(t) U^\dagger(T) \bar{\Psi}(t)$$

$$\Rightarrow \bar{\Psi}(t) = U^\dagger(T) \bar{\Psi}(t) = \bar{\Psi}(t-T)$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T = 0$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T = 0$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T = 0$$

\Rightarrow

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T = 0$$

$$\Rightarrow \delta \frac{1}{T} \int_0^T dt' Q(t') = 0$$

$$\delta \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle + i \frac{\partial}{\partial t} \langle \bar{\Psi}(t) | \delta \Psi(t) \rangle = 0$$

\Downarrow

(FTC)

$$\int_0^T dt' \delta \langle \bar{\Psi}(t') | H - i \frac{\partial}{\partial t} | \Psi(t') \rangle + i \langle \bar{\Psi}(t') | \delta \Psi(t') \rangle \Big|_0^T = 0$$

$$\Rightarrow \underbrace{\delta \frac{1}{T} \int_0^T dt' Q(t')}_Q = 0$$

time-argd. Q.E.:

time-avgd. Q.E.:

$$\delta Q = 0$$

time-avgd. Q.E.:

$$\delta Q = 0 \quad \text{for exact } \bar{\Psi}(t)$$

time-avgd. Q.E.:

$$\delta Q = 0 \quad \text{for exact } \bar{\Psi}(t)$$

$$Q \equiv \left\{ \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \bar{\Psi}(t) \rangle \right\}_T$$

time-avgd. Q.E.:

$$\delta Q = 0 \quad \text{for exact } \bar{\Psi}(t)$$

$$Q \equiv \left\{ \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle \right\}_T$$

Hellmann-Feynman:

time-avgd. Q.E.:

$$\delta Q = 0 \quad \text{for exact } \bar{\Psi}(t)$$

$$Q \equiv \left\{ \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle \right\}_T$$

Hellmann-Feynman:

$$\frac{dQ}{d\xi}$$

time-avgd. Q.E.:

$$\delta Q = 0 \quad \text{for exact } \bar{\Psi}(t)$$

$$Q \equiv \left\{ \langle \bar{\Psi}(t) | H(t) - i \frac{\partial}{\partial t} | \Psi(t) \rangle \right\}_T$$

Hellmann-Feynman:

$$\frac{dQ}{d\xi} = \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial \xi} | \Psi(t) \rangle \right\}_T$$

$$\frac{dQ}{df_q(w)}$$

$$\frac{dQ}{df_{\alpha}(\omega)} = \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_{\alpha}(\omega)} | \Psi(t) \rangle \right\}_T$$

$$\begin{aligned} \frac{dQ}{df_{\alpha}(\omega)} &= \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_{\alpha}(\omega)} | \Psi(t) \rangle \right\}_T \\ &= \left\{ \langle \Psi(t) | V_{\alpha} | \Psi(t) \rangle e^{-i\omega t} \right\}_T \end{aligned}$$

$$\frac{dQ}{df_{\alpha}(\omega)} = \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_{\alpha}(\omega)} | \Psi(t) \rangle \right\}_T$$

$$= \left\{ \langle \Psi(t) | V_{\alpha} | \Psi(t) \rangle e^{-i\omega t} \right\}_T$$

$$\frac{d^2 Q}{df_{\alpha}(\omega) df_{\beta}(\omega)} \Big|_{f=0}$$

$$\begin{aligned}\frac{dQ}{df_\alpha(\omega)} &= \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_\alpha(\omega)} | \Psi(t) \rangle \right\}_T \\ &= \left\{ \langle \Psi(t) | V_\alpha | \Psi(t) \rangle e^{-i\omega t} \right\}_T\end{aligned}$$

$$\left. \frac{d^2 Q}{df_\alpha(\omega) df_\beta(\omega)} \right|_{f=0} = \langle\langle V_\alpha; V_\beta \rangle\rangle_\omega$$

$$\frac{dQ}{df_{\alpha}(\omega')} = \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_{\alpha}(\omega')} | \Psi(t) \rangle \right\}_T$$

$$= \left\{ \langle \Psi(t) | V_{\alpha} | \Psi(t) \rangle e^{-i\omega' t} \right\}_T$$

$$\frac{d^2 Q}{df_{\alpha}(\omega') df_{\beta}(\omega)} \Big|_{f=0} = \langle\langle V_{\alpha}; V_{\beta} \rangle\rangle_{\omega} \left\{ e^{-i(\omega' + \omega)t} \right\}_T$$

$$\frac{dQ}{df_\alpha(\omega')} = \left\{ \langle \Psi(t) | \frac{\partial H(t)}{\partial f_\alpha(\omega')} | \Psi(t) \rangle \right\}_T$$

$$= \left\{ \langle \Psi(t) | V_\alpha | \Psi(t) \rangle e^{-i\omega't} \right\}_T$$

$$\frac{d^2 Q}{df_\alpha(\omega') df_\beta(\omega)} \Big|_{f=0} = \langle\langle V_\alpha; V_\beta \rangle\rangle_\omega \left\{ e^{-i(\omega'+\omega)t} \right\}_T$$



freqs have
period T , so
we need $\omega' + \omega = 0$

$$\langle\langle V_\alpha; V_\beta \rangle\rangle_\omega$$

$$\langle\langle V_\alpha; V_\beta \rangle\rangle_\omega = \frac{d^2 Q}{df_\alpha(\omega') df_\beta(\omega)} \bigg|_{f=0}^{\omega' = -\omega}$$

$$\langle\langle V_\alpha; V_\beta \rangle\rangle_\omega = \frac{d^2 Q}{df_\alpha(\omega') df_\beta(\omega)} \bigg|_{f=0}^{\omega'=-\omega}$$

$$= \frac{d^2 Q}{df_\alpha^*(\omega) df_\beta(\omega)} \bigg|_{f=0}$$

$$\langle\langle V_\alpha; V_\beta \rangle\rangle_\omega = \frac{d^2 Q}{df_\alpha(\omega') df_\beta(\omega)} \Big|_{f=0}^{\omega' = -\omega}$$

$$= \frac{d^2 Q}{df_\alpha^*(\omega) df_\beta(\omega)} \Big|_{f=0}$$