

p. t.

model Hamiltonian:

model Hamiltonian:

$$H_0 \approx H_c$$

model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0$$

model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0 \quad (\text{known e.f.s})$$

model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0 \quad (\text{known e.f.s})$$

$\Rightarrow$

model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0 \quad (\text{known e.f.s})$$

$$\Rightarrow H_c$$

model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0 \quad (\text{known e.f.s})$$

$$\Rightarrow H_c = [\langle \Phi_0 | H_c | \Phi_\tau \rangle]$$



model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_0 = \mathcal{E}_0 \Phi_0 \quad (\text{known e.f.s})$$

$$\Rightarrow H_c = [\langle \Phi_0 | H_c | \Phi_\tau \rangle]$$

=


model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_\sigma = \mathcal{E}_\sigma \Phi_\sigma \quad (\text{known e.f.s})$$

$$\Rightarrow H_c = [\langle \Phi_\sigma | H_c | \Phi_\tau \rangle]$$

||




model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_\sigma = \mathcal{E}_\sigma \Phi_\sigma \quad (\text{known e.f.s})$$

$$\Rightarrow H_c = [\langle \Phi_\sigma | H_c | \Phi_\tau \rangle]$$

"



sparse


model Hamiltonian:

$$H_0 \approx H_c$$

$$H_0 \Phi_\sigma = \mathcal{E}_\sigma \Phi_\sigma \quad (\text{known e.f.s})$$

$$\Rightarrow H_c = [\langle \Phi_\sigma | H_c | \Phi_\tau \rangle]$$

=



sparse,  
diagonally  
dominant

for weakly correlated systems

for weakly correlated systems

$$(\Phi \approx \Psi)$$

for weakly correlated systems

$$(\Phi \approx \Psi)$$

$$H_0 = f_p^p \tilde{a}_p^p$$

for weakly correlated systems  $(\Phi \approx \Psi)$

$$H_0 = f_p^p \tilde{a}_p^p \quad \text{works fine}$$



for weakly correlated systems  $(\Phi \approx \Psi)$

$$H_0 = f_p^p \tilde{a}_p^p \quad \text{works fine}$$

$$H_0 \Phi = 0 \cdot \Phi$$

for weakly correlated systems  $(\Phi \approx \Psi)$

$$H_0 = f_p^p \tilde{a}_p^p \quad \text{works fine}$$

$$H_0 \Phi = 0 \cdot \Phi$$

$$H_0 \Phi_{ij\dots}^{ab\dots} = \mathcal{E}_{ij\dots}^{ab\dots} \Phi_{ij\dots}^{ab\dots}$$

for weakly correlated systems

$$(\Phi \approx \Psi)$$

$$H_0 = f_p^p \tilde{a}_p^p \quad \text{works fine}$$

$$H_0 \Phi = 0 \cdot \Phi$$

$$H_0 \Phi_{ij\dots}^{ab\dots} = \mathcal{E}_{ij\dots}^{ab\dots} \Phi_{ij\dots}^{ab\dots}$$

super convenient



Subspace

Projection  
operator

Subspace

Projection  
operator

"model space"

Subspace

Projection  
operator

"model space"  
 $\text{span}\{\Phi\}$

## Subspace

↳ contains qualitative  
approx. for  $\Psi$   
"model space"  
 $\text{span} \{ \Phi \}$

## Projection operator



Subspace

↳ contains qualitative  
approx. for  $\Psi$

"model space"

$\text{span} \{ \Phi \}$

"orthogonal space"

Projection  
operator

## Subspace

↳ contains qualitative  
approx. for  $\Psi$

"model space"

$$\text{span} \{ \Phi \}$$

## Projection operator

"orthogonal space"

$$\text{span} \{ \Phi_i^a \} \cup \{ \Phi_{ij}^{ab} \} \cup \dots$$

## Subspace

↳ contains qualitative  
approx. for  $\Psi$

"model space"

$$\text{span} \{ \Phi \}$$

↳ contains last little bit  
of  $\Psi$  to eek out quant. accuracy

"orthogonal space"

$$\text{span} \{ \Phi_i^a \} \cup \{ \Phi_{ij}^b \} \cup \dots$$

## Projection operator

## Subspace

↪ contains qualitative approx. for  $\Psi$

"model space"

$$\text{span} \{ \Phi \}$$

↪ contains last little bit of  $\Psi$  to eek out quant. accuracy

"orthogonal space"

$$\text{span} \{ \Phi_i^a \} \cup \{ \Phi_{ij}^{\perp} \} \cup \dots$$

## Projection operator

$$P = |\Phi\rangle\langle\Phi|$$

## Subspace

↪ contains qualitative approx. for  $\Psi$

"model space"

$$\text{span} \{ \Phi \}$$

↪ contains last little bit of  $\Psi$  to eek out quant. accuracy

"orthogonal space"

$$\text{span} \{ \Phi_i^a \} \cup \{ \Phi_{ij}^{\perp} \} \cup \dots$$

## Projection operator

$$P = |\Phi\rangle\langle\Phi|$$

$$Q = 1 - P$$

$$P + Q = 1$$

$$P + Q = 1 \quad P^2 = P$$

$$P + Q = 1 \quad P^2 = P \quad Q^2 = Q$$



$$P + Q = I \quad P^2 = P \quad Q^2 = Q \quad PQ = QP = 0$$

$$P + Q = 1 \quad P^2 = P \quad Q^2 = Q \quad PQ = QP = 0$$

with intermediate normalization:

$$P + Q = 1 \quad P^2 = P \quad Q^2 = Q \quad PQ = QP = 0$$

with intermediate normalization:

$$P\Psi = \Phi$$

$$P + Q = 1 \quad P^2 = P \quad Q^2 = Q \quad PQ = QP = 0$$

with intermediate normalization:

$$P\Psi = \Phi$$

$$Q\Psi = \Psi - \Phi$$



Q

$$Q = \sum_k \left(\frac{1}{k!}\right)^2 \sum_{\substack{a_1 \dots a_k \\ i_1 \dots i_k}} |\Phi_{i_1 \dots i_k}^{a_1 \dots a_k}\rangle \langle \Phi_{i_1 \dots i_k}^{a_1 \dots a_k}|$$

$$Q = \sum_{i^a} |\Phi_i^a\rangle \langle \Phi_i^a|$$

$$+ \left(\frac{1}{2!}\right)^2 \sum_{\substack{a,b \\ i,j}} |\Phi_{ij}^{ab}\rangle \langle \Phi_{ij}^{ab}|$$

$$+ \left(\frac{1}{3!}\right)^2 \sum_{\substack{a,b,c \\ i,j,k}} |\Phi_{ijk}^{abc}\rangle \langle \Phi_{ijk}^{abc}|$$

$$+ \dots$$





the resolvent:

the resolvent:

$$R_0 = -H_0^{-1}$$


the resolvent:

$$R_0 = -H_0^{-1} \big|_{\text{ortho. space}}$$

the resolvent:

$$R_0 = -H_0^{-1} \big|_{\text{ortho. space}}$$

why?



the resolvent:

$$R_0 = -H_0^{-1} \big|_{\text{ortho. space}}$$

why?

$$H_0 \big|_{\text{model space}} = 0$$

the resolvent:

$$R_0 = -H_0^{-1} \big|_{\text{ortho. space}}$$

why?

$$H_0 \big|_{\text{model space}} = 0$$

singular  $\iff$  non-invertible





$R_0$

$$R_o = -H_o^{-1}Q$$

$$R_0 = -H_0^{-1}Q$$

$$= \sum_k \left(\frac{1}{k!}\right)^2 \sum_{\substack{a_1 \dots a_k \\ i_1 \dots i_k}} (-H_0)^{-1} |\Phi_{i_1 \dots i_k}^{a_1 \dots a_k}\rangle \langle \Phi_{i_1 \dots i_k}^{a_1 \dots a_k}|$$

$$R_0 = -H_0^{-1} Q$$

$$= \sum_k \left(\frac{1}{k!}\right)^2 \sum_{\substack{a_1 \dots a_k \\ i_1 \dots i_k}} (-\varepsilon_{i_1 \dots i_k}^{a_1 \dots a_k})^{-1} |\Phi_{i_1 \dots i_k}^{a_1 \dots a_k}\rangle \langle \Phi_{i_1 \dots i_k}^{a_1 \dots a_k}|$$

$$R_0 = -H_0^{-1} Q$$

$$= \sum_k \left(\frac{1}{k!}\right)^2 \sum_{\substack{a_1 \dots a_k \\ i_1 \dots i_k}} \left( + \mathcal{E}_{a_1 \dots a_k}^{i_1 \dots i_k} \right)^{-1} | \Phi_{i_1 \dots i_k}^{a_1 \dots a_k} \rangle \langle \Phi_{i_1 \dots i_k}^{a_1 \dots a_k} |$$

$$R_0 = -H_0^{-1}Q$$

$$= \sum_i^a \frac{|\Phi_i^a\rangle\langle\Phi_i^a|}{\mathcal{E}_a^i}$$

$$+ \left(\frac{1}{2!}\right)^2 \sum_{ij}^{ab} \frac{|\Phi_{ij}^{ab}\rangle\langle\Phi_{ij}^{ab}|}{\mathcal{E}_{ab}^{ij}}$$

$$+ \left(\frac{1}{3!}\right)^2 \sum_{ijk}^{abc} \frac{|\Phi_{ijk}^{abc}\rangle\langle\Phi_{ijk}^{abc}|}{\mathcal{E}_{abc}^{ijk}}$$

$$+ \dots$$



note that



note that

$$R_0 \Phi^{ab \dots}_{ij \dots}$$

note that

$$R_0 \Phi_{ij\dots}^{ab\dots} = (\mathcal{E}_{ab\dots}^{ij\dots})^{-1} \Phi_{ij\dots}^{ab\dots}$$

note that

$$R_0 \Phi_{ij\dots}^{ab\dots} = (\mathcal{E}_{ab\dots}^{ij\dots})^{-1} \Phi_{ij\dots}^{ab\dots}$$

and

note that

$$R_0 \Phi^{ab\dots}_{ij\dots} = (\mathcal{E}^{ij\dots}_{ab\dots})^{-1} \Phi^{ab\dots}_{ij\dots}$$

and

$$R_0 \Phi$$

note that

$$R_0 \Phi^{ab\dots}_{ij\dots} = (\mathcal{E}^{ij\dots}_{ab\dots})^{-1} \Phi^{ab\dots}_{ij\dots}$$

and

$$R_0 \Phi = 0 \cdot \Phi$$



consider

consider

$$R_0 X_1 \cdots X_n | \Phi \rangle$$



consider

$$R_0 X_1 \cdots X_n |\Phi\rangle$$

$$= R_0 : \overline{X_1 \cdots X_n} : |\Phi\rangle$$

consider

$$R_0 X_1 \cdots X_n | \Phi \rangle$$

$$= R_0 : \overline{X_1 \cdots X_n} : | \Phi \rangle$$

observations:

consider

$$R_0 X_1 \cdots X_n |\Phi\rangle$$

$$= R_0 : \overline{X_1 \cdots X_n} : |\Phi\rangle$$

observations:

1. complete contractions vanish


consider

$$R_0 X_1 \cdots X_n |\Phi\rangle$$

$$= R_0 : \overline{X_1 \cdots X_n} : |\Phi\rangle$$

observations:

each term must be able to fully contract  
one of these


$$|\Phi_{ij \dots}^{ab \dots}\rangle \langle \Phi | \tilde{a}_{ab \dots}^{ij \dots}$$

such terms will have the form

such terms will have the form

$$\text{const} \times \tilde{a}_{j_1 \dots j_m}^{b_1 \dots b_m} | \Phi \rangle$$

such terms will have the form

$$\text{const} \times \tilde{a}_{j_1 \dots j_m}^{b_1 \dots b_m} |\Phi\rangle$$

$$= \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$

such terms will have the form

$$\text{const} \times \tilde{a}_{j_1 \dots j_m}^{b_1 \dots b_m} |\Phi\rangle$$

$$= \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$





such terms will have the form

$$\text{const} \times \tilde{a}_{j_1 \dots j_m}^{b_1 \dots b_m} |\Phi\rangle$$

$$= \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$

$$\Rightarrow R_0 \times \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$

such terms will have the form

$$\text{const} \times a_{j_1 \dots j_m}^{b_1 \dots b_m} |\Phi\rangle$$

$$= \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$

$$\Rightarrow R_0 \times \text{const} \times |\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle$$

$$= \text{const} \times \frac{|\Phi_{j_1 \dots j_m}^{b_1 \dots b_m}\rangle}{\sum_{j_1 \dots j_m} \sum_{b_1 \dots b_m}}$$



$$R_0 Y_1 \cdots R_0 Y_{n-2} R_0 Y_{n-1} R_0 Y_n | \mathbb{E} \rangle$$

$$R_0 Y_1 \cdots R_0 Y_{n-2} R_0 Y_{n-1} R_0 Y_n | \mathbb{E} \rangle$$

$$= R_0 Y_1 \cdots R_0 Y_{n-2} R_0 : \overline{Y_{n-1}} | Y_n : | \mathbb{E} \rangle$$

$$R_0 Y_1 \cdots R_0 Y_{n-2} R_0 Y_{n-1} R_0 Y_n | \mathbb{E} \rangle$$

$$= R_0 Y_1 \cdots R_0 Y_{n-2} R_0 : \overline{Y_{n-1} Y_n} : | \mathbb{E} \rangle$$

$$= R_0 Y_1 \cdots R_0 : \overline{Y_{n-2} Y_{n-1} Y_n} : | \mathbb{E} \rangle$$

$$R_0 Y_1 \cdots R_0 Y_{n-2} R_0 Y_{n-1} R_0 Y_n | \Xi \rangle$$

$$= R_0 Y_1 \cdots R_0 Y_{n-2} R_0 : \overline{Y_{n-1}} | Y_n : | \Xi \rangle$$

$$= R_0 Y_1 \cdots R_0 : \overline{Y_{n-2}} Y_{n-1} | Y_n : | \Xi \rangle$$

$$= \dots$$

$$R_0 Y_1 \cdots R_0 Y_{n-2} R_0 Y_{n-1} R_0 Y_n | \mathbb{E} \rangle$$

$$= R_0 Y_1 \cdots R_0 Y_{n-2} R_0 : \overline{Y_{n-1} Y_n} : | \mathbb{E} \rangle$$

$$= R_0 Y_1 \cdots R_0 : \overline{Y_{n-2} Y_{n-1} Y_n} : | \mathbb{E} \rangle$$

$$= \dots$$

$$= R_0 : \overline{Y_1 \cdots Y_{n-2} Y_{n-1} Y_n} : | \mathbb{E} \rangle$$





Perturbation theory:

Perturbation theory:

analyze dependence of  $\Psi$  or

$$0 = \langle \Psi | \hat{O} | \Psi \rangle \text{ on } V_c \equiv H_c - H_0$$

Perturbation theory:

analyze dependence of  $\Psi$  or

$$O = \langle \Psi | \hat{O} | \Psi \rangle \text{ on } V_c \equiv H_c - H_0$$

"fluctuation potential"

Perturbation theory:

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

Perturbation theory:

↙ on/off switch

$$H(\lambda) \equiv H_0 + \lambda V_c$$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

on/off switch  
 $\lambda = 1$



Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

on/off switch  
 $\lambda = 1$  "on"

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

on/off switch

$\lambda = 1$  "on"

$\lambda = 0$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch

$\lambda = 1$  "on"

$\lambda = 0$  "off"

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch  
 $\lambda = 1$  "on"  
 $\lambda = 0$  "off"

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch  
 $\lambda = 1$  "on"  
 $\lambda = 0$  "off"

$$H(\lambda) \psi(\lambda) = E(\lambda) \psi(\lambda)$$

$$\psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \psi^{(n)}$$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch  
 $\lambda = 1$  "on"  
 $\lambda = 0$  "off"

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \Psi^{(n)}$$

$$E(\lambda) = \sum_{n=0}^{\infty} \lambda^n E_c^{(n)}$$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch  
 $\lambda = 1$  "on"  
 $\lambda = 0$  "off"

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \Psi^{(n)} \rightarrow \frac{1}{n!} \left. \frac{\partial^n \Psi(\lambda)}{\partial \lambda^n} \right|_0$$

$$E(\lambda) = \sum_{n=0}^{\infty} \lambda^n E_c^{(n)}$$

Perturbation theory:

$$H(\lambda) \equiv H_0 + \lambda V_c$$

↖ on/off switch  
 $\lambda = 1$  "on"  
 $\lambda = 0$  "off"

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \Psi^{(n)} \rightarrow \frac{1}{n!} \left. \frac{\partial^n \Psi(\lambda)}{\partial \lambda^n} \right|_0$$

$$E(\lambda) = \sum_{n=0}^{\infty} \lambda^n E_c^{(n)} \rightarrow \frac{1}{n!} \left. \frac{\partial^n E(\lambda)}{\partial \lambda^n} \right|_0$$



$$H(\lambda) \psi(\lambda) = E(\lambda) \psi(\lambda)$$

$$H(\lambda) \psi(\lambda) = E(\lambda) \psi(\lambda)$$

project by  $\Phi$ :

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

project by  $\Phi$ :

$$\lambda \langle \Phi | V_c | \Psi(\lambda) \rangle = E(\lambda)$$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

project by  $\Phi$ :

$$\lambda \langle \Phi | V_c | \Psi(\lambda) \rangle = E(\lambda)$$

$\Rightarrow$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

project by  $\Phi$ :

$$\lambda \langle \Phi | V_c | \Psi(\lambda) \rangle = E(\lambda)$$

$$\Rightarrow \langle \Phi | V_c | \Psi^{(n)} \rangle = E^{(n+1)}$$

$$H(\lambda) \psi(\lambda) = E(\lambda) \psi(\lambda)$$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$R_0 H_0 \Psi(\lambda) + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$



$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$\underbrace{R_0 H_0 \Psi(\lambda)}_{-Q \Psi(\lambda)} + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$\underbrace{R_0 H_0 \Psi(\lambda)} + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$

$$-Q \Psi(\lambda) = -\Psi(\lambda) + \Phi$$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$\underbrace{R_0 H_0 \Psi(\lambda)} + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$

$$-Q \Psi(\lambda) = -\Psi(\lambda) + \Phi$$

$\Rightarrow$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$\underbrace{R_0 H_0 \Psi(\lambda)} + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$

$$-Q \Psi(\lambda) = -\Psi(\lambda) + \Phi$$

$$\Rightarrow \Psi(\lambda)$$

$$H(\lambda) \Psi(\lambda) = E(\lambda) \Psi(\lambda)$$

operate  $R_0$  on both sides:

$$\underbrace{R_0 H_0 \Psi(\lambda)} + R_0 \lambda V_c \Psi(\lambda) = E(\lambda) R_0 \Psi(\lambda)$$

$$-Q \Psi(\lambda) = -\Psi(\lambda) + \Phi$$

$$\Rightarrow \Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$



$$\Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$

$$E^{(n+1)} = \langle \Phi | V_c | \Psi^{(n)} \rangle$$

$$\Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$

$$E^{(n+1)} = \langle \Phi | V_c | \Psi^{(n)} \rangle$$

$$1. E^{(1)}$$



$$\Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$

$$E^{(n+1)} = \langle \Phi | V_c | \Psi^{(n)} \rangle$$

$$1. E^{(1)} \quad 2. \Psi^{(1)}$$

$$\Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$

$$E^{(n+1)} = \langle \Phi | V_c | \Psi^{(n)} \rangle$$

$$1. E^{(1)} \quad 2. \Psi^{(1)} \quad 3. E^{(2)}$$

$$\Psi(\lambda) = \Phi + R_0 (\lambda V_c - E(\lambda)) \Psi(\lambda)$$

$$E^{(n+1)} = \langle \Phi | V_c | \Psi^{(n)} \rangle$$

$$1. E^{(1)} \quad 2. \Psi^{(1)} \quad 3. E^{(2)} \quad 4. \Psi^{(2)}$$

end.