What Sets Bayes Apart?

Models for Socio-Environmental Data

Chris Che-Castaldo, Mary B. Collins, N. Thompson Hobbs

June 02, 2019




Housekeeping

- Introductions
- GitHub / Website for course materials
- Getting notes just in time
- Daily schedule
- Lecture / exercise mix
- Individual modeling projects

Pace

- Challenge
- Solutions
 - Working in groups
 - Questions, questions, questions
 - Advanced problems
 - A flexible schedule

Readings




Errata and explanations can be found here



Exercise

What do statements made by journalists, attorneys, and scientists have in common? What sets the statements of scientists apart?

What is this course about?



What is this course about?

Gaining insight about socio-ecological systems by building models

$$[z_i \mid \theta_p]$$

and fitting those models to data

$$[y_i \mid z_i, \theta_d]$$

using Bayesian methods.



3 A	5 B	1 B	4 B	2 A	1 A	4 A	3 B	2 B	5 A	Block 1	
2 A	5 B	4 B	2 B	4 A	3 A	1 A	1 B	3 B	5 A	Block 2	Factorial Arrangemer Treatments Randomized Complete B Design
1 A	3 B	4 B	5 B	3 A	4 A	2 A	2 B	1 B	5 A	Block 3	
5 A	2 A	1 A	4 A	3 A	1 B	3 B	5 B	4 B	2 B	Block 1	
5 B	3 B	1 B	2 B	4 B	4 A	3 A	2 A	1 A	5 A	Block 2	Factorial Arrangemen Treatments Split-Plot De

Problems poorly suited to traditional approaches:

- Multiple sources of data
- Multiple sources of uncertainty
- Missing data
- Inference across scales
- Unobservable quantities
- Multimodal data
- Derived quantities
- Forecasting
- Synthesis

SESYNC is dedicated to fostering synthetic, actionable science related to the structure, functioning, and sustainability of socio-environmental systems.



1	Design or Purpose Measurement Variables Ranked Variables Attributes									
-	Design or Purpose	Measurement variables	Ranked variables	Attributes						
1 arriable 1 arriable 1 arriable 2 arriable	Examination of a single sample	Procedure for grouping a frequency distribution, Box 2.1; stem and leaf display, Section 2.5; esting for outliers, Section 13.4 Computing median of frequency distribution, Box 4.1 Computing artifaction of the company		Confidence limits for a percentage, Section 17.1 Runs test for randomness in dichotomized data, Box 18.3						
	Comparison of a single sample with an expected frequency distribution	Normal expected frequencies, Box 6.1 Goodniess off it tests parameters from an extrinsic hypothesis, Box 17.1; from an intrinsic hypothesis, Box 17.2 Kolmogorov-Smirrov itst of goodness of fit, Box 17.3 Graphic Tests: for normality: large sample sizes, Box 6.3, small sample sizes (rankit test), Box 6.4 Test of sample statistic against expected value, Box 7.4		Binomial expected frequencies, Box 5.1 Poisson expected frequencies, Box 5.2 Goodness of fit tests: parameters from an extrinsic hypothesis, Box 17.1; from an intrinsic hypothesis, Box 17.2						
	Single classification	Single Classification anyse: suncqual sample sizes, Box 91; equal sample sizes, Box 9.4 Planned comparison of means in anova, Box 9.8; single degree of freedom comparisons of means, Box 14.10 Unplanned comparison of means. Therethod, equal sample sizes, Box 9.9; T., GT2, and Tiduey-Farmer, ungual sample sizes, Box 9.9; T., GT2, and Tiduey-Farmer, ungual sample sizes, Box 9.10; Schriff, GT2, Box 9.12; multiple confidence limits, Section 14.10 Estimate variance components: unequal sample sizes, Box 9.2; equal sample sizes, Box 9.3 Secting confidence limits to a variance component. Box 9.3 Tests of homogeneity of variances, Box 13.1 Tests of quality of means when variances are heterogeneous, Box 13.2	Kruskal-Wallis test. Bos 13.5 Unplanned comparison of opportunity of the comparison ononparametric STP, Box 17.5	Great for homogeneity of ercorntages. Boxes 17.3 and 17.8 Comparison of several samples with an expected frequency distribution, Box 17. unplanned analysis of replicated tests of goodness of fit, Box 17.5.						
	Nested classification	Two level nested anova: equal sample sizes. Box 10.1; unequal sample sizes. Box 10.4 Three-level nested anova: equal sample sizes. Box 10.3; unequal sample sizes. Box 10.5								
	Two-way or multi-way classification	Two way arous with replication, Box 11.1; without replication, Box 11.2; unequal but proportional subcless sizes, Box 11.4; with a single missing observation, Box 11.5; with a single missing observation, Box 11.2. Three way anowa, Box 12.1 More than othere way classification, Section 12.3 and Box 12.2 Test for nonadiativity in at two way anova. Box 13.4	Friedman's method for randomized blocks, Box 13.9	Three-way log-linear model, Box 17.9 Randomized blocks for frequency data (repeated testing of the same individuals Box 17.11						


Goals

- Provide principles based understanding
- Enhance intellectual satisfaction
- Foster collaboration
- Build a foundation for self-teaching

Learning outcomes

- Explain basic principles of Bayesian inference.
- Diagram and write out mathematically correct posterior and joint distributions for Bayesian models.
- Explain basics of the Markov chain Monte Carlo (MCMC) algorithm.
- Use software for implementing MCMC.
- Develop and implement hierarchical models.
- Evaluate model fit.
- Understand papers and proposals using Bayesian methods.

Learning outcomes



B. Model specification

Diagram relationship between observed and unobserved.

Write out posterior and joint distributions using general probability notation.

Choose appropriate

probability distributions.

.

C. Model implementation

Write full conditional distributions.
Write MCMC sampling algorithm.
Or
Software

Implement MCMC on simulated data.

Implement MCMC on real data.



D. Model evaluation and inference

Posterior predictive checks

Probabilistic inference from single model

Model selection, model averaging

Topics

Day 1 - 2

Principles

- Rules of probability
- Distribution theory
- Likelihood
- Moment matching
- Bayes' theorem
- Dayes incorein
- Conjugate priors

Day 3 - 8

Implementation


- MCMC
 - JAGS
 - Regression
 - Hierarchical models
- Model checking

Day 9 - 11

Advanced topics

- Model selection
 - Designed experiments
- Mixture models
- Ordinal regression
- Dynamic models
- Spatial models
- Individual problems

Cross cutting theme



$$\mu_{i} = \frac{mx_{i}^{a}}{h^{a} + x_{i}^{a}}$$

$$[a, h, m, \sigma^{2} | \mathbf{y}] \propto \prod_{i=1}^{n} [y_{i} | \mu_{i}, \sigma^{2}][a][h][m][\sigma^{2}]$$

```
model{
    a ~ dnorm(0, .0001)
    m ~ dgamma(.01, .01)
    h ~ dgamma(.01, .01)
    sigma ~ dunif(0, 5)
    for (i in 1:length(y)){
        mu[i] <- (m * x[i]^a) / (h^a + x[i]^a)
        y[i] ~ dgamma(mu[i]^2 / sigma^2, mu[i] / sigma^2)
    }
}</pre>
```

Exercise

Describe how Bayesian analysis differs from other types of statistical analysis.

Some notation

- y data
- ullet heta a parameter or other unknown quantity of interest
- $[y \mid \theta]$ The probability distribution of y conditional on θ
- $[\theta \mid y]$ The probability distribution of θ conditional on y
- $P(y \mid \theta) = p(y \mid \theta) = [y \mid \theta] = f(y \mid \theta) = f(y, \theta)$, different notation that means the same thing.

Confidence envelopes

What sets Bayes apart? An illustration using confidence envelopes.

Notes for this are in the board notes folder.

What do we do in Bayesian modeling?

- We divide the world into things that are observed (y) and things that unobserved (θ) .
- The unobserved quantities (θ) are random variables.
- The data are random variables before they are observed and fixed after they have been observed.
- We seek to understand the probability distribution of θ using fixed observations, i.e., $[\theta \mid y]$.
- Those distributions quantify our uncertainty about θ .

You can understand it

- Rules of probability
 - Conditioning and independence
 - Law of total probability
 - The chain rule of probability
- Distribution theory
- Markov chain Monte Carlo

