

Probability Concepts and Distributions

Models for Socio-Environmental Data

Chris Che-Castaldo, Mary B. Collins, N. Thompson Hobbs

May 23, 2019




Housekeeping

Errors in text:

```
http://www.stat.colostate.edu/~hooten/papers/pdf/  
Hobbs_Hooten_Bayesian_Models_2015_errata.pdf
```

Note in particular that the plot for a cumulative distribution function is wrong.

Motivation: A general approach to scientific research



Roadmap

- ▶ The rules of probability
 - ▶ conditional probability and independence
 - ▶ the law of total probability
 - ▶ the chain law of probability
- ▶ Directed acyclic graphs (Bayesian networks)
- ▶ Probability distributions for discrete and continuous random variables
- ▶ Marginal distributions
- ▶ Moment matching

What you must know and why


| Concept to be taught | Why do you need to understand this concept? |
|-------------------------------|---|
| Conditional probability | It is the foundation for Bayes' Theorem and all inferences we will make. |
| The law of total probability | Basis for the denominator of Bayes' Theorem $[y]$ |
| Factoring joint distributions | This is the procedure we will use to build models. |
| Independence | Allows us to simplify fully factored joint distributions. |
| Probability distributions | Our toolbox for fitting models to data and representing uncertainty |
| Moments | The way we summarize distributions |
| Marginal distributions | Bayesian inference is based on marginal distributions of unobserved quantities. |
| Moment matching | Allows us to embed the predictions of models into any statistical distribution |

Motivation: The essence of Bayes

Bayesian analysis is the *only* branch of statistics that treats all unobserved quantities as random variables. We seek to understand the characteristics of the probability distributions governing the behavior of these random variables.


Motivation: models of data

$$\mu_i = g(\boldsymbol{\theta}, x_i)$$



A model of the data describes our ideas about how the data arise.

Motivation: flexibility in analysis




Motivation: flexibility in analysis

| Probability model | Support for random variable |
|---------------------|------------------------------------|
| normal | real numbers |
| multivariate normal | real numbers (vectors) |
| lognormal | non-negative real numbers |
| gamma | non-negative real numbers |
| beta | 0 to 1 real numbers |
| Bernoulli | 0 or 1 |
| binomial | counts in 2 categories |
| Poisson | counts |
| multinomial | counts in > 2 categories |
| negative binomial | counts |
| Dirichlet | proportions in ≥ 2 categories |
| Cauchy | real numbers |

Motivation: flexibility in analysis






$p = 0.5$



Work flow: probability distributions

- ▶ General properties and definitions (today)
 - ▶ discrete random variables
 - ▶ continuous random variables
- ▶ Specific distributions (cheat sheet and Probability Lab 2)
- ▶ Marginal distributions (Probability Lab 3)
- ▶ Moment matching (Probability Lab 4)

How will we use probability distributions?



Used to fit models to data, to represent uncertainty in processes and parameters, and to portray prior information

Used to make inference

Key points for today

1. What makes a function a probability mass function or a probability density function?
2. How to compute moments of distributions?
3. How to approximate moments of distributions from random draws?
4. Relationships among:
 - 4.1 probability mass function
 - 4.2 probability density function
 - 4.3 cumulative distribution function
 - 4.4 quantile function

Board work on general concepts of probability distributions with previous slide on screens.

Key points for today

1. What makes a function a probability mass function or a probability density function?
2. How to compute moments of distributions?
3. How to approximate moments of distributions from random draws.
4. Relationships among:
 - 4.1 probability mass function
 - 4.2 probability density function
 - 4.3 cumulative distribution function
 - 4.4 quantile function

Break here for marginal distributions and moment matching


Work flow: probability distributions

- ▶ General properties and definitions
 - ▶ discrete random variables
 - ▶ continuous random variables
- ▶ Specific distributions (cheat sheet and Probability Lab 2)
- ▶ Marginal distributions (Probability Lab 3)
- ▶ Moment matching (Probability Lab 4)

Key points

1. Marginal distributions summarize multivariate, joint distributions as univariate distributions.
2. Moment matching allows us to compute moments of distributions in terms of parameters and parameters of distributions in terms of moments.

How will we use marginal distributions?



Marginal distributions of discrete random variables

| | a_1 | a_2 | a_3 | a_4 | $\Pr(B) \downarrow$ |
|----------------------|----------------|----------------|----------------|-----------------|---------------------|
| b_1 | $\frac{2}{20}$ | 0 | $\frac{1}{20}$ | $\frac{3}{20}$ | $\frac{6}{20}$ |
| b_2 | 0 | $\frac{4}{20}$ | 0 | $\frac{6}{20}$ | $\frac{10}{20}$ |
| b_3 | $\frac{1}{20}$ | $\frac{1}{20}$ | $\frac{1}{20}$ | 0 | $\frac{3}{20}$ |
| b_4 | 0 | 0 | 0 | $\frac{1}{20}$ | $\frac{1}{20}$ |
| $\Pr(A) \rightarrow$ | $\frac{3}{20}$ | $\frac{5}{20}$ | $\frac{2}{20}$ | $\frac{10}{20}$ | $\frac{20}{20}$ |

Marginal distributions of discrete random variables

If we have a function $[A, B]$ specifying the joint probability of the discrete random variables A and B, then

$\sum_A [A, B]$ is the marginal probability of B
and


$\sum_B [A, B]$ is the marginal probability of A.

This same idea applies to any number of jointly distributed random variables. We simply sum over all but one.

Marginal distributions of discrete random variables

Willow establishment example on board

Joint distribution of continuous random variables



Marginal distributions of continuous random variables

Exercise: If A and B are continuous random variables and we have a function $[A,B]$ that gives their joint probability density, what is the marginal distribution of A ? Of B ?

Write two equivalent equations for these marginal distributions.

Marginal distributions of continuous random variables

If we have a function $[A, B]$ specifying the joint probability of the discrete random variables A and B, then

$\int_A [A, B] dA = \int_A [B|A][A] dA$ is the marginal probability of B


and

$\int_B [A, B] dB = \int_B [A|B][B] dB$ is the marginal probability of A.

This same idea applies to any number of jointly distributed random variables. We simply integrate over all but one.


Integrating over all but one random variable is often referred to as “integrating out.”

Marginal distributions of continuous random variables



By Bscan - Own work, CC0, <https://commons.wikimedia.org/w/index.php?curid=25235145>

Motivation: flexibility in analysis



Motivation: flexibility in analysis


| Probability model | Support for random variable |
|-------------------|------------------------------------|
| normal | all real numbers |
| lognormal | non-negative real numbers |
| gamma | non-negative real numbers |
| beta | 0 to 1 real numbers |
| Bernoulli | 0 or 1 |
| binomial | counts in 2 categories |
| Poisson | counts |
| multinomial | counts in > 2 categories |
| negative binomial | counts |
| Dirichlet | proportions in ≥ 2 categories |
| Cauchy | real numbers |

$$\mu_i = g(\theta, x_i)$$

A familiar approach


$$\boldsymbol{\theta} = (\beta_0, \beta_1)'$$

$$\mu_i = g(\boldsymbol{\theta}, x_i) = \beta_0 + \beta_1 x_i$$



The problem

All distributions have parameters:



α and β are parameters of the distribution of the random variable z .

Types of parameters

| Parameter name | Function |
|---------------------------------|------------------------------|
| intensity, centrality, location | sets position on x axis |
| shape | controls dispersion and skew |
| scale, dispersion parameter | shrinks or expands width |
| rate | scale ⁻¹ |

The problem

The normal and the Poisson are the only distributions for which the parameters of the distribution are the *same* as the moments. For all other distributions, the parameters are *functions* of the moments.

$$\alpha = m_1(\mu, \sigma^2)$$

$$\beta = m_2(\mu, \sigma^2)$$

We can use these functions to “match” the moments to the parameters.

Moment matching

$$\begin{aligned}\mu_i &= g(\theta, x_i) \\ \alpha &= m_1(\mu_i, \sigma^2) \\ \beta &= m_2(\mu_i, \sigma^2) \\ &\quad [y_i | \alpha, \beta]\end{aligned}$$

Moment matching the gamma distribution

The gamma distribution: $[z|\alpha, \beta] = \frac{\beta^\alpha z^{\alpha-1} e^{-\beta z}}{\Gamma(\alpha)}$

The mean of the gamma distribution is

$$\mu = \frac{\alpha}{\beta}$$

and the variance is

$$\sigma^2 = \frac{\alpha}{\beta^2}.$$

Discover functions for α and β in terms of μ and σ^2 .

Note: $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} \frac{dt}{t}$

Answer

$$1) \mu = \frac{\alpha}{\beta}$$

$$2) \sigma^2 = \frac{\alpha}{\beta^2}$$

Solve 1 for β , substitute for β in 2), solve for α :

$$3) \alpha = \frac{\mu^2}{\sigma^2}$$

Substitute rhs 3) for α in 2), solve for β :

$$4) \beta = \frac{\mu}{\sigma^2}$$

Moment matching the beta distribution

The beta distribution gives the probability density of random variables with support on $0, \dots, 1$.

$$[z|\alpha, \beta] = \frac{z^{\alpha-1}(1-z)^{\beta-1}}{B(\alpha, \beta)}$$

$$B = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}$$

$$\mu = \frac{\alpha}{\alpha + \beta}$$

$$\sigma^2 = \frac{\alpha \beta}{(\alpha + \beta)^2 (\alpha + \beta + 1)}$$

$$\alpha = \frac{\mu^2 - \mu^3 - \mu \sigma^2}{\sigma^2}$$

$$\beta = \frac{\mu - 2\mu^2 + \mu^3 - \sigma^2 + \mu \sigma^2}{\sigma^2}$$

You need some functions...

```
#BetaMomentMatch.R
# Function for parameters from moments
shape_from_stats <- function(mu, sigma){
  a <- (mu^2-mu^3-mu*sigma^2)/sigma^2
  b <- (mu-2*mu^2+mu^3-sigma^2+mu*sigma^2)/sigma^2
  shape_ps <- c(a,b)
  return(shape_ps)
}
# Functions for moments from parameters
beta.mean=function(a,b)a/(a+b)
beta.var = function(a,b)a*b/((a+b)^2*(a+b+1))
```

Moment matching for a single parameter

We can solve for α in terms of μ and β ,

$$\mu = \frac{\alpha}{\alpha + \beta} \quad (1)$$

$$\alpha = \frac{\mu\beta}{1 - \mu}, \quad (2)$$

which allows us to use

$$\mu_i = g(\theta, x_i) \quad (3)$$

$$y_i \sim \text{beta}\left(\frac{\mu_i\beta}{1 - \mu_i}, \beta\right) \quad (4)$$

to moment match the mean alone.

Moment matching for a single parameter

The first parameter of the lognormal = α , the mean of the random variable on the log scale. The second parameter = σ_{\log}^2 , the variance of the random variable on the log scale

We often moment match the median the lognormal distribution:

$$\text{median} = \mu_i = g(\theta, x_i) \quad (5)$$

$$\mu = e^{\alpha} \quad (6)$$

$$\alpha = \log(\mu_i) \quad (7)$$

$$y_i \sim \text{lognormal}(\log(\mu_i), \sigma_{\log}^2) \quad (8)$$

In this case, σ^2 remains on log scale.

Problems continued

Do Probability Lab #4