

Community Supported Quasi-Monte Carlo (QMC) Software

Fred Hickernell (hickernell@iit.edu), Sou-Cheng Choi (schoi32@iit.edu), Aleksei Sorokin (asorokin@hawk.iit.edu)

Illinois Institute of Technology

Software Objectives

To provide QMC software that is:

- Comprised of free open source tools
- Designed for development and support
- Easy to use for non-experts
- The recognized standard

The QMC Problem

Original Problem

$$\int_T g(t) \lambda(t) dt \quad (1)$$

$g : T \rightarrow \mathbb{R}$ = original integrand
 λ = original weight

Convenient Form

$$\int_X f(x) \rho(x) dx = \int_X f(x) \gamma(dx) \quad (2)$$

γ = well defined probability measure
 $\phi : X \rightarrow T$ = change of variables
 $f : X \rightarrow \mathbb{R}$ = integrand after change of variables

(quasi-)Monte Carlo Approximation

$$\frac{1}{n} \sum_{i=1}^n f(x_i) w_i \quad (3)$$

$\hat{\gamma}_n \approx \gamma$ = discrete probability distribution
 n guarantees

$$\left| \int_X f(x) \gamma(dx) - \frac{1}{n} \sum_{i=1}^n f(x_i) w_i \right| \leq \epsilon \quad (4)$$

Integrate

Specify and generate values $f(\hat{x})$ for $\hat{x} \in \hat{\gamma}$

Arguments

- funObj
- measureObj
- distribObj
- stopcritObj

Stopping Criterion

Finds n such that Equation (4) holds

Concrete Classes

- Central Limit Theorem (IID)
- Mean Variance (Mesh)

Measure

Specify components of a general sampling method

Implemented Functions

- Standard Uniform
- Standard Gaussian
- IID Zero Mean Gaussian
- Brownian Motion
- Lattice base 2
- Sobol base 2

Function

Specify and generate values $f(\hat{x})$ for $\hat{x} \in \hat{\gamma}$

Concrete Classes

- Keister
- Asian Call

Discrete Distribution

Specify and generate $a_n \sum_{i=1}^n w_i \delta_{\hat{x}_i}(\cdot)$

Concrete Classes


- IID
- Mesh

Accumulate Data

Accumulated data required in the computation of the integral

Results

Integration Time Comparison



Future Work

- Enhance testing and examples library
- Incorporate existing components
- Expand community of contributors

References

- S.-C. T. Choi, Y. Ding, F. J. Hickernell, L. Jiang, Ll. A. JimÁlnez Rugama, D. Li, R. Jagadeeswaran, X. Tong, K. Zhang, Y. Zhang, and X. Zhou, "GAIL: Guaranteed Automatic Integration Library (versions 1.0-2.2)," http://gailgithub.github.io/GAIL_Dev/, MATLAB software, 2013-2019.

Acknowledgements

github.com/QMCSoftware/QMCSoftware.git

Other References and Acknowledgements

Larger γ and ϵ symbols

Colors and other styling

Better captions on figures

IIT and QMC logo

Spacing of Original Problem

Fix python example to match fixed figure

Integrating over unit cube?

Python Example

```
dim = 3
stopObj = CLT_Rep(
    nInit=4, nMax=2**15, absTol=.05, J=3)
measureObj = measure().IIDZMeanGaussian(
    dimension=[dim], variance=[1/2])
distribObj = Mesh(trueD=measure().mesh(
    dimension=[dim], meshType='lattice'))
sol, out = integrate(KeisterFun(),
    measureObj, distribObj, stopObj)
```

