

## 2.5 Conditional distributions

### I. Discrete random variables:

$$P(\xi = x_i, \eta = y_j) = p_{ij}, i, j = 1, 2, \dots.$$

$$\begin{aligned} P(\eta = y_j | \xi = x_i) &= \frac{P(\eta = y_j, \xi = x_i)}{P(\xi = x_i)} \\ &= \frac{p_{ij}}{p_{i\cdot}}, \end{aligned}$$

where  $j = 1, 2, \dots$ . This is the conditional distribution of  $\eta$  conditioning on  $\xi = x_i$ .

## Definition

称 $P(\eta = y_j | \xi = x_i)$ 为在 $\xi = x_i$ 的条件下 $\eta$ 的条件概率分布列, 简称为条件分布, 记为 $p_{\eta|\xi}(y_j | x_i)$ . 称

$$P(\eta \leq y | \xi = x_i) = \sum_{j: y_j \leq y} p_{\eta|\xi}(y_j | x_i)$$

为在 $\xi = x_i$ 的条件下 $\eta$ 的条件分布函数.

从条件分布的定义和 $\xi, \eta$ 的独立性的定义可知,  $\xi, \eta$ 独立的充分必要条件是对任何 $i, j \geq 1$  有

$$P(\eta = y_j | \xi = x_i) = P(\eta = y_j).$$

## Example

在独立重复伯努里试验中, 记 $p$ 为每次试验“成功”的概率,  $S_n$ 表示第 $n$ 次成功时的试验次数. 求(1) 在 $S_n = t$ 的条件下,  $S_{n+1}$ 的条件概率分布列; (2) 在 $S_{n+1} = w$ 的条件下,  $S_n$ 的条件概率分布列.

## Example

在独立重复伯努里试验中, 记 $p$ 为每次试验“成功”的概率,  $S_n$ 表示第 $n$ 次成功时的试验次数. 求(1) 在 $S_n = t$ 的条件下,  $S_{n+1}$ 的条件概率分布列; (2) 在 $S_{n+1} = w$ 的条件下,  $S_n$ 的条件概率分布列.

解 对 $t \leq w$ , 事件 $\{S_n = t, S_{n+1} = w\}$ 意味着在 $w$ 次试验中, 第 $t, w$ 次出现“成功”, 在第1次到第 $t-1$ 次中出现 $n-1$ 次“成功”, 其余均出现“失败”. 所以

$$\begin{aligned} P(S_n = t, S_{n+1} = w) &= p \cdot p \cdot \binom{t-1}{n-1} p^{n-1} q^{w-(n+1)} \\ &= \binom{t-1}{n-1} p^{n+1} q^{w-(n+1)}. \end{aligned}$$

而

$$P(S_n = t) = \binom{t-1}{n-1} p^n q^{t-n}.$$

而

$$P(S_n = t) = \binom{t-1}{n-1} p^n q^{t-n}.$$

从而在  $S_n = t$  的条件下,  $S_{n+1}$  的条件概率分布列为

$$P(S_{n+1} = w | S_n = t) = \frac{P(S_n = t, S_{n+1} = w)}{P(S_n = t)} = pq^{w-t-1}.$$

这意味着, 在  $S_n = t$  的条件下,  $S_{n+1} - S_n$  服从几何分布.

而在  $S_{n+1} = w$  的条件下,  $S_n$  的条件概率分布列为

$$\begin{aligned} P(S_n = t | S_{n+1} = w) &= \frac{P(S_n = t, S_{n+1} = w)}{P(S_{n+1} = w)} \\ &= \frac{\binom{t-1}{n-1} p^{n+1} q^{w-(n+1)}}{\binom{w-1}{n} p^{n+1} q^{w-(n+1)}} \\ &= \frac{\binom{t-1}{n-1}}{\binom{w-1}{n}}, \quad t = n, \dots, w-1. \end{aligned}$$

这一条件分布不依赖于  $p$ .

**II. Continuous case:**  $P(\xi = x) = 0$ . Given  $\xi = x$ , the conditional distribution function of  $\eta$  can be understood as

$$P(\eta \leq y | \xi = x)$$



**II. Continuous case:**  $P(\xi = x) = 0$ . Given  $\xi = x$ , the conditional distribution function of  $\eta$  can be understood as

$$\begin{aligned} & P(\eta \leq y | \xi = x) \\ &= \lim_{\Delta x \rightarrow 0} P(\eta \leq y | x < \xi \leq x + \Delta x) \\ &= \lim_{\Delta x \rightarrow 0} \frac{P(x < \xi \leq x + \Delta x, \eta \leq y)}{P(x < \xi \leq x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{F(x + \Delta x, y) - F(x, y)}{F_{\xi}(x + \Delta x) - F_{\xi}(x)}. \end{aligned}$$

$$\begin{aligned} P(\eta \leq y | \xi = x) &= \frac{\partial F / \partial x}{F'_\xi(x)} \\ &= \frac{\int_{-\infty}^y p(x, v) dv}{p_\xi(x)} = \int_{-\infty}^y \frac{p(x, v)}{p_\xi(x)} dv. \end{aligned}$$

When  $p_\xi(x) > 0$ , conditioning on  $\xi = x$ , the density of  $\eta$  is

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_\xi(x)}.$$

## Definition

设随机向量 $(\xi, \eta)$ 有联合密度函数 $p(x, y)$ ,  $\xi$ 有边际密度函数 $p_\xi(x) = \int_{-\infty}^{\infty} p(x, y)dy$ . 若在 $x$  处,  $p_\xi(x) > 0$ , 则称

$$P(\eta \leq y | \xi = x) = \int_{-\infty}^y \frac{p(x, v)}{p_\xi(x)} dv, \quad y \in \mathbf{R}$$

为在 $\xi = x$ 的条件下,  $\eta$ 的条件分布函数, 简称为条件分布, 记作 $F_{\eta|\xi}(y|x)$ . 称

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_\xi(x)}, \quad y \in \mathbf{R} \quad (1)$$

为在 $\xi = x$ 的条件下,  $\eta$ 的条件密度函数, 简称为条件密度.

若  $p_{\xi}(x) = \int_{-\infty}^{\infty} p(x, y) dy = 0$ , 则对所有的  $y$ ,  $p(x, y) = 0$ , (1)式右边是  $\frac{0}{0}$  型不定式, 通常定义  $p_{\eta|\xi}(y|x)$  的值为 0.

若  $p_{\xi}(x) = \int_{-\infty}^{\infty} p(x, y) dy = 0$ , 则对所有的  $y$ ,  $p(x, y) = 0$ , (1)式右边是  $\frac{0}{0}$  型不定式, 通常定义  $p_{\eta|\xi}(y|x)$  的值为 0.

同理, 若  $p_{\eta}(y) > 0$ , 则在  $\eta = y$  的条件下,  $\xi$  的密度函数为

$$p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

## Example

Suppose that  $(\xi, \eta) \sim N(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, r)$ , calculate the conditional density  $p_{\eta|\xi}(y|x)$ .

## Example

Suppose that  $(\xi, \eta) \sim N(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, r)$ , calculate the conditional density  $p_{\eta|\xi}(y|x)$ .

解  $(\xi, \eta)$  的联合密度为

$$p(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-r^2}} \exp \left\{ -\frac{1}{2(1-r^2)} \left[ \frac{(x-\mu_1)^2}{\sigma_1^2} - 2r \frac{(x-\mu_1)(y-\mu_2)}{\sigma_1\sigma_2} + \frac{(y-\mu_2)^2}{\sigma_2^2} \right] \right\}.$$



## Example

Suppose that  $(\xi, \eta) \sim N(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, r)$ , calculate the conditional density  $p_{\eta|\xi}(y|x)$ .

解  $(\xi, \eta)$  的联合密度为

$$p(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-r^2}} \exp \left\{ -\frac{1}{2(1-r^2)} \left[ \frac{(x-\mu_1)^2}{\sigma_1^2} - 2r \frac{(x-\mu_1)(y-\mu_2)}{\sigma_1\sigma_2} + \frac{(y-\mu_2)^2}{\sigma_2^2} \right] \right\}.$$

下面我们推导在  $\xi = x$  的条件下,  $\eta$  的条件密度. 为此, 我们不断把不含  $y$  的因子提出来, 用常数  $C_i$  表示. 最后的常数通过  $\int_{-\infty}^{\infty} p_{\eta|\xi}(y|x) dy = 1$  求得.

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{\int_{-\infty}^{\infty} p(x, y) dy} = C_1 p(x, y)$$

$$\begin{aligned} p_{\eta|\xi}(y|x) &= \frac{p(x,y)}{\int_{-\infty}^{\infty} p(x,y)dy} = C_1 p(x,y) \\ &= C_2 \exp \left\{ -\frac{1}{2(1-r^2)} \left[ \frac{(y-\mu_2)^2}{\sigma_2^2} - 2r \frac{(x-\mu_1)(y-\mu_2)}{\sigma_1\sigma_2} \right] \right\} \end{aligned}$$

$$\begin{aligned} p_{\eta|\xi}(y|x) &= \frac{p(x,y)}{\int_{-\infty}^{\infty} p(x,y)dy} = C_1 p(x,y) \\ &= C_2 \exp \left\{ -\frac{1}{2(1-r^2)} \left[ \frac{(y-\mu_2)^2}{\sigma_2^2} - 2r \frac{(x-\mu_1)(y-\mu_2)}{\sigma_1\sigma_2} \right] \right\} \\ &= C_3 \exp \left\{ -\frac{1}{2(1-r^2)} \left( \frac{y-\mu_2}{\sigma_2} - r \frac{x-\mu_1}{\sigma_1} \right)^2 \right\} \end{aligned}$$

## 2.5 Conditional distributions

$$\begin{aligned} p_{\eta|\xi}(y|x) &= \frac{p(x, y)}{\int_{-\infty}^{\infty} p(x, y) dy} = C_1 p(x, y) \\ &= C_2 \exp \left\{ -\frac{1}{2(1-r^2)} \left[ \frac{(y-\mu_2)^2}{\sigma_2^2} - 2r \frac{(x-\mu_1)(y-\mu_2)}{\sigma_1 \sigma_2} \right] \right\} \\ &= C_3 \exp \left\{ -\frac{1}{2(1-r^2)} \left( \frac{y-\mu_2}{\sigma_2} - r \frac{x-\mu_1}{\sigma_1} \right)^2 \right\} \\ &= C_3 \exp \left\{ -\frac{\left[ y - \mu_2 - r \frac{\sigma_2}{\sigma_1} (x - \mu_1) \right]^2}{2\sigma_2^2(1-r^2)} \right\}. \end{aligned}$$

上述过程可以简写为

$$p_{\eta|\xi}(y|x) \propto_y p(x, y) \propto_y \dots \\ \propto_y \exp \left\{ -\frac{\left[ y - \mu_2 - r \frac{\sigma_2}{\sigma_1} (x - \mu_1) \right]^2}{2\sigma_2^2(1 - r^2)} \right\}.$$

上述过程可以简写为

$$\begin{aligned} p_{\eta|\xi}(y|x) &\propto_y p(x, y) \propto_y \dots \\ &\propto_y \exp \left\{ -\frac{\left[ y - \mu_2 - r \frac{\sigma_2}{\sigma_1} (x - \mu_1) \right]^2}{2\sigma_2^2(1 - r^2)} \right\}. \end{aligned}$$

回顾正态分布的密度函数知 $p_{\eta|\xi}(y|x)$ 为正态密度函数

$$p_{\eta|\xi}(y|x) = \frac{1}{\sqrt{2\pi}\sigma_2\sqrt{1-r^2}} \exp \left\{ -\frac{\left[ y - \mu_2 - r \frac{\sigma_2}{\sigma_1} (x - \mu_1) \right]^2}{\underbrace{2\sigma_2^2(1-r^2)}} \right\}. \quad (2)$$


即在 $\xi = x$ 的条件下, 二维正态分布的条件分布是正态分布 $N(\mu_2 + r \frac{\sigma_2}{\sigma_1}(x - \mu_1), (1 - r^2)\sigma_2^2)$ , 记作

$$\eta|_{\xi=x} \sim N(\underbrace{\mu_2 + \frac{r\sigma_2}{\sigma_1}(x - \mu_1)}, \underbrace{(1 - r^2)\sigma_2^2}),$$

其中第一个参数 $m = \mu_2 + r \frac{\sigma_2}{\sigma_1}(x - \mu_1)$ 是 $x$ 的线性函数, 第二个参数与 $x$ 无关.



## 2.5 Conditional distributions



$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

If  $\xi$  and  $\eta$  are independent, then

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

If  $\xi$  and  $\eta$  are independent, then

$$p_{\eta|\xi}(y|x) = p_{\eta}(y), \quad p_{\xi|\eta}(x|y) = p_{\xi}(x).$$

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

$$p(x, y) = p_{\eta|\xi}(y|x)p_{\xi}(x); \quad p(x, y) = p_{\xi|\eta}(x|y)p_{\eta}(y).$$

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

$$\underline{p(x, y) = p_{\eta|\xi}(y|x)p_{\xi}(x)}; \quad \underline{p(x, y) = p_{\xi|\eta}(x|y)p_{\eta}(y)}.$$

$$\begin{aligned} \underline{p_{\xi|\eta}(x|y)} &= \frac{p(x, y)}{p_{\eta}(y)} = \frac{p(x, y)}{\int p(u, y) du} \\ &= \frac{\underline{p_{\eta|\xi}(y|x)p_{\xi}(x)}}{\int \underline{p_{\eta|\xi}(y|u)p_{\xi}(u)} du}. \end{aligned}$$

$$p_{\eta|\xi}(y|x) = \frac{p(x, y)}{p_{\xi}(x)}; \quad p_{\xi|\eta}(x|y) = \frac{p(x, y)}{p_{\eta}(y)}.$$

$$p(x, y) = p_{\eta|\xi}(y|x)p_{\xi}(x); \quad p(x, y) = p_{\xi|\eta}(x|y)p_{\eta}(y).$$

$$\begin{aligned} p_{\xi|\eta}(x|y) &= \frac{p(x, y)}{p_{\eta}(y)} = \frac{p(x, y)}{\int p(u, y) du} \\ &= \frac{p_{\eta|\xi}(y|x)p_{\xi}(x)}{\int p_{\eta|\xi}(y|u)p_{\xi}(u) du}. \end{aligned}$$

Bayesian formula



### III. The general case:

In general, suppose that the joint distribution function of  $(\xi, \eta)$  is  $F(x, y)$ . If

$$\begin{aligned} & \lim_{\Delta y \rightarrow 0} P(\xi \leq x | \eta \in (y, y + \Delta y]) \\ &= \lim_{\Delta y \rightarrow 0} \frac{P(\xi \leq x, \eta \in (y, y + \Delta y])}{P(\eta \in (y, y + \Delta y])} \\ &= \lim_{\Delta y \rightarrow 0} \frac{F(x, y + \Delta y) - F(x, y)}{F_{\eta}(y + \Delta y) - F_{\eta}(y)} \end{aligned}$$

exists for any  $x$ , we call the limit function  $F_{\xi|\eta}(x|y)$  be the conditional distribution function of  $\xi$  for given  $\eta = y$ .

If there exists  $\{x_i\}$  such that

$$F_{\xi|\eta}(x|y) = \sum_{i:x_i \leq x} p_{\xi|\eta}(x_i|y), \quad x \in \mathbf{R},$$

then we call  $p_{\xi|\eta}(x_i|y)$ ,  $i = 1, 2, \dots$ , the conditional mass function (条件分布列). If  $F_{\xi|\eta}(x|y)$  can be represented as the form

$$F_{\xi|\eta}(x|y) = \int_{-\infty}^x p_{\xi|\eta}(v|y) dv, \quad x \in \mathbf{R},$$

then we call  $p_{\xi|\eta}(x|y)$  the conditional density function.

## Example

设 $\Lambda$ 服从伽玛分布 $\Gamma(b, a)$ , 在条件 $\Lambda = \lambda$ 下,  $X$ 服从参数为 $\lambda$  的泊松分布. 求在 $X = x$ 的条件下 $\Lambda$  的分布.

## Example

设 $\Lambda$ 服从伽玛分布 $\Gamma(b, a)$ , 在条件 $\Lambda = \lambda$ 下,  $X$ 服从参数为 $\lambda$  的泊松分布. 求在 $X = x$ 的条件下 $\Lambda$  的分布.

**解**  $\Lambda$ 为连续型随机变量,  $X$ 为离散型随机变量. 对 $x = 0, 1, \dots$ , 有

$$P(X = x | \Lambda = \lambda) = \frac{\lambda^x}{x!} e^{-\lambda}.$$

### Example

设 $\Lambda$ 服从伽玛分布 $\Gamma(b, a)$ , 在条件 $\Lambda = \lambda$ 下,  $X$ 服从参数为 $\lambda$  的泊松分布. 求在 $X = x$ 的条件下 $\Lambda$  的分布.

解  $\Lambda$ 为连续型随机变量,  $X$ 为离散型随机变量. 对 $x = 0, 1, \dots$ , 有

$$P(X = x | \Lambda = \lambda) = \frac{\lambda^x}{x!} e^{-\lambda}. \quad + o(o(\lambda))$$

这意味着

$$P(X = x | \Lambda = \lambda) = \lim_{\Delta\lambda \rightarrow 0} \frac{P(X = x, \Lambda \in (\lambda, \lambda + \Delta\lambda))}{P(\Lambda \in (\lambda, \lambda + \Delta\lambda))}$$

即

$$\begin{aligned} & P(X = x, \Lambda \in (\lambda, \lambda + \Delta\lambda)) \\ &= P(X = x | \Lambda = \lambda) P(\Lambda \in (\lambda, \lambda + \Delta\lambda)) + o(\Delta\lambda) \\ &= P(X = x | \Lambda = \lambda) \underbrace{p_\Lambda(\lambda)}_{\Delta\lambda} + o(\Delta\lambda). \end{aligned}$$

即

$$\begin{aligned} & P(X = x, \Lambda \in (\lambda, \lambda + \Delta\lambda)) \\ &= P(X = x | \Lambda = \lambda) P(\Lambda \in (\lambda, \lambda + \Delta\lambda)) + o(\Delta\lambda) \\ &= P(X = x | \Lambda = \lambda) p_{\Lambda}(\lambda) \Delta\lambda + o(\Delta\lambda). \end{aligned}$$

所以

$$P(X = x, \Lambda \leq y) = \int_{-\infty}^y P(X = x | \Lambda = \lambda) p_{\Lambda}(\lambda) d\lambda.$$

即

$$\begin{aligned} & P(X = x, \Lambda \in (\lambda, \lambda + \Delta\lambda)) \\ &= P(X = x | \Lambda = \lambda) P(\Lambda \in (\lambda, \lambda + \Delta\lambda)) + o(\Delta\lambda) \\ &= P(X = x | \Lambda = \lambda) p_{\Lambda}(\lambda) \Delta\lambda + o(\Delta\lambda). \end{aligned}$$

所以

$$P(X = x, \Lambda \leq y) = \int_{-\infty}^y P(X = x | \Lambda = \lambda) p_{\Lambda}(\lambda) d\lambda.$$

从而

$$\begin{aligned} P(\Lambda \leq y | X = x) &= \frac{P(X = x, \Lambda \leq y)}{P(X = x)} \\ &= \int_{-\infty}^y \frac{P(X = x | \Lambda = \lambda) p_{\Lambda}(\lambda)}{P(X = x)} d\lambda. \end{aligned}$$



因此在  $X = x$  的条件下,  $\Lambda$  的密度函数为

$$\begin{aligned} p_{\Lambda|\xi}(\lambda|x) &= \frac{P(X = x|\Lambda = \lambda)p_{\Lambda}(\lambda)}{P(X = x)} \\ &= \frac{P(X = x|\Lambda = \lambda)p_{\Lambda}(\lambda)}{\int_{-\infty}^{\infty} P(X = x|\Lambda = \lambda)p_{\Lambda}(\lambda)d\lambda} \\ &\propto \lambda^x e^{-\lambda} \lambda^{b-1} e^{-\lambda a} = \lambda^{x+b-1} e^{-(a+1)\lambda}, \quad \lambda > 0. \end{aligned}$$

因此在 $X = x$ 的条件下,  $\Lambda$ 的密度函数为

$$\begin{aligned}
 p_{\Lambda|\xi}(\lambda|x) &= \frac{P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)}{P(X=x)} \\
 &= \frac{P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)}{\int_{-\infty}^{\infty} P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)d\lambda} \\
 &\propto_{\lambda} \lambda^x e^{-\lambda} \lambda^{b-1} e^{-a\lambda} = \lambda^{x+b-1} e^{-(a+1)\lambda}, \quad \lambda > 0.
 \end{aligned}$$

将上式右边添加正则化常数因子使得其积分为1, 得

$$p_{\Lambda|\xi}(\lambda|x) = \frac{(a+1)^{x+b}}{\Gamma(x+b)} \lambda^{x+b-1} e^{-(a+1)\lambda}, \quad \lambda > 0.$$

即在 $X = x$ 的条件下,  $\Lambda$ 服从伽玛分布 $\Gamma(x+b, a+1)$ .

$$\begin{aligned} p_{\Lambda|X}(\lambda|x) &= \frac{P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)}{P(X=x)} \\ &= \frac{P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)}{\int_{-\infty}^{\infty} P(X=x|\Lambda=\lambda)p_{\Lambda}(\lambda)d\lambda} \end{aligned}$$

Bayes 公式.

## IV. Multi-dimensional case:

Suppose that the joint distribution function of random vectors  $\xi$  and  $\eta$  is  $F(x, y)$ . If

$$\begin{aligned}
 & \lim_{\Delta y \rightarrow 0} P(\xi \leq x | \eta \in (y, y + \Delta y]) \\
 &= \lim_{\Delta y \rightarrow 0} \frac{P(\xi \leq x, \eta \in (y, y + \Delta y])}{P(\eta \in (y, y + \Delta y])} \\
 &= \lim_{\Delta y \rightarrow 0} \frac{F(x, y + \Delta y) - F(x, y)}{F_\eta(y + \Delta y) - F_\eta(y)}
 \end{aligned}$$

exists for any  $x$ , we call the limit function  $F_{\xi|\eta}(x|y)$  be the conditional distribution function of  $\xi$  for given  $\eta = y$ .

$P(\xi \leq x | \eta = y)$

When  $(\boldsymbol{\xi}, \boldsymbol{\eta})$  is a continuous random vector with probability density function  $p(\boldsymbol{x}, \boldsymbol{y})$ , the conditional probability density function of  $\boldsymbol{\xi}$  for given  $\boldsymbol{\eta} = \boldsymbol{y}$  is

$$p_{\boldsymbol{\xi}|\boldsymbol{\eta}}(\boldsymbol{x}|\boldsymbol{y}) = \frac{p(\boldsymbol{x}, \boldsymbol{y})}{p_{\boldsymbol{\eta}}(\boldsymbol{y})} = \frac{p(\boldsymbol{x}, \boldsymbol{y})}{\int p(\boldsymbol{u}, \boldsymbol{y}) d\boldsymbol{u}},$$

if  $p_{\boldsymbol{\eta}}(\boldsymbol{y}) > 0$ .

When  $(\boldsymbol{\xi}, \boldsymbol{\eta})$  is a discrete random vector with probability mass function  $P(\boldsymbol{\xi} = \boldsymbol{x}_i, \boldsymbol{\eta} = \boldsymbol{y}_j) = p(\boldsymbol{x}_i, \boldsymbol{y}_j)$ , the conditional probability mass function of  $\boldsymbol{\xi}$  for given  $\boldsymbol{\eta} = \boldsymbol{y}_j$  is

$$p_{\boldsymbol{\xi}|\boldsymbol{\eta}}(\boldsymbol{x}_i|\boldsymbol{y}_j) = \frac{p(\boldsymbol{x}_i, \boldsymbol{y}_j)}{p_{\boldsymbol{\eta}}(\boldsymbol{y}_j)} = \frac{P(\boldsymbol{\xi} = \boldsymbol{x}_i, \boldsymbol{\eta} = \boldsymbol{y}_j)}{P(\boldsymbol{\eta} = \boldsymbol{y}_j)},$$

$\underbrace{\hspace{1.5cm}}_{\text{if } P(\boldsymbol{\eta} = \boldsymbol{y}_j) > 0.}$