

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

§4.2 Convergence in probability and weak law of large numbers

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Consider the event A in random trial E . Suppose the probability of occurring A is p ($0 < p < 1$). Now we experiment independently n times— n -fold Bernoulli trial. Let

$$\xi_i = \begin{cases} 1, & A \text{ occurs at the } i\text{-th trial,} \\ 0, & A \text{ does not occur at the } i\text{-th trial,} \end{cases}$$

$1 \leq i \leq n$. Then $P(\xi_i = 1) = p$, $P(\xi_i = 0) = 1 - p$. Let $S_n = \sum_{i=1}^n \xi_i$. Then

$$\frac{S_n}{n} = F_n(A) \text{ --- the frequency of } A.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

For any $\varepsilon > 0$ we can not expect that $|S_n/n - p| \leq \varepsilon$ holds for all the trials even if n is big enough.

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

For any $\varepsilon > 0$ we can not expect that $|S_n/n - p| \leq \varepsilon$ holds for all the trials even if n is big enough.

It is nature to hope that the probability to appear $\{|S_n/n - p| \geq \varepsilon\}$ could be as smaller as possible when n is large enough.

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Weak laws of large numbers

Theorem

(Bernoulli) Let $\{\xi_n, n \geq 1\}$ be a sequence of *independent and identically distributed* random variables with $P(\xi_n = 1) = p$, $P(\xi_n = 0) = 1 - p$, $0 < p < 1$. Put $S_n = \sum_{i=1}^n \xi_i$. Then we have

$$P\left(\left|\frac{S_n}{n} - p\right| \geq \epsilon\right) \rightarrow 0, \quad \forall \epsilon > 0,$$

i.e., for any $\epsilon > 0$ and $\delta > 0$, there is a $N = N(\epsilon, \delta)$ such that

$$P\left(\left|\frac{S_n}{n} - p\right| \geq \epsilon\right) < \delta, \quad \text{for all } n \geq N.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof. By the Chebyshev inequality,

$$P \left(\left| \frac{S_n}{n} - p \right| \geq \epsilon \right)$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof. By the Chebyshev inequality,

$$P\left(\left|\frac{S_n}{n} - p\right| \geq \epsilon\right) \leq \frac{1}{\epsilon^2} \text{Var}\left(\frac{S_n}{n}\right)$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof. By the Chebyshev inequality,

$$\begin{aligned} P\left(\left|\frac{S_n}{n} - p\right| \geq \epsilon\right) &\leq \frac{1}{\epsilon^2} \text{Var}\left(\frac{S_n}{n}\right) \\ &= \frac{1}{\epsilon^2} \frac{np(1-p)}{n^2} = \frac{1}{\epsilon^2} \frac{p(1-p)}{n} \rightarrow 0. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Theorem

(Chebyshev) Let $\{\xi_n, n \geq 1\}$ be a sequence of *independent* (or *pairwise correlated*) random variables defined on the probability space (Ω, \mathcal{F}, P) with $E\xi_n = \mu_n$ and $\text{Var}\xi_n = \sigma_n^2$. If $\sum_{k=1}^n \sigma_k^2 / n^2 \rightarrow 0$, then $\{\xi_n, n \geq 1\}$ obeys the weak law of large numbers, i.e.,

$$P \left(\left| \frac{1}{n} \sum_{k=1}^n \xi_k - \frac{1}{n} \sum_{k=1}^n \mu_k \right| \geq \epsilon \right) \rightarrow 0, \quad \forall \epsilon > 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Using the Chebyshev inequality, we have

$$\begin{aligned} & P\left(\left|\frac{1}{n} \sum_{k=1}^n (\xi_k - \mu_k)\right| \geq \varepsilon\right) \\ & \leq P\left(\left|\frac{1}{n} \sum_{k=1}^n \xi_k - E \frac{1}{n} \sum_{k=1}^n \xi_k\right| \geq \varepsilon\right) \leq \frac{1}{\varepsilon^2} \text{Var}\left(\frac{1}{n} \sum_{k=1}^n \xi_k\right) \\ & = \frac{1}{\varepsilon^2 n^2} \sum_{k=1}^n \sigma_k^2 \longrightarrow 0 ; \text{ as } n \longrightarrow \infty. \end{aligned}$$

The proof is complete.

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$, where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.
Proof.

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$, where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.

Proof. We have $E\xi_k = 0$, $Var\xi_k = k^{2s}$. When $s < 1/2$,

$$\frac{1}{n^2} \sum_{k=1}^n Var\xi_k = \frac{1}{n^2} \sum_{k=1}^n k^{2s} < \frac{1}{n^2} \sum_{k=1}^n n^{2s} = n^{2s-1} \longrightarrow 0.$$

In addition, $\{\xi_k, k \geq 1\}$ is also independent,

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$, where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.

Proof. We have $E\xi_k = 0$, $Var\xi_k = k^{2s}$. When $s < 1/2$,

$$\frac{1}{n^2} \sum_{k=1}^n Var\xi_k = \frac{1}{n^2} \sum_{k=1}^n k^{2s} < \frac{1}{n^2} \sum_{k=1}^n n^{2s} = n^{2s-1} \rightarrow 0.$$

In addition, $\{\xi_k, k \geq 1\}$ is also independent, so $\{\xi_k, k \geq 1\}$ obeys the Chebyshev LLN, i.e.,

$$P\left(\left|\frac{1}{n} \sum_{k=1}^n \xi_k\right| \geq \epsilon\right) \rightarrow 0, \quad \forall \epsilon > 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Corollary

Let $\{\xi_n, n \geq 1\}$ be a sequence of *independent and identically distributed* random variables defined (Ω, \mathcal{F}, P) with $\text{Var}(\xi_1) < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then $\{\xi_n, n \geq 1\}$ obeys the weak LLN, i.e.,

$$P\left(\left|\frac{S_n}{n} - \mu\right| \geq \epsilon\right) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad \forall \epsilon > 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Theorem

(Khinchine) Let $\{\xi_n, n \geq 1\}$ be a sequence of *independent and identically distributed* random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then $\{\xi_n, n \geq 1\}$ obeys the weak LLN, i.e.,

$$P\left(\left|\frac{S_n}{n} - \mu\right| \geq \epsilon\right) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad \forall \epsilon > 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof : Let $M_k = \sqrt{k}$,

$$\eta_k = \xi_k I\{|\xi_k| \leq M_k\} = \begin{cases} \xi_k, & \text{if } |\xi_k| \leq M_k, \\ 0, & \text{if } |\xi_k| > M_k, \end{cases}$$

$$\zeta_k = \xi_k - \eta_k = \xi_k I\{|\xi_k| > M_k\} = \begin{cases} 0, & \text{if } |\xi_k| \leq M_k, \\ \xi_k, & \text{if } |\xi_k| > M_k. \end{cases}.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Then

$$\begin{aligned} & P\left(\frac{|\sum_{k=1}^n(\eta_k - E\eta_k)|}{n} \geq \epsilon/2\right) \\ & \leq \frac{4}{\epsilon^2 n^2} \sum_{k=1}^n \text{Var}(\eta_k) \leq \frac{4}{\epsilon^2 n^2} \sum_{k=1}^n E\eta_k^2 \\ & = \sum_{k=1}^n \frac{4}{\epsilon^2 n^2} E[\xi_k^2 I\{|\xi_k| \leq M_k\}] \leq \frac{4}{\epsilon^2} E\left[\frac{\xi_1^2 I\{|\xi_1| \leq M_n\}}{n}\right] \\ & \leq \frac{4M_n}{n\epsilon^2} E[|\xi_1|] \rightarrow 0. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

$$\begin{aligned} P\left(\frac{|\sum_{k=1}^n(\zeta_k - E\zeta_k)|}{n} \geq \epsilon/2\right) &\leq \frac{2}{\epsilon n} E\left|\sum_{k=1}^n(\zeta_k - E\zeta_k)\right| \\ &\leq \frac{2}{\epsilon n} \sum_{k=1}^n E|\zeta_k - E\zeta_k| \leq 2\frac{2}{\epsilon n} \sum_{k=1}^n E|\zeta_k| \leq \frac{4}{\epsilon} \frac{1}{n} \sum_{k=1}^n E[|\xi_1| I\{|\xi_1| > M_k\}] \rightarrow 0. \end{aligned}$$

Hence,

$$\begin{aligned} &P\left(\frac{|\sum_{k=1}^n(\xi_k - E\xi_k)|}{n} \geq \epsilon\right) \\ &\leq P\left(\frac{|\sum_{k=1}^n(\zeta_k - E\zeta_k)|}{n} \geq \epsilon/2\right) + P\left(\frac{|\sum_{k=1}^n(\eta_k - E\eta_k)|}{n} \geq \epsilon/2\right) \\ &\rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Corollary

Let $\{\xi_n, n \geq 1\}$ be a sequence of *pairwise independent* and *identically distributed* random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then $\{\xi_n, n \geq 1\}$ obeys the weak LLN, i.e.,

$$P\left(\left|\frac{S_n}{n} - \mu\right| \geq \epsilon\right) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad \forall \epsilon > 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

4.2.1 Convergence in probability

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

4.2.1 Convergence in probability

Definition

Suppose ξ and $\{\xi_n, n \geq 1\}$, are defined on the same probability space (Ω, \mathcal{F}, P) . If for any $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} P(|\xi_n - \xi| \geq \varepsilon) = 0,$$

or equivalently $\lim_{n \rightarrow \infty} P(|\xi_n - \xi| < \varepsilon) = 1$, then we say that ξ_n converges to ξ in probability, written $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Throwing a dot in $[0, 1]$ randomly, the dot is located any point in $[0, 1]$ with the same possibility. Let ω denote the location of dot and define

$$\xi(\omega) = \begin{cases} 1, & \omega \in [0, 0.5], \\ 0, & \omega \in (0.5, 1], \end{cases} \quad \eta(\omega) = \begin{cases} 0, & \omega \in [0, 0.5], \\ 1, & \omega \in (0.5, 1]. \end{cases}$$

Then ξ and η have the same distribution function

$$F(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{2}, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

If we define $\xi_n = \xi$, for $n \geq 1$, then $\xi_n \xrightarrow{d} \eta$, but $|\xi_n - \eta| \equiv 1$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

- 1 Suppose ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on the probability space (Ω, \mathcal{F}, P) .
- (1) If $\xi_n \xrightarrow{P} \xi$, then $\xi_n \xrightarrow{d} \xi$.
- (2) If $\xi_n \xrightarrow{d} c$, where c is a **constant**, then $\xi_n \xrightarrow{P} c$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

1 Suppose ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on the probability space (Ω, \mathcal{F}, P) .

(1) If $\xi_n \xrightarrow{P} \xi$, then $\xi_n \xrightarrow{d} \xi$.

(2) If $\xi_n \xrightarrow{d} c$, where c is a **constant**, then $\xi_n \xrightarrow{P} c$.

Proof. (1) Let F and F_n be the cdfs of ξ and ξ_n respectively, and let x be a continuity point of F .

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

For any $\varepsilon > 0$,

$$\begin{aligned}(\xi_n \leq x) &= (\xi_n \leq x, |\xi_n - \xi| < \varepsilon) \\ &\quad + (\xi_n \leq x, |\xi_n - \xi| \geq \varepsilon) \\ &\subset (\xi \leq x + \varepsilon) \cup (|\xi_n - \xi| \geq \varepsilon).\end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

For any $\varepsilon > 0$,

$$\begin{aligned}(\xi_n \leq x) &= (\xi_n \leq x, |\xi_n - \xi| < \varepsilon) \\ &\quad + (\xi_n \leq x, |\xi_n - \xi| \geq \varepsilon) \\ &\subset (\xi \leq x + \varepsilon) \cup (|\xi_n - \xi| \geq \varepsilon).\end{aligned}$$

Thus

$$F_n(x) \leq F(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Since $\xi_n \xrightarrow{P} \xi$ as $n \rightarrow \infty$, we obtain

$$P(|\xi_n - \xi| \geq \varepsilon) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Since $\xi_n \xrightarrow{P} \xi$ as $n \rightarrow \infty$, we obtain

$$P(|\xi_n - \xi| \geq \varepsilon) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus

$$\limsup_{n \rightarrow \infty} F_n(x) \leq F(x + \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

So

$$F(x - \varepsilon) \leq F_n(x) + P(|\xi_n - \xi| \geq \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

So

$$F(x - \varepsilon) \leq F_n(x) + P(|\xi_n - \xi| \geq \varepsilon).$$

Thus

$$F(x - \varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x).$$

We conclude that

$$F(x - \varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x) \leq \limsup_{n \rightarrow \infty} F_n(x) \leq F(x + \varepsilon).$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

We conclude that

$$F(x - \varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x) \leq \limsup_{n \rightarrow \infty} F_n(x) \leq F(x + \varepsilon).$$

Letting $\varepsilon \rightarrow 0$ yields

$$\lim_{n \rightarrow \infty} F_n(x) = F(x).$$

That is

$$\xi_n \xrightarrow{d} \xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

(2) If $\xi_n \xrightarrow{d} c$, then

$$\lim_{n \rightarrow \infty} F_n(x) = \begin{cases} 0, & x < c, \\ 1, & x > c. \end{cases}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

(2) If $\xi_n \xrightarrow{d} c$, then

$$\lim_{n \rightarrow \infty} F_n(x) = \begin{cases} 0, & x < c, \\ 1, & x > c. \end{cases}$$

Hence for any $\varepsilon > 0$,

$$\begin{aligned} & P(|\xi_n - c| > \varepsilon) \\ &= P(\xi_n > c + \varepsilon) + P(\xi_n < c - \varepsilon) \\ &= 1 - P(\xi_n \leq c + \varepsilon) + P(\xi_n < c - \varepsilon) \\ &= 1 - F_n(c + \varepsilon) + F_n(c - \varepsilon - 0) \\ &\rightarrow 0 \quad (n \rightarrow \infty). \end{aligned}$$

Example

(Khinchine LLN) Let $\{\xi_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then

$$\frac{S_n}{n} \xrightarrow{P} \mu.$$

Example

(Khinchine LLN) Let $\{\xi_n, n \geq 1\}$ be a sequence of **independent and identically distributed** random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then

$$\frac{S_n}{n} \xrightarrow{P} \mu.$$

Proof. Since the limit μ is a constant, it is sufficient to show that

$$\frac{S_n}{n} \xrightarrow{d} \mu.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \longrightarrow 0,$$

since $E\xi_1 = \mu$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \longrightarrow 0,$$

since $E\xi_1 = \mu$. For every $t \in \mathbf{R}$,

$$f(t/n) = 1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty,$$

$$f_n(t) = \left(1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right)\right)^n \rightarrow e^{i\mu t}.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \rightarrow 0,$$

since $E\xi_1 = \mu$. For every $t \in \mathbf{R}$,

$$f(t/n) = 1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty,$$

$$f_n(t) = \left(1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right)\right)^n \rightarrow e^{i\mu t}.$$

By the inverse limit theorem in we know that $S_n/n \xrightarrow{d} \mu$. So, we have $S_n/n \xrightarrow{P} \mu$. The proof is complete.

- 2 Let $\{\xi, \xi_n, n \geq 1\}$ be a sequence of random variables defined on the probability space (Ω, \mathcal{F}, P) . Prove that
- (1) If $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$, then $P(\xi = \eta) = 1$.
 - (2) If $\xi_n \xrightarrow{P} \xi$, f is the continuous function on $(-\infty, \infty)$, then $f(\xi_n) \xrightarrow{P} f(\xi)$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

- 2 Let $\{\xi, \xi_n, n \geq 1\}$ be a sequence of random variables defined on the probability space (Ω, \mathcal{F}, P) . Prove that
- (1) If $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$, then $P(\xi = \eta) = 1$.
 - (2) If $\xi_n \xrightarrow{P} \xi$, f is the continuous function on $(-\infty, \infty)$, then $f(\xi_n) \xrightarrow{P} f(\xi)$.

In general, if $\boldsymbol{\xi}_n =: (\xi_{n,1}, \dots, \xi_{n,m}) \xrightarrow{P} \boldsymbol{\xi} := (\xi_1, \dots, \xi_m)$

(i.e., $\|\boldsymbol{\xi}_n - \boldsymbol{\xi}\| \rightarrow 0$, or equivalently, $\xi_{n,k} \rightarrow \xi_k$, $k = 1, \dots, m$)

and $f(\boldsymbol{x})$ is a m -dimensional continuous function, then

$$f(\boldsymbol{\xi}_n) \xrightarrow{P} f(\boldsymbol{\xi}).$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Note that the left side of the above inequality is independent of n and $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$ as $n \rightarrow \infty$. Therefore $P(|\xi - \eta| \geq \varepsilon) = 0$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Note that the left side of the above inequality is independent of n and $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$ as $n \rightarrow \infty$. Therefore $P(|\xi - \eta| \geq \varepsilon) = 0$. Furthermore,

$$\begin{aligned} P(|\xi - \eta| > 0) &= P\left(\bigcup_{n=1}^{\infty} (|\xi - \eta| \geq \frac{1}{n})\right) \\ &\leq \sum_{n=1}^{\infty} P(|\xi - \eta| \geq \frac{1}{n}) = 0, \end{aligned}$$

i.e., $P(\xi = \eta) = 1$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

(2). For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever $\|\mathbf{x} - \mathbf{y}\| < \delta$, $\|\mathbf{x}\| \leq M$, $\|\mathbf{y}\| \leq M$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

(2). For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever $\|\mathbf{x} - \mathbf{y}\| < \delta$, $\|\mathbf{x}\| \leq M$, $\|\mathbf{y}\| \leq M$. Then

$$\begin{aligned} & \{|f(\mathbf{x}) - f(\mathbf{y})| \geq \epsilon\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{x}\| > M\} \cup \{\|\mathbf{y}\| > M\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{y}\| > M/2\}. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

(2). For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever $\|\mathbf{x} - \mathbf{y}\| < \delta$, $\|\mathbf{x}\| \leq M$, $\|\mathbf{y}\| \leq M$. Then

$$\begin{aligned} & \{|f(\mathbf{x}) - f(\mathbf{y})| \geq \epsilon\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{x}\| > M\} \cup \{\|\mathbf{y}\| > M\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{y}\| > M/2\}. \end{aligned}$$

So,

$$\begin{aligned} & P(|f(\boldsymbol{\xi}_n) - f(\boldsymbol{\xi})| \geq \epsilon) \\ & \leq P(\|\boldsymbol{\xi}_n - \boldsymbol{\xi}\| \geq \delta) + P(\|\boldsymbol{\xi}\| > M/2) \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$ and then $M \rightarrow \infty$.

3 We have

1 If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

③ We have

① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;

② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

③ We have

① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;

② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;

③ If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, both η_n and c are not 0, then $\xi_n / \eta_n \xrightarrow{P} \xi / c$;

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

③ We have

- ① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;
- ② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;
- ③ If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, both η_n and c are not 0, then $\xi_n / \eta_n \xrightarrow{P} \xi / c$;
- ④ If $\xi_n \xrightarrow{d} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, then $\xi_n + \eta_n \xrightarrow{d} \xi + c$, $\eta_n \xi_n \xrightarrow{d} c\xi$.

Proof. We only give a proof of (3) and (4).

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$. For any $\epsilon > 0$, let $\delta = \min\{\epsilon \frac{1}{2} c^2, \frac{1}{2} |c|\}$. If $|\eta_n - c| < \delta$, then $|\eta_n| > |c| - \delta > \frac{1}{2} |c|$, and so

$$|\eta_n^{-1} - c^{-1}| = \frac{|\eta_n - c|}{|\eta_n||c|} < \frac{\epsilon \frac{1}{2} c^2}{\frac{1}{2} |c| \cdot |c|} = \epsilon.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$. For any $\epsilon > 0$, let $\delta = \min\{\epsilon \frac{1}{2} c^2, \frac{1}{2} |c|\}$. If $|\eta_n - c| < \delta$, then $|\eta_n| > |c| - \delta > \frac{1}{2} |c|$, and so

$$|\eta_n^{-1} - c^{-1}| = \frac{|\eta_n - c|}{|\eta_n||c|} < \frac{\epsilon \frac{1}{2} c^2}{\frac{1}{2} |c| \cdot |c|} = \epsilon.$$

It follows that

$$P(|\eta_n^{-1} - c^{-1}| \geq \epsilon) \leq P(|\eta_n - c| \geq \delta) \rightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

For (4), it suffices to show that for any bounded continuous function $g(x, y)$ we have

$$Eg(\xi_n, \eta_n) \rightarrow Eg(\xi, c). \quad (*)$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

For (4), it suffices to show that for any bounded continuous function $g(x, y)$ we have

$$Eg(\xi_n, \eta_n) \rightarrow Eg(\xi, c). \quad (*)$$

If fact, choosing $g(x, y) = e^{it(x+y)}$ and $g(x, y) = e^{itxy}$ yields

$$Ee^{it(\xi_n + \eta_n)} \rightarrow Ee^{it(\xi + c)}, Ee^{it(\xi_n \eta_n)} \rightarrow Ee^{it(c\xi)},$$

respectively, which completes the proof by the inverse limit theorem.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Now, suppose $g(x, y)$ is a continuous function with $|g(x, y)| \leq M$, then it is uniformly continuous in any bounded area. So for any given $\epsilon > 0$ and any $A > 0$ there exist a $\delta = \delta(A, \epsilon, g) > 0$ such that $|g(\xi_n, \eta_n) - g(\xi_n, c)| \leq \epsilon$ whenever $|\eta_n - c| \leq \delta$ and $|\xi_n| \leq A$.

Then

$$\begin{aligned} & |Eg(\xi_n, \eta_n) - Eg(\xi, c)| \\ \leq & |Eg(\xi_n, \eta_n) - Eg(\xi_n, c)| + |Eg(\xi_n, c) - Eg(\xi, c)| \\ \leq & E[|g(\xi_n, \eta_n) - g(\xi_n, c)|] + |Eg(\xi_n, c) - Eg(\xi, c)| \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Then

$$\begin{aligned} & |Eg(\xi_n, \eta_n) - Eg(\xi, c)| \\ \leq & |Eg(\xi_n, \eta_n) - Eg(\xi_n, c)| + |Eg(\xi_n, c) - Eg(\xi, c)| \\ \leq & E[|g(\xi_n, \eta_n) - g(\xi_n, c)|] + |Eg(\xi_n, c) - Eg(\xi, c)| \\ \leq & \epsilon + 2MP(|\eta_n - c| > \delta) \\ & + |Eg(\xi_n, c) - Eg(\xi, c)| + 2MP(|\xi_n| > A). \end{aligned}$$

The second term will converge to zero because $\eta_n \xrightarrow{P} c$. The third will also converge to zero because $\xi_n \xrightarrow{d} \xi$ and $g(x, c)$ is a continuous function of x .

For the fourth term, we can choose A such that $\pm A$ is continuous points of the distribution function of ξ . Then $2MP(|\xi_n| > A)$ will converges to

$$2MP(|\xi| > A),$$

which can be smaller than the given $\epsilon > 0$ if A is large enough. Finally, by the arbitrariness of ϵ , $(*)$ is proved.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

④ $\xi_n \xrightarrow{P} \xi$ if and only if

$$E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} \longrightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$P(|\xi_n - \xi| > \varepsilon) = \int_{|x| > \varepsilon} dF_n(x) = \int_{|x| > \varepsilon} \frac{1 + |x|}{|x|} \frac{|x|}{1 + |x|} dF_n(x)$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$\begin{aligned} P(|\xi_n - \xi| > \varepsilon) &= \int_{|x| > \varepsilon} dF_n(x) = \int_{|x| > \varepsilon} \frac{1 + |x|}{|x|} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \int_{|x| > \varepsilon} \frac{1 + \varepsilon}{\varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$\begin{aligned} P(|\xi_n - \xi| > \varepsilon) &= \int_{|x| > \varepsilon} dF_n(x) = \int_{|x| > \varepsilon} \frac{1 + |x|}{|x|} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \int_{|x| > \varepsilon} \frac{1 + \varepsilon}{\varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \frac{1 + \varepsilon}{\varepsilon} E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

That is $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} &= \int_{-\infty}^{+\infty} \frac{|x|}{1 + |x|} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} &= \int_{-\infty}^{+\infty} \frac{|x|}{1 + |x|} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \frac{\varepsilon}{1 + \varepsilon} + \int_{|x| \geq \varepsilon} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} &= \int_{-\infty}^{+\infty} \frac{|x|}{1 + |x|} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \frac{\varepsilon}{1 + \varepsilon} + \int_{|x| \geq \varepsilon} dF_n(x) \\ &= \frac{\varepsilon}{1 + \varepsilon} + P(|\xi_n - \xi| \geq \varepsilon). \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} &= \int_{-\infty}^{+\infty} \frac{|x|}{1 + |x|} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{|x|}{1 + |x|} dF_n(x) \\ &\leq \frac{\varepsilon}{1 + \varepsilon} + \int_{|x| \geq \varepsilon} dF_n(x) \\ &= \frac{\varepsilon}{1 + \varepsilon} + P(|\xi_n - \xi| \geq \varepsilon). \end{aligned}$$

Since $\xi_n \xrightarrow{P} \xi$, first letting $n \rightarrow \infty$ and then letting $\varepsilon \rightarrow 0$ yield

$$E \frac{|\xi_n - \xi|}{1 + |\xi_n - \xi|} \longrightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let

$$\rho(\xi, \eta) = E \frac{|\xi - \eta|}{1 + |\xi - \eta|}.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let

$$\rho(\xi, \eta) = E \frac{|\xi - \eta|}{1 + |\xi - \eta|}.$$

Theorem

$\rho(\cdot, \cdot)$ satisfies

- $\rho(\xi, \eta) = 0$ if and only if $P(\xi = \eta) = 1$;
- $\rho(\xi, \eta) = \rho(\eta, \xi)$;
- $\rho(\xi, \tau) \leq \rho(\xi, \eta) + \rho(\eta, \tau)$.

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;
- $(\mathfrak{R}, \rho) = (\mathfrak{R}, \xrightarrow{P})$;

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;
- $(\mathfrak{R}, \rho) = (\mathfrak{R}, \xrightarrow{P})$;
- (\mathfrak{R}, ρ) is complete, i.e., $\xi_n - \xi_m \xrightarrow{P} 0$ as $n, m \rightarrow \infty$ if and only if there exists a random variable ξ such that $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

5 (*Dominated convergence theorem*) Suppose $\xi_n \xrightarrow{P} \xi$,
 $P(|\xi_n| \leq \eta) = 1$ and $E\eta < \infty$. Then

$$E\xi_n \rightarrow E\xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

5 (*Dominated convergence theorem*) Suppose $\xi_n \xrightarrow{P} \xi$,
 $P(|\xi_n| \leq \eta) = 1$ and $E\eta < \infty$. Then

$$E\xi_n \rightarrow E\xi.$$

Proof. First, we have $P(|\xi| \leq \eta) = 1$. In fact, for any $\epsilon > 0$,

$$\begin{aligned} P(|\xi| > \eta + \epsilon) &= P(|\xi| > \eta + \epsilon, |\xi_n - \xi| < \epsilon) \\ &\quad + P(|\xi| > \eta + \epsilon, |\xi_n - \xi| \geq \epsilon) \\ &\leq P(|\xi_n - \xi| \geq \epsilon) \rightarrow 0, \end{aligned}$$

which implies $P(|\xi| \leq \eta) = 1$.

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Now, for any $\epsilon > 0$ and $M > 0$, we have

$$\begin{aligned} |\xi_n - \xi| &\leq \epsilon + |\xi_n - \xi|I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2\eta I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2MI\{|\xi_n - \xi| \geq \epsilon\} + 2\eta I\{\eta \geq M\} \quad a.s.. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Now, for any $\epsilon > 0$ and $M > 0$, we have

$$\begin{aligned} |\xi_n - \xi| &\leq \epsilon + |\xi_n - \xi|I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2\eta I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2MI\{|\xi_n - \xi| \geq \epsilon\} + 2\eta I\{\eta \geq M\} \quad a.s.. \end{aligned}$$

For any $\epsilon > 0$, choose $M > 0$ large enough such that

$$E\eta I\{\eta \geq M\} = \int_{y \geq M} y dF_\eta(y) < \epsilon/4.$$

Then choose N large enough such that

$$P(|\xi_n - \xi| \geq \epsilon) < \epsilon/(4M), \quad n \geq N.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in probability

Then for $n \geq N$,

$$\begin{aligned} |E\xi_n - E\xi| &\leq E|\xi_n - \xi| \\ &\leq \epsilon + 2MP(|\xi_n - \xi| \geq \epsilon) + 2E\eta I\{\eta \geq M\} < 2\epsilon. \end{aligned}$$

The applications of LLN

Example

Let $\{\xi_k, k \geq 1\}$ be a sequence of i.i.d. random variables with $E\xi_k = \mu$ and $Var\xi_k = \sigma^2$. Let

$$\bar{\xi}_n = \frac{1}{n} \sum_{k=1}^n \xi_k, \quad \hat{\sigma}_n^2 = \frac{1}{n} \sum_{k=1}^n (\xi_k - \bar{\xi}_n)^2.$$

Prove that $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$ and find the asymptotic distribution of $\sqrt{n} \frac{\bar{\xi}_n - \mu}{\hat{\sigma}_n}$.

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Proof.

$$\begin{aligned}\hat{\sigma}_n^2 &= \frac{1}{n} \sum_{k=1}^n (\xi_k - \bar{\xi}_n)^2 \\ &= \frac{1}{n} \sum_{k=1}^n ((\xi_k - \mu) - (\bar{\xi}_n - \mu))^2 \\ &= \frac{1}{n} \sum_{k=1}^n (\xi_k - \mu)^2 - (\bar{\xi}_n - \mu)^2.\end{aligned}$$

By the Khinchine weak LLN, we have $\bar{\xi}_n \xrightarrow{P} \mu$. Thus $\bar{\xi}_n - \mu \xrightarrow{P} 0$.

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Moreover, since $\{(\xi_k - \mu)^2, k \geq 1\}$ is i.i.d. and $E(\xi_k - \mu)^2 = Var \xi_k = \sigma^2$, $\{(\xi_k - \mu)^2, k \geq 1\}$ also obeys the Khinchine weak LLN, i.e. $\sum_{k=1}^n (\xi_k - \mu)^2 / n \xrightarrow{P} \sigma^2$. Hence $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$.

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Moreover, since $\{(\xi_k - \mu)^2, k \geq 1\}$ is i.i.d. and $E(\xi_k - \mu)^2 = Var \xi_k = \sigma^2$, $\{(\xi_k - \mu)^2, k \geq 1\}$ also obeys the Khinchine weak LLN, i.e. $\sum_{k=1}^n (\xi_k - \mu)^2 / n \xrightarrow{P} \sigma^2$. Hence $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$. By the Lindeberg-Lévy central limit theorem,

$$\sqrt{n} \frac{\bar{\xi}_n - \mu}{\sigma} = \frac{\sum_{k=1}^n (\xi_k - \mu)}{\sqrt{n\sigma^2}} \xrightarrow{d} N(0, 1).$$

Hence

$$\sqrt{n} \frac{\bar{\xi}_n - \mu}{\hat{\sigma}_n} = \frac{\sigma}{\hat{\sigma}_n} \cdot \sqrt{n} \frac{\bar{\xi}_n - \mu}{\sigma} \xrightarrow{d} N(0, 1).$$

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Example Prove that for any $q > p > 0$,

$$\lim_{n \rightarrow \infty} \int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = \frac{p+1}{q+1}.$$

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Example Prove that for any $q > p > 0$,

$$\lim_{n \rightarrow \infty} \int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = \frac{p+1}{q+1}.$$

Proof. Let $\{\xi_i\}$ i.i.d. $\sim U(0, 1)$, and let

$$\eta_n = \frac{\xi_1^q + \cdots + \xi_n^q}{\xi_1^p + \cdots + \xi_n^p}.$$

Then $0 \leq \eta_n \leq 1$ and

$$\int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = E\eta_n.$$

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

On the other hand, by WLLN,

$$\frac{1}{n} \sum_{k=1}^n \xi_k^q \xrightarrow{P} E\xi_1^q = \frac{1}{q+1}$$

$$\frac{1}{n} \sum_{k=1}^n \xi_k^p \xrightarrow{P} E\xi_1^p = \frac{1}{p+1}.$$

So,

$$\eta_n \xrightarrow{P} \frac{E\xi_1^q}{E\xi_1^p} = \frac{p+1}{q+1}.$$

Hence

$$\int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = E\eta_n \rightarrow \frac{p+1}{q+1}.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r Convergence in mean of order r :

Definition

Let $r > 0$, ξ and $\{\xi_n, n \geq 1\}$ be random variables defined on (Ω, \mathcal{F}, P) with $E|\xi|^r < \infty$ and $E|\xi_n|^r < \infty$. If

$$E|\xi_n - \xi|^r \longrightarrow 0,$$

then we say that $\{\xi_n, n \geq 1\}$ converges in mean of order r to ξ , denoted by $\xi_n \xrightarrow{L_r} \xi$.

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

$$\xi_n \xrightarrow{L_r} \xi \Rightarrow \xi_n \xrightarrow{P} \xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

$$\xi_n \xrightarrow{L_r} \xi \Rightarrow \xi_n \xrightarrow{P} \xi.$$

$$\xi_n \xrightarrow{L_r} \xi \not\Rightarrow \xi_n \xrightarrow{P} \xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Definition

Define ξ_n by $P(\xi_n = n) = 1/\log(n+3)$,

$P(\xi_n = 0) = 1 - 1/\log(n+3)$, $n = 1, 2, \dots$. It is easy to know

$\xi_n \xrightarrow{P} 0$, but for any $0 < r < \infty$,

$$E|\xi_n|^r = \frac{n^r}{\log(n+3)} \longrightarrow \infty.$$

That is, $\xi_n \xrightarrow{L_r} 0$ does not hold true.

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Theorem

Suppose $\xi_n \xrightarrow{P} \xi$, $P(|\xi_n| \leq \eta) = 1$, and $E\eta^r < \infty$. Then

$$\xi_n \xrightarrow{L_r} \xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Theorem

Suppose $\xi_n \xrightarrow{P} \xi$, $P(|\xi_n| \leq \eta) = 1$, and $E\eta^r < \infty$. Then

$$\xi_n \xrightarrow{L_r} \xi.$$

Proof. Since $\xi_n \xrightarrow{P} \xi$, so $P(|\xi| \leq \eta) = 1$. Then

$$P(|\xi_n - \xi|^r \leq 2^r \eta^r) = 1, \quad |\xi_n - \xi|^r \xrightarrow{P} 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Theorem

Suppose $\xi_n \xrightarrow{P} \xi$, $P(|\xi_n| \leq \eta) = 1$, and $E\eta^r < \infty$. Then

$$\xi_n \xrightarrow{L_r} \xi.$$

Proof. Since $\xi_n \xrightarrow{P} \xi$, so $P(|\xi| \leq \eta) = 1$. Then

$$P(|\xi_n - \xi|^r \leq 2^r \eta^r) = 1, \quad |\xi_n - \xi|^r \xrightarrow{P} 0.$$

By the dominated convergence theorem,

$$E[|\xi_n - \xi|^r] \rightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

附录:

Theorem

Suppose $E[|\xi_n|^r] < \infty$, $E[|\xi|^r] < \infty$. Then $\xi_n \xrightarrow{L_r} \xi$ if and only if

$$\xi_n \xrightarrow{P} \xi$$

and

$$\lim_{M \rightarrow \infty} \sup_n E[|\xi_n|^r I\{|\xi_n| \geq M\}] = 0.$$

(uniformly integrable).

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Proof. For the "if part",

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Proof. For the "if part", for any $M > 0$, let

$$\xi_{n,M} = (-M) \vee \xi_n \wedge M, \quad \xi_M = (-M) \vee \xi \wedge M.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Proof. For the "if part", for any $M > 0$, let

$$\xi_{n,M} = (-M) \vee \xi_n \wedge M, \quad \xi_M = (-M) \vee \xi \wedge M.$$

$$\xi_n \xrightarrow{P} \xi \implies \xi_{n,M} \xrightarrow{P} \xi_M.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Proof. For the "if part", for any $M > 0$, let

$$\xi_{n,M} = (-M) \vee \xi_n \wedge M, \quad \xi_M = (-M) \vee \xi \wedge M.$$

$$\xi_n \xrightarrow{P} \xi \implies \xi_{n,M} \xrightarrow{P} \xi_M.$$

By the dominated convergence theorem,

$$E[|\xi_{n,M}|^r] \xrightarrow{n \rightarrow \infty} E[|\xi_M|^r], \quad E[|\xi_{n,M} - \xi_M|^r] \xrightarrow{n \rightarrow \infty} 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Hence

$$\begin{aligned} E[|\xi|^r] &= \lim_{M \rightarrow \infty} E[|\xi_M|^r] \\ &= \lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} E[|\xi_{n,M}|^r] \leq \limsup_n E[|\xi_n|^r] < \infty. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Hence

$$\begin{aligned} E[|\xi|^r] &= \lim_{M \rightarrow \infty} E[|\xi_M|^r] \\ &= \lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} E[|\xi_{n,M}|^r] \leq \limsup_n E[|\xi_n|^r] < \infty. \end{aligned}$$

Note $|\xi - \xi_M| = 0$ if $|\xi| \leq M$, and $= |\xi| - M$ if $|\xi| > M$. So

$$\begin{aligned} |\xi_n - \xi| &\leq |\xi_{n,M} - \xi_M| + |\xi_n - \xi_{n,M}| + |\xi - \xi_M| \\ &\leq |\xi_{n,M} - \xi_M| + |\xi_n| I\{|\xi_n| \geq M\} + |\xi| I\{|\xi| \geq M\}. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

$$|\xi_n - \xi|^r \leq 3^r (|\xi_{n,M} - \xi_M|^r + |\xi_n|^r I\{|\xi_n| \geq M\} + |\xi|^r I\{|\xi| \geq M\}).$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

$$|\xi_n - \xi|^r \leq 3^r (|\xi_{n,M} - \xi_M|^r + |\xi_n|^r I\{|\xi_n| \geq M\} + |\xi|^r I\{|\xi| \geq M\}).$$

So,

$$\begin{aligned} & \limsup_{n \rightarrow \infty} E[|\xi_n - \xi|^r] \\ & \leq \lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} 3^r \left\{ E[|\xi_{n,M} - \xi_M|^r] \right. \\ & \quad \left. + E[|\xi_n|^r I\{|\xi_n| \geq M\}] + E[|\xi|^r I\{|\xi| \geq M\}] \right\} \\ & = 0. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

For the "only if part", it is obvious that $\xi_n \xrightarrow{L_r} \xi \implies \xi_n \xrightarrow{P} \xi$.

Hence

$$E[|\xi_{n,M} - \xi_M|^r] \rightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

For the "only if part", it is obvious that $\xi_n \xrightarrow{L_r} \xi \implies \xi_n \xrightarrow{P} \xi$.

Hence

$$E[|\xi_{n,M} - \xi_M|^r] \rightarrow 0.$$

It follows that

$$E[|(\xi_n - \xi_{n,M}) - (\xi - \xi_M)|^r] \rightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

For the "only if part", it is obvious that $\xi_n \xrightarrow{L_r} \xi \implies \xi_n \xrightarrow{P} \xi$.

Hence

$$E[|\xi_{n,M} - \xi_M|^r] \rightarrow 0.$$

It follows that

$$E[|(\xi_n - \xi_{n,M}) - (\xi - \xi_M)|^r] \rightarrow 0.$$

Note $|\xi - \xi_M| = 0$ if $|\xi| \leq M$, and $= |\xi| - M$ if $|\xi| > M$. So

$$\begin{aligned} & |\xi_n - \xi_{n,M}| \\ & \leq |\xi - \xi_M| + |(\xi_n - \xi_{n,M}) - (\xi - \xi_M)| \\ & \leq |\xi| I\{|\xi| \geq M\} + |(\xi_n - \xi_{n,M}) - (\xi - \xi_M)|, \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

$$\begin{aligned} |\xi_n - \xi_{n,M}|^r &\leq 2^r |\xi| I\{|\xi| \geq M\}^r \\ &\quad + 2^r |(\xi_n - \xi_{n,M}) - (\xi - \xi_M)|^r. \end{aligned}$$

It follows that

$$\begin{aligned} &\limsup_{n \rightarrow \infty} E[|\xi_n - \xi_{n,M}|^r] \\ &\leq 2^r E[|\xi|^r I\{|\xi| \geq M\}] \rightarrow 0, \quad M \rightarrow \infty. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Note again $|\xi_n - \xi_{n,M}| = 0$ if $|\xi_n| \leq M$, and $= |\xi_n| - M$ if $|\xi_n| > M$.

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Note again $|\xi_n - \xi_{n,M}| = 0$ if $|\xi_n| \leq M$, and $= |\xi_n| - M$ if $|\xi_n| > M$.

$$|\xi_n - \xi_{n,M}| \geq \frac{1}{2}|\xi_n| \quad \text{if } |\xi_n| \geq 2M.$$

That is

$$|\xi_n| I\{|\xi_n| \geq 2M\} \leq 2|\xi_n - \xi_{n,M}|.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Note again $|\xi_n - \xi_{n,M}| = 0$ if $|\xi_n| \leq M$, and $= |\xi_n| - M$ if $|\xi_n| > M$.

$$|\xi_n - \xi_{n,M}| \geq \frac{1}{2}|\xi_n| \quad \text{if } |\xi_n| \geq 2M.$$

That is

$$|\xi_n| I\{|\xi_n| \geq 2M\} \leq 2|\xi_n - \xi_{n,M}|.$$

It follows that

$$\lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} E[|\xi_n|^r I\{|\xi_n| \geq 2M\}] = 0.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Hence, for any $\epsilon > 0$, there exist $M_0 > 0$ and $N \geq 1$ such that

$$E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M_0, n \geq N.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Hence, for any $\epsilon > 0$, there exist $M_0 > 0$ and $N \geq 1$ such that

$$E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M_0, n \geq N.$$

For each $n = 1, \dots, N$, there exists an M_n such that

$$E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M_n.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Hence, for any $\epsilon > 0$, there exist $M_0 > 0$ and $N \geq 1$ such that

$$E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M_0, n \geq N.$$

For each $n = 1, \dots, N$, there exists an M_n such that

$$E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M_n.$$

Choose $M^* = \max\{M_0, M_1, \dots, M_N\}$, then

$$\sup_n E[|\xi_n|^r I\{|\xi_n| \geq M\}] \leq \epsilon \quad \text{for all } M \geq M^*.$$