

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

The expectation of a function of a discrete random variable:

ξ has the pmf

$$\begin{pmatrix} x_1 & x_2 & \cdots & x_k & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

Then for $\eta = f(\xi)$,

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

The expectation of a function of a discrete random variable:

ξ has the pmf

$$\begin{pmatrix} x_1 & x_2 & \cdots & x_k & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

Then for $\eta = f(\xi)$,

$$\begin{pmatrix} f(x_1) & f(x_2) & \cdots & f(x_k) & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\sum_i |y_i| p_i^*$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\sum_i |y_i| p_i^* = \sum_i \sum_{j:f(x_j)=y_i} |f(x_j)| p_j$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\begin{aligned} \sum_i |y_i| p_i^* &= \sum_i \sum_{j:f(x_j)=y_i} |f(x_j)| p_j \\ &= \sum_k |f(x_k)| p_k < \infty. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$E\eta = \sum_i y_i p_i^* = \sum_i \sum_{j: f(x_j)=y_i} f(x_j) p_j = \sum_k f(x_k) p_k.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$E\eta = \sum_i y_i p_i^* = \sum_i \sum_{j: f(x_j)=y_i} f(x_j) p_j = \sum_k f(x_k) p_k.$$

$$Ef(\xi) = \sum_k f(x_k) p_k = \sum_k f(x_k) P(\xi = x_k).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Theorem

Suppose that ξ is a discrete random variable with the distribution $F_\xi(x)$ and

$$P(\xi = x_k) = p_k, \quad k = 1, 2, \dots,$$

$f(x)$ a Borel function on the real line. Let $\eta = f(\xi)$. Then $Ef(\xi)$ exists if and only if

$$\int_{-\infty}^{\infty} |f(x)| dF_\xi(x) = \sum_k |f(x_k)| P(\xi = x_k) < \infty$$

and

$$Ef(\xi) = \sum_k f(x_k) P(\xi = x_k) = \int_{-\infty}^{+\infty} f(x) dF_\xi(x).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, we have

Theorem

Suppose that ξ is a random variable with the distribution $F_\xi(x)$, $f(x)$ a Borel function on the real line. Let $\eta = f(\xi)$. Then

$$Ef(\xi) = \int_{-\infty}^{+\infty} y dF_\eta(y) = \int_{-\infty}^{+\infty} f(x) dF_\xi(x).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

When ξ has density $p(x)$, then

$$Ef(\xi) = \int_{-\infty}^{+\infty} f(x)p(x)dx.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

When ξ is a random variable of the general type and $f(x)$ is a general Borel function, the proof of this theorem is due to measure theory. Next, we give a proof for the case when ξ is a continuous type random variable. Let $p(x)$ be the pdf of ξ . Then

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

When ξ is a random variable of the general type and $f(x)$ is a general Borel function, the proof of this theorem is due to measure theory. Next, we give a proof for the case when ξ is a continuous type random variable. Let $p(x)$ be the pdf of ξ . Then

$$\begin{aligned} E|\eta| &= \int_0^\infty P(|\eta| > y) dy = \int_0^\infty P(|f(\xi)| > y) dy \\ &= \int_0^\infty \int_{x: |f(x)| > y} p(x) dx dy \\ &= \int_{-\infty}^\infty \int_{y: |f(x)| > y, y \geq 0} p(x) dy dx \\ &= \int_{-\infty}^\infty |f(x)| p(x) dx. \end{aligned}$$

So $Ef(\xi)$ exists if and only if $\int_{-\infty}^\infty |f(x)| p(x) dx < \infty$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$\begin{aligned} E\eta &= \int_0^\infty P(\eta > y)dy - \int_0^\infty P(-\eta > y)dy \\ &= \int_0^\infty \int_{x:f(x)>y} p(x)dx dy - \int_0^\infty \int_{x:-f(x)>y} p(x)dx dy \\ &= \int_{-\infty}^\infty \left[\int_{y:f(x)>y, y\geq 0} dy - \int_{y:-f(x)>y, y\geq 0} dy \right] p(x)dx \\ &= \int_{-\infty}^\infty f(x)p(x)dx. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$\begin{aligned} E\eta &= \int_0^\infty P(\eta > y)dy - \int_0^\infty P(-\eta > y)dy \\ &= \int_0^\infty \int_{x:f(x)>y} p(x)dx dy - \int_0^\infty \int_{x:-f(x)>y} p(x)dx dy \\ &= \int_{-\infty}^\infty \left[\int_{y:f(x)>y, y\geq 0} dy - \int_{y:-f(x)>y, y\geq 0} dy \right] p(x)dx \\ &= \int_{-\infty}^\infty f(x)p(x)dx. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Theorem 1 tells us that if ξ and η have the same distribution function, then

$$Ef(\xi) = Ef(\eta).$$

On the contrary, if the above equality holds for any bounded continuous function f , then ξ and η have the same distribution function.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

$$\begin{aligned} F_{\xi}(z) &= \int_{-\infty}^z f(x) dF_{\xi}(x) \leq \int_{-\infty}^{\infty} f(x) dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} f(x) dF_{\eta}(x) = \int_{-\infty}^{z+\epsilon} f(x) dF_{\eta}(x) \\ &\leq \int_{-\infty}^{z+\epsilon} dF_{\eta}(x) = F_{\eta}(z + \epsilon). \end{aligned}$$

Letting $\epsilon \rightarrow 0$ yields $F_{\xi}(z) \leq F_{\eta}(z)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

$$\begin{aligned} F_{\xi}(z) &= \int_{-\infty}^z f(x) dF_{\xi}(x) \leq \int_{-\infty}^{\infty} f(x) dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} f(x) dF_{\eta}(x) = \int_{-\infty}^{z+\epsilon} f(x) dF_{\eta}(x) \\ &\leq \int_{-\infty}^{z+\epsilon} dF_{\eta}(x) = F_{\eta}(z + \epsilon). \end{aligned}$$

Letting $\epsilon \rightarrow 0$ yields $F_{\xi}(z) \leq F_{\eta}(z)$. Similarly, $F_{\eta}(z) \leq F_{\xi}(z)$. So ξ and η have the same distribution function.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example

(Stein's Lemma) (i) Let $\xi \sim N(0, 1)$, and g be differentiable function satisfying $|g(x)| \leq c_1 e^{c_2|x|}$ and $|g'(x)| \leq c_1 e^{c_2|x|}$ for some $c_1 > 0, c_2 > 0$. Prove

$$E[g(\xi)\xi] = E g'(\xi).$$

(ii)* On the contrary, if the above equality holds for any bounded continuous function $g(x)$ with bounded derivation, then $\xi \sim N(0, 1)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Proof. For (i), we have

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x)x \exp\left\{-\frac{x^2}{2}\right\} dx \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Proof. For (i), we have

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x)x \exp\left\{-\frac{x^2}{2}\right\} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x) d\left[-\exp\left\{-\frac{x^2}{2}\right\}\right]. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Use integration by parts to get

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \left[-g(x) \exp\left\{-\frac{x^2}{2}\right\} \Big|_{-\infty}^{\infty} \right. \\ & \quad \left. + \int_{-\infty}^{\infty} g'(x) \exp\left\{-\frac{x^2}{2}\right\} dx \right] \\ &= \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Use integration by parts to get

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \left[-g(x) \exp\left\{-\frac{x^2}{2}\right\} \Big|_{-\infty}^{\infty} \right. \\ &\quad \left. + \int_{-\infty}^{\infty} g'(x) \exp\left\{-\frac{x^2}{2}\right\} dx \right] \\ &= E g'(\xi). \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$. Suppose $|h(x)| \leq M$, then $0 \leq \frac{h(x)+M}{2M} \leq 1$. So without loss of generality, we assume $0 \leq h(x) \leq 1$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$. Suppose $|h(x)| \leq M$, then $0 \leq \frac{h(x)+M}{2M} \leq 1$. So without loss of generality, we assume $0 \leq h(x) \leq 1$. Let $g(x)$ satisfy

$$h(x) - Eh(\eta) = g'(x) - xg(x).$$

The above equation is called Stein's equation.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$. Suppose $|h(x)| \leq M$, then $0 \leq \frac{h(x)+M}{2M} \leq 1$. So without loss of generality, we assume $0 \leq h(x) \leq 1$. Let $g(x)$ satisfy

$$h(x) - Eh(\eta) = g'(x) - xg(x).$$

The above equation is called Stein's equation. It can be verified that

$$g(x) = e^{\frac{x^2}{2}} \int_{-\infty}^x [h(u) - Eh(\eta)] e^{-\frac{u^2}{2}} du$$

is a solution of the equation, and, both $g(x)$ and $g'(x)$ are bounded continuous function. (verify! Note $g(\infty) = 0$, $\Phi(x) \leq \frac{1}{|x|\sqrt{2\pi}} e^{-x^2/2}$, $\Phi(x) \leq e^{-x^2/2}$ for all $x < 0$.)

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So

$$\begin{aligned} Eh(\xi) &= \int_{-\infty}^{\infty} [g'(x) - xg(x) + Eh(\eta)] dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} g'(x) dF_{\xi}(x) - \int_{-\infty}^{\infty} xg(x) dF_{\xi}(x) + Eh(\eta) \int_{-\infty}^{\infty} dF_{\xi}(x) \\ &= Eg'(\xi) - E[\xi g(\xi)] + Eh(\eta) = Eh(\eta). \end{aligned}$$

The proof is completed. In the second equality above, the linearity of the Stieltjes integral.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

$$E[\xi g(\xi - 1)] = E[\lambda g(\xi)], \forall g(\text{bounded}) \Leftrightarrow \xi \sim P(\lambda).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

$$E[\xi g(\xi - 1)] = E[\lambda g(\xi)], \quad \forall g(\text{bounded}) \Leftrightarrow \xi \sim P(\lambda).$$

Stein-Chen method.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, suppose $(\xi_1, \dots, \xi_n) \sim F(x_1, \dots, x_n)$. Also, assume that $g(x_1, \dots, x_n)$ is a Borel function, then

$$Eg(\xi_1, \dots, \xi_n) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, \dots, x_n) dF(x_1, \dots, x_n).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

If $(\xi_1, \xi_2, \dots, \xi_n)$ has pmf

$P(\xi_1 = x_1(i_1), \xi_2 = x_2(i_2), \dots, \xi_n = x_n(i_n)) = p_{i_1 i_2 \dots i_n}$, then

$$Eg(\xi_1, \xi_2, \dots, \xi_n) = \sum_{i_1, i_2, \dots, i_n} g(x_1(i_1), x_2(i_2), \dots, x_n(i_n)) p_{i_1 i_2 \dots i_n};$$

If $(\xi_1, \xi_2, \dots, \xi_n)$ has pdf $p(x_1, x_2, \dots, x_n)$, then

$$\begin{aligned} & Eg(\xi_1, \xi_2, \dots, \xi_n) \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, x_2, \dots, x_n) p(x_1, x_2, \dots, x_n) dx_1 \cdots dx_n. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Here, the multi-variable Stieltjes integral is defined similarly as in the one-variable case. For example

$$\begin{aligned} & \int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} g(x_1, \dots, x_n) dF(x_1, \dots, x_n) \\ &= \lim \sum_{k_1, \dots, k_n} g(x_1(k_1), \dots, x_n(k_n)) \Delta F(x_1(k_1), \dots, x_n(k_n)), \end{aligned}$$

where $x_i(1), x_i(2), \dots$ is a partition of $(a_i, b_i]$, $\Delta F(x_1(k_1), \dots, x_n(k_n))$ is the probability that (ξ_1, \dots, ξ_n) falls in $(x_1(k_1), x_1(k_1 + 1)] \times \cdots \times (x_n(k_n), x_n(k_n + 1)]$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In particular, we have

$$E\xi_i = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} x_i dF(x_1, \cdots, x_n) = \int_{-\infty}^{\infty} x dF_i(x),$$

where $F_i(x)$ is the distribution function of ξ_i .

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For $F(x, y)$ it follows

$$E\xi\eta = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy dF(x, y)$$

and

$$E\xi^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x^2 dF(x, y),$$

etc.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example

Suppose R and Θ are indept., $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$.

Find $Ee^{R \sin \Theta}$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

$$Ee^{R \sin \Theta}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

$$Ee^{R \sin \Theta} = \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} r e^{-r^2/2} \frac{1}{2\pi} d\theta dr$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

$$\begin{aligned} Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \color{red}{r} e^{-r^2/2} \color{blue}{\frac{1}{2\pi}} d\theta dr \\ &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

$$\begin{aligned} Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \color{red}{r} \color{red}{e^{-r^2/2}} \color{blue}{\frac{1}{2\pi}} d\theta dr \\ &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \\ &= \int_{-\infty}^\infty e^x \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Solution.

$$\begin{aligned} Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \color{red}{r} \color{red}{e^{-r^2/2}} \color{blue}{\frac{1}{2\pi}} d\theta dr \\ &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \\ &= \int_{-\infty}^\infty e^x \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \\ &= e^{\frac{1}{2}} \int_{-\infty}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{(x-1)^2}{2}} dx = e^{\frac{1}{2}}. \end{aligned}$$