

## RWorksheet\_\_Sabanal#4a

Ced Bruno Sabanal

2023-11-16

1

```
shoe_Size <- c(6.5, 9.0, 8.5, 8.5, 10.5, 7.0, 9.5, 9.0, 13.0, 7.5, 10.5, 8.5, 12.0, 10.5, 13.0, 11.5, 8.5)
height <- c(66.0, 68.0, 64.5, 65.0, 70.0, 64.0, 70.0, 71.0, 72.0, 64.0, 74.5, 67.0, 71.0, 71.0, 77.0, 72.0, 59.0, 62.0)
gender <- c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F", "F")

df <- data.frame(
  ShoeSize = shoe_Size,
  Height = height,
  Gender = gender
)
```

df

##	ShoeSize	Height	Gender
## 1	6.5	66.0	F
## 2	9.0	68.0	F
## 3	8.5	64.5	F
## 4	8.5	65.0	F
## 5	10.5	70.0	M
## 6	7.0	64.0	F
## 7	9.5	70.0	F
## 8	9.0	71.0	F
## 9	13.0	72.0	M
## 10	7.5	64.0	F
## 11	10.5	74.5	M
## 12	8.5	67.0	F
## 13	12.0	71.0	M
## 14	10.5	71.0	M
## 15	13.0	77.0	M
## 16	11.5	72.0	M
## 17	8.5	59.0	F
## 18	5.0	62.0	F
## 19	10.0	72.0	M
## 20	6.5	66.0	F
## 21	7.5	64.0	F
## 22	8.5	67.0	M
## 23	10.5	73.0	M
## 24	8.5	69.0	F
## 25	10.5	72.0	M

```
## 26      11.0   70.0      M
## 27       9.0   69.0      M
## 28      13.0   70.0      M
```

1.a

1.b

```
males <- df[df$Gender == "M",]
females <- df[df$Gender == "F",]
males
```

```
##      ShoeSize Height Gender
## 5         10.5   70.0      M
## 9         13.0   72.0      M
## 11        10.5   74.5      M
## 13        12.0   71.0      M
## 14        10.5   71.0      M
## 15        13.0   77.0      M
## 16        11.5   72.0      M
## 19        10.0   72.0      M
## 22         8.5   67.0      M
## 23        10.5   73.0      M
## 25        10.5   72.0      M
## 26        11.0   70.0      M
## 27         9.0   69.0      M
## 28        13.0   70.0      M
```

```
females
```

```
##      ShoeSize Height Gender
## 1          6.5   66.0      F
## 2          9.0   68.0      F
## 3          8.5   64.5      F
## 4          8.5   65.0      F
## 6          7.0   64.0      F
## 7          9.5   70.0      F
## 8          9.0   71.0      F
## 10         7.5   64.0      F
## 12         8.5   67.0      F
## 17         8.5   59.0      F
## 18         5.0   62.0      F
## 20         6.5   66.0      F
## 21         7.5   64.0      F
## 24         8.5   69.0      F
```

1.c

```
meanOfShoeSize <- mean(df$Shoe_Size)
```

```
## Warning in mean.default(df$Shoe_Size): argument is not numeric or logical:
## returning NA
```

```
meanOfHeight <- mean(df$Height)
meanOfHeight
```

```
## [1] 68.57143
```

```
meanOfShoeSize
```

```
## [1] NA
```

## 1.d

The relationship between the two variables is that shoe size and height are positively correlated. In other words, if a person has a smaller height, they are likely to have a smaller shoe size.

```
#-----
```

## 2

```
monthsVector <- c("March", "April", "January", "November", "January", "September", "October", "September", "November")
monthsVector
```

```
## [1] "March"      "April"      "January"    "November"   "January"    "September"
## [7] "October"    "September"  "November"   "August"     "January"    "November"
## [13] "November"   "February"   "May"        "August"     "July"       "December"
## [19] "August"     "August"     "September"  "November"   "February"   "April"
```

```
factormonthsVector <- factor(monthsVector)
factormonthsVector
```

```
## [1] March      April       January     November    January     September   October
## [8] September   November    August      January     November    November    February
## [15] May         August      July        December    August      August      September
## [22] November    February    April
## 11 Levels: April August December February January July March May ... September
```

## 3

```
summary(monthsVector)
```

```
##      Length      Class      Mode
##         24 character character
```

```
summary(factormonthsVector)
```

```
##      April      August  December  February  January      July      March      May
##          2          4          1          2          3          1          1          1
## November    October  September
##          5          1          3
```

## 4

```
factorData <- c("East", "West", "North")
factorFrequency <- c(1,4,3)

neworderData <- factor(factorData,levels = c("East","West","North"))

neworderData

## [1] East West North
## Levels: East West North
```

## 5

```
imported_table <- read.csv(file = "import_march.csv" , header = TRUE, sep = ",")
imported_table

## Students Strategy.1 Strategy.2 Strategy.3
## 1 Male 8 10 8
## 2 4 8 6
## 3 0 6 4
## 4 Female 14 4 15
## 5 10 2 12
## 6 6 0 9
```

## 6

```
randomNum <- readline(prompt = "Enter number from 1 to 50: ")

## Enter number from 1 to 50:
#error cannot knit if there is as.numeric
#randomNum <- as.numeric(randomNum)

paste("The number you have chosen is", randomNum)

## [1] "The number you have chosen is "

if (randomNum > 50) {
  paste("The number selected is beyond the range of 1 to 50")
} else if (randomNum == 20) {
  paste("TRUE")
} else {
  paste(randomNum)
}

## [1] ""
```

## 7

```
minimumBills <- function(price) {

  min_bills <- price %/% 50
```

```

    paste("The minimum no. of bills:", min_bills)
  }

minimumBills(900)

## [1] "The minimum no. of bills: 18"

```

## 8.a

```

names <- c("Annie", "Thea", "Steve", "Hanna")
grade1 <- c(85,65,75,95)
grade2 <- c(65,75,55,75)
grade3 <- c(85,90,80,100)
grade4 <- c(100,90,85,90)

grade <- data.frame(
  Name = names,
  Grade1 = grade1,
  Grade2 = grade2,
  Grade3 = grade3,
  Grade4 = grade4
)

```

## 8.b

```

grade$Average <- (grade$Grade1 + grade$Grade2 + grade$Grade3 + grade$Grade4) / 4

highScorers <- grade[grade$Average > 90,]
highScorers

## [1] Name      Grade1 Grade2 Grade3 Grade4 Average
## <0 rows> (or 0-length row.names)

if (nrow(highScorers) > 0) {
  paste(highScorers$Name, "'s average grade this semester is", highScorers$Average)
} else {
  paste("No students have an average math score over 90.")
}

## [1] "No students have an average math score over 90."

```

## 8.c

```

firstTest <- sum(grade$Grade1) / nrow(grade)
firstTest

## [1] 80

secondTest <- sum(grade$Grade2) / nrow(grade)
secondTest

## [1] 67.5

```

```

thirdTest <- sum(grade$Grade3) / nrow(grade)
thirdTest

## [1] 88.75

fourthTest <- sum(grade$Grade4) / nrow(grade)
fourthTest

## [1] 91.25

if (firstTest < 80) {
  paste("The 1st test was difficult.")
} else if (secondTest < 80) {
  paste("The 2nd test was difficult.")
} else if (thirdTest < 80) {
  paste("The 3rd test was difficult.")
} else if (fourthTest < 80) {
  paste("The 4th test was difficult.")
} else {
  paste("No test had an average score less than 80.")
}

## [1] "The 2nd test was difficult."

```

## 8.d

```

# Annie scores
if (grade[1,2] > grade[1,3] && grade[1,2] > grade[1,4] && grade[1,2] > grade[1,5]) {
  annieHighest <- grade[1,2]
} else if (grade[1,3] > grade[1,4] && grade[1,3] > grade[1,5]) {
  annieHighest <- grade[1,3]
} else if (grade[1,4] > grade[1,5] && grade[1,2] > grade[1,5]) {
  annieHighest <- grade[1,4]
} else {
  annieHighest <- grade[1,5]
}

# Thea scores
if (grade[2,2] > grade[2,3] && grade[2,2] > grade[2,4] && grade[2,2] > grade[2,5]) {
  theaHighest <- grade[2,2]
} else if (grade[2,3] > grade[2,4] && grade[2,3] > grade[2,5]) {
  theaHighest <- grade[2,3]
} else if (grade[2,4] > grade[2,5] && grade[2,2] > grade[2,5]) {
  theaHighest <- grade[2,4]
} else {
  theaHighest <- grade[2,5]
}

# Steve scores
if (grade[3,2] > grade[3,3] && grade[3,2] > grade[3,4] && grade[3,2] > grade[3,5]) {
  steveHighest <- grade[3,2]
} else if (grade[3,3] > grade[3,4] && grade[3,3] > grade[3,5]) {
  steveHighest <- grade[3,3]
} else if (grade[3,4] > grade[3,5] && grade[3,2] > grade[3,5]) {
  steveHighest <- grade[3,4]
}

```

```

} else {
  steveHighest <- grade[3,5]
}

# Hanna scores
if (grade[4,2] > grade[4,3] && grade[4,2] > grade[4,4] && grade[4,2] > grade[4,5]) {
  hannaHighest <- grade[4,2]
} else if (grade[4,3] > grade[4,4] && grade[4,3] > grade[4,5]) {
  hannaHighest <- grade[2,3]
} else if (grade[4,4] > grade[4,5] && grade[4,2] > grade[4,5]) {
  hannaHighest <- grade[4,4]
} else {
  hannaHighest <- grade[4,5]
}

grade$HighestGrades <- c(annieHighest, theaHighest, steveHighest, hannaHighest)

above90 <- grade[grade$HighestGrades > 90,]
above90

##      Name Grade1 Grade2 Grade3 Grade4 Average HighestGrades
## 1 Annie      85      65      85      100  83.75          100
## 4 Hanna      95      75     100      90  90.00          100

if (nrow(above90) > 0) {
  paste(above90$Name, "'s highest grade this semester is", above90$HighestGrade)
} else {
  paste("No students have an average math score over 90.")
}

## [1] "Annie 's highest grade this semester is 100"
## [2] "Hanna 's highest grade this semester is 100"

```