

Testing whether the observed I<sub>ss</sub> is significantly lower than I<sub>ss.c</sub>.

Part I. For a symmetrical tree.

=====	
Prop. invar. sites	0.0000
Mean H	0.1489
Standard Error	0.0088
Hmax	1.8140
I <sub>ss</sub>	0.0821
I <sub>ss.c</sub>	0.7655
T	77.3685
DF	1034
Prob (Two-tailed)	0.0000
95% Lower Limit	0.0647
95% Upper Limit	0.0994

Part II. For an extreme asymmetrical (and generally very unlikely) tree.

=====	
I <sub>ss.c</sub>	0.5790
T	56.2634
DF	1034
Prob (Two-tailed)	0.0000
95% Lower Limit	0.0647
95% Upper Limit	0.0994

Interpretation of results:

Significant Difference		
-----		
	Yes	No
-----		
I <sub>ss</sub> < I <sub>ss.c</sub>	Little saturation	Substantial saturation
-----		
I <sub>ss</sub> > I <sub>ss.c</sub>	Useless sequences	Very poor for phylogenetics
-----		

Please cite:

Xia, X., Z. Xie, M. Salemi, L. Chen, Y. Wang. 2003. An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26:1-7.

Xia, X. and Lemey, P. 2009. Assessing substitution saturation with DAMBE. Pp. 615-630 in Philippe Lemey, Marco Salemi and Anne-Mieke Vandamme, eds. *The Phylogenetic Handbook: A Practical Approach to DNA and Protein Phylogeny*. 2nd edition Cambridge University Press.