

Testing whether the observed I<sub>ss</sub> is significantly lower than I<sub>ss.c</sub>.

Part I. For a symmetrical tree.

=====		
Prop. invar. sites	0.0000	
Mean H	1.0408	
Standard Error	0.0287	
Hmax	1.8578	
I <sub>ss</sub>	0.5602	
I <sub>ss.c</sub>	0.7951	
T	8.1797	
DF	2072	
Prob (Two-tailed)	0.0000	
95% Lower Limit	0.5039	
95% Upper Limit	0.6165	

Part II. For an extreme asymmetrical (and generally very unlikely) tree.

=====		
I <sub>ss.c</sub>	0.5975	
T	1.2999	
DF	2072	
Prob (Two-tailed)	0.1938	
95% Lower Limit	0.5039	
95% Upper Limit	0.6165	

Interpretation of results:

Significant Difference		
-----		
	Yes	No
-----		
I <sub>ss</sub> < I <sub>ss.c</sub>	Little saturation	Substantial saturation
-----		
I <sub>ss</sub> > I <sub>ss.c</sub>	Useless sequences	Very poor for phylogenetics
-----		

Please cite:

Xia, X., Z. Xie, M. Salemi, L. Chen, Y. Wang. 2003. An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26:1-7.

Xia, X. and Lemey, P. 2009. Assessing substitution saturation with DAMBE. Pp. 615-630 in Philippe Lemey, Marco Salemi and Anne-Mieke Vandamme, eds. *The Phylogenetic Handbook: A Practical Approach to DNA and Protein Phylogeny*. 2nd edition Cambridge University Press.