

Testing whether the observed Iss is significantly lower than Iss.c.

Part I. For a symmetrical tree.

Prop. invar. sites	0.0000
Mean H	0.9628
Standard Error	0.0273
Hmax	1.8525
Iss	0.5197
Iss.c	0.8039
T	10.3903
DF	2381
Prob (Two-tailed)	0.0000
95% Lower Limit	0.4661
95% Upper Limit	0.5734

Part II. For an extreme asymmetrical (and generally very unlikely) tree.

Iss.c	0.6296
T	4.0167
DF	2381
Prob (Two-tailed)	0.0001
95% Lower Limit	0.4661
95% Upper Limit	0.5734

Interpretation of results:

Significant Difference

Yes No

Iss < Iss.c	Little saturation	Substantial saturation
Iss > Iss.c	Useless sequences	Very poor for phylogenetics

Please cite:

Xia, X., Z. Xie, M. Salemi, L. Chen, Y. Wang. 2003. An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26:1-7.

Xia, X. and Lemey, P. 2009. Assessing substitution saturation with DAMBE. Pp. 615-630 in Philippe Lemey, Marco Salemi and Anne-Mieke Vandamme, eds. *The Phylogenetic Handbook: A Practical Approach to DNA and Protein Phylogeny*. 2nd edition Cambridge University Press.