

Please complete the following exercises. Feel free to work with classmates, but each student must turn in **UNIQUE** work, not photocopies or identical replicates. When applicable, use **APA format** in communicating your results in text. **Show your work!** If any question involves any math at all, show your work. When in doubt, write it out. Always show more than you think you need.

### 1) WRITE-UP - Textbook Problems

Cohen Chap	Exercises	Pts	Off
2	A	1	
	C	2, 3, 4 (no output, tables, or plots need to be included)	2
3	A	*1, 2, 3, 4, *5	5
4	A	*1, 2, 4, 9a, 10a, 11, *12, 13, *14	11
	B	3, 4, 5, 6, 7	5

### 2) SUMMARY – Your Journal Article

Do NOT re-submit your HW 0 article!		Pts	Off
Half Page	Re-read your article you selected for Unit 0 HW (do not re-submit it). Summarize any mention or use of the concepts in the above chapters.	10	

### 3) R SYNTAX – Section C: Ihno's data set – add to the skeleton R notebook and knit to .pdf & upload

Cohen Chap	Exercises	Pts	Off
2	C	1, 2, 3, 4, 6, 9, 10	8
3	C	1, 2, 3, 4, 5, 6	6
4	C	1,	2

Grading		Earned	Possible
<b>CORRECTNESS</b>	a subset of spot-checked items: must show work, especially items from back of book or done in class		50
<b>COMPLETENESS</b>	more than one item is missing or skipped: 25/50 roughly half the assignment is completed: 10/50		50
		<div style="border: 2px solid black; width: 100px; height: 20px;"></div>	100

## 2 A 1. Histograms


A psychotherapist has rated all 20 of her patients in terms of their progress in therapy, using a 7-point scale. The results are shown in the following table: (please use TWO decimal places)

	F	rf	cf	crf	cpf
Greatly improved	5				
Moderately improved	4				
Slightly improved	6				
Unchanged	2				
Slightly worse	2				
Moderately worse	1				
Greatly worse	0				

To answer the questions below, fill out the columns in the table above, such that:

- rf = relative frequency
- cf = cumulative frequency
- crf = cumulative relative frequency
- cpf = cumulative percentage frequency

a) Draw a bar graph to represent the above results.



b) What **proportion** of the patients was greatly improved?

c) How **many** patients did not improve?

d) What is the **percentile rank** of a patient who improved slightly?

What is **the percentile rank** of a patient who becomes slightly worse?

e) Which **category** of improvement corresponds to the third quartile?

Which **category** of improvement corresponds to the first quartile?

2	C	2. Distributions & Bar Plots
---	---	------------------------------

Request a frequency distribution and a bar chart for the *prevmath* and *phobia* variables.

**write code in R syntax file**

*(no output, tables, or plots need to be included)*

Would it **make sense** to request a histogram instead of a bar chart for *phobia*?

**Discuss.**

☐ yes ☐ no

2	C	3. Distributions & Bar Plots
---	---	------------------------------

Request a frequency distribution and a histogram for the *statquiz* variables.

**write code in R syntax file**

*(no output, tables, or plots need to be included)*

**Describe the shape of this distribution.**

<b>3</b>	<b>A</b>	<b>*1. Measures of central tendency</b>
----------	----------	---

Select the **measure of central tendency** that would be most appropriate for describing each of the following hypothetical sets of data:

a. Religious preferences of delegates to the United Nations	<input type="checkbox"/> Mean <input type="checkbox"/> Median <input type="checkbox"/> Mode
b. Heart rates for a group of women before they start their first aerobics class	<input type="checkbox"/> Mean <input type="checkbox"/> Median <input type="checkbox"/> Mode
c. Types of phobias exhibited by patients attending a phobia clinic	<input type="checkbox"/> Mean <input type="checkbox"/> Median <input type="checkbox"/> Mode
d. Amounts of time participants spend solving a classic cognitive problem, with some of the participants unable to solve it	<input type="checkbox"/> Mean <input type="checkbox"/> Median <input type="checkbox"/> Mode
e. Height in inches for a group of boys in the first grade	<input type="checkbox"/> Mean <input type="checkbox"/> Median <input type="checkbox"/> Mode

<b>3</b>	<b>A</b>	<b>2. Distribution descriptives</b>
----------	----------	-------------------------------------

Describe a **realistic situation** in which you would expect to obtain each of the following:

a)	A negatively skewed distribution
b)	A positively skewed distribution
c)	A bimodal distribution

<b>3</b>	<b>A</b>	<b>*3. Distribution descriptives</b>
----------	----------	--------------------------------------

A midterm exam was given in a large introductory psychology class. The **median** score was 85, the **mean** was 81, and the **mode** was 87.

What kind of distribution would you expect from these exam scores?

### 3 A 4. Measures of central tendency

A veterinarian is interested in the life span of golden retrievers. She recorded the age at death (in years) of the retrievers treated in her clinic. The ages were **12, 9, 11, 10, 8, 14, 12, 1, 9, 12.**

a) Calculate the mean, median, and mode of life span:

Mean

Median

Mode




b) After examining her records, the veterinarian determined that the dog that had died at 1 year was killed by a car. **Recalculate** the mean, median, and mode without that dog's data.

Mean

Median

Mode




c) Which measure of central tendency in part b **changed the most**, compared to the values originally calculated in part a?

☐ Mean ☐ Median ☐ Mode

Explain.

<b>3</b>	<b>A</b>	<b>5. Measures of variability</b>
----------	----------	-----------------------------------

Which of the three most popular **measures of variability** would you choose in each of the following situations?

a. The distribution is badly skewed with a few extreme outliers in one direction.	<input type="checkbox"/> Range <input type="checkbox"/> SIQ range <input type="checkbox"/> SD
b. You are planning to perform advanced statistical procedures (e.g., draw inferences about population parameters).	<input type="checkbox"/> Range <input type="checkbox"/> SIQ range <input type="checkbox"/> SD
c. You need to know the maximum width taken up by the distribution.	<input type="checkbox"/> Range <input type="checkbox"/> SIQ range <input type="checkbox"/> SD
d. You need a statistic that takes into account every score in the population.	<input type="checkbox"/> Range <input type="checkbox"/> SIQ range <input type="checkbox"/> SD
e. The highest score in the distribution is “more than 10.”	<input type="checkbox"/> Range <input type="checkbox"/> SIQ range <input type="checkbox"/> SD

<b>4</b>	<b>A</b>	<b>*1. Z-scores</b>
----------	----------	---------------------

If you **convert each score in a set of scores to a z score**, which of the following will be **true** about the resulting set of z scores?

a. The mean will equal 1.	<input type="checkbox"/> TRUE
b. The variance will equal 1.	<input type="checkbox"/> TRUE
c. The distribution will be normal in shape.	<input type="checkbox"/> TRUE
d. All of the above.	<input type="checkbox"/> TRUE
e. None of the above.	<input type="checkbox"/> TRUE

<b>4</b>	<b>A</b>	<b>2. Z-scores</b>
----------	----------	--------------------

The distribution of body weights for adults is somewhat **positively skewed**— there is much more room for people to be above average than below.

If you take **the mean weights for random groups of 10 adults** each and form a new distribution, how will this new distribution **compare** to the distribution of individuals?

a. The new distribution will be <b>more symmetrical</b> than the distribution of individuals.	<input type="checkbox"/> TRUE
b. The new distribution will <b>more closely resemble the normal</b> distribution.	<input type="checkbox"/> TRUE
c. The new distribution will be <b>narrower</b> (i.e., have a smaller standard deviation) than the distribution of individuals.	<input type="checkbox"/> TRUE
d. All of the above.	<input type="checkbox"/> TRUE
e. None of the above.	<input type="checkbox"/> TRUE

- a) Calculate  $\mu$  and  $\sigma$  for the following set of scores and then convert each score to a z score:  
64, 45, 58, 51, 53, 60, 52, 49.

Mean:  $\mu$

Standard Deviation:  $\sigma$

$\mu =$

$\sigma =$

z-scores:

--	--	--	--	--	--	--	--

- b) Calculate the mean and standard deviation of these z scores.

Mean:  $\mu$

Standard Deviation:  $\sigma$

$\mu =$

$\sigma =$

Did you obtain the values you **expected**?  
**Explain.**

☐ yes ☐ no





4 A 13. Standard error for the mean

- a) In one college, freshman English classes always contain exactly **20 students**. An English teacher wonders how much these classes are **likely to vary** in terms of their verbal scores on the SAT. What would you expect for the standard deviation (i.e., **standard error**) of class means on the verbal SAT?

$SE_{\mu} =$

- b) Suppose that a crew for the space shuttle consists of **seven people**, and we are interested in the average weights of all possible shuttle crews. If the **standard deviation for weight is 30 pounds**, what is the **standard deviation for the mean weights** of shuttle crews (i.e., the standard error of the mean)?

$SE_{\mu} =$

4 A \*14. Standard error for the mean

If for a particular sampling distribution of the mean we know that the **standard error is 4.6**, and we also know that  $\sigma = 32.2$ , what is the sample size (  $n$  )?

$n =$

4 B \*3. Area under a normal curve

Assume that the resting heart rate in humans is normally distributed with  $\mu = 72$  bpm (i.e., beats per minute) and  $\sigma = 8$  bpm.

a) What proportion of the population has resting heart rates **above** 82 bpm?





**Above 70 bpm?**




b) What proportion of the population has resting heart rates **below** 75 bpm?




**Below 50 bpm?**





c) What proportion of the population has resting heart rates **between** 80 and 85 bpm?




**Between 60 and 70 bpm?**





**Between 55 and 75 bpm?**




**4 B 4. Area under a normal curve**

Assume that the resting heart rate in humans is normally distributed with  $\mu = 72$  bpm (i.e., beats per minute) and  $\sigma = 8$  bpm.


- a) **Above** what heart rate do you find the upper 25% of the people? (That is, what heart rate is at the 75th percentile, or third quartile?)




- b) **Below** what heart rate do you find the lowest 15% of the people? (That is, what heart rate is at the 15th percentile?)





- c) **Between** which two heart rates do you find the middle 75% of the people?




**4 B \*5. Area under a normal curve**

A new preparation course for the math SAT is open to those who have already taken the test once and scored in the **middle 90%** of the population.

In what **range** must a testtaker's previous score have fallen for the test-taker to be eligible for the new course?




4 B 6. Area under a normal curve

A teacher thinks her class has an unusually high IQ, because her 36 students have an average IQ ( $\bar{X}$ ) of 108. If the population mean is 100 and  $\sigma = 15$ .

a) What is the **z score** for this class?

b) What **percentage** of classes ( $n = 36$ , randomly selected) would be even higher on IQ?




4 B \*7. Area under a normal curve

An aerobics instructor thinks that his class has an unusually low resting heart rate. If  $\mu = 72$  bpm and  $\sigma = 8$  bpm, and his class of 14 pupils has a mean heart rate ( $\bar{X}$ ) of 66,

a) What is the **z score** for the aerobics class?

b) What is the **probability** of randomly selecting a group of 14 people with a mean resting heart rate **lower** than the mean for the aerobics class?



Re-read your article you selected for Unit 0 HW (do not re-submit it).  
Summarize any mention or use of the concepts in the above chapters.

*(you may choose to **type** this summary and include a printed copy here instead of hand writing  
OR upload a typed document to CANVAS)*