

Monte Carlo simulation

2026-08-28

Summary

Estimating an expectation by simulating years, and how far off the estimate is.

- An expected annual damage is an integral that usually has no closed form, so we average simulated years instead.
- The law of large numbers says the average converges, given independent draws, identical distribution, and a finite mean, and says nothing about how fast.
- The estimator is unbiased: on average it lands on the right answer for any n .
- Its variance falls like $1/n$, so the error falls like $1/\sqrt{n}$ and halving the error costs four times the draws.
- Independence is what licenses splitting the variance of the sum, so draws that covary make the estimator noisier than n independent ones.
- The relative error depends on the coefficient of variation of the quantity being averaged, which is a property of that quantity and not of how the simulation is run.

Key results.

$$\theta = E[g(X)], \quad \hat{\theta}_n = \frac{1}{n} \sum_{i=1}^n g(X_i), \quad E[\hat{\theta}_n] = \theta$$

$$SE(\hat{\theta}_n) = \frac{sd(g(X))}{\sqrt{n}}, \quad \frac{SE(\hat{\theta}_n)}{\theta} = \frac{CV}{\sqrt{n}}$$

Lab 1 breaks the finite-mean assumption and shows what happens, and has you find the coefficient of variation for several distributions to see which converge.

Bibliography