

Mathematics Courses
Framework Series

Real Analysis

Fourth Edition,
by *H.L. Royden* and *P.M. Fitzpatrick*

aytony

2024 -- 04 -- 17



Contents

Preliminaries on Sets, Mappings, and Relations	3
1 The Real Numbers: Sets, Sequences, and Functions	4
1.1 The Field, Positivity, and Completeness Axioms	4
1.2 The Natural and Rational Numbers	4
1.3 Countable and Uncountable Sets	4
1.4 Open Sets, Closed Sets, and Borel Sets of Real Numbers	4
1.5 Sequences of Real Numbers	5
1.6 Continuous Real-valued Functions of a Real Variable	6
2 Lebesgue Measure	8
2.1 Intruduction	8
2.2 Lebesgue Outer Measure	8
2.3 The σ -Algebra of Lebesgue Measurable Sets	8
2.4 Outer and Inner Approximation of Lebesgue Measurable Sets	9
2.5 Countable Additivity, Continuity, and the Borel-Cantelli Lemma	9
2.6 Nonmeasurable Sets	10
2.7 The Cantor Set and the Cantor-Lebesgue Function	10
3 Lebesgue Measurable Functions	11
3.1 Sums, Products, and Compositions	11
3.2 Sequential Pointwise Limits and Simple Approximation	12
3.3 Littlewood's Three Principles, Egoroffs's Theorem, and Lusin's Theorem	12
4 Lebesgue Integration	14
4.1 The Rimemann Integral	14
4.2 The Lebesgue Integral of a Bounded Measurable Function over a Set of Finite Measure	14
4.3 The Lebesgue Integral of a Measurable Nonnegatice Function	16
4.4 The General Lebesgue Integral	17
4.5 Contable Addativity and Continuity of Integration	19
4.6 Uniform Integrability: The Vitali Convergence Theorem	19

Preliminaries on Sets, Mappings, and Relations

Definition: A relation R on a set X is called an **equivalence relation** provided it is reflexive, symmetric, and transitive.

Definition: Let \mathcal{F} be a nonempty family of nonempty sets. A **choice function** on \mathcal{F} is a function f from \mathcal{F} to $\cup_{F \in \mathcal{F}} F$ with the property that for each set F in \mathcal{F} , $f(F)$ is a member of F .

Zermelo's Axiom of Choice: Let \mathcal{F} be a nonempty collection of nonempty sets. Then there is a choice function on \mathcal{F} .

Definition: A relation R on a set nonempty X is called a **partial ordering** provided it is reflexive, transitive and for x, x' in X ,

$$\text{if } xRx' \text{ and } x'R x, \text{ then } x = x'.$$

A subset E of X is said to be **totally ordered** provided for x, x' in E , either xRx' or $x'R x$. A member x of X is said to be an **upper bound** for a subset E of X provided $x'R x$ for all $x' \in E$, and is said to be **maximal** provided the only member x' of X for which xRx' is $x' = x$.

Zorn's Lemma: Let X be a partially ordered set for which every totally ordered subset has an upper bound. Then X has a maximal member.

1 | The Real Numbers: Sets, Sequences, and Functions

1.1 | The Field, Positivity, and Completeness Axioms

The Completeness Axiom: Let E be a nonempty set of real numbers that is bounded above. Then among the set of upper bounds for E there is a smallest, or least, upper bound.

1.2 | The Natural and Rational Numbers

Definition: A set E of real numbers is said to be **inductive** provided it contains 1 and if the number x belongs to E , the number $x + 1$ also belongs to E .

Principle of Mathematical Induction: For each natural number n , let $S(n)$ be some mathematical assertion. Suppose $S(1)$ is true. Also suppose that whenever k is a natural number for which $S(k)$ is true, then $S(k + 1)$ is also true. Then $S(n)$ is true for every natural number n .

Theorem 1.1: Every nonempty set of natural numbers has a smallest member.

Archimedean Property: For each pair of positive real numbers a and b , there is a natural number n for which $na > b$.

Definition: A set E of real numbers is said to be **dense** in \mathbb{R} provided between any two real numbers there lies a member of E .

Theorem 1.2: The rational numbers are dense in \mathbb{R} .

1.3 | Countable and Uncountable Sets

Definition: A set E is said to be **finite** provided either it is empty or there is a natural number n for which E is equipotent to $\{1, \dots, n\}$. We say that E is **countably infinite** provided E is equipotent to the set \mathbb{N} of natural numbers. A set that is either finite or countably infinite is said to be **countable**. A set that is not countable is called **uncountable**.

Theorem 1.3: A subset of a countable set is countable. In particular, every set of natural numbers is countable.

Corollary 1.4: The following sets are countably infinite:

1. For each natural number n , the Cartesian product $\overbrace{\mathbb{N} \times \dots \times \mathbb{N}}^{n \text{ times}}$.
2. The set of rational numbers \mathbb{Q} .

Theorem 1.5: A nonempty set is countable if and only if it is the image of a function whose domain is a nonempty countable set.

Corollary 1.6: The union of a countable collection of countable sets is countable.

Theorem 1.7: A nondegenerate interval of real numbers is uncountable.

1.4 | Open Sets, Closed Sets, and Borel Sets of Real Numbers

Definition: A set \mathcal{O} of real numbers is called **open** provided for each $x \in \mathcal{O}$, there is a $r > 0$ for which the interval $(x - r, x + r)$ is contained in \mathcal{O} .

Proposition 1.8: The set of real numbers \mathbb{R} and the empty-set \emptyset is open; the intersection of any finite collection of open sets is open and the union of any collection of open sets is open.

Proposition 1.9: Every nonempty open set is the disjoint union of a countable collection of open intervals.

Definition: For a set E of real numbers, a real number x is called a **point of closure** of E provided every open interval that contains x also contains a point in E . The collection of points of closure of E is called the **closure** of E and denoted by \overline{E} .

Proposition 1.10: For a set of real numbers E , its closure \overline{E} is closed. Moreover, \overline{E} is the smallest closed set that contains E in the sense that if F is closed and $E \subseteq F$, then $\overline{E} \subseteq F$.

Proposition 1.11: A set of real numbers is open if and only if its complement in \mathbb{R} is closed.

Proposition 1.12: The empty-set \emptyset are closed; the union of any finite collection of closed sets is closed; and the intersection of any collection of closed sets is closed.

The Heine-Borel Theorem: Let F be a closed and bounded set of real numbers. Then every open cover of F has a finite subcover.

The Nested Set Theorem: Let $\{F_n\}_{n=1}^{\infty}$ be a descending countable collection of nonempty closed sets of real numbers for which F_1 bounded. Then

$$\bigcap_{n=1}^{\infty} F_n \neq \emptyset.$$

Definition: Given a set X , a collection \mathcal{A} of subsets of X is called a σ -algebra (of subsets of X) provided:

1. The empty-set, \emptyset , belongs to \mathcal{A} ;
2. The complement in X of a set in \mathcal{A} also belongs to \mathcal{A} ;
3. The union of a countable collection of sets in \mathcal{A} also belongs to \mathcal{A} .

Proposition 1.13: Let \mathcal{F} be a collection of subsets of a set X . Then the intersection \mathcal{A} of all σ -algebras of subsets of X that contain \mathcal{F} is a σ -algebra that contains \mathcal{F} . Moreover, it is the smallest σ -algebra of subsets of X that contains \mathcal{F} in the sense that any σ -algebra that contains \mathcal{F} also contains \mathcal{A} .

Definition: The collection \mathcal{B} of Borel sets of real numbers is the smallest σ -algebra of sets of real numbers that contains all of the open sets of real numbers.

1.5 | Sequences of Real Numbers

Definition: A sequence $\{a_n\}$ is said to **converge** to the number a provided for every $\varepsilon > 0$, there is an index N for which

$$\text{if } n \geq N, \text{ then } |a - a_n| < \varepsilon.$$

We call a the **limit** of the sequence and denote the convergence of $\{a_n\}$ by writing

$$\{a_n\} \rightarrow a \text{ or } \lim_{n \rightarrow \infty} a_n = a.$$

Proposition 1.14: Let the sequence of real numbers $\{a_n\}$ converge to the real number a . Then the limit is unique, the sequence is bounded, and, for a real number c ,

$$\text{if } a_n \leq c \text{ for all } n, \text{ then } a \leq c.$$

Theorem 1.15 (the Monotone Convergence Criterion for Real Sequences): A monotone sequence of real numbers converges if and only if it is bounded.

Theorem 1.16 (the Volzano-Weierstrass Theorem): Every bounded sequence of real numbers has a convergent subsequence.

Definition: A sequence of real numbers $\{a_n\}$ is said to be **Cauchy** provided for each $\varepsilon > 0$, there is an index N for which

$$\text{if } n, m \geq N, \text{ then } |a_m - a_n| < \varepsilon.$$

Theorem 1.17 (the Cauchy Convergence Criterion for Real Sequences): A sequence of real numbers converges if and only if it is Cauchy.

Theorem 1.18 (Linearity and Monotonicity of Convergence of Real Sequences): Let $\{a_n\}$ and $\{b_n\}$ be convergent sequences of real numbers. Then for each pair of real numbers α and β , the sequence $\{\alpha \cdot a_n + \beta \cdot b_n\}$ is convergent and

$$\lim_{n \rightarrow \infty} [\alpha \cdot a_n + \beta \cdot b_n] = \alpha \cdot \lim_{n \rightarrow \infty} a_n + \beta \cdot \lim_{n \rightarrow \infty} b_n.$$

Moreover,

$$\text{if } a_n \leq b_n \text{ for all } n, \text{ then } \lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} b_n.$$

Definition: Let $\{a_n\}$ be sequence of real numbers. The limit superior of $\{a_n\}$, denoted by $\limsup\{a_n\}$, is defined by

$$\limsup\{a_n\} = \lim_{n \rightarrow \infty} [\sup\{a_k \mid k \geq n\}].$$

The limit inferior of $\{a_n\}$, denoted by $\liminf\{a_n\}$, is defined by

$$\liminf\{a_n\} = \lim_{n \rightarrow \infty} [\inf\{a_k \mid k \geq n\}].$$

Proposition 1.19: Let $\{a_n\}$ and $\{b_n\}$ be sequences of real numbers.

1. $\limsup\{a_n\} = \ell \in \mathbb{R}$ if and only if for each $\varepsilon > 0$, there are infinitely many indices n for which $a_n > \ell - \varepsilon$ and only finitely many indices n for which $a_n > \ell + \varepsilon$.
2. $\limsup\{a_n\} = \infty$ if and only if $\{a_n\}$ is not bounded above.
3. $\limsup\{a_n\} = -\liminf\{-a_n\}$.
4. A sequence of real numbers $\{a_n\}$ converges to an extended real number a if and only if

$$\liminf\{a_n\} = \limsup\{a_n\} = a.$$

5. If $a_n \leq b_n$ for all n , then

$$\limsup\{a_n\} \leq \limsup\{b_n\}.$$

Proposition 1.20: Let $\{a_n\}$ be a sequence of real numbers.

1. The series $\sum_{k=1}^{\infty} a_k$ is summable if and only if for each $\varepsilon > 0$, there is an index N for which

$$\left| \sum_{k=n}^{n+m} a_k \right| < \varepsilon \text{ for } n \geq N \text{ and any natural number } m.$$

2. If the series $\sum_{k=1}^{\infty} |a_k|$ is summable, then $\sum_{k=1}^{\infty} a_k$ also is summable.
3. If each term a_k is nonnegative, then the series $\sum_{k=1}^{\infty} a_k$ is summable if and only if the sequence of partial sums is bounded.

1.6 | Continuous Real-valued Functions of a Real Variable

Proposition 1.21: A real-valued function f defined on a set E of real numbers is continuous at the point $x_* \in E$ if and only if whenever a sequence $\{x_n\}$ in E converges to x_* , its image sequence $\{f(x_n)\}$ converges to $f(x_*)$.

Proposition 1.22: Let f be a real-valued function defined on a set E of real numbers. Then f is continuous on E if and only if for each open set \mathcal{O} ,

$$f^{-1}(\mathcal{O}) = E \cap \mathcal{U} \text{ where } \mathcal{U} \text{ is an open set.}$$

The Extreme Value Theorem: A continuous real-valued function on a nonempty closed, bounded set of real numbers takes a minimum and maximum value.

The Intermediate Value Theorem: Let f be a continuous real-valued function on the closed, bounded interval $[a, b]$ for which $f(a) < c < f(b)$. Then there is a point x_0 in (a, b) at which $f(x_0) = c$.

Definition: A real-valued function f defined on a set E of real numbers is said to be **uniformly continuous** provided for each $\varepsilon > 0$, there is a $\delta > 0$ such that for all $x, x' \in E$,

$$\text{if } |x - x'| < \delta, \text{ then } |f(x) - f(x')| < \varepsilon.$$

Theorem 1.23: A continuous real-valued function on a closed, bounded set of real numbers is uniformly continuous.

Definition: A real-valued function f defined on a set E of real numbers is said to be **increasing** provided $f(x) \leq f(x')$ whenever x, x' belong to E and $x \leq x'$, and **decreasing** provided $-f$ is increasing. It is called **monotone** if it is either increasing or decreasing.

2 | Lebesgue Measure

2.1 | Intruduction

2.2 | Lebesgue Outer Measure

Example: A countable set has outer measure zero. Indeed, let C be a countable set enumerated as $C = \{c_k\}_{k=1}^{\infty}$. Let $\varepsilon > 0$. For each natural number k , define $I_k = (c_k - \varepsilon/2^{k+1}, c_k + \varepsilon/2^{k+1})$. The countable collection of open intervals $\{I_k\}_{k=1}^{\infty}$ covers C . Therefore

$$0 \leq m^*(C) \leq \sum_{k=1}^{\infty} \ell(I_k) = \sum_{k=1}^{\infty} \varepsilon/2^k = \varepsilon.$$

This inequality holds for each $\varepsilon > 0$. Hence $m^*(E) = 0$.

Proposition 2.1: The outer measure of an interval is its length.

Proposition 2.2: Outer measure is translation invariant, that is, for any set A and number y ,

$$m^*(A + y) = m^*(A).$$

Proposition 2.3: Outer measure is countably subadditive, that is, if $\{E_k\}_{k=1}^{\infty}$ is any countable collection of sets, disjoint or not, then

$$m^*\left(\bigcup_{k=1}^{\infty} E_k\right) \leq \sum_{k=1}^{\infty} m^*(E_k).$$

2.3 | The σ -Algebra of Lebesgue Measurable Sets

Definition: A set E is said to be **measurable** provided for any set A ,

$$m^*(A) = m^*(A \cap E) + m^*(A \cap E^C).$$

Proposition 2.4: Any set of outer measure zero is measurable. In particular, any countable set is measurable.

Proposition 2.5: The union of a finite collection of measurable sets is measurable.

Proposition 2.6: Let A be any set and $\{E_k\}_{k=1}^n$ a finite disjoint collection of measurable sets. Then

$$m^*\left(A \cap \left[\bigcup_{k=1}^n E_k\right]\right) = \sum_{k=1}^n m^*(A \cap E_k).$$

In particular,

$$m^*\left(\bigcup_{k=1}^n E_k\right) = \sum_{k=1}^n m^*(E_k).$$

Proposition 2.7: The union of a countable collection of measurable sets is measurable.

Proposition 2.8: Every interval is measurable.

Theorem 2.9: The collection \mathcal{M} of measurable sets is a σ -algebra that contains the σ -algebra \mathcal{B} of Borel sets. Each interval, each open set, each closed set, each G_δ set, and each F_δ set is measurable.

Proposition 2.10: The translate of a measurable set is measurable.

2.4 | Outer and Inner Approximation of Lebesgue Measurable Sets

Theorem 2.11: Let E be any set of real numbers. Then each of the following four assertions is equivalent to the measurability of E .

(Outer Approximation by Open Sets and G_δ Sets)

1. For each $\varepsilon > 0$, there is an open set \mathcal{O} containing E for which $m^*(\mathcal{O} \setminus E) < \varepsilon$.
2. There is a G_δ set G containing E for which $m^*(G \setminus E) = 0$.

(Inner Approximation by Closed Sets and F_δ Sets)

1. For each $\varepsilon > 0$, there is a closed set F contained in E for which $m^*(E \setminus F) < \varepsilon$.
2. There is an F_δ set F contained in E for which $m^*(E \setminus F) = 0$.

Theorem 2.12: Let E be a measurable set of finite outer measure. Then for each $\varepsilon > 0$, there is a finite disjoint collection of open intervals $\{I_k\}_{k=1}^n$ for which if $\mathcal{O} = \bigcup_{k=1}^n I_k$, then

$$m^*(E \setminus \mathcal{O}) + m^*(\mathcal{O} \setminus E) < \varepsilon.$$

Remark: A comment regarding assertion in Theorem 2.12 is in order. By the definition of outer measure, for any bounded set E , regardless of whether or not it is measurable, and any $\varepsilon > 0$, there is an open set \mathcal{O} such that $E \subseteq \mathcal{O}$ and $m^*(\mathcal{O}) < m^*(E) + \varepsilon$ and therefore $m^*(\mathcal{O}) - m^*(E) < \varepsilon$. This does not imply that $m^*(\mathcal{O} \setminus E) < \varepsilon$, because the excision property

$$m^*(\mathcal{O} \setminus E) = m^*(\mathcal{O}) - m^*(E)$$

is false unless E is measurable.

2.5 | Countable Additivity, Continuity, and the Borel-Cantelli Lemma

Definition: The restriction of the set function outer measure to the class of measurable sets is called **Lebesgue measure**. It is denoted by m , so that if E is a measurable set, its Lebesgue measure, $m(E)$, is defined by

$$m(E) = m^*(E).$$

Proposition 2.13: *Lebesgue measure is countably additive*, that is, if $\{E_k\}_{k=1}^\infty$ is a countable disjoint collection of measurable sets, then its union $\bigcup_{k=1}^\infty E_k$ also is measurable and

$$m\left(\bigcup_{k=1}^\infty E_k\right) = \sum_{k=1}^\infty m(E_k).$$

Theorem 2.14: The set function Lebesgue measure, defined on the σ -algebra of Lebesgue measurable sets, assigns length to any interval, is translation invariant, and is countable additive.

Theorem 2.15 (the Continuity of Measure): Lebesgue measure possesses the following continuity properties:

1. If $\{A_k\}_{k=1}^\infty$ is an ascending collection of measurable sets, then

$$m\left(\bigcup_{k=1}^{\infty} A_k\right) = \lim_{k \rightarrow \infty} m(A_k).$$

1. If $\{B_k\}_{k=1}^{\infty}$ is a descending collection of measurable sets and $m(B_1) < \infty$, then

$$m\left(\bigcap_{k=1}^{\infty} B_k\right) = \lim_{k \rightarrow \infty} m(B_k).$$

The Borel-Cantelli Lemma: Let $\{E_k\}_{k=1}^{\infty}$ be a countable collection of measurable sets for which $\sum_{k=1}^{\infty} m(E_k) < \infty$. Then almost all $x \in \mathbb{R}$ belong to at most finitely many of the E_k 's.

Remark: In our forthcoming study of Lebesgue integration it will be apparent that it is the countable additivity of Lebesgue measure that provides the Lebesgue integral with its decisive advantage over the Riemann integral.

2.6 | Nonmeasurable Sets

Lemma 2.16: Let E be a bounded measurable set of real numbers. Suppose there is a bounded, countably infinite set of real numbers Λ for which the collection of translates of E , $\{\lambda + E\}_{\lambda \in \Lambda}$, is disjoint. Then $m(E) = 0$.

Theorem 2.17 (Vitali): Any set E of real numbers with positive outer measure contains a subset that fails to be measurable.

Theorem 2.18: There are disjoint sets of real numbers A and B for which

$$m^*(A \cup B) < m^*(A) + m^*(B).$$

2.7 | The Cantor Set and the Cantor-Lebesgue Function

Proposition 2.19: The Cantor set C is a closed, uncountable set of measure zero.

Proposition 2.20: The Cantor-Lebesgue function φ is an increasing continuous function that maps $[0, 1]$ onto $[0, 1]$. Its derivative exists on the open set \mathcal{O} , the complement in $[0, 1]$ of the Cantor set,

$$\varphi' = 0 \text{ on } \mathcal{O} \text{ while } m(\mathcal{O}) = 1.$$

Proposition 2.21: Let φ be the Cantor-Lebesgue function and define the function ψ on $[0, 1]$ by

$$\psi(x) = \varphi(x) + x \text{ for all } x \in [0, 1].$$

Proposition 2.22: There is a measurable set, a subset of the Cantor set, that is not a Borel set.

3 | Lebesgue Measurable Functions

3.1 | Sums, Products, and Compositions

Proposition 3.1: Let the function f have a measurable domain E . Then the following statements are equivalent:

1. For each real number c , the set $\{x \in E \mid f(x) > c\}$ is measurable.
2. For each real number c , the set $\{x \in E \mid f(x) \geq c\}$ is measurable.
3. For each real number c , the set $\{x \in E \mid f(x) < c\}$ is measurable.
4. For each real number c , the set $\{x \in E \mid f(x) \leq c\}$ is measurable.

Each of these properties implies that for each extended real number c ,

the set $\{x \in E \mid f(x) = c\}$ is measurable.

Definition: An extended real-valued function f defined on E is said to be **Lebesgue measurable**, or simply **measurable**, provided its domain E is measurable and it satisfies one of the four statements of Proposition 3.1.

Proposition 3.2: Let the function f be defined on a measurable set E . Then f is measurable if and only if for each open set \mathcal{O} , the inverse image of \mathcal{O} under f , $f^{-1}(\mathcal{O}) = \{x \in E \mid f(x) \in \mathcal{O}\}$, is measurable.

Proposition 3.3: A real-valued function that is continuous on its measurable domain is measurable.

Proposition 3.4: A monotone function that is defined on an interval is measurable.

Proposition 3.5: Let f be an extended real-valued function on E .

1. If f is measurable on E and $f = g$ a.e. on E , then g is measurable on E .
2. For a measurable subset D of E , f is measurable on E if and only if the restrictions of f to D and $E \sim D$ are measurable.

Theorem 3.6: Let f and g be measurable functions on E that are finite a.e. on E .

(Linearity) For any α and β ,

$$\alpha f + \beta g \text{ is measurable on } E.$$

(Products)

$$fg \text{ is measurable on } E.$$

Example: There are two measurable real-valued functions, each defined on all of \mathbb{R} , whose composition fails to be measurable. By Proposition 2.21, there is a continuous, strictly increasing function ψ defined on $[0, 1]$ and a measurable subset A of $[0, 1]$ for which $\psi(A)$ is nonmeasurable. Extend ψ to a continuous, strictly increasing function that maps \mathbb{R} onto \mathbb{R} . The function ψ^{-1} is continuous and therefore is measurable. On the other hand, A is a measurable set and so its characteristic function χ_A is a measurable function. We claim that the composition $f = \chi_A \circ \psi^{-1}$ is not measurable. Indeed, if I is any open interval containing 1 but not 0, then its inverse image under f is the nonmeasurable set $\psi(A)$.

Proposition 3.7: Let g be a measurable real-valued function defined on E and f a continuous real-valued function defined on all of \mathbb{R} . Then the composition $f \circ g$ is a measurable function on E .

Proposition 3.8: For a finite family $\{f_k\}_{k=1}^n$ of measurable functions with common domain E , the functions $\max\{f_1, \dots, f_n\}$ and $\min\{f_1, \dots, f_n\}$ also are measurable.

3.2 | Sequential Pointwise Limits and Simple Approximation

Definition: For a sequence $\{f_n\}$ of functions with common domain E , a function f on E and a subset A of E , we say that

1. The sequence $\{f_n\}$ converges to f pointwise on A provided

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \text{ for all } x \in A.$$

2. The sequence $\{f_n\}$ converges to f pointwise a.e. on A provided it converges to f pointwise on $A \sim B$, where $m(B) = 0$.
3. The sequence $\{f_n\}$ converges to f uniformly on A provided for each $\varepsilon > 0$, there is an index N for which

$$|f - f_n| < \varepsilon \text{ on } A \text{ for all } n \geq N.$$

Proposition 3.9: Let $\{f_n\}$ be a sequence of measurable functions on E that converges pointwise a.e. on E to the function f . Then f is measurable.

Definition: A real-valued function φ defined on a measurable set E is called **simple** provided it is measurable and takes only a finite number of values.

The Simple Approximation Lemma: Let f be a measurable real-valued function on E .

Assume f is bounded on E , that is, there is an $M \geq 0$ for which $|f| \leq M$ on E . Then for each $\varepsilon > 0$, there are simple functions φ_ε and ψ_ε defined on E which have the following approximation properties:

$$\varphi_\varepsilon \leq f \leq \psi_\varepsilon \text{ and } 0 \leq \psi_\varepsilon - \varphi_\varepsilon < \varepsilon \text{ on } E.$$

The Simple Approximation Theorem: An extended real-valued function f on a measurable set E is measurable if and only if there is a sequence $\{\varphi_n\}$ of simple functions on E which converges pointwise on E to f and has the property that

$$|\varphi_n| \leq |f| \text{ on } E \text{ for all } n.$$

If f is nonnegative, we may choose $\{\varphi_n\}$ to be increasing.

3.3 | Littlewood's Three Principles, Egoroff's Theorem, and Lusin's Theorem

Egoroff's Theorem: Assume E has finite measure. Let $\{f_n\}$ be a sequence of measurable functions on E that converges pointwise on E to the real-valued function f . Then for each $\varepsilon > 0$, there is a closed set F contained in E for which

$$\{f_n\} \rightarrow f \text{ uniformly on } F \text{ and } m(E \sim F) < \varepsilon.$$

Lemma 3.10: Under the assumptions of Egoroff's Theorem, for each $\eta > 0$ and $\delta > 0$, there is a measurable subset A of E and an index N for which

$$|f_n - f| < \eta \text{ on } A \text{ for all } n \geq N \text{ and } m(E \sim A) < \delta.$$

Proposition 3.11: Let f be a simple function defined on E . Then for each $\varepsilon > 0$, there is a continuous function g on \mathbb{R} and a closed set F contained in E for which

$$f = g \text{ on } F \text{ and } m(E \sim F) < \varepsilon.$$

Lusin's Theorem: Let f be a real-valued measurable function on E . Then for each $\varepsilon > 0$, there is a continuous function g on \mathbb{R} and a closed set F contained in E for which

$$f = g \text{ on } F \text{ and } m(E \sim F) < \varepsilon.$$

4 | Lebesgue Integration

4.1 | The Riemann Integral

Example (Dirichlet's Function): Define f on $[0, 1]$ by setting $f(x) = 1$ if x is rational and 0 if x is irrational. Let P be any partition of $[0, 1]$. By the density of the rationals and the irrationals,

$$L(f, P) = 0 \text{ and } U(f, P) = 1.$$

Thus

$$(R) \int_0^1 f = 0 < 1 = (R) \int_0^1 \bar{f},$$

so f is not Riemann integrable. The set of rational numbers in $[0, 1]$ is countable. Let $\{q_k\}_{k=1}^{\infty}$ be an enumeration of the rational numbers in $[0, 1]$. For a natural number n , define f_n on $[0, 1]$ by setting $f_n(x) = 1$, if $x = q_k$ for some q_k with $1 \leq k \leq n$, and $f(x) = 0$ otherwise. Then each f_n is a step function, so it is Riemann integrable. Thus, $\{f_n\}$ is an increasing sequence of Riemann integrable functions on $[0, 1]$,

$$|f_n| \leq 1 \text{ on } [0, 1] \text{ for all } n$$

and

$$\{f_n\} \rightarrow f \text{ pointwise on } [0, 1].$$

However, the limit function f fails to be Riemann integrable on $[0, 1]$.

4.2 | The Lebesgue Integral of a Bounded Measurable Function over a Set of Finite Measure

Definition: For a simple function ψ defined on a set of finite measure E , we define the integral of ψ over E by

$$\int_E \psi = \sum_{i=1}^n a_i \cdot m(E_i),$$

where ψ has the canonical representation.

Lemma 4.1: Let $\{E_i\}_{i=1}^n$ be a finite disjoint collection of measurable subsets of a set of finite measure E . For $1 \leq i \leq n$, let a_i be a real number.

$$\text{If } \varphi = \sum_{i=1}^n a_i \cdot \chi_{E_i} \text{ on } E, \text{ then } \int_E \varphi = \sum_{i=1}^n a_i \cdot m(E_i).$$

Proposition 4.2 (Linearity and Monotonicity of Integration): Let φ and ψ be simple functions defined on a set of finite measure E . Then for any α and β ,

$$\int_E (\alpha\varphi + \beta\psi) = \alpha \int_E \varphi + \beta \int_E \psi.$$

Moreover,

$$\text{if } \varphi \leq \psi \text{ on } E, \text{ then } \int_E \varphi \leq \int_E \psi.$$

Definition: A bounded function f on a domain E of finite measure is said to be **Lebesgue integrable** over E provided its upper and lower Lebesgue integrals over E are equal. The common value of the upper and lower integrals is called the **Lebesgue integral**, or simply the integral, of f over E and is denoted by $\int_E f$.

Theorem 4.3: Let f be a bounded function defined on the closed, bounded interval $[a, b]$. If f is Riemann integrable over $[a, b]$, then it is Lebesgue integrable over $[a, b]$ and the two integrals are equal.

Example: The set E of rational numbers in $[0, 1]$ is a measurable set of measure zero. The Dirichlet function f is the restriction to $[0, 1]$ of the characteristic function of E , χ_E . Thus f is integrable over $[0, 1]$ and

$$\int_{[0,1]} f = \int_{[0,1]} 1 \cdot \chi_E = 1 \cdot m(E) = 0.$$

We have shown that f is not Riemann integrable over $[0, 1]$.

Theorem 4.4: Let f be a bounded measurable function on a set of finite measure E . Then f is integrable over E .

Theorem 4.5 (Linearity and Monotonicity of Integration): Let f and g be bounded measurable functions on a set of finite measure E . Then for any α and β ,

$$\int_E (\alpha f + \beta g) = \alpha \int_E f + \beta \int_E g.$$

Moreover,

$$\text{if } f \leq g \text{ on } E, \text{ then } \int_E f \leq \int_E g.$$

Corollary 4.6: Let f be a bounded measurable function on a set of finite measure E . Suppose A and B are disjoint measurable subsets of E . Then

$$\int_{A \cup B} f = \int_A f + \int_B f.$$

Corollary 4.7: Let f be a bounded measurable function on a set of finite measure E . Then

$$\left| \int_E f \right| \leq \int_E |f|.$$

Proposition 4.8: Let $\{f_n\}$ be a sequence of bounded measurable functions on a set of finite measure E .

$$\text{If } \{f_n\} \rightarrow f \text{ uniformly on } E, \text{ then } \lim_{n \rightarrow \infty} \int_E f_n = \int_E f.$$

Example: For each natural number n , define f_n on $[0, 1]$ to have the value 0 if $x \geq 2/n$, have $f(1/n) = n$, $f(0) = 0$ and to be linear on the intervals $[0, 1/n]$ and $[1/n, 2/n]$. Observe that $\int_0^1 f_n = 1$ for each n . Define $f \equiv 0$ on $[0, 1]$. Then

$$\{f_n\} \rightarrow f \text{ pointwise on } [0, 1], \text{ but } \lim_{n \rightarrow \infty} \int_0^1 f_n \neq \int_0^1 f.$$

Thus, pointwise convergence alone is not sufficient to justify passage of the limit under the integral sign.

The Bounded Convergence Theorem: Let $\{f_n\}$ be a sequence of measurable functions on a set of finite measure E . Suppose $\{f_n\}$ is uniformly pointwise bounded on E , that is, there is a number $M \geq 0$ for which

$$|f_n| \leq M \text{ on } E \text{ for all } n.$$

$$\text{If } \{f_n\} \rightarrow f \text{ pointwise on } E, \text{ then } \lim_{n \rightarrow \infty} \int_E f_n = \int_E f.$$

Remark: Prior to the proof of the Bounded Convergence Theorem, no use was made of the countable additivity of Lebesgue measure on the real line. Only finite additivity was used, and it was used just once, in the proof of Lemma 1. But for the proof of the Bounded Convergence Theorem we used Egoroff's Theorem. The proof of Egoroff's Theorem needed the continuity of Lebesgue measure, a consequence of countable additivity of Lebesgue measure.

4.3 | The Lebesgue Integral of a Measurable Nonnegative Function

Definition: For f a nonnegative measurable function on E , we define the integral of f over E by

$$\int_E f = \sup \left\{ \int_E h \mid h \text{ bounded, measurable, of finite support and } 0 \leq h \leq f \text{ on } E \right\}.$$

Chebychev's Inequality: Let f be a nonnegative measurable function on E . Then for any $\lambda > 0$,

$$m\{x \in E \mid f(x) \geq \lambda\} \leq \frac{1}{\lambda} \cdot \int_E f.$$

Proposition 4.9: Let f be a nonnegative measurable function on E . Then

$$\int_E f = 0 \text{ if and only if } f = 0 \text{ a.e. on } E.$$

Theorem 4.10 (Linearity and Monotonicity of Integration): Let f and g be nonnegative measurable functions on a set of finite measure E . Then for any $\alpha > 0$ and $\beta > 0$,

$$\int_E (\alpha f + \beta g) = \alpha \int_E f + \beta \int_E g.$$

Moreover,

$$\text{if } f \leq g \text{ on } E, \text{ then } \int_E f \leq \int_E g.$$

Theorem 4.11 (Additivity over Domains of Integration): Let f be a nonnegative measurable function on E . If A and B are disjoint measurable subsets of E , then

$$\int_{A \cup B} f = \int_A f + \int_B f.$$

In particular, if E_0 is a subset of E of measure zero, then

$$\int_E f = \int_{E \sim E_0} f.$$

Fatou's Lemma: Let $\{f_n\}$ be a sequence of nonnegative measurable functions on E .

$$\text{If } \{f_n\} \rightarrow f \text{ pointwise a.e. on } E, \text{ then } \int_E f \leq \liminf \int_E f_n.$$

Example: Let $E = (0, 1]$ and for a natural number n , define $f_n = n \cdot \chi_{(0, 1/n)}$. Then $\{f_n\}$ converges pointwise on E to $f \equiv 0$ on E . However,

$$\int_E f = 0 < 1 = \lim_{n \rightarrow \infty} \int_E f_n.$$

As another example of strict inequality in Fatou's Lemma, let $E = \mathbb{R}$ and for a natural number n , define $g_n = \chi_{(n, n+1)}$. Then $\{g_n\}$ converges pointwise on E to $g \equiv 0$ on E . However,

$$\int_E g = 0 < 1 = \lim_{n \rightarrow \infty} \int_E g_n.$$

The Monotone Convergence Theorem: Let $\{f_n\}$ be an increasing sequence of nonnegative measurable functions on E .

$$\text{If } \{f_n\} \rightarrow f \text{ pointwise a.e. on } E, \text{ then } \lim_{n \rightarrow \infty} \int_E f_n = \int_E f.$$

Corollary 4.12: Let $\{u_n\}$ be a sequence of nonnegative measurable functions on E .

$$\text{If } f = \sum_{n=1}^{\infty} u_n \text{ pointwise a.e. on } E, \text{ then } \int_E f = \sum_{n=1}^{\infty} \int_E u_n.$$

Definition: A nonnegative measurable function f on a measurable set E is said to be **integrable** over E provided

$$\int_E f < +\infty.$$

Proposition 4.13: Let the nonnegative function f be integrable over E . Then f is finite a.e. on E .

Beppo Levi's Lemma: Let $\{f_n\}$ be an increasing sequence of nonnegative measurable functions on E . If the sequence of integrals $\left\{ \int_E f_n \right\}$ is bounded, then $\{f_n\}$ converges pointwise on E to a measurable function f that is finite a.e. on E and

$$\lim_{n \rightarrow \infty} \int_E f_n = \int_E f < +\infty.$$

4.4 | The General Lebesgue Integral

Proposition 4.14: Let f be a measurable function on E . Then f^+ and f^- are integrable over E if and only if $|f|$ is integrable over E .

Definition: A measurable function f on E is said to be **integrable** over E provided $|f|$ is integrable over E . When this is so we define the integral of f over E by

$$\int_E f = \int_E f^+ - \int_E f^-.$$

Proposition 4.15: Let f be integrable over E . Then f is finite a.e. on E and

$$\int_E f = \int_{E \sim E_0} f \text{ if } E_0 \subseteq E \text{ and } m(E_0) = 0.$$

Proposition 4.16 (The Integral Comparison Test): Let f be a measurable function on E . Suppose there is a nonnegative function g that is integrable over E and dominates f in the sense that

$$|f| \leq g \text{ on } E.$$

Then f is integrable over E and

$$\left| \int_E f \right| \leq \int_E |f|.$$

Theorem 4.17 (Linearity and Monotonicity of Integration): Let f and g be integrable over E . Then for any $\alpha > 0$ and $\beta > 0$, the function $\alpha f + \beta g$ is integrable over E and

$$\int_E (\alpha f + \beta g) = \alpha \int_E f + \beta \int_E g.$$

Moreover,

$$\text{if } f \leq g \text{ on } E, \text{ then } \int_E f \leq \int_E g.$$

Theorem 4.18 (Additivity over Domains of Integration): Let f be integrable over E . Assume A and B are disjoint measurable subsets of E . Then

$$\int_{A \cup B} f = \int_A f + \int_B f.$$

The Lebesgue Dominated Convergence Theorem: Let $\{f_n\}$ be a sequence of measurable functions on E . Suppose there is a function g that is integrable over E and dominates $\{f_n\}$ on E in the sense that $|f_n| \leq g$ on E for all n .

If $\{f_n\} \rightarrow f$ pointwise a.e. on E , then f is integrable over E and $\lim_{n \rightarrow \infty} \int_E f_n = \int_E f$.

Theorem 4.19 (General Lebesgue Dominated Convergence Theorem): Let $\{f_n\}$ be a sequence of measurable functions on E that converges pointwise a.e. on E to f . Suppose there is a sequence $\{g_n\}$ of nonnegative measurable functions on E that converges pointwise a.e. on E to g and dominates $\{f_n\}$ on E in the sense that

$$|f_n| \leq g_n \text{ on } E \text{ for all } n.$$

$$\text{If } \lim_{n \rightarrow \infty} \int_E g_n = \int_E g < +\infty, \text{ then } \lim_{n \rightarrow \infty} \int_E f_n = \int_E f.$$

Remark: If Fatou's Lemma and the Lebesgue Dominated Convergence Theorem, the assumption of pointwise convergence a.e. on E rather than on all of E is not a decoration pinned on to honor generality. It is necessary for future applications of these results. We provide one illustration of this necessity. Suppose f is an increasing function on all of \mathbb{R} . A forthcoming theorem of Lebesgue (Lebesgue's Theorem of Chapter 6) tells us that

$$\lim_{n \rightarrow \infty} \frac{f(x + 1/n) - f(x)}{1/n} = f'(x) \text{ for almost all } x.$$

From this and Fatou's Lemma we will show that for any closed, bounded interval $[a, b]$,

$$\int_a^b f'(x) dx \leq f(b) - f(a).$$

In general, given a nondegenerate closed, bounded interval $[a, b]$ and a subset A of $[a, b]$ that has measure zero, there is an increasing function f on $[a, b]$ for which the limit in previous formula fails to exist at each point in A .

4.5 | Countable Additivity and Continuity of Integration

Theorem 4.20 (The Countable Additivity of Integration): Let f be integrable over E and $\{E_n\}_{n=1}^{\infty}$ a disjoint countable collection of measurable subsets of E whose union is E . Then

$$\int_E f = \sum_{n=1}^{\infty} \int_{E_n} f.$$

Theorem 4.21 (The Continuity of Integration): Let f be integrable over E .

1. If $\{E_n\}_{n=1}^{\infty}$ is an ascending countable collection of measurable subsets of E , then

$$\int_{\bigcup_{n=1}^{\infty} E_n} f = \lim_{n \rightarrow \infty} \int_{E_n} f.$$

2. If $\{E_n\}_{n=1}^{\infty}$ is a descending countable collection of measurable subsets of E , then

$$\int_{\bigcap_{n=1}^{\infty} E_n} f = \lim_{n \rightarrow \infty} \int_{E_n} f.$$

4.6 | Uniform Integrability: The Vitali Convergence Theorem

Lemma 4.22: Let E be a set of finite measure and $\delta > 0$. Then E is the disjoint union of a finite collection of sets, each of which has measure less than δ .

Proposition 4.23: Let f be a measurable function on E . If f is integrable over E , then for each $\varepsilon > 0$, there is a $\delta > 0$ for which

$$\text{if } A \subseteq E \text{ is measurable and } m(A) < \delta, \text{ then } \int_A |f| < \varepsilon.$$

Conversely, in the case $m(E) < \infty$, if for each $\varepsilon > 0$, there is a $\delta > 0$ for which previous formula holds, then f is integrable over E .

Definition: A family \mathcal{F} of measurable functions on E is said to be **uniformly integrable over E** provided for each $\varepsilon > 0$, there is a $\delta > 0$ such that for each $f \in \mathcal{F}$,

$$\text{if } A \subseteq E \text{ is measurable and } m(A) < \delta, \text{ then } \int_A |f| < \varepsilon.$$

Proposition 4.24: Let $\{f_k\}_{k=1}^n$ be a finite collection of functions, each of which is integrable over E . Then $\{f_k\}_{k=1}^\infty$ is uniformly integrable.

Proposition 4.25: Assume E has finite measure. Let the sequence of functions $\{f_n\}$ be uniformly integrable over E . If $\{f_n\} \rightarrow f$ pointwise a.e. on E , then f is integrable over E .

The Vitali Convergence Theorem: Let E be of finite measure. Suppose the sequence of functions $\{f_n\}$ is uniformly integrable over E .

$$\text{If } \{f_n\} \rightarrow f \text{ pointwise a.e. on } E, \text{ then } f \text{ is integrable over } E \text{ and } \lim_{n \rightarrow \infty} \int_E f_n = \int_E f.$$

Theorem 4.26: Let E be of finite measure. Suppose $\{h_n\}$ is a sequence of nonnegative integrable functions that converges pointwise a.e. on E to $h \equiv 0$. Then

$$\lim_{n \rightarrow \infty} \int_E h_n = 0 \text{ if and only if } \{h_n\} \text{ is uniformly integrable over } E.$$