

A CONDITIONAL SKEIN-LASAGNA OBSTRUCTION FOR A TRACE-73 CAPPELL–SHANESON CANDIDATE, PENDING GEOMETRIC IDENTIFICATION

ANONYMOUS REPOSITORY AUDIT

ABSTRACT. We study a proposed trace-73 Cappell–Shaneson construction associated with the matrix

$$A = \begin{pmatrix} 0 & 269 & 1240 \\ 0 & 41 & 189 \\ 1 & 0 & 32 \end{pmatrix}.$$

A public finite calculation produces the nonzero cubic Burau scalar -59072 in the grading convention corresponding to quantum degree 494. An exact Artin–Magnus certificate and the pure-braid Andreadakis theorem establish a third-order property of the public braid. We prove an abstract obstruction: if the compact word ledger is realized by the required framed Kirby presentation, if the Burau functional is identified with the specified rational $N = 2$ Manolescu–Walker–Wedrich coefficient trace in the required symmetric-monoidal form, then the candidate is a homotopy sphere not diffeomorphic to S^4 , subject also to the stated final identifications. A relative complete-sphere argument shows that the three-handle closure then follows without an independent sphere-movie hypothesis. The Kirby and coefficient-trace identifications remain open. The Lean development checks the finite and quotient algebra; it does not construct the external geometry or formalize four-dimensional differential topology.

CONTENTS

1. Introduction	2
Contributions	3
2. The trace-73 Cappell–Shaneson candidate	3
3. Precise statements	4
4. The finite computational layer	5
4.1. Integral lattice calculations	5
4.2. The public Burau computation	6
5. Lasagna modules and the intended obstruction	6

Date: September 2, 2026.

2020 Mathematics Subject Classification. Primary 57K40, 57R55; Secondary 57K18, 68V20.

Key words and phrases. smooth Poincaré conjecture, Cappell–Shaneson sphere, skein lasagna module, Khovanov homology, formal verification.

6. Compact product-ribbon presentation	7
7. The coefficient-trace comparison	8
8. Three-handle sphere closure	9
9. Final identifications	10
10. Formalization boundary	11
11. Status table and reproducibility ledger	11
12. Related work	12
13. Reproducibility and review	12
14. Conclusion	13
Appendix A. Retired historical objects	13
Appendix B. Correspondence with the Lean fields	14
References	15

1. INTRODUCTION

The smooth four-dimensional Poincare conjecture asks whether every smooth closed manifold homotopy equivalent to S^4 is diffeomorphic to S^4 . Freedman’s work proves the topological counterpart: a homotopy 4-sphere is homeomorphic to S^4 [8]. The remaining question is therefore a smooth classification problem, and any proposed counterexample must establish two logically separate conclusions:

- (1) the candidate is a homotopy 4-sphere;
- (2) the candidate is not diffeomorphic to the standard smooth sphere.

Cappell and Shaneson constructed a distinguished family of potential homotopy 4-spheres from torus automorphisms [5]. Their handle structures were studied by Aitchison and Rubinstein [1]. Many subfamilies were subsequently shown to be standard, through work including Akbulut [2], Gompf [9], Kim–Yamada [12], and Iwaki [11]. This history makes geometric identification especially important: agreement of a matrix, a relator, or a trace parameter is not a substitute for identifying a complete framed handle presentation.

Skein lasagna modules extend Khovanov–Rozansky link homology to smooth four-manifolds [15]. Manolescu and Neithalath gave a two-handlebody description [13], and Manolescu, Walker, and Wedrich gave formulas compatible with general handle decompositions [14]. These results provide an attractive route to conclusion (ii): a nonzero class in a nonzero bidegree cannot be carried by a diffeomorphism to the lasagna module of S^4 , which is concentrated in bidegree zero.

We investigate whether the public finite class can be constructed for the standard-form Cappell–Shaneson parameter (41, 189, 73). Compact product-ribbon, point-push, owner-lift and Nielsen ledgers replace parts of the unavailable historical data, but they do not by themselves give an embedded framed Kirby equivalence or the candidate-specific quantum-trace map. Once

those are supplied in a relative monoidal form, a standard sphere system disjoint from the detector gives the three-handle closure. The paper therefore separates the conditional theorem from the remaining comparison gaps.

Contributions. The main contributions are as follows.

- (1) We give a compact word-level Aitchison–Rubinstein ledger and a six-sweep construction of the public point-push braid, while isolating the missing framed-embedding/Kirby equivalence.
- (2) We identify the local checked R-matrix with the public Burau blocks and prove a full-entry third-order statement, while isolating the missing candidate-specific MWW-to-endpoint comparison and simultaneous transport.
- (3) We use the relative uniqueness of complete sphere systems and MWW’s intrinsic local module action to derive the three-handle closure from P0 and the symmetric-monoidal form of C, without TH-sized sphere movies.
- (4) We provide executable ledgers, mutation tests and a kernel-checked Lean algebraic core.

Historical files referenced by earlier versions of the project are not publicly available. Their hashes are retained as provenance. The compact scripts regenerate useful projections of the proposed geometry but do not yet regenerate every load-bearing candidate-specific construction.

2. THE TRACE-73 CAPPELL–SHANESON CANDIDATE

Let

$$(1) \quad A = \begin{pmatrix} 0 & 269 & 1240 \\ 0 & 41 & 189 \\ 1 & 0 & 32 \end{pmatrix}.$$

This is Iwaki’s standard form $X_{41,189,73}$; the triple occurs among the trace-73 representatives in his table [11, Table 2]. Direct integer arithmetic gives

$$(2) \quad \det A = 1, \quad \det(A - I) = 1.$$

For $A \in \mathrm{SL}(3, \mathbb{Z})$, let $f_A : T^3 \rightarrow T^3$ be the induced diffeomorphism, made equal to the identity near a chosen point. Surgery on the corresponding circle in the mapping torus gives a Cappell–Shaneson manifold Σ_A^ε , where $\varepsilon \in \mathbb{Z}/2$ records the framing choice. The standard criterion says that Σ_A^ε is a homotopy 4-sphere if and only if $\det(A - I) = \pm 1$; see Iwaki’s Proposition 2.1 and the original sources cited there [11, 5].

Consequently, (2) proves that the standard construction Σ_A^ε is a homotopy 4-sphere. It does not prove that a separately generated large Kirby diagram is a diagram of that construction. That identification is the first and most load-bearing hypothesis below.

Iwaki proves standardness for many infinite families. The appearance of (41, 189, 73) in a representative table is not a standardness or exoticness

theorem for this individual sphere. Likewise, the trace bound in Kim–Yamada’s work stops below 73 [12]. These observations explain the choice of candidate but do not decide its diffeomorphism type.

3. PRECISE STATEMENTS

We first state the abstract theorem in the form actually proved by the Lean development. This prevents geometric intuition from silently strengthening the result.

Fix a collection of manifold labels, a graded family of rational vector spaces $G(M, q)$, labels X and S^4 , and predicates $\text{HS}(M)$ and $\text{Diff}(M, N)$.

Theorem 3.1 (Abstract nonstandardness theorem). *Suppose there are rational vector spaces W_0, W_1, W_2, W_3 , an element $x_0 \in W_0$, linear maps*

$$W_0 \xrightarrow{q_{01}} W_1 \xrightarrow{q_{12}} W_2, \quad \ell_i : W_i \longrightarrow \mathbb{Q} \quad (i = 0, 1, 2),$$

and linear isomorphisms

$$T : W_2 \xrightarrow{\cong} W_3, \quad F : W_3 \xrightarrow{\cong} G(X, 494),$$

such that

$$(3) \quad \ell_0(x_0) = -59072,$$

$$(4) \quad \ell_1 q_{01} = \ell_0,$$

$$(5) \quad \ell_2 q_{12} = \ell_1.$$

Assume also that every element of $G(S^4, 494)$ is zero and that every diffeomorphism $X \rightarrow S^4$ induces a linear isomorphism $G(X, 494) \cong G(S^4, 494)$. Then X is not diffeomorphic to S^4 .

Proof. Set $v = F(T(q_{12}(q_{01}(x_0))))$. If $v = 0$, injectivity of F and T implies $q_{12}(q_{01}(x_0)) = 0$. Applying ℓ_2 and using (4)–(5) gives $-59072 = 0$, a contradiction. Thus $v \neq 0$. If X were diffeomorphic to S^4 , the induced isomorphism would send v to an element of $G(S^4, 494)$, hence to zero, contradicting injectivity. $\square$

The Lean theorem corresponding to Theorem 3.1 accepts the input structures `ExternalGeometry` and `CSExternalGeometry`; no inhabitants of these structures are constructed in the repository. We name the candidate-specific assumptions needed to instantiate them.

Hypothesis 3.2 (Compact geometric presentation, P0). The public compact Aitchison–Rubinstein product-ribbon ledger is a labelled and framed handle presentation of Σ_A^0 . Its two product cancellations give the registered genus-two words, and its six-sweep tubular point-push induces the public 44- and 88-strand braids on all oriented cable states. The detector collar is contained in a specified boundary 3-ball B , and the presentation has three 3-handles followed by one final 4-handle.

Hypothesis 3.3 (Coefficient comparison, C). The completed MWW coefficient quantum trace admits a grading-preserving, symmetric-monoidal natural map to the BPW/BHPW weight-one endpoint representation, natural also for boundary cobordisms supported away from B . Under this map the point-push action is the public Burau representation, and the divided cubic row descends through every beta and psi relation. Its selected value is -59072 up to an invertible convention unit, and hence is nonzero in quantum degree 494.

Theorem 3.4 (Relative three-handle consequence, S). *Under Hypotheses 3.2 and 3.3, the three 3-handles may be attached along a complete sphere system disjoint from B . For each sphere the detector kills the undotted MWW relation and identifies the once-dotted map with the source term. Hence the divided detector descends through the complete three-handle coequalizer.*

Hypothesis 3.5 (Candidate-level final identifications, P3). MWW's three-/four-handle formulas identify the final quotient with $\mathcal{S}_{0,(0,494),0}^2(\Sigma_A^0; \mathbb{Q})$. The corresponding summand for S^4 is zero, and diffeomorphisms induce grading-preserving equivalences. Moreover, $\det(A - I) = 1$ makes Σ_A^0 a homotopy sphere. The general published theorems supporting these statements are not the candidate-specific identifications asserted here.

Theorem 3.6 (Conditional trace-73 theorem). *Assume Hypotheses 3.2, 3.3, and 3.5. Then the resulting trace-73 candidate satisfies*

$$\Sigma_A^0 \simeq S^4 \quad \text{and} \quad \Sigma_A^0 \not\cong_{\text{diff}} S^4.$$

Proof of Theorem 3.6. Hypotheses 3.2 and 3.5 and the Cappell–Shaneson criterion give the homotopy-sphere assertion. Hypothesis 3.3 and Theorem 3.4 instantiate the maps and rows in Theorem 3.1; its selected class is nonzero in quantum degree 494. MWW's four-handle isomorphism preserves this degree, while the same degree of the standard S^4 module vanishes. Diffeomorphism invariance and Theorem 3.1 rule out a diffeomorphism to S^4 . $\square$

4. THE FINITE COMPUTATIONAL LAYER

The public finite layer has three independent parts: lattice arithmetic, a chosen-sphere coordinate determinant, and a truncated Burau calculation.

4.1. Integral lattice calculations. Besides (2), the repository records the proposed sphere coordinate columns

$$(6) \quad C = \begin{pmatrix} -1311 & -189 & 41 \\ 8608 & 1241 & -269 \\ -1 & 0 & 1 \end{pmatrix}, \quad \det C = 1.$$

Thus the three recorded coordinate vectors form a basis of $\mathbb{Z}^3$. This is only the last arithmetic step in a geometric-basis argument. It does not construct embedded spheres, prove disjointness, or identify $\mathbb{Z}^3$ with $H_2^{\text{sph}}(\partial W_2)$.

The Lean files also exhibit integral inverses for $A - I$ and for the induced map on $H_2(T^3; \mathbb{Z})$. These computations eliminate the candidate-specific lattice algebra from the homotopy-sphere branch. Van Kampen, Wang–Mayer–Vietoris, Poincare duality, and Hurewicz–Whitehead arguments remain supplied through a topology structure rather than formalized as four-manifold topology.

4.2. The public Burau computation. Let $\varepsilon = t - 1$. The script `scripts/recompute_t73_delta3.py` reads a result-free JSON input and performs the following finite operations:

- (1) parse 252 primitive crossing rows and reconstruct an 11,340-letter Artin word on 44 strands;
- (2) cable it to a 45,360-letter word on 88 strands;
- (3) apply the unreduced Burau representation over $\mathbb{Z}[\varepsilon]/(\varepsilon^7)$;
- (4) substitute $t = q^{-2}$ with $q = 1 + h$, equivalently $\varepsilon = (1 + h)^{-2} - 1$;
- (5) extract the coefficient of h^3 after the specified source vector and covector are applied.

Denote the resulting scalar by δ_3^{Bur} . The independently reproduced public result is

$$(7) \quad \delta_3^{\text{Bur}} = -59072 = (-2)^3 \cdot 7384.$$

The public input does not contain either decimal literal 7384 or -59072 ; the former is, however, stored as the constant `cubicBase` in the Lean finite layer. Lean checks only the arithmetic $(-2)^3 \cdot 7384 = -59072$ and never evaluates the 45,360-letter word.

Proposition 4.1 (Exact scope of the computation). *Equation (7) is a statement about the specified truncated Burau model. The script and JSON alone do not define the MWW invariant. Under the coefficient-trace comparison of Theorem 3.3, however,*

$$\delta_3^{\text{MWW}} = \delta_3^{\text{Bur}} = -59072,$$

up to the harmless global convention unit normalized in that theorem.

Proof. The script defines vector and covector operations in an 88-dimensional truncated polynomial representation. The displayed MWW equality is exactly the additional content of Hypothesis 3.3; it does not follow from the finite script or from the local Burau-block calculation. $\square$

Thus the finite recomputation and the categorical comparison have distinct roles: the former supplies the integer, while the latter makes it a four-manifold invariant.

5. LASAGNA MODULES AND THE INTENDED OBSTRUCTION

We recall only the features needed for the comparison. For a smooth compact oriented four-manifold W and a framed link $L \subset \partial W$, the skein

lasagna module $\mathcal{S}_0^N(W; L)$ is generated by decorated surfaces in W , modulo local relations coming from KhR_N cobordism maps [15, 14].

For a handle decomposition

$$W_1 \subset W_2 \subset W_3 \subset W_4,$$

where W_1 is a one-handlebody, W_2 adds two-handles along a framed link, W_3 adds three-handles along embedded spheres, and W_4 adds four-handles, the published results have the following roles.

- MWW Theorem 3.2 identifies the two-handle stage with a cabled skein lasagna quotient, whose relations include beta and psi operations.
- MWW Theorems 3.7 and 3.10 describe each three-handle as a co-equalizer of the two hemisphere maps, and give the complete handle formula.
- MWW Proposition 3.4 says that a four-handle attachment induces an isomorphism for the empty boundary link.
- MWW Corollary 3.5 gives $\mathcal{S}_0^N(S^4) \cong \mathbb{Z}$, concentrated in bidegree $(0, 0)$.

At $N = 2$, a nonzero class in bidegree $(0, 494)$ would therefore obstruct a diffeomorphism to S^4 . The intended proof constructs a functional at the one-handle stage, shows that it annihilates the two-handle and three-handle relations, and transports the resulting nonzero class through the four-handle isomorphism.

The formal algebra correctly captures the last sentence. The next sections record partial constructions and the exact candidate-specific inputs still needed by that algebra.

6. COMPACT PRODUCT-RIBBON PRESENTATION

Start with the Aitchison–Rubinstein product-annulus handle construction [1]. For the j th coordinate circle, order the integer-face crossings of the straight segment from 0 to Ae_j by their rational times. A fixed product perturbation resolves simultaneous events. This gives the mapping-torus attaching word

$$m_j = t\phi_A(x_j)t^{-1}x_j^{-1}$$

with its product-annulus normal.

The section handle cancels the base one-handle t without depositing a twist. Since $Ae_1 = e_3$, the remaining word $m_1 = zx^{-1}$ gives a second product cancellation and replaces x by z . The resulting public ledger has

$$\begin{aligned} |m_2| &= 311, & [m_2]_{(y,z)} &= (40, 269), \\ |m_3| &= 1460, & [m_3]_{(y,z)} &= (189, 1271). \end{aligned}$$

The word $r_{zx} = [z, z]$ formally contracts through two product bigons. If this word projection lifts to the claimed product-annulus embedding, it gives the desired framed genus-two presentation of Σ_A^0 . The current generator does not output an embedded framed link or a labelled ambient isotopy/Kirby-move

ledger, so this lift remains Hypothesis 3.2. This is not a merely cosmetic omission: tying a local knot into a component inside a ball preserves its free-group word and integer framing but can change the framed surgery manifold. Thus no word/framing-name argument can recover the required whole-link embedding.

The point-push collar is generated by six affine sweeps, each with 42 pure crossing factors. The generator regenerates all 252 public rows and the 11,340-letter braid. Its return wicket closes a framed loop of the r_{xy} product connector through the complement of the m_2 passages. The rows alone do not construct a tubular ambient motion in the actual handlebody. Such a motion Φ_t , together with its simultaneous action on all parallel copies, must be extracted from a solution of P0 before the public 88-strand cabling can be identified with the candidate’s MWW state.

7. THE COEFFICIENT-TRACE COMPARISON

The compact words give

$$p_y = 42 + 2 = 44, \quad p_z = 269 + 2 = 271.$$

The counts suggest product subrectangles pairing the 44 y wickets with 44 z passages and 227 remaining circles. The desired cut coefficient would then have the two-sided Hattori form

$$M_R(T, T') \cong \text{Hom}(B_{act}T, B_{act}T')\{-44\} \otimes A^{\otimes 227}, \quad A = \mathbb{Q}[X]/(X^2).$$

MWW Theorem 4.7 identifies the one-handle module with the coefficient trace of the cut tangle categories and describes cobordism maps component-wise [14]. BPW’s functorial quantum vertical-to-horizontal trace and annular functor give

$$q \text{Tr}(\mathcal{C}; M_R) \longrightarrow hTr_q(BN) \xrightarrow{F_{A_q}} \text{gRep}(U_q(\mathfrak{sl}_2))$$

as constructed by BPW [4]. BHPW supplies strict integral tangle/foam functoriality [3]. The base change $q \mapsto 1 + h$ is a rational localization followed by a Noetherian adic completion and is flat.

The endpoint type must be stated carefully. Put

$$E_{86} = (V^{\otimes 86})_{\lambda=86}, \quad E_{88} = (V^{\otimes 88})_{\lambda=86}.$$

Then E_{86} has rank one and E_{88} has rank 88; after restriction to weight spaces the cup lies in $\text{Hom}_{\mathbb{Q}[[h]]}(E_{86}, E_{88}) \cong E_{88}$. The tangle maps are $U_q(\mathfrak{sl}_2)$ intertwiners before this restriction. It is therefore incorrect to place a “weight-86 summand” on an intertwiner space itself.

There is also no abstract obstruction at the representable-coefficient step. For a rational-linear autofunctor B , the coefficient bimodule $(T, T') \mapsto \text{Hom}(BT, BT')$ is two-sided equivalent to the regular coefficient bimodule, and hence their coefficient HH_0 groups are linearly equivalent. This quotient statement is kernel checked in `Smooth4PC/RepresentableCoefficient`.

lean. Its use here still requires the missing candidate-specific, action-compatible Hattori equivalence from M_R to that representable model, as well as its q -deformed completed lift.

On the weight-one subspace, the normalized checked R-matrix is

$$\begin{pmatrix} 1 - q^{-2} & q^{-1} \\ q^{-1} & 0 \end{pmatrix}.$$

After conjugation by $\text{diag}(1, q^{-1})$ and setting $t = q^{-2}$, this becomes the public Burau block. This is a valid local comparison. It does not construct a map from the candidate's MWW coefficient trace to the specified qHH0/Chern summand, nor the single coordinate change P that must transport the actual operator, source vector, and covector simultaneously.

The compact word has zero writhe and identity permutation. Direct evaluation of all 88 basis vectors proves

$$\rho_h(W) - I \in h^3 \text{End}(E_{88}).$$

An exact noncommutative Magnus expansion of the Artin action on F_{44} shows that the automorphism is the identity through degrees one, two and three. All intermediate integer coefficients have absolute value at most four. Darné's Andreadakis equality for the pure braid group then gives

$$W \in \Gamma_3(P_{44}).$$

See [6, Theorem 6.2]. Every physical cabling is a group homomorphism and hence preserves the lower central series. Since each local pure generator acts as $I + O(h)$, every statewise cabling satisfies $\rho_h(W_s) - I = O(h^3)$. If the missing candidate-specific map and simultaneous transport exist, BPW's tangle action gives statewise equations $Q_e W_s = W_t Q_e$ for the beta and psi edges. General functoriality transports an identified action; it does not construct this missing arrow. Finally, the core identities

$$(\epsilon \otimes \epsilon)\Delta(1) = 0, \quad (\epsilon \otimes \epsilon)\Delta(X) = 1$$

give the required Frobenius equations after the actual maps have been bound. Thus the public results prove the order-three and local algebraic parts of C, while the coefficient-trace comparison and complete beta/psi descent remain Hypothesis 3.3.

8. THREE-HANDLE SPHERE CLOSURE

We prove Theorem 3.4. Remove the final 4-handle and call the remaining handlebody W_3 . Its boundary is S^3 . Reversing the three 3-handles attaches three three-dimensional 1-handles to S^3 , and hence

$$\partial W_2 \cong \#^3(S^1 \times S^2).$$

The actual attaching sphere system has connected complement, since sphere surgery on it gives the connected boundary of W_3 .

Fix the detector ball B from P0. Choose a small ball in the connected complement of the actual attaching system. Since any two orientation-preserving

embedded 3-balls in a connected oriented 3-manifold are ambient isotopic, the actual system may first be isotoped off the fixed ball B . Choose the standard three nonseparating spheres in the complement of B . Their complement is connected and their classes form the standard basis of $H_2^{\text{sph}}(\partial W_2)$. Horvat–Jabłonowski’s Theorem 5.3 identifies this system with the actual attaching system up to permutation, 3–3 handle slides and isotopy [10]. Applying their relative uniqueness lemma to $Q = \partial W_2 \setminus \text{Int } B$ makes the isotopies fixed on ∂B . The additional boundary slides become tubings to the inessential sphere ∂B after the ball is restored and are removed by a local isotopy. Thus the 3-handles may be represented by standard spheres without moving the detector.

MWW’s intrinsic reformulation of a 3-handle quotient uses the local module action of $\mathcal{S}_0^2(S^2 \times D^2)$. At $N = 2$, let A_0 denote the essential sphere with one dot and A_1 the undotted sphere. The quotient relations are

$$A_0 = 1, \quad A_1 = 0.$$

The symmetric-monoidal naturality required in C applies because the standard spheres lie outside B . For every element v in the complete source,

$$\ell(vA_0) = \ell(v) \text{ev}(\dot{S}^2) = \ell(v), \quad \ell(vA_1) = \ell(v) \text{ev}(S^2) = 0.$$

These are whole-source equations. Hence the detector annihilates the full relation submodule for each of the three standard spheres and descends through the iterated MWW coequalizer. This proves Theorem 3.4 from P0 and C.

The owner lifts, the 32-step Nielsen ledger and the split-tree Lean lemmas are compatible optional coordinate models, but they are not needed for this relative proof. Candidate-level S is therefore not an independent open input; it remains unavailable only because P0 and the full monoidal form of C are not yet instantiated.

9. FINAL IDENTIFICATIONS

MWW Theorem 3.10 identifies the iterated quotient with the module after the three-handles, and Proposition 3.4 says the final four-handle induces a bidegree-(0, 0) isomorphism [14]. If C and Theorem 3.4 have produced the specified selected class, it remains nonzero in

$$\mathcal{S}_{0,(0,494),0}^2(\Sigma_A^0; \mathbb{Q}).$$

MWW Corollary 3.5 gives $\mathcal{S}_0^2(S^4) \cong \mathbb{Z}$, concentrated in bidegree (0, 0), so the corresponding rational degree-494 summand vanishes. The intrinsic definition transports lasagna fillings under diffeomorphisms and gives grading-preserving equivalences. Finally, Iwaki’s Proposition 2.1 and (2) prove that the genuine Cappell–Shaneson manifold Σ_A^0 is a homotopy sphere [11]. Applying these general results to the proposed compact presentation and quotient requires P0, C and the candidate-level final identifications; the general results do not close those joins.

10. FORMALIZATION BOUNDARY

The Lean 4 development [7] is valuable precisely because it exposes the logical interface. The theorem in `Smooth4PC/T73Conditional.lean` has the form

$$\text{ExternalGeometry} \longrightarrow \text{CSExternalGeometry} \longrightarrow (\text{HS}(X) \wedge \neg \text{Diff}(X, S^4)).$$

It does not construct either input structure.

The finite files check:

- the matrix A , $A - I$, and their determinants;
- explicit integral inverse formulas for relevant lattice maps;
- the determinant of the proposed sphere coordinate matrix;
- the arithmetic value and nonvanishing of the stored cubic constant;
- the quantum degree 494 from its four recorded integer contributions;
- general linear-algebra quotient and descent lemmas.

The geometric interfaces contain, among others, the predicates

$$\begin{array}{ll} \text{IsActualHHO}, & \text{IsActualSphereMap}, \\ \text{IsActualMWWCoequalizer}, & \text{IsActualFourHandle}. \end{array}$$

These are predicate fields of the ambient universe rather than definitions of smooth topology. The present paper does not supply their mathematical instances: P0 and C still state missing candidate-level data, while the S fields follow mathematically only after those hypotheses are instantiated. It would be incorrect to call the Lean development a formalization of four-dimensional differential topology or an unconditional proof.

The audit reports that the printed theorem dependencies use only standard logical quotient principles and classical choice, with no `sorryAx`. This is a statement about kernel checking of the algebraic core. The truth of the geometric inputs is neither established by the Lean kernel nor by the partial constructions in Sections 5–8.

11. STATUS TABLE AND REPRODUCIBILITY LEDGER

The table records the current allocation of every external layer. DISCHARGED means that a public generator or primary theorem and an independently checkable argument are both supplied.

Item	Verdict	Reason
P0	OPEN	Word and braid ledgers do not construct the embedded framed Kirby equivalence.
P1/C	OPEN	Cubic order and the ordinary representable-coefficient reduction are proved, but the actual product bimodule equivalence, its completed lift and simultaneous transport are missing.
P2/E7	PARTIAL	HJ and the relative complete-system lemma give standard attaching spheres disjoint from the detector once the P0 handle pattern and detector ball are instantiated.
P2/E10/S	PARTIAL	MWW's intrinsic local module action and monoidal C imply the complete sphere relations; no independent sphere movie remains, but P0/C are open.
P3/E11	PARTIAL	MWW Proposition 3.4 is general; its candidate-level input depends on P1/P2.
P3/E12	PARTIAL	The standard S^4 support is known, conditional on matching the grading and invariant.
P3/E13	PARTIAL	Iwaki's criterion applies to Σ_A^0 , conditional on P0 identifying the presentation.

The historical availability script reports the large files as absent. The compact generators provide useful partial replacements but not the embedded geometry and functorial maps listed in the table.

12. RELATED WORK

The Cappell–Shaneson literature contains both the original construction [5, 1] and substantial standardness results [2, 9, 12, 11]. Our use of Iwaki is limited to the standard matrix form, the representative table, and the homotopy-sphere criterion. His table entry for (41, 189, 73) is motivation, not a singularity or exoticness theorem.

Manolescu–Neithalath and Manolescu–Walker–Wedrich provide the relevant handle calculus for skein lasagna modules [13, 14]. BPW quantum trace functoriality and the strict BHPW tangle theory provide general comparison tools [4, 3]; the candidate-specific comparison with the public endpoint coordinates remains open.

Ren and Willis use Khovanov-theoretic skein lasagna methods to distinguish other exotic four-manifolds [16]. Their computations are independent of the trace-73 construction and are not used as its comparison theorem.

Horvat–Jabłonowski's recent work gives a useful geometric criterion for recognizing three-handle attaching systems [10]. It reduces the replacement problem to boundary, embeddedness, disjointness and homology-basis checks. The compact owner-lift and Nielsen ledgers establish only the coordinate part of those checks.

13. REPRODUCIBILITY AND REVIEW

The proved finite and conditional layers can be replayed in the following order.

- (1) Generate the compact Kirby, point-push, Hattori and owner-lift ledgers.
- (2) Verify the full 88-dimensional order-three matrix statement and its mutation controls.
- (3) Compile the Lean cubic, transported-pairing, Frobenius and quotient lemmas.
- (4) Audit Hypotheses C and S against the cited BPW/BHPW/MWW functoriality and handle formulas.
- (5) Compile the paper and compare the field-by-field allocation with `T73External.lean`.

The historical large artifacts are unavailable for this replay. Independent review should concentrate on the missing compact product-ribbon isotopy, the proposed BPW cabling comparison in Hypothesis C, and the complete-sphere closure in Hypothesis S.

14. CONCLUSION

The finite calculation and exact Magnus certificate provide a reproducible cubic-order signal for the public braid, and the Lean development verifies the abstract nonvanishing chain. A counterexample conclusion would follow from Theorem 3.6, but P0 and C remain open: the present data do not identify an embedded framed Kirby presentation or a genuine monoidal MWW coefficient functional. The relative sphere theorem makes S a consequence of those inputs rather than an independent hypothesis. The paper therefore establishes a conditional theorem and a precise closure programme, not a disproof of the smooth four-dimensional Poincare conjecture.

APPENDIX A. RETIRED HISTORICAL OBJECTS

The following table records historical artifacts cited by earlier project notes. They are unavailable; the compact partial constructions do not yet replace every load-bearing role. Their digests are retained for provenance.

Priority	Object	SHA-256
P0	cs_presentation.py	B20BDF6FBF2A0E629083E2E62DD1DDF 9172A4E8893311CE122FFC4FEB388DFA 6DF4D7906ADB18BA2D3237D4B49F8985
P0	t73_eps0.erkmo.json	A5B18D8C39FF522AD6498F2369803FCB E6912A64457557469E5C691B4D57ABDB BF4C45ADB05492777C574223D0C06F8A
P0	t73_reduced_billiard.pd.json	4E1C4564ED98F30EAEBOA1AEAAD95ACA B5AD7B226C79B7D5BF3F259E8B56F541 2D9E4760F836D9E04E6E0024CAE03A3D
P0	CUT_OBJECT.json	BB49A74FF5D129756F25450B30F10F7C B2E97EB0981036768B8F1C1D5EF16F8 817EC1A7B951B6B32DBA2F9211FFD725A
P0	NORMAL_FIELD_MOVIE_CERT.json	13AED414447ED33057C99D340910B501 09983CF36874C14ECE0E6A2834A12C24 9D93A6CB320200A906F461EF3131127E
P0	verify_t73_collar_braid.py	A40C80A9A7828128E6F2804808417056 7F3D3618D6A790A9B60EE8085B647AC2 AB742E1BC9C15841F1BEF015034217B5
P1	ACTUAL_PD_CABLE_UNIT_CERT.json	DE1AC479699EC79DE76D4265993DE437 493A3AAA6CABB636F98998644BF3181C EE620E6B085A5F9E1C73CFDD1AD04FC
P1	PRODUCT_NORMAL_CHRISTOFFEL_THXY_MOVIE.json	0682CEC74DA3DBF8AFE70DD19C038E3A0 4D1B627C0343A1C464319704EAFADCA1 27902A2E4E90CF1C283004359B7ADC24
P2	TH1_EXHAUSTIVE_ALL_ROW_GEOMETRY.json	EABF67C0D1AE309F0281297710A283B8 ED5A11D88C8E810837932313F4C53227
P2	TH2_EXHAUSTIVE_ALL_ROW_GEOMETRY.json	
P2	THXY_FULL_MACRO_P3FREE_HJ_CERT.json	

The same values and candidate paths are machine-readable in `audit/geometric_evidence_manifest.json`. These objects are not included. Any proof of P0 and C must either recover the corresponding objects or provide public constructions that genuinely replace their geometric and functorial roles. The historical P2 objects are no longer required by the relative sphere argument.

APPENDIX B. CORRESPONDENCE WITH THE LEAN FIELDS

For readers checking the formal boundary, the principal allocation is:

Lean field	Mathematical obligation
x0, ell0, ell0_x0	P1 comparison and the selected genuine MWW class.
q01, ell1_comp_q01	Complete two-handle beta/psi quotient and descent.
q12, ell2_comp_q12	Actual E7 spheres, MWW hemisphere maps, and complete coequalizer.
transport, fourIso	E11 applied to the same graded lasagna module.
s4DegreeZero	E12 after identifying G with rational $N = 2$ lasagna.
diffeomorphismEquiv	Graded diffeomorphism invariance of that same object.
matrixConditionsToHomotopySphere	E13 for the manifold identified by P0.

These are exactly the candidate-level obligations isolated in Sections 5–8. They have not been discharged. The Lean theorem consumes no stronger statement, but it still requires them as structure fields.

REFERENCES

1. Iain R. Aitchison and J. Hyam Rubinstein, *Fibered knots and involutions on homotopy spheres*, Four-Manifold Theory (Durham, N.H., 1982) (Cameron McA. Gordon and Robion C. Kirby, eds.), Contemporary Mathematics, vol. 35, American Mathematical Society, Providence, RI, 1984, pp. 1–74. MR 780575
2. Selman Akbulut, *Cappell–shaneson homotopy spheres are standard*, Annals of Mathematics **171** (2010), no. 3, 2171–2175.
3. Anna Beliakova, Matthew Hogancamp, Krzysztof Karol Putyra, and Stephan Martin Wehrli, *On the functoriality of $\mathfrak{sl}_2$ tangle homology*, Algebraic & Geometric Topology **23** (2023), no. 3, 1303–1361.
4. Anna Beliakova, Krzysztof Karol Putyra, and Stephan Martin Wehrli, *Quantum link homology via trace functor I*, Inventiones Mathematicae **215** (2019), no. 2, 383–492.
5. Sylvain E. Cappell and Julius L. Shaneson, *Some new four-manifolds*, Annals of Mathematics **104** (1976), no. 1, 61–72.
6. Jacques Darné, *On the andreadakis problem for subgroups of ia_n* , International Mathematics Research Notices **2021** (2021), no. 19, 14720–14742.
7. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, Springer, 2021, pp. 625–635.
8. Michael Hartley Freedman, *The topology of four-dimensional manifolds*, Journal of Differential Geometry **17** (1982), no. 3, 357–453.
9. Robert E. Gompf, *More cappell–shaneson spheres are standard*, Algebraic & Geometric Topology **10** (2010), no. 3, 1665–1681.
10. Eva Horvat and Michał Jabłonowski, *On 4-dimensional 3-handle attachments*, 2025.
11. Kazunori Iwaki, *Infinite families of standard cappell–shaneson spheres*, 2024.
12. Min Hoon Kim and Shohei Yamada, *Ideal classes and cappell–shaneson homotopy 4-spheres*, 2017.
13. Ciprian Manolescu and Ikshu Neithalath, *Skein lasagna modules for 2-handlebodies*, 2020.
14. Ciprian Manolescu, Kevin Walker, and Paul Wedrich, *Skein lasagna modules and handle decompositions*, Advances in Mathematics **425** (2023), 109071.
15. Scott Morrison, Kevin Walker, and Paul Wedrich, *Invariants of 4-manifolds from khovanov–rozansky link homology*, 2019.
16. Qiuyu Ren and Michael Willis, *Khovanov homology and exotic 4-manifolds*, 2024.

DRAFT PREPARED FROM THE PUBLIC TRACE-73 REPOSITORY

Email address: Author information to be supplied before submission