



Started on	Sunday, 2 November 2025, 9:06 PM
State	Finished
Completed on	Sunday, 16 November 2025, 9:02 PM
Time taken	13 days 23 hours
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

**Question 1** | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

**For example:**

Input	Result
3 1 3 5 4	1

**Answer:** (penalty regime: 0 %)

```

1  #include<stdio.h>
2  #include<stdlib.h>
3  int main(){
4      int n,v;
5      scanf("%d",&n);
6      int arr[n];
7      for(int i=0;i<n;i++)
8          scanf("%d",&arr[i]);
9      scanf("%d",&v);
10     int f=0;
11     for(int i=0;i<n;i++){
12         for(int j=i+1;j<n;j++){
13             if(abs(arr[i]-arr[j])==v)
14             {
15                 printf("%d",1);
16                 f=1;
17                 break;
18             }
19         }
20     }
21     if(f==1)
22         break;
23 }
24 if(f==0)
25     printf("%d",0);
26
27 }
28

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)