



CHANDRU G S 2024-CSE ▾

C2

Started on	Thursday, 23 October 2025, 10:22 PM
State	Finished
Completed on	Thursday, 23 October 2025, 10:24 PM
Time taken	2 mins 6 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

**Question 1** | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

**For example:**

Input	Result
3 1 3 5 4	1

**Answer:** (penalty regime: 0 %)

```

1 | #include<stdio.h>
2 | int main(){
3 |     int n,k;
4 |     scanf("%d",&n);
5 |     int arr[n];
6 |     for(int i=0;i<n;i++){
7 |         scanf("%d",&arr[i]);
8 |     }
9 |     scanf("%d",&k);
10 |    int f=0;
11 |    for(int j=n-1;j>=0;j--){
12 |        for(int i=0;i!=j;i++){
13 |            if(arr[j]-arr[i]==k)
14 |            {
15 |                printf("1");
16 |                f=1;
17 |                break;
18 |            }
19 |            if(f==1)
20 |                break;
21 |        }
22 |    }
23 |    if(f==0)
24 |        printf("0");
25 | }
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)