

141. (b) $[H^+] = 1.00 \times 10^{-6}$ mole/litre

$$pH = -\log [1.00 \times 10^{-6}]; pH = 6.$$

142. (a) $[H^+]$ is in moles per litre.

143. (d) As the solution is acidic, $pH < 7$. This is because $[H^+]$ from H_2O ($10^{-7}M$) cannot be neglected in comparison to 10^{-8} .

145. (b) pH of $0.001 M HCl = 10^{-3}M[H^+]$, $pH = 3$.

146. (d) Because it can furnish H^+ ions in solutions.

147. (c) Because it is a strong acid.

$$H^+ = 10^{-1}$$

$$pH = -\log [H^+] = -\log [10^{-1}]; pH = 1.$$

148. (b) Buffer solution is a combination of weak acid and conjugate base. $NaCl$ is a salt and $NaOH$ is the base.

149. (a) $[H^+] = \sqrt{Kc} = \sqrt{10^{-5} \times 0.1} = 10^{-3}$, $pH = 3$.

150. (d) In $\frac{N}{10}NaOH$ have $[OH^-] = 10^{-1}M$ means $pOH = 1$ and then $pH + pOH = 14$

$$pH = 14 - pOH = 13.$$

151. (b) Borate ions are hydrolyzed to develop alkaline nature in solution.

152. (d) Less the pH , more acidic is the solution.

153. (b) Acidic



Explanation (Simple & Word-Friendly):

$pH = 7 \rightarrow$ Neutral (e.g., pure water)

$pH > 7 \rightarrow$ Basic (e.g., NaOH solution)

$pH < 7 \rightarrow$ Acidic (e.g., vinegar, lemon juice)

Since $pH = 6$, it is less than 7, so the solution is slightly acidic.

154. (b) The equal conc. of salt and acid.

155. (c) $pH = -\log K_a + \log \frac{[KCN]}{[HCN]}$

$$pH = -\log[5 \times 10^{-10}] + \log\left(\frac{0.15}{1.5}\right) = 8.302$$

156. (d) $NH_4OH + NH_4Cl$

Explanation (Simple & Word-Friendly):

Type of Buffer Example pH

Acidic buffer Weak acid + its salt $pH < 7$

Basic buffer Weak base + its salt $pH > 7$

(a) $CH_3COOH + CH_3COONa \rightarrow$ weak acid + salt \rightarrow acidic buffer

(b) $HCOOH + HCOOK \rightarrow$ weak acid + salt \rightarrow acidic buffer

(c) $CH_3COONH_4 \rightarrow$ not a buffer

(d) $NH_4OH + NH_4Cl \rightarrow$ weak base + its salt \rightarrow basic buffer $\rightarrow pH > 7$

157. (c) $pH = pK_a + \log \frac{[Salt]}{[Acid]}$ equimolar means

$$\frac{[Salt]}{[Acid]} = 1; pH = 4.74 + 0 = 4.74$$

158. (a) Because of $NaCl$ is a salt of strong acid and strong base. So that it is neutral.

159. (c) When strong acid and strong base are react neutral salt are formed. So that

$NaCl$ is a neutral salt.

160. (b) Addition of 25 mL of 0.02 M HCl



**Explanation:**

pH decreases when the **concentration of H^+ increases**.

Adding **0.02 M HCl** increases the H^+ concentration \rightarrow lowers the pH.

Adding 0.005 M HCl **dilutes the solution slightly** \rightarrow pH increases slightly.

Mg reacts with HCl, but this does not decrease pH significantly in a buffered way.

