


Graph Signal Processing - Graph Laplacian

Prof. Luis Gustavo Nonato


August 23, 2017

Adjacency Matrix

Adjacency Matrix



Adjacency Matrix



$$\mathbf{A} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

The Graph Laplacian

The Graph Laplacian

$$\mathbf{L} = \mathbf{D} - \mathbf{A}$$


\mathbf{A} is the adjacency matrix and \mathbf{D} is a diagonal matrix with

$$d_{ii} = \sum_j a_{ij}$$

The Graph Laplacian

$$\mathbf{L} = \mathbf{D} - \mathbf{A}$$


\mathbf{A} is the adjacency matrix and \mathbf{D} is a diagonal matrix with $d_{ii} = \sum_j a_{ij}$



The Graph Laplacian

$$\mathbf{L} = \mathbf{D} - \mathbf{A}$$

\mathbf{A} is the adjacency matrix and \mathbf{D} is a diagonal matrix with $d_{ii} = \sum_j a_{ij}$




$$\underbrace{\begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ -1 & 4 & -1 & -1 & -1 \\ 0 & -1 & 2 & 0 & -1 \\ 0 & -1 & 0 & 2 & -1 \\ 0 & -1 & -1 & -1 & 3 \end{bmatrix}}_{\mathbf{L}} = \underbrace{\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 3 \end{bmatrix}}_{\mathbf{D}} - \underbrace{\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{bmatrix}}_{\mathbf{A}}$$

The Graph Laplacian

If the graph has weights w_{ij} associated to the edges, then the graph Laplacian can incorporate such weights.

The Graph Laplacian


If the graph has weights w_{ij} associated to the edges, then the graph Laplacian can incorporate such weights.



$$\mathbf{L} = \mathbf{D} - \mathbf{W}$$

The Graph Laplacian

If the graph has weights w_{ij} associated to the edges, then the graph Laplacian can incorporate such weights.




$$\mathbf{L} = \mathbf{D} - \mathbf{W}$$

$$\begin{bmatrix} w_{12} & -w_{12} & 0 & 0 & 0 \\ -w_{21} & \sum_{j \neq 2} w_{2j} & -w_{23} & -w_{24} & -w_{25} \\ 0 & -w_{32} & \sum_{j \neq 3} w_{3j} & 0 & -w_{35} \\ 0 & -w_{42} & 0 & \sum_{j \neq 4} w_{4j} & -w_{45} \\ 0 & -w_{52} & -w_{53} & -w_{54} & \sum_{j \neq 5} w_{5j} \end{bmatrix}$$


Let $G = (V, E)$ be a graph with node set V and edge set E .
We can define a scalar function $f : V \rightarrow \mathbb{R}$ that associates a scalar to each node of the graph.

Let $G = (V, E)$ be a graph with node set V and edge set E .
We can define a scalar function $f : V \rightarrow \mathbb{R}$ that associates a scalar to each node of the graph.




Graph Laplacian

Let $G = (V, E)$ be a graph with node set V and edge set E .
We can define a scalar function $f : V \rightarrow \mathbb{R}$ that associates a scalar to each node of the graph.



Graph Laplacian

Let $G = (V, E)$ be a graph with node set V and edge set E .
We can define a scalar function $f : V \rightarrow \mathbb{R}$ that associates a scalar to each node of the graph.




f satisfies the Graph Laplacian equation if:

$$\mathbf{L}f = 0$$

Graph Laplacian

Let $G = (V, E)$ be a graph with node set V and edge set E .
We can define a scalar function $f : V \rightarrow \mathbb{R}$ that associates a scalar to each node of the graph.



f satisfies the Graph Laplacian equation if:

$$\mathbf{L}\mathbf{f} = \mathbf{0}$$

or equivalently

$$f_i = \frac{1}{l_{ii}} \sum_{j \neq i} f_j$$

(the value in each node is the average of values in neighbor nodes)

What is so interesting about about the Graph Laplacian (GL)?

What is so interesting about about the Graph Laplacian (GL)?

- The GL is a symmetric matrix (real eigenvalues and eigenvectors, the latter forming an orthogonal basis).

What is so interesting about about the Graph Laplacian (GL)?

- The GL is a symmetric matrix (real eigenvalues and eigenvectors, the latter forming an orthogonal basis).
- The GL is positive semidefinite (assuming non-negative edge weights)

What is so interesting about about the Graph Laplacian (GL)?

- The GL is a symmetric matrix (real eigenvalues and eigenvectors, the latter forming an orthogonal basis).
- The GL is positive semidefinite (assuming non-negative edge weights)

$$\mathbf{x}^\top \mathbf{L} \mathbf{x} = \sum_{(i,j) \in E} w_{ij} (x_i - x_j)^2 \geq 0$$

What is so interesting about about the Graph Laplacian (GL)?

- The GL is a symmetric matrix (real eigenvalues and eigenvectors, the latter forming an orthogonal basis).
- The GL is positive semidefinite (assuming non-negative edge weights)

$$\mathbf{x}^\top \mathbf{L} \mathbf{x} = \sum_{(i,j) \in E} w_{ij} (x_i - x_j)^2 \geq 0$$

Eigenvalues are non-negative

What is so interesting about about the Graph Laplacian (GL)?

- The GL is a symmetric matrix (real eigenvalues and eigenvectors, the latter forming an orthogonal basis).
- The GL is positive semidefinite (assuming non-negative edge weights)

$$\mathbf{x}^\top \mathbf{L} \mathbf{x} = \sum_{(i,j) \in E} w_{ij} (x_i - x_j)^2 \geq 0$$

Eigenvalues are non-negative

- The eigenvectors are "nice" functions defined on the graph.

Graph Laplacian: Spectral Properties

Graph Laplacian: Spectral Properties

Property 1

GL has an eigenpair $(\lambda_0 = 0, \mathbf{u}_0 = \mathbf{1} = (1, 1, \dots, 1)^\top)$.

Graph Laplacian: Spectral Properties

Property 1

GL has an eigenpair $(\lambda_0 = 0, \mathbf{u}_0 = \mathbf{1} = (1, 1, \dots, 1)^\top)$.

$$\mathbf{L}\mathbf{u}_0 = \begin{bmatrix} \vdots \\ -w_{i1} & 0 & -w_{i3} & \cdots & \sum_j w_{ij} & \cdots & 0 & -w_{3(n-1)} & 0 \\ \vdots \end{bmatrix} \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} = 0\mathbf{u}_0$$

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Number the vertices in G such that:

- vertices $1, \dots, k$ are in one component
- vertices $k + 1, \dots, n$ are in the other component.

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Number the vertices in G such that:

- vertices $1, \dots, k$ are in one component
- vertices $k + 1, \dots, n$ are in the other component.

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{L}_1 & 0 \\ \hline 0 & \mathbf{L}_2 \end{array} \right]$$

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Number the vertices in G such that:

- vertices $1, \dots, k$ are in one component
- vertices $k + 1, \dots, n$ are in the other component.

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{L}_1 & 0 \\ \hline 0 & \mathbf{L}_2 \end{array} \right]$$

\mathbf{L}_1 and \mathbf{L}_2 are GL by themselves, so both has an eigenvalue zero.

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Number the vertices in G such that:

- vertices $1, \dots, k$ are in one component
- vertices $k + 1, \dots, n$ are in the other component.

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{L}_1 & 0 \\ \hline 0 & \mathbf{L}_2 \end{array} \right]$$

\mathbf{L}_1 and \mathbf{L}_2 are GL by themselves, so both has an eigenvalue zero. Since the spectrum of block diagonal matrix is the union of the spectrum in each block, L will have $\lambda_0 = 0$ with multiplicity 2.

Graph Laplacian: Spectral Properties

Property 2

The multiplicity of the eigenvalue zero is equal to the number of connected components of the graph

Suppose G has 2 connected components.

Number the vertices in G such that:

- vertices $1, \dots, k$ are in one component
- vertices $k + 1, \dots, n$ are in the other component.

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{L}_1 & 0 \\ \hline 0 & \mathbf{L}_2 \end{array} \right]$$

\mathbf{L}_1 and \mathbf{L}_2 are GL by themselves, so both has an eigenvalue zero. Since the spectrum of block diagonal matrix is the union of the spectrum in each block, L will have $\lambda_0 = 0$ with multiplicity 2. The proof can easily be generalized to any number of components.

Graph Laplacian: Spectral Properties

Property 3

Supposing G is connected, the eigenvector associated to the smallest non-zero eigenvalue (Fiedler Vector) provides the best one-dimensional embedding of G .

Graph Laplacian: Spectral Properties

Property 3

Supposing G is connected, the eigenvector associated to the smallest non-zero eigenvalue (Fiedler Vector) provides the best one-dimensional embedding of G .

In mathematical terms, we want to find $f : V \rightarrow \mathbb{R}, f_i = f(v_i)$ for each node v_i in G , so as to minimize

$$\sum_{(i,j) \in E} w_{ij}(f_i - f_j)^2, \quad \mathbf{f}^\top \mathbf{1} = 0$$

Graph Laplacian: Spectral Properties

Property 3

Supposing G is connected, the eigenvector associated to the smallest non-zero eigenvalue (Fiedler Vector) provides the best one-dimensional embedding of G .

In mathematical terms, we want to find $f : V \rightarrow \mathbb{R}, f_i = f(v_i)$ for each node v_i in G , so as to minimize

$$\sum_{(i,j) \in E} w_{ij}(f_i - f_j)^2, \quad \mathbf{f}^\top \mathbf{1} = 0$$

$$\mathbf{f}^\top \mathbf{L} \mathbf{f} = \sum_{(i,j) \in E} w_{ij}(f_i - f_j)^2$$

Graph Laplacian: Spectral Properties

Property 3

Supposing G is connected, the eigenvector associated to the smallest non-zero eigenvalue (Fiedler Vector) provides the best one-dimensional embedding of G .

In mathematical terms, we want to find $f : V \rightarrow \mathbb{R}, f_i = f(v_i)$ for each node v_i in G , so as to minimize

$$\sum_{(i,j) \in E} w_{ij} (f_i - f_j)^2, \quad \mathbf{f}^\top \mathbf{1} = 0$$

$$\mathbf{f}^\top \mathbf{L} \mathbf{f} = \sum_{(i,j) \in E} w_{ij} (f_i - f_j)^2$$

Imposing the additional constraint $\|\mathbf{f}\|^2 = 1$, the Courant-Fiecher theorem ensures that the minimum is reached when \mathbf{f} is the eigenvector associated to the second smallest eigenvalue.

Graph Laplacian: Spectral Properties

Property 3 can be generalized to embed a graph in \mathbb{R}^d .

Graph Laplacian: Spectral Properties

Property 3 can be generalized to embed a graph in \mathbb{R}^d .

$\mathbf{f} : V \rightarrow \mathbb{R}^d, \mathbf{f}(v_i) = (f_1(v_i), f_2(v_i), \dots, f_d(v_i))$ that minimizes

Graph Laplacian: Spectral Properties

Property 3 can be generalized to embed a graph in \mathbb{R}^d .

$\mathbf{f} : V \rightarrow \mathbb{R}^d$, $\mathbf{f}(v_i) = (f_1(v_i), f_2(v_i), \dots, f_d(v_i))$ that minimizes

$$\sum_{(i,j) \in E} w_{ij} \|\mathbf{f}(v_i) - \mathbf{f}(v_j)\|^2, \quad \mathbf{F}^\top \mathbf{1} = 0$$

where $\mathbf{F} = \left[\begin{array}{c|c|c|c} | & | & & | \\ \mathbf{f}_1 & \mathbf{f}_2 & \cdots & \mathbf{f}_d \\ | & | & & | \end{array} \right]$

Graph Laplacian: Spectral Properties

Property 3 can be generalized to embed a graph in \mathbb{R}^d .

$\mathbf{f} : V \rightarrow \mathbb{R}^d$, $\mathbf{f}(v_i) = (f_1(v_i), f_2(v_i), \dots, f_d(v_i))$ that minimizes

$$\sum_{(i,j) \in E} w_{ij} \|\mathbf{f}(v_i) - \mathbf{f}(v_j)\|^2, \quad \mathbf{F}^\top \mathbf{1} = 0$$

where $\mathbf{F} = \begin{bmatrix} | & | & & | \\ \mathbf{f}_1 & \mathbf{f}_2 & \cdots & \mathbf{f}_d \\ | & | & & | \end{bmatrix}$ Imposing $\|\mathbf{f}_i\| = 1$ the solution is given by:

$$\mathbf{F} = \begin{bmatrix} | & | & & | \\ \mathbf{u}_1 & \mathbf{u}_2 & \cdots & \mathbf{u}_d \\ | & | & & | \end{bmatrix}$$

where \mathbf{u}_i are the eigenvectors of \mathbf{L} .

Graph Laplacian: Spectral Properties


- A positive (or negative) *strong nodal domain* of a function f defined on V is a maximal connected subgraph of G where $f(v) > 0$ (or $f(v) < 0$). The number of strong nodal domains of f is $\mathcal{S}(f)$.

Graph Laplacian: Spectral Properties

- A positive (or negative) *strong nodal domain* of a function f defined on V is a maximal connected subgraph of G where $f(v) > 0$ (or $f(v) < 0$). The number of strong nodal domains of f is $\mathcal{S}(f)$.
- A positive (or negative) *weak nodal domain* of f is a maximal connected subgraph of G where $f(v) \geq 0$ (or $f(v) \leq 0$) which contains at least one vertex v such that $f(v) \neq 0$. The number of weak nodal domains of f is denoted by $\mathcal{W}(f)$.

Graph Laplacian: Spectral Properties

- A positive (or negative) *strong nodal domain* of a function f defined on V is a maximal connected subgraph of G where $f(v) > 0$ (or $f(v) < 0$). The number of strong nodal domains of f is $\mathcal{S}(f)$.
- A positive (or negative) *weak nodal domain* of f is a maximal connected subgraph of G where $f(v) \geq 0$ (or $f(v) \leq 0$) which contains at least one vertex v such that $f(v) \neq 0$. The number of weak nodal domains of f is denoted by $\mathcal{W}(f)$.



$$\begin{aligned}\mathcal{S}(f) &= \{\{1\}, \{2\}, \{4\}, \{5\}\} \\ \mathcal{W}(f) &= \{\{1, 2, 3\}, \{3, 4, 5\}\}\end{aligned}$$

Graph Laplacian: Spectral Properties

Property 4: Discrete Courant's Nodal Theorem

Let G be a connected graph with n vertices. Any Graph Laplacian eigenvector \mathbf{u}_k with corresponding eigenvalue λ_k with multiplicity r has at most $k + 1$ weak nodal domains and $k + r$ strong nodal domains, i.e.,

$$\mathcal{W}(\mathbf{u}_k) \leq k + 1, \quad \mathcal{S}(\mathbf{u}_k) \leq k + r$$

where $k \in [0, n - 1]$

Graph Laplacian: Spectral Properties

Property 4: Discrete Courant's Nodal Theorem





Let G be a connected graph with n vertices. Any Graph Laplacian eigenvector \mathbf{u}_k with corresponding eigenvalue λ_k with multiplicity r has at most $k + 1$ weak nodal domains and $k + r$ strong nodal domains, i.e.,

$$\mathcal{W}(\mathbf{u}_k) \leq k + 1, \quad \mathcal{S}(\mathbf{u}_k) \leq k + r$$

where $k \in [0, n - 1]$

This theorem was proved by Davies, Gladwell, Leydold, Stadler in 2001 and it is the discrete version of the Courant's Nodal Theorem for the Laplace operator on manifolds.

Graph Laplacian: Spectral Properties



Graph Laplacian: Normalized Version

There are other versions of Graph Laplacian:

Graph Laplacian: Normalized Version

There are other versions of Graph Laplacian:

- Normalized Graph Laplacian:

$$\mathcal{L} = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{1/2}$$

Graph Laplacian: Normalized Version

There are other versions of Graph Laplacian:

- Normalized Graph Laplacian:

$$\mathcal{L} = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{1/2}$$

$0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1} \leq 2$, with $\lambda_{n-1} = 2$ if and only if G is bipartite.

- In the case of normalized graph Laplacian, there are a multitude of results giving upper and lower bounds for the eigenvalues, specially related to the diameter of G .

Graph Laplacian: Normalized Version

There are other versions of Graph Laplacian:

- Normalized Graph Laplacian:

$$\mathcal{L} = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{1/2}$$

$0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1} \leq 2$, with $\lambda_{n-1} = 2$ if and only if G is bipartite.

- In the case of normalized graph Laplacian, there are a multitude of results giving upper and lower bounds for the eigenvalues, specially related to the diameter of G .
- Normalized Graph Laplacian is also closely related with random walks on graphs.

$$\mathbf{P} = \mathbf{D}^{-1/2} (\mathbf{I} - \mathcal{L}) \mathbf{D}^{1/2}$$