


Correction

Information theory broadens the spectrum of molecular ecology and evolution

W.B. Sherwin,* A. Chao, L. Jost, and P.E. Smouse

*Correspondence: w.sherwin@unsw.edu.au (W.B. Sherwin).
DOI of original article: <https://doi.org/10.1016/j.tree.2017.09.012>
(Trends in Ecology and Evolution 32:12, p:948–963, 2017)

In Sherwin *et al.* [1], several corrections are required, having been noticed when assisting other researchers to use the methods. In the main text, in Figure III in Box 2, nine subscripts were incorrect (reversing localities 1 and 2). The correct figure is shown below.



Trends in Ecology & Evolution

Figure III. Data for mutual information between alleles and location.

On pages 5 and 6 of the supplement, it should be explained that $\overline{1D_\alpha} = e^{\overline{1H_\alpha}}$, that is, the averaging happens before conversion to the D scale (see Equations 10 and 11 of Jost [2]). Similarly, on page 8 of the supplement, $\overline{2D_\alpha} = 1/(1 - \overline{2H_\alpha})$.

On Page 6 of the supplement, the equation from Dewar *et al.* [3] is incorrectly transcribed; the correct equation is:

$$I_{SNP} = \frac{0.25 + (1 - 1/N)m^2}{1 + 4(N-1)\left\{\lambda + m(1-4\lambda)\left[(1-m)\left(1 + \frac{1}{8N\lambda}\right) + \frac{(m-0.5)}{N}\right]\right\}}.$$

The authors and publisher apologise for any confusion.



References

1. Sherwin, W.B. *et al.* (2017) Information theory broadens the spectrum of molecular ecology and evolution. *Trends Ecol. Evol.* 32, 948–963
2. Jost, L. (2007) Partitioning diversity into independent alpha and beta components. *Ecology* 88, 2427–2439
3. Dewar, R.C. *et al.* (2011) Predictions of single-nucleotide polymorphism differentiation between two populations in terms of mutual information. *Mol. Ecol.* 20, 3156–3166