

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
26 October 2017 (26.10.2017)

(10) International Publication Number  
**WO 2017/182561 A1**

## (51) International Patent Classification:

*G01N 33/68* (2006.01)      *A61K 39/395* (2006.01)  
*G01N 33/574* (2006.01)

KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

## (21) International Application Number:

PCT/EP2017/059377

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

## (22) International Filing Date:

20 April 2017 (20.04.2017)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

16166476.8      21 April 2016 (21.04.2016)      EP

(71) Applicant: SPHINGOTEC THERAPEUTICS GMBH [DE/DE]; Neuendorfstrasse 15A, 16761 Hennigsdorf (DE).

(72) Inventor: BERGMANN, Dr. Andreas; Am Rosenanger 78, 13465 Berlin (DE).


(74) Agent: BOEHMERT & BOEHMERT; Dr. Ute Kilger, Hollerallee 32, 28209 Bremen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,

## Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))
- with sequence listing part of description (Rule 5.2(a))

## (54) Title: METHODS FOR DETERMINING DPP3 AND THERAPEUTIC METHODS



(57) Abstract: The present invention is directed to methods for determining active DPP3 in a bodily fluid sample, an assay or kit for determining active DPP3 in a bodily fluid sample, a method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes and methods of treating or preventing said disease.

## Methods for determining DPP3 and therapeutic methods

The present invention is directed to methods for determining active DPP3 in a bodily fluid sample, an assay or kit for determining active DPP3 in a bodily fluid sample, a method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes and methods of treating said disease.

### State of the Art

Dipeptidyl peptidase 3 – also known as Dipeptidyl aminopeptidase III, Dipeptidyl arylamidase III, Dipeptidyl peptidase III, Enkephalinase B or red cell angiotensinase; short name: DPP3, DPPIII – is a metallopeptidase that removes dipeptides from physiologically active peptides, such as enkephalins and angiotensins.

DPP3 was first identified and its activity measured in extracts of purified bovine anterior pituitary by Ellis & Nuenke 1967. The enzyme, which is listed as EC 3.4.14.4, has a molecular mass of about 83 kDa and is highly conserved in prokaryotes and eukaryotes (Prajapati & Chauhan 2011). The amino acid sequence of the human variant is depicted in SEQ ID NO 1. Dipeptidyl peptidase III is a mainly cytosolic peptidase which is ubiquitously expressed. Despite lacking a signal sequence, a few studies reported membranous activity (Lee & Snyder 1982).

DPP3 is a zinc-depending exo-peptidase belonging to the peptidase family M49. It has a broad substrate specificity for oligopeptides from three/ four to ten amino acids of various compositions and is also capable of cleaving after proline. DPP3 is known to hydrolyze dipeptides from the N-terminus of its substrates, including angiotensin II, III and IV; Leu- and Met-enkephalin; endomorphin 1 and 2. The metallopeptidase DPP3 has its activity optimum at pH 8.0-9.0 and can be activated by addition of divalent metal ions, such as Co<sup>2+</sup> and Mg<sup>2+</sup>. Structural analysis of DPP3 revealed the catalytic motifs HELLGH (hDPP3 450-455) and EECRAE (hDPP3 507-512), as well as following amino acids, that are important for substrate binding and hydrolysis: Glu316, Tyr, 318, Asp366, Asn391, Asn394, His568, Arg572, Arg577, Lys666 and Arg669 (Prajapati & Chauhan 2011; Kumar et al. 2016; numbering refers to the sequence of human DPP3, see SEQ ID No. 1). Considering all known amino

acids or sequence regions that are involved in substrate binding and hydrolysis, the active site of human DPP3 can be defined as the area between amino acids 316 and 669.

The activity of DPP3 can be inhibited unspecifically by different general protease inhibitors  
5 (e.g. PMSF, TPCK), sulphydryl reagents (e.g. pHMB, DTNB) and metal chelators (EDTA, o-phenanthroline; Abramić et al. 2000).

DPP3 activity can be inhibited specifically by different kinds of compounds: an endogenous DPP3 inhibitor is the peptide spinorphin. Several synthetic derivatives of spinorphin, e.g.  
10 tynorphin, have been produced and shown to inhibit DPP3 activity to varying extents (Yamamoto et al. 2000). Other published peptide inhibitors of DPP3 are propioxatin A and B (US4804676) and propioxatin A analogues (Inaoka et al. 1988).

DPP3 can also be inhibited by small molecules such as fluostatins and benzimidazol  
15 derivatives. Fluostatins A and B are antibiotics produced in *Streptomyces sp.* TA-3391 that are non-toxic and strongly inhibit DPP3 activity. So far 20 different derivatives of benzimidazol have been synthesized and published (Agić et al. 2007; Rastija et al. 2015), of which the two compounds 1' and 4' show the strongest inhibitory effect (Agić et al. 2007).  
For a complete list of DPP3 inhibitors see table 2.

20 The exact biological function of DPP3 in cellular physiology is not understood, but recent findings indicate its role not only in protein metabolism but also in blood pressure regulation, pain modulation and inflammatory processes (Prajapati & Chauhan 2011).

25 DPP3 has been shown to be a promising biomarker in several publications, which all refer to intracellular DPP3. It has been shown that DPP3 activity is elevated in homogenates of ovarian and endometrial tumors. DPP3 activity even increases with the severity/ malignancy of said tumors (Šimaga et al. 1998 and 2003). Immune histology and western blot analysis of glioblastoma cell lines also revealed elevated DPP3 levels (Singh et al. 2014).

30 Intracellular or membranous DPP3 was also proposed to be a potential arteriorisk marker (US2011008805) and marker for rheumatoid arthrosis (US2006177886). The patent application WO2005106486 claims: DPP3 expression and activity as diagnostic marker and

DPP3 as therapeutic target in all kinds of diseases, due to ubiquitous expression of DPP3 in or at surface of cell. EP1498480 mentions the potential diagnostic and therapeutic use of hydrolytic enzymes, including DPP3.

- 5 Not only has DPP3 been proposed as potential biomarker but also as potential therapeutic target due to its ability to cleave several bioactive peptides. Overexpression of DPP3 protects neuroblastoma cells from oxidative stress (Liu et al. 2007). Influenca A virus changes host DPP3 levels for own replication (cell culture studies, Meliopoulos et al. 2012). Enkephalin and/ or angiotensin degrading enzymes in general, including DPP3, have a therapeutic potential as targets for treatment of pain, cardiovascular diseases (CVD) and cancer, and the corresponding inhibitors as potential treatments of pain, mental illnesses and CVD (Khaket et al. 2012, Patel et al. 1993, Igic et al. 2007).
- 10

15 Although DPP3 is known as an intracellular protein, DPP3 activity was detected in some bodily fluids: retroplacental serum (Shimamori et al. 1986), seminal plasma (Vanha-Perttula et al. 1988) and CSF (Aoyagi et al. 1993). In CSF there were elevated DPP3 activity levels measured in patients suffering from Alzheimer's disease (AD, Aoyagi et al. 1993). Wattiaux et al. (2007) proposed the release of intracellular DPP3 as a marker for dead and/ or dying cells in a cell culture system. It has also been proposed that the release of DPP3 from necrotic 20 cells influences immune response in a mouse model (Gamrekelashvili et al. 2013).

The object of the present invention is the provision of a method for determining active DPP3 in a sample of bodily fluid specifically, i.e. that active DPP3 is determined but not any other aminopeptidase than DPP3.

25

It is an object of the invention to provide respective assays and kits.

Another object of the invention is the provision of a method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes and methods of treating 30 said disease.

Subject of the present invention is a method for determining active DPP3 in a bodily fluid sample of a subject comprising the steps

- Contacting said sample with a capture-binder that binds specifically to full-length DPP3,
- Separating DPP3 bound to said capture-binder
- Adding a substrate of DPP3 to said separated DPP3
- Quantifying DPP3 activity by measuring the conversion of the substrate of DPP3.

10 In a specific embodiment of the present invention said method is an enzyme capture assay (ECA, see e.g. US5612186A, US5601986A).

All definitions and specific embodiments specified in the description shall apply for all aspects and objects of the invention. It shall be understood that definitions and specific 15 embodiments that are outlined in detail for one aspect or object of the invention shall be definitions and specific embodiments for other aspects and objects of the invention too, e.g. the definition of capture binder or specific capture is applicable to all embodiments of the invention: e.g. methods of determining DPP3, assays and kits, diagnostic methods, therapeutic methods. Such definitions may not be repeated throughout the description.

20 Binding specifically to full-length DPP3 means that said capture-binder does not bind to any other protein than DPP3. Binding specifically to full-length DPP3 means that said capture-binder does not bind to any other aminopeptidase than DPP3.

25 A binder that binds to full-length DPP3 is binder that binds to the protein of SEQ ID No. 1. A binder that binds to full-length DPP3 is binder that binds to SEQ ID No. 1.

In a specific embodiment said capture-binder inhibits DPP3 activity in a liquid phase assay less than 50 %, preferably less than 40 %, preferably less than 30 %. A liquid phase assay is 30 an assay where the cleavage of substrates by DPP3 is taking place in the liquid phase.

Inhibition of DPP3 activity in a liquid phase assay by a binder may be determined as followed according to the present invention: Possible DPP3 capture-binders are incubated with

recombinant or purified native DPP3 and specific DPP3 substrates in a liquid phase assay. Preferably, as capture-binder for the ECA is chosen the one with the least inhibitory ability. The capture-binder should inhibit DPP3 activity less than 50 %, preferably less than 40 %, preferably less than 30 %. The specific liquid phase DPP3 activity assay to determine the 5 inhibitory ability of possible capture-binders is described in detail in example 1 and comprises the following steps:

- Incubation of 25 ng/ ml recombinant GST-hDPP3 with 5 µg/ ml of the respective capture-binder and buffer control in 50 mM Tris-HCl, pH 7,5 and 100 µM ZnCl<sub>2</sub> for 1 10 hour at room temperature.
  - Addition of the fluorogenic substrate Arg-Arg-βNA (20 µl, 2 mM).
  - Incubation at 37 °C and monitoring the generation of free βNA in a Twinkle LB 970 microplate fluorometer (Berthold Technologies GmbH) over 1 hour. Fluorescence of βNA is detected by exciting at 340 nm and measuring emission at 410 nm.
  - Slopes (in RFU/ min) of increasing fluorescence of the different samples are calculated. The slope of GST-hDPP3 with buffer control is appointed as 100 % activity. The inhibitory ability of a possible capture-binder is defined as the decrease 15 of GST-hDPP3 activity by incubation with said capture-binder in percent.
- 20 In contrast thereto, a solid phase assay is an assays where the respective binding events take place at the solid phase (see example 4 and 5).

In order to be clear, the method for determining active DPP3 may be conducted as liquid 25 phase assay and as solid phase assay. Inhibition of DPP3 activity may be determined in a liquid phase assay nevertheless according to the above-described procedure.

Thus, a solid phase assay is an embodiment of the present invention. Contacting said sample with a capture-binder that binds specifically to full-length DPP3 may occur in liquid phase (liquid phase capture assay) and thereafter the separation step may comprise immobilization 30 of said capture-binder-DPP3-complex. Alternatively the capture-binder may be immobilized on a surface and the binding event – capture-binder to DPP3 - takes place on the solid phase (solid phase capture assay).

In one specific embodiment, to prevent total inhibition of DPP3 and to inhibit DPP3 activity less than 50 %, preferably less than 40 %, preferably less than 30 % in the afore explained liquid phase assay, the capture-binder should, preferably, not bind the DPP3 in the region at or to the region of amino acids 316 – 669 of SEQ ID No. 1. The region of amino acids 316 – 5 669 of SEQ ID No. 1 includes the active center of DPP3 and the area of substrate binding (Prajapati & Chauhan 2011; Kumar et al. 2016).

DPP3 activity can be measured by detection of cleavage products of DPP3 specific substrates.

10 Known peptide hormone substrates include angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH (Adrenocorticotrophic hormone) and MSH (melanocyte-stimulating hormone; Abramić et al. 2000, Baršun et al. 2007, Dhanda et al. 2008). The cleavage of mentioned peptide hormones as well as other untagged oligopeptides (e.g. Ala-Ala-Ala-Ala, Dhanda et al. 2008) can be 15 monitored by detection of the respective cleavage products. Detection methods include, but are not limited to, HPLC analysis (e.g. Lee & Snyder 1982), mass spectrometry (e.g. Abramić et al. 2000), H<sub>1</sub>-NMR analysis (e.g. Vandenberg et al. 1985), capillary zone electrophoresis (CE; e.g. Baršun et al. 2007), thin layer chromatography (e.g. Dhanda et al. 2008) or reversed phase chromatography (e.g. Mazocco et al. 2006).

20 Detection of fluorescence due to hydrolysis of fluorogenic substrates by DPP3 is a standard procedure to monitor DPP3 activity. Those substrates are specific di- or tripeptides (Arg-Arg, Ala-Ala, Ala-Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe) coupled to a fluorophore. Fluorophores include but are not 25 limited to  $\beta$ -naphthylamide (2-naphthylamide,  $\beta$ NA, 2NA), 4-methoxy- $\beta$ -naphthylamide (4-methoxy-2-naphthylamide) and 7-amido-4-methylcoumarin (AMC, MCA; Abramić et al. 2000, Ohkubo et al. 1999). Cleavage of these fluorogenic substrates leads to the release of fluorescent  $\beta$ -naphthylamine or 7-amino-4-methylcoumarin respectively. In a liquid phase assay or an ECA substrate and DPP3 are incubated in for example a 96 well plate format and 30 fluorescence is measured using a fluorescence detector (Ellis & Nuenke 1967). Additionally, DPP3 carrying samples can be immobilized and divided on a gel by electrophoresis, gels stained with fluorogenic substrate (e.g. Arg-Arg- $\beta$ NA) and Fast Garnet GBC and fluorescent protein bands detected by a fluorescence reader (Ohkubo et al. 1999).

The same peptides (Arg-Arg, Ala-Ala, Ala-Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe) can be coupled to chromophores, such as p-nitroanilide diacetate. Detection of color change due to hydrolysis of 5 chromogenic substrates can be used to monitor DPP3 activity.

Another option for the detection of DPP3 activity is a Protease-Glo<sup>TM</sup> Assay (commercially available at Promega). In this embodiment of said method DPP3 specific di- or tripeptides (Arg-Arg, Ala-Ala, Ala-Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-10 Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe) are coupled to aminoluciferin. Upon cleavage by DPP3, aminoluciferin is released and serves as a substrate for a coupled luciferase reaction that emits detectable luminescence.

In a preferred embodiment DPP3 activity is measured by addition of the fluorogenic substrate 15 Arg-Arg-βNA and monitoring fluorescence in real time.

In a specific embodiment of said method for determining inhibitory effect of DPP3 binders and/ or DPP3 activity in a bodily fluid sample of a subject said binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

An antibody according to the present invention is a protein including one or more polypeptides substantially encoded by immunoglobulin genes that specifically bind an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha (IgA), gamma (IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, IgG<sub>4</sub>), delta (IgD), epsilon (IgE) and mu (IgM) constant region 20 genes, as well as the myriad immunoglobulin variable region genes. Full-length immunoglobulin light chains are generally about 25 kD or 214 amino acids in length. Full-length immunoglobulin heavy chains are generally about 50 kD or 446 amino acid in length. Light chains are encoded by a variable region gene at the NH<sub>2</sub>-terminus (about 110 amino 25 acids in length) and a kappa or lambda constant region gene at the COOH-terminus. Heavy chains are similarly encoded by a variable region gene (about 116 amino acids in length) and one of the other constant region genes.

The basic structural unit of an antibody is generally a tetramer that consists of two identical pairs of immunoglobulin chains, each pair having one light and one heavy chain. In each pair,

the light and heavy chain variable regions bind to an antigen, and the constant regions mediate effector functions. Immunoglobulins also exist in a variety of other forms including, for example, Fv, Fab, and (Fab')<sub>2</sub>, as well as bifunctional hybrid antibodies and single chains (e.g., Lanzavecchia et al. 1987; Huston et al. 1988; Bird et al. 1988; Hood et al. 1984; Hunkapiller & Hood, 1986). An immunoglobulin light or heavy chain variable region includes a framework region interrupted by three hypervariable regions, also called complementarity determining regions (CDR's) (see, Kabat et al. 1983). As noted above, the CDRs are primarily responsible for binding to an epitope of an antigen. An immune complex is an antibody, such as a monoclonal antibody, chimeric antibody, humanized antibody or human antibody, or functional antibody fragment, specifically bound to the antigen.

Chimeric antibodies are antibodies whose light and heavy chain genes have been constructed, typically by genetic engineering, from immunoglobulin variable and constant region genes belonging to different species. For example, the variable segments of the genes from a mouse monoclonal antibody can be joined to human constant segments, such as kappa and gamma 1 or gamma 3. In one example, a therapeutic chimeric antibody is thus a hybrid protein composed of the variable or antigen-binding domain from a mouse antibody and the constant or effector domain from a human antibody, although other mammalian species can be used, or the variable region can be produced by molecular techniques. Methods of making chimeric antibodies are well known in the art, e.g., see U.S. Patent No. 5,807,715. A "humanized" immunoglobulin is an immunoglobulin including a human framework region and one or more CDRs from a non-human (such as a mouse, rat, or synthetic) immunoglobulin. The non-human immunoglobulin providing the CDRs is termed a "donor" and the human immunoglobulin providing the framework is termed an "acceptor." In one embodiment, all the CDRs are from the donor immunoglobulin in a humanized immunoglobulin. Constant regions need not be present, but if they are, they must be substantially identical to human immunoglobulin constant regions, i.e., at least about 85-90%, such as about 95% or more identical. Hence, all parts of a humanized immunoglobulin, except possibly the CDRs, are substantially identical to corresponding parts of natural human immunoglobulin sequences. A "humanized antibody" is an antibody comprising a humanized light chain and a humanized heavy chain immunoglobulin. A humanized antibody binds to the same antigen as the donor antibody that provides the CDRs. The acceptor framework of a humanized immunoglobulin or antibody may have a limited number of substitutions by amino acids taken from the donor

framework. Humanized or other monoclonal antibodies can have additional conservative amino acid substitutions which have substantially no effect on antigen binding or other immunoglobulin functions. Exemplary conservative substitutions are those such as gly, ala; val, ile, leu; asp, glu; asn, gln; ser, thr; lys, arg; and phe, tyr. Humanized immunoglobulins  
5 can be constructed by means of genetic engineering (*e.g.*, see U.S. Patent No. 5,585,089). A human antibody is an antibody wherein the light and heavy chain genes are of human origin. Human antibodies can be generated using methods known in the art. Human antibodies can be produced by immortalizing a human B cell secreting the antibody of interest. Immortalization can be accomplished, for example, by EBV infection or by fusing a human B cell with a  
10 myeloma or hybridoma cell to produce a trioma cell. Human antibodies can also be produced by phage display methods (see, *e.g.*, PCT Publication No. WO91/17271; PCT Publication No. WO92/001047; PCT Publication No. WO92/20791, which are herein incorporated by reference), or selected from a human combinatorial monoclonal antibody library (see the Morphosys website). Human antibodies can also be prepared by using transgenic animals  
15 carrying a human immunoglobulin gene (for example, see PCT Publication No. WO93/12227; PCT Publication No. WO91/10741, which are herein incorporated by reference).

Thus, the DPP3 antibody may have the formats known in the art. Examples are human  
20 antibodies, monoclonal antibodies, humanized antibodies, chimeric antibodies, CDR-grafted antibodies. In a preferred embodiment antibodies according to the present invention are recombinantly produced antibodies as *e.g.* IgG, a typical full-length immunoglobulin, or antibody fragments containing at least the F-variable domain of heavy and/or light chain as *e.g.* chemically coupled antibodies (fragment antigen binding) including but not limited to  
25 Fab-fragments including Fab minibodies, single chain Fab antibody, monovalent Fab antibody with epitope tags, *e.g.* Fab-V5Sx2; bivalent Fab (mini-antibody) dimerized with the CH3 domain; bivalent Fab or multivalent Fab, *e.g.* formed via multimerization with the aid of a heterologous domain, *e.g.* via dimerization of dHLX domains, *e.g.* Fab-dHLX-FSx2; F(ab')2-fragments, scFv-fragments, multimerized multivalent or/and multispecific scFv-  
30 fragments, bivalent and/or bispecific diabodies, BITE® (bispecific T-cell engager), trifunctional antibodies, polyvalent antibodies, *e.g.* from a different class than G; single-domain antibodies, *e.g.* nanobodies derived from camelid or fish immunoglobulins and numerous others.

In addition to anti-DPP3 antibodies other biopolymer scaffolds are well known in the art to complex a target molecule and have been used for the generation of highly target specific biopolymers. Examples are aptamers, spiegelmers, anticalins and conotoxins.

5 Non-Ig scaffolds may be protein scaffolds and may be used as antibody mimics as they are capable to bind to ligands or antigens. Non-Ig scaffolds may be selected from the group comprising tetranectin-based non-Ig scaffolds (e.g. described in US 2010/0028995), fibronectin scaffolds (e.g. described in EP 1266 025; lipocalin-based scaffolds ((e.g. described in WO 2011/154420); ubiquitin scaffolds (e.g. described in WO 2011/073214), transferring scaffolds (e.g. described in US 2004/0023334), protein A scaffolds (e.g. described in EP 2231860), ankyrin repeat based scaffolds (e.g. described in WO 2010/060748), microprotein (preferably microproteins forming a cystine knot) scaffolds (e.g. described in EP 2314308), Fyn SH3 domain based scaffolds (e.g. described in WO 2011/023685) EGFR-A-domain based scaffolds (e.g. described in WO 2005/040229) and Kunitz domain based scaffolds (e.g. described in EP 1941867). Non-Ig scaffolds may be peptide or oligonucleotide aptamers. Aptamers are usually created by selecting them from a large random sequence pool and are either short strands of oligonucleotides (DNA, RNA or XNA; Xu et al. 2010, Deng et al. 10 2014) or short variable peptide domains attached to a protein scaffold (Li et al. 2011).

15 20 In one embodiment of the invention anti-DPP3 antibodies according to the present invention may be produced as follows:

Mice are immunized with either recombinant DPP3 (e.g. GST-hDPP3 from USBio, Salem, USA), peptides comprising parts of the DPP3 amino acid sequence, e.g. conjugated to BSA or native purified DPP3 (e.g. from human erythrocytes, Abramić et al. 1988).

25 30 Balb/c mice were intraperitoneally (i.p.) injected with 100 µg recombinant GST-hDPP3, native purified hDPP3 or DPP3-peptide-BSA-conjugates at day 0 (emulsified in TiterMax Gold Adjuvant), 100 µg at day 14 (emulsified in complete Freund's adjuvant) and 50 µg at day 21 and 28 (in incomplete Freund's adjuvant). At day 49 the animal received an intravenous (i.v.) injection of 50 µg GST-hDPP3, native purified hDPP3 or DPP3-peptide-

BSA-conjugates dissolved in saline. Three days later the mice were sacrificed and the immune cell fusion was performed.

Splenocytes from the immunized mice and cells of the myeloma cell line SP2/0 were fused with 1 ml 50% polyethylene glycol for 30 s at 37°C. After washing, the cells were seeded in 96-well cell culture plates. Hybrid clones were selected by growing in HAT medium [RPMI 1640 culture medium supplemented with 20% fetal calf serum and HAT-Supplement]. After one week, the HAT medium was replaced with HT Medium for three passages followed by returning to the normal cell culture medium.

The cell culture supernatants were primarily screened for recombinant DPP3 binding IgG antibodies two weeks after fusion. Therefore recombinant GST-tagged DPP3 (USBiologicals, Salem, USA) was immobilized in 96-well plates (100 ng/ well) and incubated with 50 µl cell culture supernatant per well for 2 hours at room temperature. After washing of the plate, 50 µl / well POD-rabbit anti mouse IgG was added and incubated for 1 h at RT. After a next washing step, 50 µl of a chromogen solution (3,7 mM o-phenylenediamin in citrate/ hydrogen phosphate buffer, 0.012% H<sub>2</sub>O<sub>2</sub>) were added to each well, incubated for 15 minutes at RT and the chromogenic reaction stopped by the addition of 50 µl 4N sulfuric acid. Absorption was detected at 490 nm.

The positive tested microcultures were transferred into 24-well plates for propagation. After retesting the selected cultures were cloned and recloned using the limiting-dilution technique and the isotypes were determined.

Antibodies raised against GST-tagged human DPP3 or DPP3-peptides were produced via standard antibody production methods (Marx et al., 1997) and purified via Protein A. The antibody purities were ≥ 90% based on SDS gel electrophoresis analysis.

Antibodies may be produced by means of phage display according to the following procedure:

The human naive antibody gene libraries HAL7/8 were used for the isolation of recombinant single chain F-Variable domains (scFv) against DPP3 peptide. The antibody gene libraries

were screened with a panning strategy comprising the use of peptides containing a biotin tag linked via two different spacers to the DPP3 peptide sequence. A mix of panning rounds using non-specifically bound antigen and streptavidin bound antigen were used to minimize background of non-specific binders. The eluted phages from the third round of panning have  
5 been used for the generation of monoclonal scFv expressing E.coli strains. Supernatant from the cultivation of these clonal strains has been directly used for an antigen ELISA testing (see also Hust et al. 2011; Schütte et al. 2009)

Humanization of murine antibodies may be conducted according to the following procedure:  
10 For humanization of an antibody of murine origin the antibody sequence is analyzed for the structural interaction of framework regions (FR) with the complementary determining regions (CDR) and the antigen. Based on structural modeling an appropriate FR of human origin is selected and the murine CDR sequences are transplanted into the human FR. Variations in the amino acid sequence of the CDRs or FRs may be introduced to regain structural interactions,  
15 which were abolished by the species switch for the FR sequences. This recovery of structural interactions may be achieved by random approach using phage display libraries or via directed approach guided by molecular modeling. (Almagro & Fransson 2008).

In an alternative embodiment the DPP3 antibody format is selected from the group  
20 comprising Fv fragment, scFv fragment, Fab fragment, scFab fragment, F(ab)<sub>2</sub> fragment and scFv-Fc Fusion protein. In another preferred embodiment the antibody format is selected from the group comprising scFab fragment, Fab fragment, scFv fragment and bioavailability optimized conjugates thereof, such as PEGylated fragments.

25 In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said binder is an antibody.

In one embodiment, a capture or binding assay may be performed to detect and/or quantitate the DPP3. A binder reactive with DPP3 protein, but which does not interfere with peptidase activity by more than 50 %, preferably less than 40 %, preferably less than 30 % in a liquid phase assay, may be immobilized upon a solid phase. In one embodiment to prevent inhibition of DPP3 the capture-binder should preferably not bind DPP3 in the region of amino acids 316 – 669 of SEQ ID No. 1. The region of amino acids 316 – 669 of SEQ ID No. 1  
30

includes the active center of DPP3 and the area of substrate binding (Prajapati & Chauhan 2011; Kumar et al. 2016).

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said binder may be selected from the group of antibody, antibody fragments, non-Ig scaffold or aptamers.

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said capture binder reactive with DPP3 is immobilized on a solid phase.

The test sample is passed over the immobile binder, and DPP3, if present, binds to the binder and is itself immobilized for detection. A substrate may then be added, and the reaction product may be detected to indicate the presence or amount of DPP3 in the test sample. For the purposes of the present description, the term "solid phase" may be used to include any material or vessel in which or on which the assay may be performed and includes, but is not limited to: porous materials, nonporous materials, test tubes, wells, slides, agarose resins (e.g. Sepharose from GE Healthcare Life Sciences), magnetic particals (e.g. Dynabeads<sup>TM</sup> or Pierce<sup>TM</sup> magnetic beads from Thermo Fisher Scientific), etc.

Binders of protein or peptide origin (e.g. antibody, antibody fragments, non-Ig scaffold) are immobilized onto the solid phase by methods comprising: physical adsorption (e.g. by electrostatic interaction or hydrophobic interaction), bioaffinity immobilization (e.g. avidin-biotin, protein A/ G/ L, His-tag and Ni<sup>2+</sup>-NTA, GST-tag and glutathione, DNA hybridization, aptamers), covalent bond (e.g. amine and N-hydroxysuccinimide) or a combination of said immobilization methods (Kim & Herr 2013). Binders of oligonucleotide origin (e.g. aptamers) may be immobilized onto the solid phase by utilization of the (strept)avidin-biotin system (Müller et al. 2012, Deng et al. 2014).

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3. That separation step can be any other step that separates DPP3 bound to said capture-binder from the ingredients of said bodily fluid sample.

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said DPP3 substrate conversion by immobilized DPP3 is measured (detected) by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, 5 luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

10 In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, 15 dynorphin, proctolin, ACTH and MSH, or di- and tri-peptides coupled to a fluorophore, a chromophore or aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay). Di- or tripeptides that are cleaved by DPP3 include, but are not limited to, Arg-Arg, Ala-Ala, Ala-Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe. Fluorophores include but are not limited to  $\beta$ -naphtylamide (2-naphtylamide,  $\beta$ NA, 2NA), 4-methoxy- $\beta$ -naphtylamide (4-methoxy-2-naphtylamide) and 7-amido-4-methylcoumarin 20 (AMC, MCA; Abramic et al. 2000, Ohkubo et al. 1999). Cleavage of these fluorogenic substrates leads to the release of fluorescent  $\beta$ -naphtylamine or 7-amino-4-methylcoumarin respectively. Chromophores include but are not limited to p-nitroanilide diacetate (pNA). The hydrolysis of a peptide-pNA bond in the chromogenic substrates results in the release of pNA which in turn changes color. Thus the change in absorbance (DA/min) is directly proportional 25 to the enzymatic activity. Using the Protease-Glo<sup>TM</sup> Assay from Promega, upon cleavage by DPP3, aminoluciferin is released and serves as a substrate for a coupled luciferase reaction that emits detectable luminescence.

30 In a preferred embodiment DPP3 activity is measured by addition of the fluorogenic substrate Arg-Arg- $\beta$ NA and monitoring fluorescence in real time. In one specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said sample is selected from the group comprising whole blood, serum, plasma, cerebrospinal fluid, urine, saliva, sputum, and pleural effusions.

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said sample is a blood sample selected from the group comprising whole blood, serum and plasma.

5 Another embodiment of the present invention is an assay or kit for determining active DPP3 in a bodily fluid sample of a subject comprising:

- A capture-binder that binds to full-length DPP3 specifically
- A substrate of DPP3.

10

The kit may additionally comprise a calibrator:

- 15
- A calibrator may be samples with known DPP3 concentration (native, purified or recombinant);
  - A calibrator may be the cleavage products themselves, such as free fluorophores (e.g. 2-naphthylamine), free chromophores (e.g. p-nitroanilide) or free luciferin.

The kit may additionally comprise washing reagents:

- 20
- Washing reagent (can be any aqueous buffer solution with or without detergent. Here we used 8 mM Tris-HCl, pH 7,5, 60 mM NaCl, 0,02 % Tween 20.).

In a specific embodiment said assay is an enzyme capture assay (ECA, e.g. US5612186A, US5601986A).

25

In a specific embodiment said capture-binder that binds to full-length DPP3 specifically inhibits less than 50% DPP3 activity in a liquid phase assay, preferably less than 40%, more preferably 30 %. For definition of liquid phase assay see above. In one specific embodiment to prevent inhibition of DPP3 the capture-binder should not bind DPP3 in the area around the active center and substrate binding region (amino acids 316 – 669 of SEQ ID No. 1).

30

In a specific embodiment of said assay or kit for determining active DPP3 in a bodily fluid sample of a subject said binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

- 5 In a specific embodiment of said assay or kit for determining active DPP3 in a bodily fluid sample of a subject said binder is an antibody.

The term "antibody" generally comprises monoclonal and polyclonal antibodies and binding fragments thereof, in particular Fc-fragments as well as so called "single-chain-antibodies" 10 (Bird et al. 1988), chimeric, humanized, in particular CDR-grafted antibodies, and dia or tetrabodies (Holliger et al. 1993). Also comprised are immunoglobulin-like proteins that are selected through techniques including, for example, phage display to specifically bind to the molecule of interest contained in a sample. In this context the term "specific binding" refers to antibodies raised against the molecule of interest or a fragment thereof. An antibody is 15 considered to be specific, if its affinity towards the molecule of interest or the aforementioned fragment thereof is at least preferably 50-fold higher, more preferably 100-fold higher, most preferably at least 1000-fold higher than towards other molecules comprised in a sample containing the molecule of interest. It is well known in the art how to make antibodies and to select antibodies with a given specificity.

20 In a specific embodiment of said assay or kit for determining active DPP3 in a bodily fluid sample of a subject said capture binder is immobilized on a surface.

In a specific embodiment of said assay or kit for determining active DPP3 in a bodily fluid 25 sample of a subject said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or di- and tri-peptides coupled to a fluorophore, a chromophore or aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay). Di- or tripeptides that are cleaved by DPP3 include, but are not limited to, Arg-Arg, Ala-Ala, Ala- 30 Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe. Fluorophores include but are not limited to  $\beta$ -naphthylamide (2-naphthylamide,  $\beta$ NA, 2NA), 4-methoxy- $\beta$ -naphthylamide (4-methoxy-2-naphthylamide) and 7-amido-4-methylcoumarin (AMC, MCA; Abramić et al. 2000, Ohkubo et al. 1999).

Cleavage of these fluorogenic substrates leads to the release of fluorescent β-naphthylamine or 7-amino-4-methylcoumarin respectively. Chromophores include but are not limited to p-nitroanilide diacetate (pNA). The hydrolysis of a peptide-pNA bond in the chromogenic substrates results in the release of pNA which in turn changes color. Thus the change in absorbance (DA/min) is directly proportional to the enzymatic activity. Using the Protease-Glo™ Assay from Promega, upon cleavage by DPP3, aminoluciferin is released and serves as a substrate for a coupled luciferase reaction that emits detectable luminescence.

In a preferred embodiment DPP3 activity is measured by addition of the fluorogenic substrate Arg-Arg-βNA and monitoring fluorescence in real time.

In a specific embodiment of said assay or kit for determining active DPP3 in a bodily fluid sample of a subject said calibrator is selected from the group comprising: A) recombinant DPP3 (e.g. GST-hDPP3 from USBio), purified native DPP3 (e.g. from human erythrocytes, Abramić et al. 1988) or DPP3 fragments (native, synthetic or recombinant), B) The cleavage products themselves: free fluorophores (e.g. 2-naphthylamine), free chromophores (e.g. p-nitroanilide) or free luciferin, for the quantification of fluorescence, color change and bioluminescence signals.

One embodiment of the present invention of using DPP3 as diagnostic biomarker comprises immunoassays in various formats such as for instance radioimmunoassay (RIA), chemiluminescence- and fluorescence-immunoassays, Enzyme-linked immunoassays (ELISA), Luminex-based bead arrays, protein microarray assays, and rapid test formats such as for instance immunochromatographic strip tests.

The assays can be homogenous or heterogeneous assays, competitive and non-competitive sandwich assays. In a particularly preferred embodiment employing the two antibodies according to the present invention, the assay is in the form of a sandwich assay, which is a non-competitive immunoassay, wherein DPP3 or a fragment thereof to be detected and/or quantified is bound to the first antibody and to the second antibody. The first antibody may be bound to a solid phase, e.g. a bead, a surface of a well or other container, a chip or a strip, and the second antibody is an antibody which is labeled, e.g. with a dye, with a radioisotope, or a reactive or catalytically active moiety. The amount of labeled antibody bound to the analyte is

then measured by an appropriate method. The general composition and procedures involved with “sandwich assays” are well-established and known to the skilled person. (The Immunoassay Handbook, Ed. David Wild, 2005; Hultschig et al. 2006).

- 5 In a particularly preferred embodiment the assay comprises a liquid reaction mixture, wherein  
a first labeling component is attached to the first antibody, wherein said first labeling  
component is part of a labeling system based on fluorescence- or chemiluminescence-  
quenching or amplification, and a second labeling component of said marking system is  
attached to the second antibody, so that upon binding of both antibodies to the analyte a  
10 measurable signal is generated that allows for the detection of the formed sandwich  
complexes in the solution comprising the sample.

In the context of the present invention, fluorescence based assays comprise the use of dyes,  
which may for instance be selected from the group comprising FAM (5-or  
15 6-carboxyfluorescein), VIC, NED, Fluorescein, Fluoresceinisothiocyanate (FITC),  
IRD-700/800, Cyanine dyes, such as CY3, CY5, CY3.5, CY5.5, Cy7, Xanthen,  
6-Carboxy-2' ,4' ,7' ,4,7-hexachlorofluorescein (HEX), TET, 6-Carboxy-4' ,  
5' -dichloro-2' ,7' -dimethoxyfluorescein (JOE), N,N,N' ,N' -Tetramethyl-6-  
20 carboxyrhodamine (TAMRA), 6-Carboxy-X-rhodamine (ROX), 5-Carboxyrhodamine-6G  
(R6G5), 6-carboxyrhodamine-6G (RG6), Rhodamine, Rhodamine Green, Rhodamine Red,  
Rhodamine 110, BODIPY dyes, such as BODIPY TMR, Oregon Green, Coumarines such as  
Umbelliferone, Benzimides, such as Hoechst 33258; Phenanthridines, such as Texas Red,  
25 Yakima Yellow, Alexa Fluor, PET, Ethidiumbromide, Acridinium dyes, Carbazol dyes,  
Phenoxazine dyes, Porphyrine dyes, Polymethin dyes, and the like.

25 In the context of the present invention, chemiluminescence based assays comprise the use of  
dyes, based on the physical principles described for chemiluminescent materials in Kirk-  
Othmer, Encyclopedia of chemical technology, 4th ed., executive editor, J. I. Kroschwitz;  
editor, M. Howe-Grant, John Wiley & Sons, 1993, vol. 15, p. 518-562, incorporated herein by  
30 reference, including citations on pages 551-562. Preferred chemiluminescent dyes are  
acridiniumesters.

One embodiment of the present invention involves a chemical assay for DPP3. The assay uses an enzyme substrate which reacts with DPP3 to form a detectable reaction product. Alternatively, the rate of the reaction of the substrate can be monitored to determine the presence or amount of DPP3 in a test sample. Suitable enzyme substrates include, but are not limited to, dipeptide substrates such as Arg-Arg-β -NA or Arg-Arg-AMC.

Assays embodying such reagents and reactions can be performed in any suitable reaction vessel, for example, a test tube or well of a microtiter plate. Alternatively, assay devices may be developed in disposable form such as dipstick or test strip device formats which are well known to those skilled-in-the-art and which provide ease of manufacture and use. Such disposable assay devices may be packaged in the form of kits containing all necessary materials, reagents and instructions for use.

Assay devices of the present invention could advantageously be formatted as dipstick or test strip devices. For example, a dipstick may be made from a piece of bibulous material containing a chromogenic substrate for DPP3. Alternatively, the dipstick could be made from a nonporous material on which the substrate is coated. Upon contacting the device with the desired test sample, the substrate and any DPP3 present in the sample would interact to form a detectable reaction upon the device.

In an alternative embodiment, the device may be a test strip, wherein the substrate is contained in one or more zones along the length of a strip of bibulous material. Upon contact of one end of the strip with the desired test sample, the liquid sample migrates along the bibulous material. The reaction of the substrate and production of a detectable signal indicates the presence of DPP3 in the test sample. In a multi-zoned device, the number of discrete or isolated zones along the length of the strip which produce a detectable signal may also indicate the quantity of DPP3 present in the test sample. Alternatively, a major portion of the test strip may contain the substrate. The length of the colored reaction formed in a test strip having such a single, elongated substrate zone may be used to indicate the presence or amount of DPP3 in the test sample.

In an alternative assay embodiment, the rate at which the reaction occurs may be detected as an indication of the quantity of DPP3 present in the test sample. For example, the rate at

which the substrate is reacted may be used to indicate the quantity of DPP3 present in the test sample. Alternatively, the rate at which the reaction product is formed may be used to indicate the quantity of DPP3 present in the test sample.

- 5 In yet another embodiment, a capture or binding assay may be performed to detect and/or quantitate the protease. For example, an antibody reactive with DPP3 protein, but which does not interfere with peptidase activity, may be immobilized upon a solid phase. The test sample is passed over the immobile antibody, and DPP3, if present, binds to the antibody and is itself immobilized for detection. A substrate may then be added, and the reaction product may be
- 10 detected to indicate the presence or amount of DPP3 in the test sample. For the purposes of the present description, the term "solid phase" may be used to include any material or vessel in which or on which the assay may be performed and includes, but is not limited to, porous materials, nonporous materials, test tubes, wells, slides, etc.
- 15 In another specific embodiment DPP3 ECA can be performed as a test strip assay. In an exemplary test strip device, a test sample application pad is optionally attached to one end of a porous strip. The strip contains an immobilized antibody which will bind to and thereby immobilize DPP3 at a predetermined site for subsequent detection. Optionally, the device may include an end of assay indicator which is positioned at the distal end of the test strip away from the test sample contact site. The end of assay indicator produces a detectable signal upon contact with the test sample or an assay reagent thereby indicating that the assay is complete.

A test sample application pad may be a portion of the porous strip itself or a material in fluid-flow contact with the end of the porous strip, referred to as the proximal end, such that the test sample can pass or migrate from the application pad to the porous strip. Fluid-flow contact can include physical contact of the application pad to the porous strip as well as the separation of the application pad from the porous strip by an intervening space or additional material which still allows fluid to flow between the application pad and the porous strip. Substantially all of the application pad can overlap the porous strip to enable the test sample to pass through substantially any part of the application pad to the proximal end of the porous strip. Alternatively, only a portion of the application pad might be in fluid-flow contact with the

porous strip. The application pad can be any material which can transfer the test sample to the porous strip.

The porous strip of the assay device can be any suitably absorbent, porous, bibulous, chromatographic or capillary possessing material through which a test sample containing the analyte can be transported by a capillary or wicking action. Natural, synthetic, or naturally occurring materials that are synthetically modified, can be used as the porous strip including, but not limited to: cellulose materials such as paper, cellulose, and cellulose derivatives such as cellulose acetate and nitrocellulose; fiberglass; cloth, both naturally occurring (e.g., cotton) and synthetic (e.g., nylon); porous gels such as silica gel, agarose, dextran, and gelatin; porous fibrous matrixes; starch based materials, such as crosslinked dextran chains; ceramic materials; films of polyvinyl chloride and combinations of polyvinyl chloride-silica; and the like. The porous strip should not interfere with the production of a detectable signal. The porous strip should have a reasonable inherent strength, or strength can be provided by means of a supplemental support.

The particular dimensions of the porous strip will be a matter of convenience, depending upon the size of the test sample involved, the assay protocol, the means for detecting and measuring the signal, and the like. For example, the dimensions may be chosen to regulate the rate of fluid migration as well as the amount of test sample to be imbibed by the porous strip.

In one possible test strip device of the present invention, a DPP3 substrate and/or DPP3 capture antibody may be immobilized on the porous strip to form at least one analyte detection site, i.e., that region of the porous strip having one or more assay reagents non-diffusively attached thereto. In another device embodiment, the measurement or detection region of the test strip may include a plurality of sites containing a DPP3 substrate and/or immobilized anti-DPP3 antibody. Optionally, the different detection sites may contain different amounts of substrate and/or immobilized anti-DPP3 antibody, i.e., a higher amount in the first detection site and lesser amounts in subsequent sites. For example, if 20 nanograms of antibody captures the equivalent of 1 nmol/min/ml of DPP3, then the first detection site of an assay device might contain 50 nanograms of anti-DPP3 antibody while the subsequent sites contain 10, 20, 30, etc. nanograms of antibody. Upon the addition of test sample, the number of sites displaying a detectable signal provides a quantitative indication of

the amount of DPP3 present in the sample. The detection sites may be configured in any suitably detectable shape and are typically in the shape of a bar spanning the width of the test strip.

- 5     Optionally, the multi-capture site device may be prepared such that if a threshold amount of DPP3 is not present in the test sample, then substantially all of the DPP3 will bind to the antibody in the first capture site and thus become immobilized at that site. If a greater than threshold amount of DPP3 is present in the test sample, the remaining DPP3 will bind to subsequent detection zones of immobilized antibody along the length of the test strip. The  
10    greater the amount of DPP3 in the test sample, the greater the number of capture sites that will display a detectable signal due to the presence of DPP3. As will be appreciated by those skilled-in-the-art, devices containing multiple DPP3 substrate sites can also be produced wherein the amount of substrate in the individual sites is designed to produce a quantitative or semiquantitative assay result.

15    Another important embodiment of the invention is a method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes comprising:

- 20    • determining the amount of total DPP3 or determining the amount of active DPP3 in a sample of bodily fluid of said subject,  
• comparing said determined amount to a predetermined threshold,  
• wherein said subject is diagnosed as having a disease or condition accompanied by or related to necrotic processes if said determined amount is above said predetermined threshold.

25    In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said amount of total or active DPP3 is determined in the unit of concentration.

30    Methods of determining the amount of total or active DPP3 are known in the art. In the context of a method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to the present invention either state-of-the art methods

and assays may be used or the above-described methods and assays for determining DPP3 may be used.

The threshold is pre-determined by measuring DPP3 concentration and or DPP3 activity in  
5 healthy controls and calculating e.g. the according 75-percentile, more preferably the 90-percentile, even more preferably the 95-percentile. The upper border of the 75-percentile, more preferably the 90-percentile, even more preferably the 95-percentile, defines the threshold for healthy versus diseased patients. In relation to said percentiles, the threshold that divides between healthy and diseased patients may be between 5 and 25 ng/ ml, more  
10 preferably between 7 and 20 ng/ ml, more preferably between 8 and 18 ng/ ml, most preferred between 10 and 15 ng/ ml in plasma using a sandwich type anti-DPP3 immunoassay (see example 3). In a DPP3 specific enzyme capture activity assay in plasma, the threshold that divides between healthy and diseased patients may be between 0.5 and 2 nmol  $\beta$  NA min<sup>-1</sup> ml<sup>-1</sup>, more preferably between 0.7 and 1.8 nmol  $\beta$  NA min<sup>-1</sup> ml<sup>-1</sup>, more preferably between  
15 0.8 and 1.5 nmol  $\beta$  NA min<sup>-1</sup> ml<sup>-1</sup>, most preferred between 1.0 and 1.3 nmol  $\beta$  NA min<sup>-1</sup> ml<sup>-1</sup> (see example 5).

The person skilled in the art knows how to determine thresholds from conducted previous studies. The person skilled in the art knows that a specific threshold value may depend on the  
20 cohort used for calculating a pre-determined threshold that can be later-on used in routine. The person skilled in the art knows that a specific threshold value may depend on the calibration used in the assay. The person skilled in the art knows that a specific threshold value may depend on the sensitivity and/or specificity that seems to be acceptable for the practitioner.

25 The sensitivity and specificity of a diagnostic test depends on more than just the analytical "quality" of the test, they also depend on the definition of what constitutes an abnormal result. In practice, Receiver Operating Characteristic curves (ROC curves), are typically calculated by plotting the value of a variable versus its relative frequency in "normal" (i.e. apparently healthy) and "disease" populations (i.e. patients suffering from an infection). Depending on  
30 the particular diagnostic question to be addressed, the reference group must not be necessarily "normals", but it might be a group of patients suffering from another disease or condition, from which the diseased group of interest shall be differentiated. For any particular marker, a

distribution of marker levels for subjects with and without a disease will likely overlap. Under such conditions, a test does not absolutely distinguish normal from disease with 100% accuracy, and the area of overlap indicates where the test cannot distinguish normal from disease. A threshold is selected, above which (or below which, depending on how a marker changes with the disease) the test is considered to be abnormal and below which the test is considered to be normal. The area under the ROC curve is a measure of the probability that the perceived measurement will allow correct identification of a condition. ROC curves can be used even when test results do not necessarily give an accurate number. As long as one can rank results, one can create a ROC curve. For example, results of a test on "disease" samples might be ranked according to degree (e.g. 1=low, 2=normal, and 3=high). This ranking can be correlated to results in the "normal" population, and a ROC curve created. These methods are well known in the art (see, e.g., Hartley et al, 1982). Preferably, a threshold is selected to provide a ROC curve area of greater than about 0.5, more preferably greater than about 0.7. The term "about" in this context refers to +/- 5% of a given measurement.

Once the threshold value is determined by using a previous study cohort and taking into consideration all the above-mentioned points the medical practitioner will use the pre-determined threshold for the methods of diagnosing a disease according to the invention and will determine whether the subject has a value above or below said pre-determined threshold value in order to make an appropriate diagnosis.

DPP3 concentrations in tissue homogenates and bodily fluids can be measured using several commercially available DPP3 ELISA kits (e.g. from LifeSpan BioSciences). Those assays are all based upon a Sandwich assay principle and for research use only.

The standard procedure to measure DPP3 levels is to determine the DPP3 activity using fluorogenic substrate (e.g. Arg-Arg- $\beta$ -naphtylamine) in a liquid phase assay (Ellis & Nuenke 1967). Commercially available kits (e.g. from BPS Bioscience) usually contain low binding black microtiter plates, recombinant DPP3, fluorogenic substrate and respective buffers. Those kits are regularly used as screening assay for DPP3 substrates and inhibitors.

In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said sample is selected from the group

comprising whole blood, serum and plasma. The bodily fluid in the context of the method of the present invention maybe also selected from the group of blood, serum, plasma, cerebrospinal fluid, urine, saliva, sputum, and pleural effusions.

- 5 Necrotic processes are herein defined by all processes in the body that lead to the death of cells and release of DPP3 from the cells cytoplasm into the extracellular space and/ or bodily fluids. Those processes include, but are not limited to, necrosis, apoptosis, necroptosis, eryptosis.
- 10 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes, said disease is selected from the group comprising: heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)) or SIRS or sepsis, cancer, acute kidney injury (AKI), central nervous system (CNS) disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension. Table 1 lists clinical symptoms/ diseases that are accompanied by or related to necrotic processes and the respective necrotic event.
- 15
- 20 In another embodiment said disease is selected from the group comprising: acute heart failure (AHF), myocardial infarction (MI), liver failure, burn injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)) or SIRS or sepsis, cancer, and acute kidney injury (AKI).
- In one embodiment said disease is hypotension.

25

<b>Clinical symptom/ disease</b>	<b>necrotic/ apoptotic event</b>	<b>reference</b>
acute heart failure	sudden death of cells/ tissue areas/ organs	Fischer et al. 2005, Zong et al. 2006
myocardial infarction		
stroke		
liver failure/ injury		Lanier et al. 2011
kidney failure/ injury		Raghupathi 2004
burn injuries		
traumatic injuries (e.g. traumatic brain injury); polytrauma		
viral + microbial + parasite infections	death of macrophages, lysis of host cells	Zong et al. 2006; Fink et al. 2005

AIDS	progressive death of immune cells	Fischer et al. 2005
Sepsis	necrosis due to immune response/ apoptosis of immune cells	Pinheiro da Silva et al. 2009
Malaria	lysis of host erythrocytes (eryptosis)	Lang et al. 2015
neurodegenerative diseases (e.g. AD, Huntington)	slowly progressive neuronal cell death	Fischer et al. 2005
other CNS disorders (e.g. seizures)	Excitotoxic cell death	Zong et al. 2006
cancer	increased inflammation	Wallach et al. 2014
autoimmune disease		
Kawasaki disease	increased inflammation and necrosis of small blood vessels	Dimitrades et al. 2014

**Table 1: Diseases accompanied by or related to necrotic processes.**

- 5 Apoptosis is the process of programmed cell death (PCD) that may occur in multicellular organisms. Biochemical events lead to characteristic cell changes (morphology) and death. These changes include blebbing, cell shrinkage, nuclear fragmentation, chromatin condensation, and chromosomal DNA fragmentation. For review see Elmore 2007.
- 10 Necrosis is a form of cell injury, which results in the premature death of cells in living tissue by autolysis (Vanlangenakker et al. 2008). Necrosis is caused by factors external to the cell or tissue, such as infection, toxins, or trauma, which result in the unregulated digestion of cell components.
- 15 While apoptosis often provides beneficial effects to the organism, necrosis is almost always detrimental and can be fatal.

20 Cellular death due to necrosis does not follow the apoptotic signal transduction pathway, but rather various receptors are activated, and result in the loss of cell membrane integrity and an uncontrolled release of products of cell death into the extracellular space. This initiates in the surrounding tissue an inflammatory response, which prevents nearby phagocytes from locating and eliminating the dead cells by phagocytosis. For this reason, it is often necessary to remove necrotic tissue surgically, a procedure known as debridement. Untreated necrosis results in a build-up of decomposing dead tissue and cell debris at or near the site of the cell 25 death.

A form of programmed necrosis, called necroptosis, has been recognized as an alternate form of programmed cell death. Necroptosis may serve as a cell-death backup to apoptosis when the apoptosis signaling is blocked by endogenous or exogenous factors such as viruses or mutations (Linkermann et al. 2014). Most recently, other types of regulated necrosis have been discovered as well, which share several signaling events with necroptosis and apoptosis (Vanden Berghe et al. 2014).

Heart failure (HF) is a cardiac condition that occurs, when a problem with the structure or function of the heart impairs its ability to supply sufficient blood flow to meet the body's needs. It can cause a large variety of symptoms, particularly shortness of breath (SOB) at rest or during exercise, signs of fluid retention such as pulmonary congestion or ankle swelling and objective evidence of an abnormality of the structure or function of the heart at rest.

Heart failure is a clinical syndrome characterized by a constellation of symptoms and signs caused by cardiac dysfunction. It is one of the major causes of morbidity and mortality in the developed countries, with a prevalence of 1-2%. Heart failure can be grouped into chronic HF and acute HF. Patients with chronic HF can be grouped into stable chronic HF, worsening signs and symptoms of chronic HF and acute decompensation of chronic HF. Acute heart failure (AHF) is defined as a rapid onset of signs and symptoms of heart failure resulting in the need for urgent therapy or hospitalization. AHF can present as acute de novo HF (new onset of AHF in a patient without previous cardiac dysfunction) or acute decompensation of chronic HF. AHF is the leading cause of hospitalization in adults older than 65 years of age. Despite marked improvements in the prognosis of chronic heart failure patients, primarily related to therapeutic advances over the past few decades, both short- and long-term outcomes remain very poor once patients are hospitalized for decompensated heart failure. Nearly 25% of patients hospitalized for AHF need readmission within 30 days of hospital discharge while <50% survive beyond 5 years after hospitalization. In addition to significantly reducing survival and quality of life of affected patients, the monetary burden of AHF on health care systems is enormous. The total cost of heart failure care was estimated to be \$31 billion in the US alone in 2012 and majority of this cost is associated with in-hospital care. This cost is projected to increase to an unprecedented \$70 billion in 2030 due to ageing of populations.

Heart failure comprises a wide range of patients, from those with normal left ventricular ejection fraction (LVEF) typically considered as  $\geq 50\%$ , also known as HF with preserved EF (HFpEF) to those with reduced LVEF, typically considered as  $< 40\%$ , also known as HF with reduced EF (HFrEF). Patients with an LVEF in the range of 40–49% represent a ‘grey area’,  
5 which is defined as HF with mid-range EF (HFmrEF) (Ponikowski et al. 2016).

Heart failure may occur as acute or chronic heart failure.

The term "acute" is used to mean rapid onset and to describe exacerbated or decompensated  
10 heart failure, referring to episodes in which a patient can be characterized as having a change in heart failure signs and symptoms resulting in a need for urgent therapy or hospitalization.

The term "chronic" refers to long duration. Chronic heart failure is a long-term condition, usually kept stable by the treatment of symptoms (stable chronic HF).

- 15 Stable chronic HF is characterized by:
- (i) the presence of structural or functional failure of the heart that impairs its ability to supply sufficient blood flow to meet body's needs,
  - (ii) the absence of volume overload (manifested by pulmonary and/ or systemic congestion) and/ or profound depression of cardiac output (manifested by hypotension, renal insufficiency and/ or a shock syndrome),
- 20 and whereas the patient is not in need of urgent therapy or therapy adjustment and does not require hospitalization.

Chronic HF with worsening signs and symptoms is characterized by:

- 30 (i) the presence of structural or functional failure of the heart that impairs its ability to supply sufficient blood flow to meet body's needs,

- (ii) volume overload (manifested by pulmonary and/ or systemic congestion) and/ or profound depression of cardiac output (manifested by hypotension, renal insufficiency and/ or a shock syndrome)

5 and whereas the patient is not in need of urgent therapy and does not require hospitalization, but is in need of therapy adjustment.

Chronic heart failure may also decompensate (termed acute decompensated heart failure or acute decompensated chronic heart failure), which is most commonly the result from an 10 intercurrent illness (such as pneumonia), myocardial infarction, arrhythmias, uncontrolled hypertension or a patient's failure to maintain fluid restriction, diet or medication. After treatment, patients with acute decompensated chronic HF may return to a stable chronic compensated status (stable chronic HF).

15 New onset acute HF and acute decompensated chronic HF are characterized by:

- (i) the presence of structural or functional failure of the heart that impairs its ability to supply sufficient blood flow to meet body's needs,
- 20 (ii) volume overload (manifested by pulmonary and/ or systemic congestion) and/ or profound depression of cardiac output (manifested by hypotension, renal insufficiency and/ or a shock syndrome)

and whereas the patient is in need of urgent therapy or therapy adjustment and does require 25 hospitalization.

Acute HF		Chronic HF	
New-onset AHF	Acute decompensated HF = acute decompensated chronic HF	Worsening signs/ symptoms of chronic HF	Stable chronic HF

The above definitions of acute heart failure that either new-onset AHF or acute decompensated HF or acute decompensated chronic HF or worsening signs/symptoms of chronic heart failure are in line with Voors et al. 2016.

5 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes DPP3 activity can be determined using a liquid phase activity assay or using an enzyme capture activity assay.

10 In a liquid phase assay samples of bodily fluids are directly subjected to fluorogenic substrates (e.g. Arg-Arg- $\beta$ -NA). Since there are many different amino peptidases in the plasma (Sanderink et al. 1988), it is possible that the used substrate is cleaved by peptidases other than DPP3. To circumvent this problem one preferred method of detecting specific DPP3 activity is the use of an enzyme capture activity assay.

15 In one specific embodiment determination of active DPP3 in an enzyme capture assay comprises the steps:

- Contacting said sample with a capture-binder that binds to full-length DPP3 but preferably inhibits DPP3 activity in a liquid phase assay less than 50%, preferably less than 40%, more preferably 30 %. To prevent inhibition of DPP3 the capture-binder should not bind DPP3 in the area around the active center and substrate binding region (amino acids 316 – 669 of SEQ ID No. 1).
- Separating DPP3 bound to said capture binder from bodily fluid sample,
- Adding substrate of DPP3 to said separated DPP3,
- Quantifying DPP3 activity by measuring the conversion of the substrate of DPP3,
- Evaluation of measured signals in comparison to non-diseased controls. The threshold may be pre-determined e.g. by measuring DPP3 concentration and or DPP3 activity in healthy controls and calculating the according 75-percentile. The upper border of the 75-percentile defines the threshold for healthy versus diseased patients.

In one specific embodiment determination of active DPP3 is conducted according to the above- described method according to the present invention.

In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

- 5 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes is an enzyme capture assay (ECA, US5612186A, US5601986A). The binder of DPP3 in this assay is an antibody.

- 10 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said capture binder is immobilized on a surface. A binder reactive with DPP3 protein, but which does not interfere with peptidase activity by more than 50 %, preferably less than 40 %, preferably less than 30 %, may be immobilized upon a solid phase. To prevent inhibition of DPP3 the capture-binder should not bind DPP3 in the area around the active center and substrate binding region (amino acids 15 316 – 669 of SEQ ID No. 1).

In a specific embodiment of said method for determining active DPP3 in a bodily fluid sample of a subject said binder may be selected from the group of antibody, antibody fragments, non-Ig scaffold or aptamers.

- 20 The test sample is passed over the immobile binder, and DPP3, if present, binds to the binder and is itself immobilized for detection. A substrate may then be added, and the reaction product may be detected to indicate the presence or amount of DPP3 in the test sample. For the purposes of the present description, the term "solid phase" may be used to include any material or vessel in which or on which the assay may be performed and includes, but is not limited to: porous materials, nonporous materials, test tubes, wells, slides, agarose resins (e.g. Sepharose from GE Healthcare Life Sciences), magnetic particals (e.g. Dynabeads<sup>TM</sup> or Pierce<sup>TM</sup> magnetic beads from Thermo Fisher Scientific), etc.

- 30 Binders of protein or peptide origin (e.g. antibody, antibody fragments, non-Ig scaffold) are immobilized onto the solid phase by methods comprising: physical adsorption (e.g. by electrostatic interaction or hydrophobic interaction), bioaffinity immobilization (e.g. avidin-biotin, protein A/ G/ L, His-tag and Ni<sup>2+</sup>-NTA, GST-tag and glutathione, DNA hybridization,

aptamers), covalent bond (e.g. amine and N-hydroxysuccinimide) or a combination said immobilization methods (Kim & Herr 2013). Binders of oligonucleotide origin (e.g. aptamers) may be immobilized onto the solid phase by utilization of the (strept)avidin-biotin system (Müller et al. 2012, Deng et al. 2014).

5 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.

10 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said DPP3 substrate conversion is measured by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg-2NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

15 20 In a specific embodiment of said method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes said substrate may be selected from the group comprising: angiotensin II and III, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or di- and tripeptides coupled to a fluorophore, a chromophore or aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay). Di- or tripeptides that are cleaved by DPP3 include, but are not limited to, Arg-Arg, Ala-Ala, Ala-Arg, Ala-Phe, Asp-Arg, Gly-Ala, Gly-Arg, Gly-Phe, Leu-Ala, Leu-Gly, Lys-Ala, Phe-Arg, Suc-Ala-Ala-Phe. Fluorophores include but are not limited to  $\beta$ -naphthylamide (2-naphthylamide,  $\beta$ NA, 2NA), 4-methoxy- $\beta$ -naphthylamide (4-methoxy-2-naphthylamide) and 7-amido-4-methylcoumarin (AMC, MCA; Abramic et al. 2000, Ohkubo et al. 1999). Cleavage of these fluorogenic substrates leads to the release of fluorescent  $\beta$ -naphthylamine or 7-amino-4-methylcoumarin respectively. Chromophores include but are not limited to p-nitroanilide diacetate (pNA). The hydrolysis of a peptide-pNA bond in the chromogenic substrates results in the release of pNA which in turn changes color. Thus the

change in absorbance (DA/min) is directly proportional to the enzymatic activity. Using the Protease-GloTM Assay from Promega, upon cleavage by DPP3, aminoluciferin is released and serves as a substrate for a coupled luciferase reaction that emits detectable luminescence.

- 5 In a preferred embodiment DPP3 activity is measured by addition of the fluorogenic substrate Arg-Arg- $\beta$ NA and monitoring fluorescence in real time.

Another important embodiment of the present invention is an inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or  
10 related to necrotic processes.

Inhibitors are molecules that preferably significantly inhibit DPP3 activity. Those molecules can be peptides and small molecules (see table 2) or antibodies (see table 3). Significantly inhibiting means inhibiting more than 80 % in a liquid phase assay as described above, preferably more than 90 %, more preferably almost or actually 100% inhibition.  
15

Peptide inhibitors of DPP3 include, but are not limited to, spinorphin, synthetic derivatives of spinorphin (tynorphin and other peptides, see table 2; Yamamoto et al. 2000), propioxatin A and B (US4804676) and synthetic propioxatin A analogues (Inaoka et al. 1988).

20 Small molecule inhibitors of DPP3 include, but are not limited to, fluostatins and benzimidazol derivatives. Fluostatins A and B are antibiotics produced in *Streptomyces* sp. TA-3391 that are non-toxic and strongly inhibit DPP3 activity. So far 20 different derivatives of benzimidazol have been synthesized and published (Agic et al. 2007; Rastija et al. 2015),  
25 of which the two compounds 1' and 4' show the strongest inhibitory effect (Agic et al. 2007).

Compound Type	Full Name/Active-oral Structure	Reference
fluostatins	Fluostatin A	Akiyama et al. 1998
	Fluostatin B	
small molecules	Compound 1	Agic et al. 2007
	Compound 2	
	Compound 3	
	Compound 4	
	Compound 1'-8'	Rastija et al. 2015
	1' Ar=phenyl 4' Ar=o-Cl-phenyl 7' Ar=2-furyl 8' Ar=2-thienyl	
	2' Ar=phenyl 5' Ar=o-Cl-phenyl	
	3' Ar=phenyl 6' Ar=o-Cl-phenyl	
	Compound 1'-16	
	1 Ar=phenyl 5 Ar=o-Cl-phenyl 9 Ar=2-furyl 13 Ar=2-thienyl	
	3 Ar=phenyl 7 Ar=o-Cl-phenyl 14 Ar=2-thienyl	
	4 Ar=phenyl 8 Ar=o-Cl-phenyl 11 Ar=2-furyl 15 Ar=2-thienyl	
	12 Ar=2-furyl 16 Ar=2-thienyl	
	2 Ar=phenyl 6 Ar=o-Cl-phenyl 10 Ar=2-furyl	

Compound Name	With N-Substituted Amino Acid Structures	Patent
	<p>Propioxatin A and B</p> <p>A: R = H B: R = CH<sub>3</sub></p>	US 4804676 A/ Inaoka et al. 1986
peptides  propioxatins and propioxatin A analogues	<p>Propioxatin A analogues (Compound 1-17)</p> <p>1</p> <p>2</p> <p>3</p> <p>4</p> <p>5</p> <p>6</p> <p>7</p> <p>8</p> <p>9</p> <p>10</p> <p>11</p> <p>12</p> <p>13</p> <p>14</p> <p>15</p> <p>16</p> <p>17</p>	Inaoka et al. 1988

<b>spinorphin and spinorphin derivatives</b>	Spinorphin (LVVYPWT)	Yamamoto et al. 2000; Chiba et al. 2003
	Tynorphin (VVYPW)	
	Spinorphin derivatives (AVYPW, FIVPW, FVAPW, FVYPW, GVYPW, IVYPW, LVVPW, LVVYP, LVVYPW, LVYPW, PWT, SVYPW, VVYP, VVYPWT, VYP, VYPW, VYPWT, WVYPW, YAIPW, YPW, YPWT, YSIPW, YSVPW, YVYPW)	

**Table 2: Peptide and small molecule inhibitors of DPP3**

- In a preferred embodiment of the invention the chosen inhibitor is pharmaceutically acceptable, selective and/ or specific for DPP3, and does not cross the cell membrane and/ or the blood brain barrier. Selective and specific inhibitors of DPP3 do not bind to other proteins/ enzymes or are bound by other proteins/ enzymes, and do not inhibit any other enzyme/ protease/ peptidase other than DPP3. Small peptides can be bound and cleaved by unspecific aminopeptidases and small molecule inhibitors easily cross the cell membrane and the blood brain barrier. Anti-DPP3 antibodies, anti-DPP3 antibody fragments or anti-DPP3 non-Ig scaffolds bind DPP3 specifically and selectively, and do not cross the cell membrane or the blood brain barrier. Therefor the preferred inhibitors of DPP3 activity are specific anti-DPP3 antibodies, antibody fragments or non-Ig scaffolds.
- In a specific embodiment of the invention the inhibitor of the activity of DPP3 is used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said inhibitor is selected from the group comprising an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold.
- In a specific embodiment of the invention the inhibitor or effector of the activity of DPP3 is used in prevention or therapy of disease or condition in a subject accompanied by or related to necrotic processes wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)) or SIRS or sepsis, cancer, acute kidney injury (AKI), central nervous system (CNS) disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension.

In another embodiment said disease is selected from the group comprising: acute heart failure (AHF), myocardial infarction (MI), liver failure, burn injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)) or SIRS or sepsis, cancer, and acute kidney injury (AKI).

5

In one specific embodiment the invention the inhibitor of the activity of DPP3 is used in prevention of a disease or condition in a subject wherein the disease or condition is acute heart failure (AHF), myocardial infarction (MI), liver failure, cancer, acute kidney injury (AKI) and hypotension.

10

In one specific embodiment the invention the inhibitor of the activity of DPP3 is used in therapy of a disease or condition in a subject wherein the disease or condition is acute heart failure (AHF), myocardial infarction (MI), liver failure, cancer, acute kidney injury (AKI) and hypotension.

15

In one specific embodiment for all embodiments of the invention said disease is not Alzheimer. In one specific embodiment for all embodiments of the invention said disease is not cancer. In one specific embodiment for all embodiments of the invention said disease is not Rheumatoid Arthritis.

20

In one specific embodiment said disease or condition is hypotension.

In a specific embodiment of the invention the inhibitor of the activity of DPP3 is used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said inhibitor is an antibody that is mono-binding or at least two-binding.

In a specific embodiment of the invention the inhibitor or effector of the activity of DPP3 is used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said inhibitor or effector is an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 1, in a specific embodiment an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 2.

In a specific embodiment of the invention the inhibitor or effector of the activity of DPP3 is used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said inhibitor or effector is an antibody or fragment or scaffold that exhibits a minimum binding affinity to DPP3 of less than  $10^{-7}$  M.

5 In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor or effector is an antibody or fragment or scaffold binds to full-length DPP3 and inhibits activity of DPP3 of at least 10%, or at least 50%, more preferred at least 60%, even more preferred more than 70 %, even more preferred more than 80 %, even more preferred more than 90 %, even more preferred more than 95 %. Activity may be determined in a liquid phase assay as described above.

10 15 In a specific embodiment of the invention the inhibitor or effector of the activity of DPP3 is used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said inhibitor or effector is an antibody or fragment or scaffold that is monospecific.

20 25 Monospecific anti-DPP3 antibody or monospecific anti-DPP3 antibody fragment or monospecific anti-DPP3 non-Ig scaffold means that said antibody or antibody fragment or non-Ig scaffold binds to one specific region encompassing at least 5 amino acids within the target DPP3. Monospecific anti-DPP3 antibody or monospecific anti-DPP3 antibody fragment or monospecific anti-DPP3 non-Ig scaffold are anti-DPP3 antibodies or anti-DPP3 antibody fragments or anti-DPP3 non-Ig scaffolds that all have affinity for the same antigen.

30 In another specific and preferred embodiment the anti-DPP3 antibody or the anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold binding to DPP3 is a monospecific antibody, antibody fragment or non-Ig scaffold, respectively, whereby monospecific means that said antibody or antibody fragment or non-Ig scaffold binds to one specific region encompassing at least 4 amino acids within the target DPP3. Monospecific antibodies or fragments or non-Ig scaffolds according to the invention are antibodies or fragments or non-Ig scaffolds that all have affinity for the same antigen. Monoclonal antibodies are monospecific, but monospecific

antibodies may also be produced by other means than producing them from a common germ cell.

In a specific embodiment of the invention the inhibitor or effector of the activity of DPP3 is  
5 used in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes wherein said subject has an elevated level of DPP3. An elevated level is a level above a pre-determined threshold.

Another embodiment of the present invention is a pharmaceutical composition comprising an  
10 inhibitor of the activity of DPP3 as described above for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes.

Another embodiment of the present invention is a method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes wherein an inhibitor  
15 of the activity of DPP3 is administered.

In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor is selected from the group comprising an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-  
20 Ig scaffold.

In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection  
25 (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)) or SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension.

30 In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor is an antibody that is mono-binding or at least two-binding.

In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor is an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 1, in a specific embodiment an anti-DPP3 antibody or anti-DPP3 antibody fragment or 5 anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 2.

In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor is an antibody or fragment or scaffold that exhibits a minimum binding affinity to DPP3 of less than  $10^{-7}$  M.

10 In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor or effector is an antibody or fragment or scaffold that is monospecific.

15 In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said inhibitor or effector is an antibody or fragment or scaffold binds to full-length DPP3 and inhibits activity of DPP3 of at least 10%, or at least 50%, more preferred at least 60%, even more preferred more than 70 %, even more preferred more than 80 %, even more preferred more than 90 %, even more 20 preferred more than 95 %.

25 In a specific embodiment of the method for preventing or treating a disease or condition in a subject accompanied by or related to necrotic processes said subject has an elevated level of DPP3. An elevated level is a level above a pre-determined threshold. Thresholds are defined above.

Another embodiment of the present invention is the clearance of DPP3 from the patient's blood. Clearance can be achieved by several apheresis techniques and/ or affinity chromatography steps (Balogun et al. 2010). Those methods include, but are not limited to, 30 filtering patient plasma through an adsorber containing specific and highly affine DPP3 antibody coupled to agarose resins, see example 12 for analysis of DPP3 binding to possible adsorber material.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Routes of administration are generally classified by the location at which the substance is applied. Common examples include oral, epicutaneous, subcutaneous, intradermal, sublingual, intramuscular, intraarterial, intravenous and 5 intraperitoneal administration.

Pharmaceutical compositions may also be applied via the central nervous system (CNS), for example epidural (synonym: peridural) by injection or infusion into the epidural space, intracerebral (into the cerebrum) direct injection into the brain, intracerebroventricular 10 (administration into the ventricular system of the brain) or intrathecal (into the spinal cord canal).

In the following specific embodiments of the invention are outlined:

- 15 1. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes comprising:
  - determining the amount of total DPP3 and/or determining the amount of active DPP3 in a sample of bodily fluid of said subject,
  - comparing said determined amount of total DPP3 or active DPP3 to a predetermined threshold,
  - wherein said subject is diagnosed as having a disease or condition accompanied by or related to necrotic processes if said determined amount is above said predetermined threshold.
- 20 2. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 1 wherein the amount of total or active DPP3 is determined in the unit of concentration.
- 25 3. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 1 or 2, wherein said sample is selected from the group comprising whole blood, serum and plasma.

4. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1-3, wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension.
5. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1-4, wherein the amount of total DPP3 and/or the amount of active DPP3 is determined in a bodily fluid sample of said subject and comprises the steps:
  - Contacting said sample with a capture-binder that binds specifically to full-length DPP3,
  - Separating DPP3 bound to said capture binder,
  - Adding substrate of DPP3 to said separated DPP3,
  - Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.
6. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 5, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.
7. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 5 or 6, wherein said capture-binder is an antibody.
8. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-7, wherein said capture binder is immobilized on a surface.

9. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-8, wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.

5

10. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-9, wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

15

11. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-10, wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or di-peptides coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.

20

12. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-11, wherein said substrate may be selected from the group comprising: a di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.

25

13. Method for monitoring a disease or condition in a subject accompanied by or related to necrotic processes wherein the method of diagnosing according to any of claims 1 to 12 is conducted at least twice.

30

14. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes.

15. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to claim 14 wherein said inhibitor is selected from the group comprising an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold.

16. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to claim 14 or 15 wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF),, myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), or SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases,vascular diseases (e.g. Kawasaki syndrome) and hypotension.

17. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claim 14 to 16 wherein said inhibitor is an antibody that is mono-binding or at least two-binding.

18. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 17 wherein said inhibitor is an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 1, in particular that binds to SEQ ID No. 2.

19. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 18 wherein said inhibitor is an antibody or fragment or scaffold that exhibits a minimum binding affinity to DPP3 of equal or less than  $10^{-7}$  M.

20. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any

claims 14 to 19 wherein said inhibitor is an antibody or fragment or scaffold that is monospecific.

21. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
5 condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 20 wherein said inhibitor is an antibody or fragment or scaffold binds to  
full-length DPP3 and inhibits activity of DPP3 of at least 10%, or at least 50%, more  
preferred at least 60%, even more preferred more than 70 %, even more preferred more  
than 80 %, even more preferred more than 90 %, even more preferred more than 95 %.

10 22. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 21 wherein said inhibitor is selective and specific to DPP3 and not cross  
cell membranes and/or the blood brain barrier.

15 23. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 22 wherein said subject has an amount of total DPP3 and/or an amount of  
20 active DPP3 in a sample of bodily fluid of said subject that is above a predetermined  
threshold.

25 24. Pharmaceutical composition comprising an inhibitor of the activity of DPP3 according  
to any of the claims 14 to 23 for use in prevention or therapy of a disease or condition in  
a subject accompanied by or related to necrotic processes.

25 25. Use of an inhibitor of the activity of DPP3 according to any of the claims 14 to 23 in a  
method of extracorporeal removal of DPP3 from plasma comprising apheresis and  
affinity chromatography.

30 26. Method for determining active DPP3 in a bodily fluid sample of a subject comprising  
the steps:

- Contacting said sample with a capture-binder that binds specifically to full-length DPP3,
- Separating DPP3 bound to said capture binder,
- Adding substrate of DPP3 to said separated DPP3,
- Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.

- 5            27. Method for determining active DPP3 in a bodily fluid sample of a subject according to claim 26 wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.
- 10            28. Method for determining active DPP3 in a bodily fluid sample of a subject according to claims 26 or 27 wherein said capture-binder is an antibody.
- 15            29. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 28 wherein said capture binder is immobilized on a surface.
- 20            30. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 29 wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.
- 25            31. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 30 wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).
- 30

32. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 31, wherein said substrate may be selected from the group comprising: A di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.

5

33. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 32, wherein said sample is a blood sample selected from the group comprising whole blood, serum and plasma.

10 34. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject comprising:

- A capture-binder that binds specifically to full-length DPP3,
- A substrate of DPP3.

15

35. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 34, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

20 36. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 34 or 35, wherein said capture-binder inhibits less than 50% DPP3 activity in a liquid phase assay, preferably less than 40%, more preferably less than 30 %.

25 37. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 36, wherein the binding region of DPP3 for the capture-binder is not within the region of amino acids 316 - 669 of SEQ Id No. 1.

38. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 37 wherein said binder is an antibody.

30

39. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 38 wherein said capture-binder is immobilized on a surface.

40. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 39 wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or a di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the preferred di-peptide is Arg-Arg.
- 5
41. Use of a method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26-33 or use of an assay or kit according to any of claims 34 to 40 in a method of for diagnosing or monitoring a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1 -13.
- 10

## EXAMPLES

15 1. **Example 1**

DPP3 activity in plasma of a variety of diseased patients (patients with acute myocardial infarction (AMI), cardiogenic shock, septic shock and liver failure) was determined using a specific DPP3 capture activity assay and compared to DPP3 plasma activities of healthy controls.

20  
1.1. **Study cohort:**

Plasma samples were obtained from 388 patients entering the emergency department directly at their first presentation. With their final diagnosis these patients could be divided into 4 subgroups: patients suffering from acute myocardial infarction (AMI), cardiogenic shock, septic shock and liver failure. The control group was a collection of 25 plasma samples from 93 healthy controls.

30  
1.2. **hDPP3 capture activity assay:**

DPP3 of plasma samples (10  $\mu$ l) is firstly enriched by an affinity purification step and secondly its activity measured by addition of the fluorogenic substrate Arg-Arg- $\beta$ NA (for a detailed description see example 4). The calculated slopes (in nmol  $\beta$ NA/ min per

ml sample [nmol βNA min<sup>-1</sup> ml<sup>-1</sup>]) of increasing fluorescence of the different samples refer to the 10 µl sample size.

### 1.3. Results:

5 Comparison of patient samples and healthy controls showed significantly higher DPP3 activity values for all patients with severe diseases or organ failure (Fig. 10)

## 2. Example 2

10 In this experiment, the effect of recombinant hDPP3 injection in healthy rats was studied by monitoring blood pressure.

### 2.1. Method:

15 2-3 months old male Wistar rats (Charles River Laboratories, Germany) were used for this study. For measurement and recording of the blood pressure (BP) a catheter (Introcan-W; 22 G/1"; B.Braun) was inserted into the Arteria carotis communis dextra (right common carotid artery). Human recombinant dipeptidyl peptidase 3 with an N-terminal GST-tag (recGST-hDPP3) was injected via the tail vain.

20 At first the animals were anesthetized with Isoflurane for weighing (whole g) and intraperitoneal (i.p.) injection of 1.2 g / kg BW Urethane (c = 0.4 g / mL) for long-term anesthesia. Then the ventral region of the neck was shorn and wiped off with ethanol. The vessels were prepared and the catheters were inserted. In the end both catheters are flushed with heparinized isotonic sodium chloride solution. Then the pressure transducers (medex logical, Medex Medical Ltd.) were connected with the patient monitoring system (Datex-Ohmeda, GE). A laptop computer via S/5 collect software, which was connected to the BP monitor, separately recorded the BP data.

25 Rats were treated with recGST-hDPP3 0.2 mg/ kg in PBS by injection into the tail vain. Blood pressure was constantly monitored before and after injection of DPP3.

## 2.2. Results:

Injection of recombinant GST-hDPP3 in healthy rats leads to an instant decrease of blood pressure (Fig. 11).

5

## 3. Example 3

Generation of antibodies and determination DPP3 binding ability: Several murine antibodies were produced and screened by their ability of binding human DPP3 in a sandwich or activity assay (see table 3).

10

## 3.1. Methods:

### - Peptides/ conjugates for immunization:

DPP3 peptides for immunization were synthesized, see table 3, (JPT Technologies, Berlin, Germany) with an additional N-terminal cystein (if no cystein is present within the selected DPP3-sequence) residue for conjugation of the peptides to Bovine Serum Albumin (BSA). The peptides were covalently linked to BSA by using Sulfolink-coupling gel (Perbio-science, Bonn, Germany). The coupling procedure was performed according to the manual of Perbio. Recombinant GST-hDPP3 was produced by USBio.

20

### - Immunization of mice, immune cell fusion and screening:

Balb/c mice were intraperitoneally (i.p.) injected with 84 µg GST-hDPP3 or 100 µg DPP3-peptide-BSA-conjugates at day 0 (emulsified in TiterMax Gold Adjuvant), 84 µg or 100 µg at day 14 (emulsified in complete Freund's adjuvant) and 25 42 µg or 50 µg at day 21 and 28 (in incomplete Freund's adjuvant). At day 49 the animal received an intravenous (i.v.) injection of 42 µg GST-hDPP3 or 50 µg DPP3-peptide-BSA-conjugates dissolved in saline. Three days later the mice were sacrificed and the immune cell fusion was performed.

30 Splenocytes from the immunized mice and cells of the myeloma cell line SP2/0 were fused with 1 ml 50% polyethylene glycol for 30 s at 37°C. After washing, the cells were seeded in 96-well cell culture plates. Hybrid clones were selected by growing in HAT medium [RPMI 1640 culture medium supplemented with 20% fetal calf serum and

HAT-Supplement]. After one week, the HAT medium was replaced with HT Medium for three passages followed by returning to the normal cell culture medium.

The cell culture supernatants were primarily screened for recombinant DPP3 binding IgG antibodies two weeks after fusion. Therefore recombinant GST-tagged DPP3 (USBiologicals, Salem, USA) was immobilized in 96-well plates (100 ng/ well) and incubated with 50 µl cell culture supernatant per well for 2 hours at room temperature. After washing of the plate, 50 µl / well POD-rabbit anti mouse IgG was added and incubated for 1 h at RT. After a next washing step, 50 µl of a chromogen solution (3,7 mM o-phenylenediamin in citrate/ hydrogen phosphate buffer, 0.012% H<sub>2</sub>O<sub>2</sub>) were added to each well, incubated for 15 minutes at RT and the chromogenic reaction stopped by the addition of 50 µl 4N sulfuric acid. Absorption was detected at 490 nm.

The positive tested microcultures were transferred into 24-well plates for propagation. After retesting the selected cultures were cloned and recloned using the limiting-dilution technique and the isotypes were determined.

- **Mouse monoclonal antibody production**

Antibodies raised against GST-tagged human DPP3 or DPP3-peptides were produced via standard antibody production methods (Marx et al., 1997) and purified via Protein A. The antibody purities were ≥ 90% based on SDS gel electrophoresis analysis.

- **Characterization of antibodies – hDPP3-inhibition analysis**

To analyze the capability of DPP3 inhibition by the different antibodies and antibody clones a DPP3 activity assay with known procedure (Jones et al., 1982) was performed. Recombinant GST-tagged hDPP3 was diluted in assay buffer (25 ng/ ml GST-DPP3 in 50 mM Tris-HCl, pH7,5 and 100 µM ZnCl<sub>2</sub>) and 200 µl of this solution incubated with 10 µg of the respective antibody at room temperature. After 1 hour of pre-incubation, fluorogenic substrate Arg-Arg-βNA (20 µl, 2mM) was added to the solution and the generation of free βNA over time was monitored using the Twinkle LB 970 microplate fluorometer (Berthold Technologies GmbH & Co. KG) at 37 °C. Fluorescence of βNA is detected by exciting at 340 nm and measuring emission at 410 nm. Slopes (in RFU/ min) of increasing fluorescence of the different samples are calculated. The slope of

GST-hDPP3 with buffer control is appointed as 100 % activity. The inhibitory ability of a possible capture-binder is defined as the decrease of GST-hDPP3 activity by incubation with said capture-binder in percent. The resulting lowered DPP3 activities are shown in figure 1a and table 3.

5

### 3.2. Results:

The following table represents a selection of obtained antibodies and their maximum inhibition rate (table 3). The monoclonal antibodies raised against the below depicted

10 DPP3 regions, were selected by their ability to bind native DPP3 (mAb-FL-DPP3\_2555 as solid phase and \_2553 as tracer for the immuno assay, for details see example 4).

For the immobilized DPP3 activity assay (see examples 6 and 7) it was necessary to select a solid phase antibody that does not inhibit DPP3 activity too strongly. As a cut off for antibody screening, solid phase antibodies should not inhibit DPP3 activity by more than 50 %. Since mAbDPP3\_2555 showed the lowest inhibition rate (table 3, figure 1 A).

15 For the generation of a strong DPP3 inhibitor that can be used therapeutically (see examples 8 - 13), it was necessary to select a DPP3 binder that shows the highest inhibition rate. The monoclonal antibody mAbDPP3\_1967, with the ability of inhibiting DPP3 activity by 70 %, was chosen as possible therapeutic antibody (see figure 1 A and table 3) and all further analysis was carried out using this antibody. Figure 1 B shows the inhibitory curve of mAbDPP3\_1967 which has an IC<sub>50</sub> of 0.2041 µg/ ml.

20  
25

Sequence number	Antigen/ Immunogen	hDPP3 region	Designation	Clone	Max inhibition of DPP3
<b>SEQ ID: 1</b>	GST tagged recombinant FL-DPP3	1-737	mAb-FL-DPP3	2552	65%
				2553	35%
				2554	30%
				2555	25%
<b>SEQ ID: 2</b>	CETVINPETGEQIQSWYRSGE	474-493	mAb-pepl-DPP3	1963	60%
				1964	60%
				1965	70%
				1966	65%
				1967	70%
				1968	65%
				1969	70%

**Table 3: Immunogen sequence, designation and characteristics of produced anti-DPP3 antibodies**

#### 4. Example 4

Determination of the antibody combination that yields high signal/ noise ratios in an hDPP3 immunoassay.

##### 4.1. Methods:

- **Monoclonal antibody production**

Antibodies raised against GST-tagged human DPP3 were produced via standard antibody production methods (Marx et al., 1997) and purified via Protein A. The antibody purities were  $\geq 90\%$  based on SDS gel electrophoresis analysis. Different clones were analyzed in their capability of binding DPP3. Resulting positive clones were used as solid phase or tracer antibodies.

- **Solid phase**

96-Well polystyrene microplates (Greiner Bio-One International AG, Austria) were coated (1 h at room temperature) with one antiDPP3 antibody clone (catching antibody; 1.5 µg antibody/ 0.25 mL 100 mmol/ L NaCl, 50 mmol/ L Tris/ HCl, pH 7.8). After blocking with 5% bovine serum albumine, the microplates were vacuum dried.

- **Labelling procedure (tracer)**

100 µg (100 µl) of a different antiDPP3 antibody (detection antibody, 1 mg/ ml in PBS, pH 7.4) were mixed with 10 µl acridinium NHS-ester (1 mg/ml in acetonitrile, InVent GmbH, Germany; EP 0 353 971) and incubated for 30 min at room temperature. Labelled antiDPP3 antibody was purified by gel-filtration HPLC on Shodex Protein 5 µm KW-803 (Showa Denko, Japan). The purified labeled antibody was diluted in assay buffer (50 mmol/l potassiumphosphate, 100 mmol/l NaCl, 10 mmol/l Na<sub>2</sub>-EDTA, 5 g/l bovine serum albumin, 1 g/l murine IgG, 1 g/l bovine IgG, 50 µmol/l amastatin, 100 µmol/l leupeptin, pH 7.4). The final concentration was approx. 7\*10<sup>6</sup> relative light units (RLU) of labelled compound (approx. 20 ng labeled antibody) per 200 µl. acridinium ester chemiluminescence was measured by using a Centro LB 960 luminometer (Berthold Technologies GmbH & Co. KG).

- **Calibrators**

A stock solution (in PBS, pH 7.4) of recombinant human GST- DPP3 (USBiological, USA) was linearly diluted using (50 mmol/ L potassium phosphate, 100 mmol/ L NaCl, 10 mmol/ L Na-EDTA, 5 g/ L bovine serum albumin, 1 g/ L murine IgG, 1 g/ L bovine IgG, 50 µmol/ L amastatin, 100 µmol/ L leupeptin, pH 7.4) The stock solution was stored at -80 °C. Calibrators were prepared before use.

- **hDPP3 immunoassay**

30 10 µl of sample (or calibrator) were pipetted into coated 96-well microplates, after adding labeled and diluted detection antibody (200 µl), the plates were incubated for 18-24 h at 2-8 °C. Unbound tracer was removed by washing 4 times with 350 µl washing solution (20 mM PBS, pH 7.4, 0.1 % Triton X-100). Well-bound

chemiluminescence was measured by using the Centro LB 960 luminometer (Berthold Technologies GmbH & Co. KG).

#### 4.2. Results:

5 All antibodies were used in a sandwich immunoassay, as coated microplate and labeled antibody, combined in the following variations (tables 4 and 5). Incubation was performed as described under hDPP3-Immunoassay. Results are given in ratio of specific signal / background signal for recombinant human GST-DPP3 (at 100, 10 and  
10 1 ng/ml) and native hDPP3 in plasma samples.

SP-AB	Tracer				100 ng/mL 10 ng/mL 1 ng/mL
	2552	2553	2554	2555	
2552	-	5.210 496 46	258	8.882 815 76	
2553	2.403 169 15	-	1.773 86 8	3.406 306 29	
2554	428 31 4	3.320 300 30	-	8.388 750 71	
2555	3.819 273 25	1.289 115 10	1.906 102 8	-	

Table 4: Signal/ noise ratio in antiDPP3 antibody pairs – measurement of recombinant GST-hDPP3 (SP – solid phase).

SP-AB	Tracer				nat. Sample 1 nat. Sample 2
	2552	2553	2554	2555	
2552	-	2.042 1.499	2	2.302 1.579	
2553	712 475	-	424 362	1.163 977	
2554	5 5	631 367	-	1.087 527	
2555	1.513 894	737 527	702 400	-	

Table 5: Signal/ noise ratio in antiDPP3 antibody pairs – measurement of human plasma samples (SP – solid phase).

All combinations displayed good signal/ noise ratios for recombinant GST-hDPP3. Furthermore, all combinations except for 2552 and 2554 yielded a good signal/ noise ratios for the native samples. Consequently, all remaining combinations can be used for further investigations. Concerning the highest absolute RLU signal, we used 2555 as solid phase antibody and 2553 as labeled antibody.

## 5. Example 5

5 DPP3 concentration in plasma of a variety of diseased patients (patients with acute heart failure (AHF), myocardial infarct (MI), sepsis, cancer, acute kidney injury (AKI) and lower respiratory tract infections (LRTI)) was determined using an hDPP3 immuno assay and compared to DPP3 plasma concentrations of healthy controls.

### 5.1. Study cohort:

10 Plasma samples were obtained from 214 patients entering the emergency department or the oncology department directly at their first presentation. With their final diagnosis these patients could be divided into 6 subgroups: patients suffering from acute heart failure (AHF), myocardial infarct (MI), sepsis, cancer, acute kidney injury (AKI) and lower respiratory tract infections (LRTI). The control group was a collection of 15 plasma samples from 93 healthy controls.

### 5.2. hDPP3 immuno assay:

20 mAbDPP3\_2555 was used as solid phase antibody and mAbDPP3\_2553 as labelled tracer antibody. Antibody immobilization, labelling and incubation were performed as described in example 4.

### 5.3. Results:

25 Together with the corresponding diagnostic details, the generated data was statistically analyzed (fig. 2 A). All patients showed significantly elevated DPP3 plasma concentrations compared to healthy controls (norm). Table 6 shows the percentage of patients with DPP3 values above the 75-percentile of the control group and their respective diagnosis. Analysis of plasma DPP3 levels indicates the disease status of a 30 patient. This revelation can be used in fields of diagnosis and also built the basis for therapeutic treatment, e.g. by inhibition of DPP3.

indication	percentage of patients with DPP3 values above 75-percentile of control group					
	AHF	MI	sepsis	cancer	AKI	LRTI
Sandwich assay	48,7	64,3	85,5	91,7	51,4	53,3

**Table 6: Comparison of DPP3 values of diseased patients and healthy controls in a sandwich type immune assay.**

The same study cohort was analyzed by their mortality. Patients who died after admission to the emergency department had significantly higher plasma DPP3 levels than emergency patients who survived in hospital. Thus elevated DPP3 concentrations indicate a bad prognosis concerning mortality (fig. 2 B).

## 6. Example 6

Not only can the amount of DPP3 in human plasma be determined by DPP3 concentrations but also by activity assays. One standard procedure is a soluble activity assay using Arg-Arg-βNA as fluorogenic substrate:

The activity of native human DPP III was determined by the hydrolysis of Arg-Arg-β-naphthylamide (Bachem Holdig AG, Switzerland) to form the fluorescent β-naphthylamine. 200 µl buffer (50 mM TRIS/HCl, pH 8.8, 0.04% NaN3, 50 µM Amastatin, 100 µM Leupeptin) and 10 µl of sample (human plasma) was pipetted into black 96-well microplates (Greiner Bio-One International GmbH, Austria)) and pre-tempered at 37 °C for 10 minutes. After adding substrate (20 µl, 2 mM), the increase of fluorescence was monitored for 1 h at 37 °C in a Twinkle LB 970 microplate fluorometer (Berthold Technologies GmbH & Co. KG) using excitation wavelengths of 340 nm an emission wavelengths of 410 nm. Slopes of increasing fluorescence of the different samples are calculated in nmol βNA/ min per ml sample [nmol βNA min<sup>-1</sup> ml<sup>-1</sup>] referring to the calibrator free βNA.

With the described standard, soluble, DPP3 activity assay it can not be determined whether one measures DPP3 activity or the activity of other amino peptidases in the plasma. In order to generate signals specific for DPP3 an enzyme capture assay is

performed, where in a first step DPP3 is immobilized to a surface via binding to a monoclonal antibody, and after a washing step, only specific DPP3 activity can be measured:

5       The solid phase was prepared as described in Example 5 using black 96-well microplates (Greiner Bio-One International GmbH, Austria). 10 µl sample (plasma or standard) and 200 µl buffer (50 mmol/l potassiumphosphate, 100 mmol/l NaCl, 5 g/l bovine serum albumin, 1 g/l murine IgG, 1 g/l bovine IgG, 50 µmol/l Amastatin, 100 µmol/l Leupeptin, pH 7.4) were pipetted into said coated microplates and incubated  
10      (18-24 h, 2-8 °C, 600 rpm). Unbound analyte was removed by washing (3 x 350 µl) with washing solution. After adding substrate (200 µl, 100 µM, in 50 mM Tris/HCl, pH (25 °C) 8.8, 0.04% NaN<sub>3</sub>), the increase of fluorescence was monitored for 1 h at 37 °C in Twinkle LB 970 microplate fluorometer (Berthold Technologies GmbH & Co. KG) using excitation wavelengths of 340 nm an emission wavelengths of 410 nm. Slopes  
15      of increasing fluorescence of the different samples are calculated in nmol βNA/ min per ml sample [nmol βNA min<sup>-1</sup> ml<sup>-1</sup>] referring to the calibrator free βNA.

In each activity assay type free βNA was used as assay calibrator. Therefor increasing concentrations of βNA (in 200 µl, in 50 mM Tris/HCl, pH (25 °C) 8.8, 0.04% NaN<sub>3</sub>; 0, 20      4, 8, 16, 32, 64, 125, 250 µM βNA) were measured in the Twinkle LB 970 microplate fluorometer (Berthold Technologies GmbH & Co. KG) at 37 °C using excitation wavelengths of 340 nm an emission wavelengths of 410 nm. All sample measurements  
15      were calibrated on this βNA standard.

25      7. Example 7

DPP3 activity in plasma of a variety of diseased patients (patients with acute heart failure (AHF), sepsis, acute kidney injury (AKI) and lower respirational tract infections (LRTI)) was determined using the specific DPP3 capture activity assay and compared to DPP3 plasma activities of healthy controls.

### 7.1. Methods:

Parts of the cohort analyzed in example 5 were subjected to a DPP3 specific enzyme capture activity assay. In this assay DPP3 of plasma samples (10 µl) is firstly enriched by an affinity purification step and secondly its activity measured by addition of the fluorogenic substrate Arg-Arg-βNA (for a detailed description see example 6). The calculated slopes (in nmol βNA/ min per ml sample [nmol βNA min<sup>-1</sup> ml<sup>-1</sup>]) of increasing fluorescence of the different samples refer to the 10 µl sample size.

### 10 7.2. Results:

Comparison of patient samples and healthy controls showed significantly higher DPP3 activity values for all patients (AHF, sepsis, AKI and LTRI, fig. 3).

15 Table 7 shows the percentage of patients with DPP3 values above the 75-percentile of the control group and their respective diagnosis. The activity shows a better division of healthy controls and diseased patients.

	percentage of patients with DPP3 values above 75-percentile of control group					
Indication	AHF	MI	sepsis	cancer	AKI	LRTI
Sandwich assay	48,7	64,3	85,5	91,7	51,4	53,3
Activity assay	77,8	n.a.	87,0	n.a.	70,0	85,7

20 **Table 7: Comparison of DPP3 values of diseased patients and healthy controls in a sandwich type immune assay and in an enzyme capture activity assay.**

To even better compare the DPP3 activity assay with the concentration assay we performed an ROC (Receiver Operating Characteristic) analysis for the division of healthy controls from AHF (fig. 4 A) or sepsis patients (fig. 4 B). The values of area under the curve (AUC) and confidence interval (CI) are depicted in table 8. Data analysis revealed a higher specificity for the activity assay compared to the sandwich immune assay.

	value	Sandwich assay	Activity assay
Normal vs. AHF	AUC	0.6747	0.8506
	95 % CI	0.5638 - 0.7853	0.7482 - 0.9530
	p value	< 0.005	< 0.0001
<hr/>			
Normal vs. SOD1G93A	AUC	0.8875	0.9384
	95 % CI	0.8270 - 0.9479	0.8865 - 0.9479
	p value	< 0.0001	< 0.0001

Table 8: Data of ROC analysis (AUC – area under the curve; CI – confidence interval).

## 8. Example 8

In this experiment, the general safety of increasing dosages of mAbDPP3 in healthy mice was monitored.

### 8.1. Method:

Female BALB/c nude (CAnN.Cg-Foxn1nu/Crl) Mice (Charles River GmbH, Sulzfeld, Germany) aged 4-5 weeks at delivery and weighing approximately 15-18 g were kept under optimum hygienic conditions, air-conditioned with 10-15 air changes per hour, and continually monitored environment with target ranges for temperature  $22 \pm 3^{\circ}\text{C}$  and for relative humidity 30-70%, 12 hours artificial fluorescent light / 12 hours dark. Maximum 4 animals were kept per individual ventilated cage (IVC) and fed with a diet consisting of M-Zucht (ssniff Spezialdiäten GmbH) and autoclaved community tab water.

After an acclimatization period of 4 days application of mAbDPP3 was started: Three different mAbDPP3 concentrations (0.65 mg/ kg, 1.9 mg/ kg and 5.75 mg/ kg) in PBS were injected into 4 mice per group. MAbDPP3 was administered intraperitoneally (i.p.) at days 1, 3, 5 and 7, and mice monitored for 14 days.

**8.2. Results:**

All mice survived the 14 day treatment with no side effects, also at the highest dosage. MAbDPP3 is safe to be used in other animal experiments, which will always be performed with a concentration of 1.9 mg/ kg.

**9. Example 9**

In this experiment, the general safety of mAbDPP3 treatment in healthy rats was studied by monitoring mean blood pressure.

10  
15  
2-3 months old male Wistar rats (Charles River Laboratories, Germany) were used for this study. For measurement and recording of the blood pressure (BP) a catheter (Introcan-W; 22 G/1"; B.Braun) was inserted into the Arteria carotis communis dextra (right common carotid artery). The administration and sampling catheter were inserted into the Vena jugularis sinistra (left jugular vein).

20  
25  
At first the animals were anesthetized with Isoflurane for weighing (whole g) and intraperitoneal (i.p.) injection of 1.2 g / kg BW Urethane ( $c = 0.4$  g / mL) for long-term anesthesia. Then the ventral region of the neck was shorn and wiped off with ethanole. The vessels were prepared and the catheters were inserted. In the end both catheters are flushed with heparinized isotonic sodium chloride solution. Then the pressure transducers (medex logical, Medex Medical Ltd.) were connected with the patient monitoring system (Datex-Ohmeda, GE). A laptop computer via S/5 collect software, which was connected to the BP monitor, separately recorded the BP data.

30  
Rats were treated with PBS, 1.9 mg/ kg and 5.75 mg/ kg mAbDPP3 in PBS ( $n = 3$  per group). The compounds were applied via the venous catheter. Blood pressure was monitored for 1 h prior to application of the compounds and over 6 h after application of compounds.

**9.2. Results:**

Rats reacted well to the mAbDPP3 treatment. Rats treated with the high dosage of mAbDPP3 showed a slight increase in mean blood pressure (fig. 5). In general the mAbDPP3 is also safe to use in a rat model even in higher dosages.

**10. Example 10**

In this study, it was analyzed how treatment with mAbDPP3 can influence sepsis mortality in a CLP mouse model.

10

**10.1. Methods:**

12-15 week old male C57Bl/6 mice (Charles River Laboratories, Germany) were used for the study. Peritonitis had been surgically induced under light isofluran anesthesia. Incisions were made into the left upper quadrant of the peritoneal cavity (normal location of the cecum). The cecum was exposed and a tight ligature was placed around the cecum with sutures distal to the insertion of the small bowel. One puncture wound was made with a 24-gauge needle into the cecum and small amounts of cecal contents were expressed through the wound. The cecum was replaced into the peritoneal cavity and the laparotomy site was closed. Finally, animals were returned to their cages with free access to food and water. 500µl saline were given s.c. as fluid replacement.

20

MAbDPP3 (1,9 mg/ kg in PBS) was tested versus vehicle (PBS). I.v. injection of the compound and vehicle was performed 5 minutes before CLP (preventive treatment) and after full development of sepsis, 2 hours after CLP (therapeutic treatment). Each group contained 10 mice and were followed over a period of 7 days.

25

**10.2. Results:**

30

It can be seen from figure 6 that the mAbDPP3 antibody reduced mortality considerably as compared to PBS application. After 4 days 75 % of the mice survived when treated with mAbDPP3. In contrast thereto almost all mice were dead after 4 days when treated with vehicle.

## 11. Example 11

A septic shock model was used to induce heart failure in rats and then to characterize mAbDPP3's influence on heart function.

5

### 11.1. Methods:

#### - **Study design**

The study flow is depicted in Figure 7 A below. After CLP or sham surgery the animals were allowed to rest for 20 hours with free access to water and food. Afterwards they were anesthetized, tracheotomy done and arterial and venous line laid. At 24 hours after CLP surgery either mAbDPP3 or vehicle (saline) were administered with 2 mg/kg. Hemodynamics were monitored invasively and continuously from  $t = 0$  till 3 h. Echocardiography of the heart was performed just after surgery, 15 minutes, 1 h, 2 h and 3 h after mAbDPP3 or saline injection.

#### - **CLP model of sepsis**

Male Wistar rats (2-3 months, 300 to 400 g, group size refer to Table 1) from the Centre d'élevage Janvier (France) were allocated randomly to one of three groups. All the animals were anesthetized using ketamine hydrochloride (90 mg/ kg) and xylazine (9 mg/ kg) intraperitoneally (i.p.). For induction of polymicrobial sepsis, cecal ligation and puncture (CLP) was performed using Rittirsch's protocol with minor modification. A ventral midline incision (1.5 cm) was made to allow exteriorization of the cecum. The cecum is then ligated just below the ileocecal valve and punctured once with an 18-gauge needle. The abdominal cavity is then closed in two layers, followed by fluid resuscitation (3 ml/ 100 g body of weight of saline injected subcutaneously) and returning the animal to its cage. Sham animals were subjected to surgery, without getting their cecum punctured.

#### - **Invasive Blood Pressure**

Hemodynamic variables were obtained using the AcqKnowledge system (BIOPAC Systems, Inc., USA). It provides a fully automated blood pressure analysis system. The catheter is connected to the BIOPAC system through a pressure sensor.

For the procedure, rats were anesthetized (ketamine and xylazine). Animals were moved to the heating pad for the desired body temperature to 37–37.5 °C. The temperature feedback probe was inserted into the rectum. The rats were placed on the operating table in a supine position. The trachea was opened and a catheter (16G) was inserted for an external ventilator without to damage carotid arteries and vagus nerves. The arterial catheter was inserted into the right carotid artery. The carotid artery is separate from vagus before ligation.

A central venous catheter was inserted through the left jugular vein allowing administration of drug.

Following surgery, the animals were allowed to rest for the stable condition prior to hemodynamic measurements. Then baseline blood pressure (BP) were recorded. During the data collection, saline infusion via arterial line was stopped.

- **Echocardiography**

Animals were anesthetized using ketamine hydrochloride. Chests were shaved and rats were placed in decubitus position.

For transthoracic echocardiographic (TTE) examination a commercial GE Healthcare Vivid 7 Ultra-sound System equipped with a high frequency (14-MHz) linear probe and 10-MHz cardiac probe was used. All examinations were recorded digitally and stored for subsequent off-line analysis.

Grey scale images were recorded at a depth of 2 cm. Two-dimensional examinations were initiated in a parasternal long axis view to measure the aortic annulus diameter and the pulmonary artery diameter. We also employed M-mode to measure left ventricular (LV) dimensions and assess fractional shortening (FS%). LVFS was calculated as LV end-diastolic diameter - LV end-systolic diameter / LV end-diastolic diameter and expressed in %. The time of end-diastole was therefore defined at the maximal diameter of the LV. Accordingly, end-systole was defined as the minimal

diameter in the same heart cycle. All parameters were measured manually. Three heart cycles were aver-aged for each measurement.

From the same parasternal long axis view, pulmonary artery flow was recorded using pulsed wave Doppler. Velocity time integral of pulmonary artery outflow was measured.

From an apical five-chamber view, mitral flow was recorded using pulsed Doppler at the level of the tip of the mitral valves.

- **Experimentation time points and animals groups**
- Baseline BP and echocardiography were recorded after surgery. Then mAbDPP3 (2 mg/kg) or vehicle (saline) was injected (i.v., 5 minutes after surgery) and saline infusion was started. Hemodynamics points (BP and echocardiography) were registered 15 minutes after mAbDPP3 or vehicle injection, and 1, 2 and 3 hours after. There were 1 control group and 2 CLP groups which are summarized in the table below (table 9). At the end of the experiment, the animal was euthanized, blood was drawn for EDTA-plasma generation and organs harvested for subsequent analysis.

Group	Group size	CLP	Treatment
1 - sham	7	no	saline
2 - CLP-saline	7	yes	saline
3 - CLP-mAbDPP3	10	yes	mAbDPP3

20 **Table 9: Experimental groups**

### 11.2. Results:

Compared to sham animals the septic rats have very low blood pressure and a decreased shortening fraction of the heart. Application of mAbDPP3 significantly increased the shortening fraction (fig. 7 B), raised the mean blood pressure (fig. 7 C) and improved the well-being of septic rats tremendously.

**12. Example 12**

The aim of the herein described study was to assess the potential anti-proliferative effect of mAbDPP3 in an *in vitro* cell culture system employing several cancer cell lines.

**5 12.1. Method:**

A stock solution of mAbDPP3 (1 mg/ ml in PBS) was diluted to cover the final concentration range between 0 – 100 µg/ ml. PBS was used as reference compound. Cancer cells stemming from established cancer cell lines (A549, HCT116, MDA-MB231) were cultured in DMEM containing 10 % FCS and Penicillin/Streptomycin.

10 A549 cells are adenocarcinomic human alveolar basal epithelial cells. This cell line was first established through the removal and culturing of cancerous lung tissue in the explanted tumor of a 58-year-old male. In nature, these cells are squamous and responsible for the diffusion of some substances, such as water and electrolytes, across the alveoli of lungs. In case said A549 cells are cultured *in vitro*, they grow as monolayer cells, adhered to the culture flask. A further characteristic is that these cells are able to synthesize lecithin and contain high level of desaturated fatty acids. The A549 cell line is widely used as *in vitro* model for a type II pulmonary epithelial cell model for drug metabolism and as a transfection host.

15 HCT116 cell line denotes human colon cancer cells. These epithelial cells have adherent culture properties, and stem from a male adult. This cell line is a suitable transfection host. This line has a mutation in codon 13 of the ras proto-oncogene, and can be used as a positive control for PCR assays of mutation in this codon.

20 MDA-MB231 cell line denotes human breast adenocarcinoma cells having epithelial morphology. These cells were isolated from pleural effusions of Caucasian breast cancer patient.

30 For each cell line 96 well suspension cell culture plates were prepared. 100 µL of the soft agar bottom layer (0.6 % final concentration in complete medium) was poured and left to solidify. 50 µL of the soft agar top layer (0.4 % final concentration) containing

the corresponding cells and cell number were then added on top, solidified and such 96 well plates incubated overnight at 37°C, 10 % CO<sub>2</sub>.

Next day, compounds were added into the inner wells of the plate. Subsequently, the assays were incubated in cell culture incubators. Finally, the assays were developed using Alamar Blue and upon 3-5 h of incubation at 37 °C fluorescence intensity was determined (excitation: 560 nm; emission: 590 nm). As low control, cells were treated with 10 - 5 M staurosporine (6-fold values). As high control, cells were treated with 0.1 % DMSO (solvent control, 6-fold values).

Raw data were converted into percent soft agar growth relative to high controls (solvent 0.1 % DMSO) and low controls (10 - 5 M staurosporine), which were set to 100 % and 0 %, respectively. IC<sub>50</sub> calculation was performed using GraphPad Prism 5 software with a variable slope sigmoidal response-fitting model using 0 % soft agar growth as bottom constraint and without bottom constraint and 100 % soft agar growth as top constraint.

## 12.2. Results:

In the cell culture system, growth of three cancer cell lines (A549, HCT116, MDA-MB231) was assessed depending on various doses of mAbDPP3 (fig. 8) applied. IC<sub>50</sub> values were determined using standard parameters based on the signal of the solvent control as top constraint (100 % soft agar growth) and the signal of the staurosporine control as bottom constraint (0 % soft agar growth). The respective IC<sub>50</sub> values are summarized in Table 10.

MAbDPP3 treatment had an antiproliferative effect on the three cell lines tested.

Cell line	Tissue source	Incubation time	IC <sub>50</sub>
A549	Lung	8 days	6.3 µg/ ml
HCT116	Colon	8 days	2.0 µg/ ml
MDA-MB231	Breast	11 days	8.7 µg/ ml

Table 10: IC<sub>50</sub> values for mAbDPP3 treatment.

**13. Example 13**

The aim of the herein described study was to assess the ability of mAbDPP3 to prevent tumor formation in xenograft models for breast cancer and colon cancer (tumor growth inhibition study).  
5

**13.1. Methods:**

Monolayer MDA-MB-231 cells (breast cancer) and HCT-116 cells (colon cancer),  
10 respectively, were grown in DMEM + 10% FCS. The cells were cultured in a humidified atmosphere of 90% air and 10% carbon dioxide at 37°C. Media was routinely changed every 3 days. Confluent cultures were split 1:3 to 1:3 every 3-4 days using Trypsin/EDTA and seeded at a density of approximately 3-4 x 10<sup>6</sup> cells/15 cm<sup>2</sup> + 25 mL medium.  
15

Female BALB/c nude (CAnN.Cg-Foxn1<sup>nu</sup>/Crl) Mice (Charles River GmbH, Sulzfeld, Germany) aged 4-5 weeks at delivery and weighing approximately 15-18 g were kept under optimum hygienic conditions, air-conditioned with 10-15 air changes per hour, and continually monitored environment with target ranges for temperature 22 ± 3°C and for relative humidity 30-70%, 12 hours artificial fluorescent light / 12 hours dark.  
20 Maximum 4 animals were kept per individual ventilated cage (IVC) and fed with a diet consisting of M-Zucht (ssniff Spezialdiäten GmbH) and autoclaved community tab water.

Human breast cancer MDA-MB-231 cells and colon cancer HCT-116 cells provided by ATCC will be used in this study. The cell subculture within 5 passages before inoculated into the mice. 3×10<sup>6</sup> cell /0.1mL will be s.c. injected into the right flank of mouse. When the tumor volume reaches about 100 - 200 mm<sup>3</sup>, the 20 mice bearing suitable size tumor will be randomized into grouping according to the tumor volume and body weight (10 mice per group). Animals will be dosed according to the tumor volume and body weight and start dosing. The groups are indicated in the table below (Table 11).  
25  
30

Group	Concen- tra-tion	Route	schema	Matrix	cell line	Observa- tion time	animal number
mAbDPP3 - breast	1,9 mg/ kg	i.v.	days 1, 3, 5, 7, 9	PBS	MDA- MB231	24 days	10
mAbDPP3 - colon					HCT116	28 days	10
PBS - breast	-	i.v.	days 1, 3, 5, 7, 9	PBS	MDA- MB231	24 days	10
PBS - colon					HCT116	28 days	10

**Table 11: Overview of treatment strategy.**

Animal behavior and welfare was observed daily and tumor growth was recorded every two days by calliper measurement over a period of 24 or days. Primary tumor sizes were measured by callipering (manual calliper, OMC Fontana). Tumor sizes were calculated according to the formula  $V = W^2 \times L / 2$  ( $L$ = length and  $W$ = the perpendicular width of the tumor,  $L>W$ ). The individual relative tumor volume (RTV) was calculated as follows:  $RTV = V_t / V_0$ , where  $V_t$  is the volume on each day, and  $V_0$  is the volume at the beginning of the treatment.

5

10

### 13.2. Results:

In the xenograft cancer models the application of mAbDPP3 reduced the formation of all tumors investigated (fig. 9 A - D). Under mAbDPP3 treatment it takes the breast cell 15 tumor 2 days longer to grow to 20-fold size (fig. 9 B) and the colon cell tumor 2 days longer to grow to 10-fold size (fig. 9 D). In this model, the tumors induced by the colon cancer cell line grew a lot slower than the ones induced by the breast cancer cell line.

### 14. Example 14

20

In order to assess the possibility of using a DPP3-adsorber to clear plasma from excess DPP3, it should be analyzed whether DPP3 would sufficiently bind to an antiDPP3 column. Affinity chromatography columns were prepared by immobilizing DPP3 binding antibodies to GlycoLink columns (Thermo Fisher). Binding of DPP3 to those columns was 25 analyzed by measuring DPP3 concentration before and after flow through these columns.

#### 14.1. Methods:

In a first step all DPP3 binding antibodies (mAbDPP3\_2552, 2553, 2554, 2555) were  
5 oxidized and immobilized on one GlycoLink column each according to the instructor's manual. Subsequently recombinant GST-hDPP3 (USBio) or patient plasma samples were loaded onto the column and binding of DPP3 monitored.

- **Oxidization of antibodies**

10 Therefore 300 µl of 3 mg/ ml of the respective antiDPP3 antibody solution were diluted in 700 µl GlycoLink Coupling Buffer to a final volume of 1 ml and pH < 6. To oxidize the carbohydrate groups of the antibodies, 2. 1 mg of sodium meta-periodate were added to the solution and incubated at room temperature for 30 minutes. The oxidizing agent was removed from the antibody solution using desalting columns.

15 - **Preparation of GlycoLink columns**

To catalyze coupling reaction 0.1 M aniline was added to the oxidized antibody. The solution was than applied onto an equilibrated GlycoLink column and incubated for 4 hours at room temperature. Unbound material was allowed to flow through the 20 column. Afterwards the column was washed, equilibrated and stored until further usage.

- **DPP3 affinity purification**

25 Either recombinant GST-hDPP3 or patient plasma was diluted in coupling buffer (final concentration of recDPP3 = 100 ng/ ml, 1 ml; 1 ml plasma + 1 ml coupling buffer), applied to the equilibrated mAbDPP3 column and incubated for 30 minutes at room temperature. The entire flow-through was saved to evaluate binding efficiency and capacity. The column was washed, bound protein eluted using GlycoLink Elution Buffer and the column equilibrated before storage.

30 - **Analysis of DPP3 content**

The DPP3 concentration of samples and recombinant GST-hDPP3 before affinity purification and of the flow through were measured using a sandwich type

luminescence immune assay (for details see example 4). 20 µl of sample (or calibrator) were pipetted into coated 96-well microplates, after adding labeled and diluted detection antibody (200 µl), the plates were incubated for 18-24 h at 2-8 °C. Unbound tracer was removed by washing 4 times with 350 µl washing solution (20 mM PBS, pH 7.4, 0.1 % Triton X-100). Well-bound chemiluminescence was measured by using the Centro LB 960 luminometer (Berthold Technologies GmbH & Co. KG).

#### 14.2. Results:

Almost all of DPP3, in plasma and in solutions of recombinant DPP3, is removed from the samples by affinity chromatography (see table 12). The different mAbDPP3 antibodies show different affinities towards DPP3. Since mAbDPP3\_2555 shows the highest binding rate of recDPP3, this antibody was chosen to be used in chromatography of plasma samples. Table 12 shows that DPP3 plasma levels can be strongly decreased by affinity chromatography. These results show, that it would be possible to use a DPP3-adsorber in apheresis to clear patients plasma off excess DPP3.

sample	immobilized antibody	c <sub>DPP3</sub> [ng/ml]	
		before GlycoLink	after GlycoLink
recombinant GST-hDPP3	2552	100	16,3
	2553	100	21,6
	2554	100	14,1
	2555	100	8,5
plasma_1	2555	18	1,2
plasma_2		26,1	4,3
plasma_3		37,5	3,7

Table 12: Concentrations of native or recombinant DPP3 in samples before and after affinity chromatography with depicted anti-DPP3 antibodies.

## REFERENCES

- Abramić, M. et al., 2000. Human and rat dipeptidyl peptidase III: Biochemical and mass spectrometric arguments for similarities and differences. *Biological Chemistry*, 381(December), pp.1233–1243.
- Abramić, M., Zubanović, M. & Vitale, L., 1988. Dipeptidyl peptidase III from human erythrocytes. *Biol Chem Hoppe Seyler*, pp.29–38.
- Agić, D. et al., 2007. Novel amidino-substituted benzimidazoles: Synthesis of compounds and inhibition of dipeptidyl peptidase III. *Bioorganic Chemistry*, 35(2), pp.153–169.
- Almagro JC, Fransson J., 2008. Humanization of antibodies. *Front Biosci.* 2008 Jan 1;13:1619-33.
- Aoyagi, T. et al., 1993. Enzymatic Changes in Cerebrospinal Fluid of Patients with Alzheimer-Type Dementia. *J Clin Biochem Nutr*, 14, pp.133–139.
- Balogun, R.A. et al., 2010. Clinical applications of therapeutic apheresis. *Journal of clinical apheresis*, 25(5), pp.250–64.
- Baršun, M. et al., 2007. Human dipeptidyl peptidase III acts as a post-proline-cleaving enzyme on endorphins. *Biological Chemistry*, 388(3), pp.343–348.
- Berghe, T. Vanden et al., 2014. Regulated necrosis: the expanding network of non-apoptotic cell death pathways. *Nature Reviews Molecular Cell Biology*, 15(2), pp.135–147.
- Bird et al., 1988. Single-chain antigen-binding proteins. *Science* 242:423-426.
- Chiba, T. et al., 2003. Inhibition of recombinant dipeptidyl peptidase III by synthetic hemorphin-like peptides. *Peptides*, 24(5), pp.773–778.
- Couston, R.G. et al., Adsorption behavior of a human monoclonal antibody at hydrophilic and hydrophobic surfaces. *mAbs*, 5(1), pp.126–39.
- Deng, B. et al., 2014. Aptamer binding assays for proteins: The thrombin example-A review. *Analytica Chimica Acta*, 837, pp.1–15.
- Dhanda, S., Singh, J. & Singh, H., 2008. Hydrolysis of various bioactive peptides by goat brain dipeptidylpeptidase-III homologue. *Cell biochemistry and function*, 26(3), pp.339–45.
- Dimitriades, V.R., Brown, A.G. & Gedalia, A., 2014. Kawasaki disease: pathophysiology, clinical manifestations, and management. *Current rheumatology reports*, 16(6), p.423.
- Ellis, S. & Nuenke, J.M., 1967. Dipeptidyl Arylamidase III of the Pituitary: Purification and Characterization. *Journal of Biological Chemistry*, 242(20), pp.4623–4629.

- Elmore, S., 2007. Apoptosis: A Review of Programmed Cell Death. *Toxicologic Pathology*, 35(4), pp.495–516.
- Fink, S.L. & Cookson, B.T., 2005. Apoptosis, Pyroptosis, and Necrosis: Mechanistic Description of Dead and Dying Eukaryotic Cells. *Infection and Immunity*, 73(4), pp.1907–1916.
- Fischer, U. & Schulze-Osthoff, K., 2005. New approaches and therapeutics targeting apoptosis in disease. *Pharmacological reviews*, 57(2), pp.187–215.
- Gamrekelashvili, J. et al., 2013. Peptidases released by necrotic cells control CD8 + T cell cross-priming. *journal of clinical investigation*, 123(11), pp.4755–4768.
- Hartley et al, 1982. *Radiology* 143: 29-36
- Holliger P. et al. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6444-8
- Hood et al., Immunology, Benjamin, N.Y., 2nd ed., 1984
- Hultschig C et al., *Curr Opin Chem Biol.* 2006 February; 10(1):4-10. PMID: 16376134)
- Hunkapiller & Hood, 1986. The growing immunoglobulin gene superfamily. *Nature* 323:15-16.
- Hust, M., Meyer, T., Voedisch, B., Rülker, T., Thie, H., El-Ghezal, A., Kirsch, M.I., Schütte, M., Helmsing, S., Meier, D., Schirrmann, T., Dübel, S., 2011. A human scFv antibody generation pipeline for proteome research. *Journal of Biotechnology* 152, 159–170
- Huston et al., 1988. Protein engineering of antibody binding sites: recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in Escherichia coli. *Proc. Natl. Acad. Sci. U.S.A.*, 85:5879-5883.
- Igic, R. & Behnia, R., 2007. Pharmacological, immunological, and gene targeting of the renin-angiotensin system for treatment of cardiovascular disease. *Current pharmaceutical design*, 13(12), pp.1199–214.
- Inaoka, Y. & Naruto, S., 1988. Propioxatins A and B, New Enkephalinase B Inhibitors, IV. Characterization of the Active Site of the Enzyme Using Synthetic Propioxatin Analogues. *J. Biochem*, 104(5), pp.706–711.
- Jones, T.H. & Kapralou, A, 1982. A rapid assay for dipeptidyl aminopeptidase III in human erythrocytes. *Analytical biochemistry*, 119(2), pp.418–23.
- Kabat et al., 1983. Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services.
- Khaket, T.P. et al., 2012. Enkephalin degrading enzymes: metalloproteases with high potential for drug development. *Current pharmaceutical design*, 18(2), pp.220–30.

- Kim, D. & Herr, A.E., 2013. Protein immobilization techniques for microfluidic assays. *Biomicrofluidics*, 7(4), p.41501.Kirk-Othmer, Encyclopedia of chemical technology, 4th ed., executive editor, J. I. Kroschwitz; editor, M. Howe-Grant, John Wiley & Sons, 1993, vol. 15, p. 518-562
- 5 Kumar, P. et al., 2016. Substrate complexes of human dipeptidyl peptidase III reveal the mechanism of enzyme inhibition. *Scientific Reports*, 6(March), p.23787.
- Lang, E. & Lang, F., 2015. Triggers, Inhibitors, Mechanisms, and Significance of Eryptosis: The Suicidal Erythrocyte Death. *BioMed Research International*, 2015, p.513518.
- Lanier, S.T. et al., Spatiotemporal progression of cell death in the zone of ischemia surrounding burns. *Wound repair and regeneration : official publication of the Wound Healing Society [and] the European Tissue Repair Society*, 19(5), pp.622–32.
- 10 Lanzavecchia, A. & Scheidegger, D., 1987. The use of hybrid hybridomas to target human cytotoxic T lymphocytes, *Eur. J. Immunol.* 17:105.
- Lee, C.M. & Snyder, S.H., 1982. Dipeptidyl-aminopeptidase III of rat brain. Selective affinity for enkephalin and angiotensin. *The Journal of biological chemistry*, 257(20), pp.12043–50.
- 15 Li, J. et al., 2011. Peptide aptamers with biological and therapeutic applications. *Current medicinal chemistry*, 18(27), pp.4215–22.
- Linkermann, A. & Green, D.R., 2014. Necroptosis. *New England Journal of Medicine*, 370(5), pp.455–465.
- 20 Liu, Y. et al., 2007. A genomic screen for activators of the antioxidant response element. *Proceedings of the National Academy of Sciences of the United States of America*, 104(12), pp.5205–10.
- Marx et al., 1997. Monoclonal Antibody Production, ATLA 25, 121.
- 25 Mazzocco, C. et al., 2006. Identification and characterization of two dipeptidyl-peptidase III isoforms in *Drosophila melanogaster*. *FEBS Journal*, 273(5), pp.1056–1064.
- Meliopoulos, V.A. et al., 2012. MicroRNA regulation of human protease genes essential for influenza virus replication. *PloS one*, 7(5), p.e37169.
- 30 Müller, J. et al., 2016. Aptamer-Based Enzyme Capture Assay for Measurement of Plasma Thrombin Levels. *Methods in molecular biology* (Clifton, N.J.), 1380, pp.179–89.
- Müller, J. et al., 2012. Monitoring of plasma levels of activated protein C using a clinically applicable oligonucleotide-based enzyme capture assay. *Journal of Thrombosis and Haemostasis*, 10(3), pp.390–398.

- Ohkubo, I. et al., 1999. Dipeptidyl peptidase III from rat liver cytosol: purification, molecular cloning and immunohistochemical localization. *Biological chemistry*, 380(12), pp.1421–1430.
- Patel, A., Smith, H.J. & Sewell, R.D., 1993. Inhibitors of enkephalin-degrading enzymes as potential therapeutic agents. *Progress in medicinal chemistry*, 30, pp.327–78.
- Pinheiro Da Silva, F. & Nizet, V., 2009. Cell death during sepsis: Integration of disintegration in the inflammatory response to overwhelming infection. *Apoptosis*, 14(4), pp.509–521.
- Ponikowski, P. et al., 2016. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *European Journal of Heart Failure*, 18(8), pp.891–975.
- 10 Prajapati, S.C. & Chauhan, S.S., 2011. Dipeptidyl peptidase III: a multifaceted oligopeptide N-end cutter. *FEBS Journal*, 278(18), pp.3256–3276.
- Raghupathi, R., 2004. Cell death mechanisms following traumatic brain injury. *Brain pathology (Zurich, Switzerland)*, 14(2), pp.215–22.
- 15 Rastija, V. et al., 2015. Synthesis, QSAR, and Molecular Dynamics Simulation of Amidino-substituted Benzimidazoles as Dipeptidyl Peptidase III Inhibitors. *Acta Chimica Slovenica*, 62, pp.867–878.
- Rittirsch, D., Huber-Lang, M., Flierl, M. Ward, P.: Immunodesign of experimental sepsis by cecal ligation and punc-ture, *Nature Protocols* 4, - 31 - 36 (2009)
- Sanderink, G.J., Artur, Y. & Siest, G., 1988. Human aminopeptidases: a review of the literature. *Journal of clinical chemistry and clinical biochemistry. Zeitschrift für klinische Chemie und klinische Biochemie*, 26(12), pp.795–807.
- 20 Schütte, M., Thullier, P., Pelat, T., Wezler, X., Rosenstock, P., Hinz, D., Kirsch, M.I., Hasenberg, M., Frank, R., Schirrmann, T., Gunzer, M., Hust, M., Dübel, S., 2009. Identification of a putative Crf splice variant and generation of recombinant antibodies for the specific detection of *Aspergillus fumigatus*. *PLoS One* 4, e6625
- Shimamori, Y., Watanabe, Y. & Fujimoto, Y., 1986. Purification and Characterization of Dipeptidyl Aminopeptidase III from Human Placenta. *Chem. Pharm. Bull.*, 34(8), pp.3333–3340.
- 25 Šimaga, Š. et al., 1998. Dipeptidyl peptidase III in malignant and non-malignant gynaecological tissue. *European Journal of Cancer*, 34(3), pp.399–405.
- Šimaga, Š. et al., 2003. Tumor cytosol dipeptidyl peptidase III activity is increased with histological aggressiveness of ovarian primary carcinomas. *Gynecologic Oncology*, 91(1), pp.194–200.

- Singh, R. et al., 2014. Transcription factor C/EBP-beta mediates downregulation of dipeptidyl-peptidase III expression by interleukin-6 in human glioblastoma cells. FEBS Journal, 281, pp.1629–1641.
- Vandenberg, I., King, F. & Kuchel, P., 1985. Enkephalin Degradation by Human Erythrocytes and Hemolysates Studied Using  $^1\text{H}$  NMR Spectroscopy. Archives of Biochemistry and Biophysics, 242(2), pp.515–522.
- Vanha-Perttula, T., 1988. Dipeptidyl peptidase III and alanyl aminopeptidase in the human seminal plasma: origin and biochemical properties. Clinica chimica acta; international journal of clinical chemistry, 177(2), pp.179–95.
- 10 Vanlangenakker, N. et al., 2008. Molecular mechanisms and pathophysiology of necrotic cell death. Current molecular medicine, 8(3), pp.207–20.
- Voors, A.A. et al., 2016. A systems BIOlogy Study to TAilored Treatment in Chronic Heart Failure: rationale, design, and baseline characteristics of BIOSTAT-CHF. European Journal of Heart Failure, 18(6), pp.716–726.
- 15 Wallach, D., Kang, T.-B. & Kovalenko, A., 2014. Concepts of tissue injury and cell death in inflammation: a historical perspective. Nature reviews. Immunology, 14(1), pp.51–9.
- Wattiaux, R. et al., 2007. Lysosomes and Fas-mediated liver cell death. The Biochemical journal, 403(1), pp.89–95.
- Wild, David (2005). The Immunoassay Handbook, Elsevier LTD, Oxford; 3rd ed., ISBN-13: 20 978-0080445267
- Xu, Y., Yang, X. & Wang, E., 2010. Review: Aptamers in microfluidic chips. Analytica chimica acta, 683(1), pp.12–20.
- Yamamoto, Y. et al., 2000. Characterization of tynorphin, a potent endogenous inhibitor of dipeptidyl peptidaseIII. Peptides, 21(4), pp.503–8.
- 25 Yamamoto, Y. et al., 1998. Inhibitory action of spinorphin, an endogenous regulator of enkephalin-degrading enzymes, on carrageenan-induced polymorphonuclear neutrophil accumulation in mouse air-pouches. Life sciences, 62(19), pp.1767–73.
- Zong, W.-X. & Thompson, C.B., 2006. Necrotic Cell Death as a Cell Fate. Genes & Development, 20, pp.1–5.
- 30 Zuk et al., Enzyme Immunoassay--A Quantitative Immunoassay Requiring No Instrumentation, Clinical Chemistry, 31 (7): 1144-1150 (1985)

**SEQUENCES****SEQ ID NO: 1**

5 hDPP3 AS 1-737  
MADTQYILPNDIGVSSLDCREAFRLSPTERLYAYHLSRAAWYGGAVLLQTSPPEAP  
YIYALLSRLFRAQDPDQLRQHALAEGLTEEEYQAFLVYAAGVYSNMGNYKSGD  
FVPNLPKEKLERVILGSEAAQQHPEEVRLWQTCGELMFSLEPRLRHLGLGKE  
FSGNCTMEDAKLAQDFLDSQNLSAYNTRLFKEVDGEGKPYYEVRLASVLGSEPL  
10 DS EVTSKLKSYFRGSPFQVTRGDYAPILKVVQELEKAKAYAANSHQGQMLAQYIES  
TQGSIEAHKRGSRFWIQDKGPIVESYIGFIESYRDPFGSRGEFEGFVAVVN  
15 KAMSAKFE RLVASAEQLLKELPWPPTFEKDKFLPDFTSLDVLTFAGSGIPAGIN  
FKNVSLGNVLAVAYATQREKLTFLEEDDKDLYILWKGPSFDVQVGLHE  
LFVQDEKGAFNFDQETVINPETGEQIQSWYRSGETWDSKFSTIASSYEECRA  
E5 LCLHPQVLEIFGFEGADAEDVIYVNWLNMVRAGLLALEFY  
TPEAFNWRQAHMQARF VILRVILLEAGEGLVTITPTGSDGRPDARV  
19 RLDRSKIRSVGKPALERFLRRLQVLKSTG DVAGGRALYEGYATV  
DAPPECFLRTLDTVLLRKESRKLIQPNTRLLEGSDVQLLEY  
EASAAGLIRSFSERFPEDGPELEEILTQLATADARFWKG  
GPSEAPSGQA

**20 SEQ ID NO: 2**

hDPP3 AS 474-493 (N-Cys)  
CETVINPETGEQIQSWYRSGE

## FIGURE DESCRIPTION

### Fig. 1: Inhibition of DPP3 activity

- 5 (A) The activity of recombinant GST-hDPP3 was measured in the presence of several different antiDPP3 antibodies. DPP3 binding antibodies, that were produced against peptides and/ or full-length (FL) native DPP3, show a strong inhibitory effect of up to 70 %.
- 10 (B) Inhibition curve of recombinant GST-hDPP3 with inhibitory mAbDPP3. Inhibition of DPP3 by a specific antibody is concentration dependent, with an IC<sub>50</sub> at ~0.2 µg/ ml.

### Fig. 2: DPP3 concentration as diagnostic marker

- 15 (A) DPP3 concentration in EDTA plasma of healthy controls and patients with various diseases (AHF – acute heart failure, MI – myocardial infarct, sepsis, cancer, AKI – acute kidney injury, LRTI – lower respirational tract infection). Medians of patient groups differ significantly from healthy controls (Mann-Whitney test p < 0.005).
- 20 (B) Comparison of plasma DPP3 concentrations of patients that died shortly within admission to the emergency department and surviving patients. Surviving patients show significantly lower DPP3 levels (Mann-Whitney test p < 0.05).

### Fig. 3: DPP3 activity as diagnostic marker

- 25 DPP3 activity in EDTA plasma of healthy controls and patients with various diseases (AHF – acute heart failure, sepsis, AKI – acute kidney injury, LRTI – lower respirational tract infection). Medians of patient groups differ significantly from healthy controls (Mann-Whitney test p < 0.0001).

### Fig. 4: ROC plot analysis of the DPP3 activity and concentration assay.

- 30 (A) ROC analysis of healthy controls and patients suffering from AHF.
- (B) ROC analysis of healthy controls and patients suffering from sepsis.

**Fig. 5: Safety of mAbDPP3 treatment (blood pressure)**

Healthy rats treated with PBS or mAbDPP3 (5,75 mg/ kg). Blood pressure (BP) was measured and recorded via a catheter inserted into the Arteria carotis communis dextra. The administration and sampling catheter were inserted into the Vena jugularis sinistra. Treatment of slightly increases relative blood pressure compared to PBS treated rats (n = 3 per group).

**Fig. 6: Influence of mAbDPP3 on mortality of septic mice**

Septic mice (CLP model) were treated with PBS or mAbDPP3 (1,9 mg/ kg) 5 minutes before and 2 h after CLP. Mortality was monitored over 7 days. The Kaplan-Meyer plot shows increased survival of septic mice after mAbDPP3 treatment.

**Fig. 7: Influence of mAbDPP3 on heart failure of septic rats**

(A) Experimental design of heart failure study of rats in septic shock.

(B) CLP induces heart failure in rats, as indicated by a decreased shortening fraction compared to sham animals. This shortening fraction is significantly increased by mAbDPP3 treatment (2 mg/ kg; n ≥ 7 per group; Mann-Whitney test p < 0.0001).

(C) Mean blood pressure of vehicle treated septic rats decreases with time whereas mAbDPP3 treatment leads to a significant increase in mBP (2 mg/ kg; n ≥ 7 per group; Mann-Whitney test p < 0.005).

**Fig. 8: Influence of mAbDPP3 on tumor growth *in vitro***

Soft-Agar Assay with tumor cell lines (lung, colon and breast cancer). Addition of antiDPP3 antibody reduces tumor cell growth.

**Fig. 9: Influence of mAbDPP3 on tumor growth *in vivo***

(A) Mice with xenograft of breast tumor cells (n = 10 per group) were treated with PBS or mAbDPP3. Growth curve of relative tumor volume over 24 days shows a decreased tumor growth in mAbDPP3 treated mice.

(B) Comparison of time the breast cell tumor needs to 20-fold increase its volume with and without mAbDPP3 treatment. Growth takes significantly longer with mAbDPP3 treatment (Mann-Whitney test,  $p < 0.05$ ).

5 (C) Mice with xenograft of colon tumor cells ( $n = 10$  per group) were treated with PBS or mAbDPP3. Growth curve of relative tumor volume over 30 days shows a decreased tumor growth in mAbDPP3 treated mice.

10 (D) Comparison of time the colon cell tumor needs to 10-fold increase its volume with and without mAbDPP3 treatment. Growth takes longer with mAbDPP3 treatment.

**Fig. 10: DPP3 activity as diagnostic marker (II)**

DPP3 activity in EDTA plasma of healthy controls and patients with various diseases (acute myocardial infarction (AMI), cardiogenic shock, septic shock and liver failure). Medians of 15 patient groups differ significantly from healthy controls (Mann-Whitney test  $p < 0.05$ ).

**Fig. 11: Effect of DPP3 on blood pressure in healthy rats**

Healthy male Wistar rats were injected with 0.2 mg/ kg recombinant GST-hDPP3. Blood 20 pressure (BP) was measured and recorded via a catheter inserted into the Arteria carotis communis dextra. DPP3 was injected i.v. via the tail vain. DPP3 injection leads to decreased BP.

**CLAIMS**

1. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes comprising:
  - determining the amount of total DPP3 and/or determining the amount of active DPP3 in a sample of bodily fluid of said subject,
  - comparing said determined amount of total DPP3 or active DPP3 to a predetermined threshold,
  - wherein said subject is diagnosed as having a disease or condition accompanied by or related to necrotic processes if said determined amount is above said predetermined threshold.
- 15 2. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 1 wherein the amount of total or active DPP3 is determined in the unit of concentration.
- 20 3. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 1 or 2, wherein said sample is selected from the group comprising whole blood, serum and plasma.
- 25 4. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1-3, wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF),, myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension.
- 30 5. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1-4, wherein the amount of total DPP3 and/or the amount of active DPP3 is determined in a bodily fluid sample of said subject and comprises the steps:

- Contacting said sample with a capture-binder that binds specifically to full-length DPP3,

- Separating DPP3 bound to said capture binder,

- Adding substrate of DPP3 to said separated DPP3,

- Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.

5           6. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 5, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

10          7. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 5 or 6, wherein said capture-binder is an antibody.

15          8. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-7, wherein said capture binder is immobilized on a surface.

20          9. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-8, wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.

25          10. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-9, wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary

zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

- 5 11. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-10, wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.
- 10 12. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 5-11, wherein said substrate may be selected from the group comprising: a di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.
- 15 13. Method for monitoring a disease or condition in a subject accompanied by or related to necrotic processes wherein the method of diagnosing according to any of claims 1 to 12 is conducted at least twice.
- 20 14. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes.
- 25 15. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to claim 14 wherein said inhibitor is selected from the group comprising an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold.
- 30 16. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to claim 14 or 15 wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), SIRS or sepsis, cancer, acute kidney injury (AKI), CNS

disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases, vascular diseases (e.g. Kawasaki syndrome) and hypotension.

- 5        17. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claim 14 to 16 wherein said inhibitor is an antibody that is mono-binding or at least two-binding.
- 10      18. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 17 wherein said inhibitor is an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 1, in particular that binds to SEQ ID No. 2.
- 15      19. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 18 wherein said inhibitor is an antibody or fragment or scaffold that exhibits a minimum binding affinity to DPP3 of equal or less than  $10^{-7}$  M.
- 20      20. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 19 wherein said inhibitor is an antibody or fragment or scaffold that is monospecific.
- 25      21. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 14 to 20 wherein said inhibitor is an antibody or fragment or scaffold binds to full-length DPP3 and inhibits activity of DPP3 of at least 10%, or at least 50%, more preferred at least 60%, even more preferred more than 70 %, even more preferred more than 80 %, even more preferred more than 90 %, even more preferred more than 95 %.
- 30      22. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any

claims 14 to 21 wherein said inhibitor is selective and/or specific to DPP3 and not cross cell membranes and/or the blood brain barrier.

23. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to any claims 14 to 22 wherein said subject has an amount of total DPP3 and/or an amount of active DPP3 in a sample of bodily fluid of said subject that is above a predetermined threshold.  
5
- 10 24. Pharmaceutical composition comprising an inhibitor of the activity of DPP3 according to any of the claims 14 to 23 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes.
- 15 25. Use of an inhibitor of the activity of DPP3 according to any of the claims 14 to 23 in a method of extracorporeal removal of DPP3 from plasma comprising apheresis and affinity chromatography.
- 20 26. Method for determining active DPP3 in a bodily fluid sample of a subject comprising the steps:
  - Contacting said sample with a capture-binder that binds specifically to full-length DPP3,
  - Separating DPP3 bound to said capture binder,
  - Adding substrate of DPP3 to said separated DPP3,
  - Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.  
25
- 30 27. Method for determining active DPP3 in a bodily fluid sample of a subject according to claim 26 wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.
28. Method for determining active DPP3 in a bodily fluid sample of a subject according to claims 26 or 27 wherein said capture-binder is an antibody.

29. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 28 wherein said capture binder is immobilized on a surface.
- 5    30. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 29 wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.
- 10    31. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 30 wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).
- 15
- 20    32. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 31, wherein said substrate may be selected from the group comprising: A di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.
- 25    33. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26 to 32, wherein said sample is a blood sample selected from the group comprising whole blood, serum and plasma.
- 30    34. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject comprising:
- A capture-binder that binds specifically to full-length DPP3,
  - A substrate of DPP3.

35. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 34, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

5  
36. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 34 or 35, wherein said capture-binder inhibits less than 50% DPP3 activity in a liquid phase assay, preferably less than 40%, more preferably less than 30 %.

10 37. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 36, wherein the binding region of DPP3 for the capture-binder is not within the region of amino acids 316 - 669 of SEQ ID No. 1.

15 38. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 37 wherein said binder is an antibody.

39. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 38 wherein said capture-binder is immobilized on a surface.

20 40. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 34 to 39 wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or a di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.

25  
30 41. Use of a method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 26-33 or use of an assay or kit according to any of claims 34 to 40 in a method of for diagnosing or monitoring a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1 - 13.

**AMENDED CLAIMS**  
**received by the International Bureau on 21 August 2017 (21.08.2017)**

- 5     1. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes comprising:
- 10    • determining the amount of total DPP3 and/or determining the amount of active DPP3 in a sample of bodily fluid of said subject,  
      • comparing said determined amount of total DPP3 or active DPP3 to a predetermined threshold,  
      • wherein said subject is diagnosed as having a disease or condition accompanied by or related to necrotic processes if said determined amount is above said predetermined threshold.
- 15    • wherein said sample is selected from the group comprising whole blood, serum and plasma.
- 20    2. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 1 wherein the amount of total or active DPP3 is determined in the unit of concentration.
- 25    3. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 1 or 2, wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF),,, myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases and vascular diseases (e.g. Kawasaki syndrome) and hypotension.
- 30    4. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 1-3, wherein the amount of total DPP3 and/or the amount of active DPP3 is determined in a bodily fluid sample of said subject and comprises the steps:

- Contacting said sample with a capture-binder that binds specifically to full-length DPP3,

- Separating DPP3 bound to said capture binder,

- Adding substrate of DPP3 to said separated DPP3,

- 5      • Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.

5. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claim 4, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

10     6. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to claims 4 or 5, wherein said capture-binder is an antibody.

15

7. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 4-6, wherein said capture binder is immobilized on a surface.

20

8. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 4-7, wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.

25

9. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 4-8, wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

10. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 4-9, wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.
- 5
11. Method for diagnosing a disease or condition in a subject accompanied by or related to necrotic processes according to any of claims 4-10, wherein said substrate may be selected from the group comprising: a di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.
- 10
12. Method for monitoring a disease or condition in a subject accompanied by or related to necrotic processes wherein the method of diagnosing according to any of claims 1 to 11 is conducted at least twice.
- 15
13. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes, wherein said disease is selected from the group comprising heart failure, chronic heart failure, acute heart failure (AHF), myocardial infarction (MI), stroke, liver failure, burn injuries, traumatic injuries, severe infection (microbial, viral (e.g. AIDS), parasitic (e.g. Malaria)), SIRS or sepsis, cancer, acute kidney injury (AKI), CNS disorders (e.g. seizures, neurodegenerative diseases), autoimmune diseases and vascular diseases (e.g. Kawasaki syndrome) and hypotension.
- 20
14. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to claim 13 wherein said inhibitor is selected from the group comprising an anti-DPP3 antibody or anti-DPP3 antibody fragment or anti-DPP3 non-Ig scaffold.
- 25
15. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes according to
- 30

claim 14 wherein said inhibitor is an antibody that is mono-binding or at least two-binding.

16. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
5 condition in a subject accompanied by or related to necrotic processes according to  
claim 14 or 15 wherein said inhibitor is an anti-DPP3 antibody or anti-DPP3 antibody  
fragment or anti-DPP3 non-Ig scaffold that binds to SEQ ID No. 1, in particular that  
binds to SEQ ID No. 2.

10 17. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to  
according to any of claims 14 to 16 wherein said inhibitor is an antibody or fragment or  
scaffold that exhibits a minimum binding affinity to DPP3 of equal or less than  $10^{-7}$  M.

15 18. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 17 wherein said inhibitor is an antibody or fragment or scaffold that is  
monospecific.

20 19. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 18 wherein said inhibitor is an antibody or fragment or scaffold binds to  
full-length DPP3 and inhibits activity of DPP3 of at least 10%, or at least 50%, more  
preferred at least 60%, even more preferred more than 70 %, even more preferred more  
25 than 80 %, even more preferred more than 90 %, even more preferred more than 95 %.

20. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any  
claims 14 to 19 wherein said inhibitor is selective and/or specific to DPP3 and not cross  
30 cell membranes and/or the blood brain barrier.

21. Inhibitor of the activity of DPP3 for use in prevention or therapy of a disease or  
condition in a subject accompanied by or related to necrotic processes according to any

claims 14 to 20 wherein said subject has an amount of total DPP3 and/or an amount of active DPP3 in a sample of bodily fluid of said subject that is above a predetermined threshold.

- 5 22. Pharmaceutical composition comprising an inhibitor of the activity of DPP3 according to any of the claims 14 to 21 for use in prevention or therapy of a disease or condition in a subject accompanied by or related to necrotic processes.
- 10 23. Use of an inhibitor of the activity of DPP3 according to any of the claims 14 to 21 in a method of extracorporeal removal of DPP3 from plasma comprising apheresis and affinity chromatography.
- 15 24. Method for determining active DPP3 in a bodily fluid sample of a subject comprising the steps:
  - Contacting said sample with a capture-binder that binds specifically to full-length DPP3,
  - Separating DPP3 bound to said capture binder,
  - Adding substrate of DPP3 to said separated DPP3,
  - 20 • Quantifying of said active DPP3 by measuring and quantifying the conversion of a substrate of DPP3.
- 25 25. Method for determining active DPP3 in a bodily fluid sample of a subject according to claim 24 wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.
- 30 26. Method for determining active DPP3 in a bodily fluid sample of a subject according to claims 24 or 25 wherein said capture-binder is an antibody.
27. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 24 to 26 wherein said capture binder is immobilized on a surface.

28. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 24 to 27 wherein said separation step is a washing step that removes ingredients of the sample that are not bound to said capture-binder from the captured DPP3.

5

29. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 24 to 28 wherein DPP3 substrate conversion is detected by a method selected from the group comprising: fluorescence of fluorogenic substrates (e.g. Arg-Arg- $\beta$ NA, Arg-Arg-AMC), color change of chromogenic substrates, luminescence of substrates coupled to aminoluciferin (Promega Protease-Glo<sup>TM</sup> Assay), mass spectrometry, HPLC/ FPLC (reversed phase chromatography, size exclusion chromatography), thin layer chromatography, capillary zone electrophoresis, gel electrophoresis followed by activity staining (immobilized, active DPP3) or western blot (cleavage products).

10

30. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 24 to 29, wherein said substrate may be selected from the group comprising: A di-peptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the di-peptide is Arg-Arg.

15

31. Method for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 24 to 30, wherein said sample is a blood sample selected from the group comprising whole blood, serum and plasma.

20

32. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject comprising:

- A capture-binder that binds specifically to full-length DPP3,
- A substrate of DPP3.

25

33. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 32, wherein said capture-binder may be selected from the group of antibody, antibody fragment or non-IgG scaffold.

34. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to claim 32 or 33, wherein said capture-binder inhibits less than 50% DPP3 activity in a liquid phase assay, preferably less than 40%, more preferably less than 30 %.

5

35. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 32 to 34, wherein the binding region of DPP3 for the capture-binder is not within the region of amino acids 316 - 669 of SEQ Id No. 1.

10 36. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 32 to 35 wherein said binder is an antibody.

37. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 32 to 36 wherein said capture-binder is immobilized on a surface.

15

38. Assay or kit for determining active DPP3 in a bodily fluid sample of a subject according to any of claims 32 to 37 wherein said substrate may be selected from the group comprising: angiotensin II, III and IV, Leu-enkephalin, Met-enkephalin, endomorphin 1 and 2, valorphin,  $\beta$ -casomorphin, dynorphin, proctolin, ACTH and MSH, or a dipeptide coupled to a fluorophore, a chromophore or aminoluciferin wherein the dipeptide is Arg-Arg.

20



**FIGURES****Fig. 1 A:****Fig. 1 B:**

Fig. 2 A:




Fig. 2 B:




Fig. 3:




Fig. 4 A:




Fig. 4 B:




Fig. 5:




Fig. 6:




Fig. 7 A:




Fig. 7 B:




Fig. 7 C:




Fig. 8:




Fig. 9 A:




Fig. 9 B:




Fig. 9 C:




Fig. 9 D:




Fig 10:





Fig 11:



# INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2017/059377

**A. CLASSIFICATION OF SUBJECT MATTER**  
 INV. G01N33/68 G01N33/574 A61K39/395  
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
**G01N A61K**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>AOYAGI T ET AL: "ENZYMATIC CHANGES IN CEREBROSPINAL FLUID OF PATIENTS WITH ALZHEIMER-TYPE DEMENTIA", JOURNAL OF CLINICAL BIOCHEMISTRY AND NUTRITION, TOKYO, JP, vol. 14, no. 2, 1 January 1993 (1993-01-01), pages 133-139, XP009037233, ISSN: 0912-0009 cited in the application abstract; tables 1-2</p> <p>-----</p> <p style="text-align: right;">-/-</p>	1,2,4, 13-16, 22-25

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
9 June 2017	21/06/2017
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  <b>Fleitmann, J</b>

## INTERNATIONAL SEARCH REPORT

International application No PCT/EP2017/059377
---

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SIMAGA S ET AL: "Dipeptidyl peptidase III in malignant and non-malignant gynaecological tissue", EUROPEAN JOURNAL OF CANCER, ELSEVIER, AMSTERDAM, NL, vol. 34, no. 3, 1 February 1998 (1998-02-01), pages 399-405, XP004284955, ISSN: 0959-8049, DOI: 10.1016/S0959-8049(97)00401-2 cited in the application abstract page 400, left-hand column, paragraph 6 - page 401, left-hand column, paragraph 2; figure 2; table 1 page 403, left-hand column, paragraph 3 -----	1,2,4,5, 14,15, 17,18, 20, 23-32, 34-41
X	US 2006/177886 A1 (HAZATO TADAHIKO [JP] ET AL) 10 August 2006 (2006-08-10) claim 1; example 1 -----	1,2,4, 13,41
X	WO 2005/106486 A2 (BAYER HEALTHCARE AG [DE]; GOLZ STEFAN [DE]; BRUEGGEMEIER ULF [DE]; GEE) 10 November 2005 (2005-11-10) page 35, line 28 - page 36, line 7; claims 18.19-21,26.; table 1 -----	1-40
X	CHIBA T ET AL: "Inhibition of recombinant dipeptidyl peptidase III by synthetic hemorphin-like peptides", PEPTIDES, ELSEVIER, AMSTERDAM, NL, vol. 24, no. 5, 1 May 2003 (2003-05-01), pages 773-778, XP002342483, ISSN: 0196-9781, DOI: 10.1016/S0196-9781(03)00119-0 abstract -----	14,23,24
X	AGIC ET AL: "Novel amidino-substituted benzimidazoles: Synthesis of compounds and inhibition of dipeptidyl peptidase III", BIOORGANIC CHEMISTRY, ACADEMIC PRESS INC., NEW YORK, NY, US, vol. 35, no. 2, 9 March 2007 (2007-03-09), pages 153-169, XP005913983, ISSN: 0045-2068, DOI: 10.1016/J.BIOORG.2006.11.002 abstract -----	14,23,24
1		

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No  
PCT/EP2017/059377

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2006177886 A1	10-08-2006	JP 2006230318 A US 2006177886 A1	07-09-2006 10-08-2006
WO 2005106486 A2	10-11-2005	NONE	