

TITLE: Kittiwake Energetics Moc Project

Fred Tremblay^{1*}

¹Department of Natural Resources, McGill University, Montreal, Canada

*Corresponding author

Abstract

Here, we investigate the effects of the two-sample method compared to the single-sample method for measuring energy expenditure using DLW in free-ranging black-legged kittiwakes (*Rissa tridactyla*).

Introduction

Energy is a common currency for any living organism, and understanding how animals use energy is critical to understanding their ecology and consequently how they might adapt to changes in their environment. Individual variation in daily energy expenditure is often linked to varying activity budgets. Doubly-labelled water (DLW) is a commonly used technique to measure energy expenditure in wild animals [Speakman1997]. The use of DLW enables us to measure daily energy expenditure by injecting a bolus of heavy water (deuterium and ^{18}O). Commonly used when using DLW is the two sample method. Yet, when working with free-ranging animals, the manipulations linked with the two sample method can be invasive and stressful for the animals. Here, we investigate the effects of the two-sample method compared to the single-sample method for measuring energy expenditure using DLW in free-ranging black-legged kittiwakes (*Rissa tridactyla*).

Methods

We measured energy expenditure in black-legged kittiwakes using doubly-labelled water using two commonly used methods: the single-sample and the double-sample method. Along with our measure of energy expenditure, we calculated time-activity budgets of free ranging kittiwake using GPS-accelerometry data. We completed all analysis and figures using Rstudio.

Results

Both DLW methods are comparable and did not have a differential impact on the kittiwakes' energetics (figure 1) and time-activity budget (figure 2).




Figure 1: Energy expenditure (kJ/day) of single-sampled (grey) and two-sampled (beige) black-legged kittiwakes.




Figure 2: Time-activity budget of single-sampled (grey) and two-sampled (beige) black-legged kittiwakes.

Discussion

The two-sample method when measuring energy expenditure has no differential impact on the kittiwakes when compared to the single sample method. Thus, as the two-sample method allows us to measure energy expenditure more precisely [Speakman1997], this method should be favored.

References

output: pdf_document: latex_engine: xelatex