

# How does tree harvest influence benthic invertebrate density?

Sam Straus<sup>1,1,\*</sup>, Bob Security<sup>b</sup>, Cat Memes<sup>b,1</sup>, Derek Zoolander<sup>a,1</sup>

<sup>a</sup>Department Street City State Zip

<sup>b</sup>Department Street City State Zip

---

## Abstract

We looked at the density of *Baetis* spp. in harvested and non-harvested stream catchments.

**Keywords:** stream ecology, entomology

---

## 1. Introduction

*Baetis* spp, belonging to the insect order Ephemeroptera, are commonly used as stream quality indicators (Wallace and Gurtz, 1986).

## 2. Methods

Researchers collected data on the number of *Baetis* spp. per square meter in the benthic environment of various catchments from 1995 to 2005. Some of the catchments were harvested, while others were not.

## 3. Results

Figure 1 is generated using an R chunk.




Figure 1: *Baetis* density in harvested and non-harvested catchments

---

\*Corresponding author

Email addresses: samantha.straus@mcgill.ca (Sam Straus), bob@example.com (Bob Security), cat@example.com (Cat Memes), derek@example.com (Derek Zoolander)

#### 4. Discussion

Benthic invertebrates are commonly used as indicators of stream water quality (Guilpart et al., 2012). *Baetis* sp. (Ephemeroptera) are used as stream quality indicators for catchments (Wallace and Gurtz, 1986).

#### References

- Alexis Guilpart, Jean-Marc Roussel, Joël Aubin, Thierry Caquet, Mickaël Marle, and Hervé Le Bris. The use of benthic invertebrate community and water quality analyses to assess ecological consequences of fish farm effluents in rivers. *Ecological Indicators*, 23:356–365, 2012. Citation Key: guilpart2012use Publisher: Elsevier.
- J Bruce Wallace and Martin E Gurtz. Response of baetis mayflies (ephemeroptera) to catchment logging. *American Midland Naturalist*, page 25–41, 1986. Citation Key: wallace1986response Publisher: JSTOR.