



Exercises for Neuronify

Andreas V. Solbrå
Milad H. Mobarhan
Svenn-Arne Dragly
Simen Tennøe

April 29, 2015

Exercise 1 Small Networks

In this exercise we will connect neurons to make simple neural networks. In Neuronify, you can connect a neuron to another neuron just like any other device, such as those we looked at in the previous exercise.


Part I

In this part we will make the simplest possible network by connecting one neuron to another.

Make a small network by doing the following:

- Create an empty canvas by pressing the menu button in the upper left corner, then press select simulation and click on the empty canvas.
- Create two neurons, and label them “A” and “B”. You can set neuron labels by pressing a neuron, pressing the gear in the lower right corner. This will open up a tab on the right side of the screen. Press the field marked “Label:” and enter the label you want. Complete the operation by pressing Enter.
- Connect neuron A to neuron B by clicking on neuron A, grabbing the connection and pulling it to neuron B.
- Connect a DC clamp to neuron A as in exercise 1. Connect a Voltmeter to both neurons by dragging the voltmeter onto the canvas, pressing it, pressing the gear in the lower right corner, and pressing the “Connect to all neurons”-button.

Your canvas should now look like this:



Note that the voltage trace from neuron B is very different from neuron A, because neuron A receives a steady current input, while neuron B receives the spikes from A.

Using the default settings for both neurons, check that 75 mA is the smallest amount of Output Current from the DC clamp that still allows neuron B to fire. Now, set the resting

potential of both neurons to -70 mV, and attempt to find the minimum current output needed to induce spiking in neuron B.


Part II

In this exercise we will look at a simple convergent connections. Create a new empty canvas. Create two neurons, label them “A1” and “B”. Set the stimulation output of Neuron A1 to 0.5. Connect neuron A1 to neuron B. Create a current generator and connect it to neuron A1.

In this exercise we will look at a simple convergent connections. Create a new empty canvas. Create two neurons, label them “A1” and “B”. Set the stimulation output of Neuron A1 to 0.5. Connect neuron A1 to neuron B. Create a DC clamp and connect it to neuron A1.

We can make add noise to a neuron by adding Poisson generators. Add two Poisson generators to the canvas (they are the green squares with question marks on them). Set their rates to 2.0 ms^{-1} and their stimulation output to 0.1. Make one of the generators inhibitory. Connect a voltmeter to both neurons.

Now your canvas should look like this:



Because of the Poisson generators, there is a chance that neuron B will eventually fire in this circuit, but the odds are astronomically small. Disconnect the Poisson generators by pressing on the wires going from the generator to the neuron and then pressing the trash can in the lower left corner.

- Make several more neurons with Stimulation output 0.5, and connect the DC generator to each of them. Label them “A2”, “A3”, and so on. How many neurons do you have to connect to neuron B in order to make it fire?
- Add one excitatory and one inhibitory spike generators to each of the neurons A1, A2, How many neurons do you need to connect to B in order to make it fire now?

Exercise 2 Lateral inhibition

In this exercise we will create simple circuits that illustrate the concept of lateral inhibition, which is the capacity of an excited neuron to reduce the activity of its neighbors through mediating inhibitory neurons. This type of inhibition occurs primarily in visual processes to increase the sharpness in visual response.

1. Create a circuit in Neuronify consisting of two passive excitatory neurons and one passive inhibitory neuron, all connected to the same DC clamp. Adjust the current output of the DC clamp and see how the spiking pattern of the neurons changes.
2. Create lateral connections from the inhibitory neuron to the excitatory neurons. How does these connections affect the activity in the excitatory neurons?
3. Make the connections in the network shown below such that the spiking pattern is the same as in the figure. You are allowed to adjust the *stimulation output* property of the cells and use DC clamp with different current output for the upper neurons.
4. The network you have created now is an example where sharpening of the signal occurs through lateral inhibition. How those the relative strength of the inputs to the upper neurons affect the spiking of the lowermost neurons?
5. Does the lateral inhibition circuit increase or decrease the contrast between strong and weak signals?




Figure 1: Lateral inhibition network.