Neuronify: a new tool for creating simple neural networks

Simen Tennøe, Svenn-Arne Dragly, Andreas V. Solbrå, Milad H. Mobarhan

May 20, 2015

In this lecture we will introduce Neuronify, a new tool for creating neural networks

```
[remember picture,overlay] [xshift=-3.5cm,yshift=1.5cm] at
```


```
(current page.center) ; [xshift=-2.cm,yshift=1.5cm, right] at (current page.center) Integrate and fire neurons; [remember picture,overlay] [xshift=-2cm,yshift=-.5cm] at (current
```


```
page.center) ; [xshift=1cm,yshift=-.5cm, left] at (current page.center) Neuronify; [remember picture,overlay] [xshift=-.5cm,yshift=-2.5cm] at
```


```
(current page.center);


[xshift=2.8cm,yshift=-2.5cm, left] at (current page.center)

Exercises;
```









Hodgkin-Huxley

Hodgkin-Huxley

$$C\frac{dV}{dt} = I_{inj} - \bar{g}_{Na}m^{3}h(V - V_{Na}) - \bar{g}_{K}n^{4}(V - V_{K}) - g_{L}(V - V_{L})$$

$$\frac{dn}{dt} = \alpha_{n}(V)(1 - n) - \beta_{n}(V)n \quad \frac{dm}{dt} = \alpha_{m}(V)(1 - m) - \beta_{m}(V)m$$

$$\frac{dh}{dt} = \alpha_{h}(V)(1 - h) - \beta_{h}(V)h \quad \alpha_{n}(V) = \frac{0.01(V + 55)}{1 - \exp[-(V + 55)/10]}$$

$$\beta_{n}(V) = 1.125 \exp[-(V + 65)/80] \quad \alpha_{m}(V) = \frac{0.1(V + 40)}{1 - \exp[-(V + 40)/10]}$$

$$\beta_{m}(V) = 4 \exp[-(V + 65)/18] \quad \alpha_{h}(V) = 0.07 \exp[-(V + 65)/20]$$

$$\beta_{n}(V) = \frac{1}{1 + \exp[-(V + 35)/10]}$$

Hodgkin-Huxley

$$C\frac{dV}{dt} = I_{inj} - \bar{g}_{Na}m^{3}h(V - V_{Na}) - \bar{g}_{K}n^{4}(V - V_{K}) - g_{L}(V - V_{L})$$

$$\frac{dn}{dt} = \alpha_{n}(V)(1 - n) - \beta_{n}(V)n \quad \frac{dm}{dt} = \alpha_{m}(V)(1 - m) - \beta_{m}(V)m$$

$$\frac{dh}{dt} = \alpha_{h}(V)(1 - h) - \beta_{h}(V)h \quad \alpha_{n}(V) = \frac{0.01(V + 55)}{1 - \exp[-(V + 55)/10]}$$

$$\beta_{n}(V) = 1.125 \exp[-(V + 65)/80] \quad \alpha_{m}(V) = \frac{0.1(V + 40)}{1 - \exp[-(V + 40)/10]}$$

$$\beta_{m}(V) = 4 \exp[-(V + 65)/18] \quad \alpha_{h}(V) = 0.07 \exp[-(V + 65)/20]$$

$$\beta_{n}(V) = \frac{1}{1 + \exp[-(V + 35)/10]}$$

Integrate and fire

Hodgkin-Huxley

$$C\frac{dV}{dt} = I_{inj} - \bar{g}_{Na}m^{3}h(V - V_{Na}) - \bar{g}_{K}n^{4}(V - V_{K}) - g_{L}(V - V_{L})$$

$$\frac{dn}{dt} = \alpha_{n}(V)(1 - n) - \beta_{n}(V)n \quad \frac{dm}{dt} = \alpha_{m}(V)(1 - m) - \beta_{m}(V)m$$

$$\frac{dh}{dt} = \alpha_{h}(V)(1 - h) - \beta_{h}(V)h \quad \alpha_{n}(V) = \frac{0.01(V + 55)}{1 - \exp[-(V + 55)/10]}$$

$$\beta_{n}(V) = 1.125 \exp[-(V + 65)/80] \quad \alpha_{m}(V) = \frac{0.1(V + 40)}{1 - \exp[-(V + 40)/10]}$$


$$\beta_{m}(V) = 4 \exp[-(V + 65)/18] \quad \alpha_{h}(V) = 0.07 \exp[-(V + 65)/20]$$

$$\beta_{n}(V) = \frac{1}{1 + \exp[-(V + 35)/10]}$$


Integrate and fire

$$C_m \frac{dV(t)}{dt} = -\frac{V(t) - E_m}{R_m} + I$$

The integrate and fire model is modeled as a simple RC circuit that is shortcircuted once a threshold is reached



The integrate and fire model is modeled as a simple RC circuit that is shortcircuted once a threshold is reached



$$C_m \frac{dV(t)}{dt} = -\frac{V(t) - E_m}{R_m} + I$$

Receptive Field

$$r(t) = r_0 + \int \int D(\mathbf{r}, \tau) s(\mathbf{r}, t - \tau) d\tau d\mathbf{r}$$
 (1)

figures/gabor.png

Receptive Field

figures/recRFs.png