

Table 1: AFP Filled Correctly

	Overall (N=17)
Number of Hospital	
0	2 (100.0%)
Dates of VPI delievered via AVS	
0	11 (73.3%)
1	4 (26.7%)
Dates of VPO delivered via AVS	
0	10 (71.4%)
1	4 (28.6%)
Source of the vaccine information	
0	9 (60.0%)
1	6 (40.0%)
Probable Diagnosis	
0	9 (52.9%)
1	8 (47.1%)
Date of Hospitalisation	
0	1 (50.0%)
1	1 (50.0%)
Doses of VPO delivered via routine immunisaiton	
0	8 (47.1%)
1	9 (52.9%)
Doses of VPI delivered via AVS	
0	8 (47.1%)
1	9 (52.9%)
Doses of VPO delivered via AVS	
0	6 (37.5%)
1	10 (62.5%)
Total number of previous vaccines	
0	6 (35.3%)
1	11 (64.7%)
Dosese of VPI delivered via routine immunisation	
0	6 (35.3%)
1	11 (64.7%)
Coordinates (often incorrect)	
0	5 (29.4%)
1	12 (70.6%)
True AFP Case?	
0	5 (29.4%)
1	12 (70.6%)
Hospitalisation recorded	
0	4 (23.5%)
1	13 (76.5%)
Date of Birth	
0	3 (17.6%)
1	14 (82.4%)
Date of stool sample 2	
0	2 (11.8%)
1	15 (88.2%)
Province	
0	1 (5.9%)
1	16 (94.1%)

	Overall (N=17)
Name of Mother/Father	
0	1 (5.9%)
1	16 (94.1%)
Sex	
0	1 (5.9%)
1	16 (94.1%)
Is paralysis progressive	
0	1 (5.9%)
1	16 (94.1%)
Asymetric Paralysis	
0	1 (5.9%)
1	16 (94.1%)
Recent IM Injection	
0	1 (5.9%)
1	16 (94.1%)
Health District Noted	
1	17 (100.0%)
Formation	
1	17 (100.0%)
Telephone or Address	
1	17 (100.0%)
Town or Village	
1	17 (100.0%)
Name of Patient	
1	17 (100.0%)
Age	
1	17 (100.0%)
Name of person notified	
1	17 (100.0%)
Date of notification	
1	17 (100.0%)
Date of investigation	
1	17 (100.0%)
Name of hospital recorded	
1	2 (100.0%)
Date of paralysation recorded	
1	17 (100.0%)
Fever present	
1	17 (100.0%)
Flaccid Paralysis	
1	17 (100.0%)
Site of paralysis	
1	17 (100.0%)
Is the limb sensitive to pain	
1	17 (100.0%)
Site of Injection	
1	3 (100.0%)
Dates of previous VPO Vaccines	
1	16 (100.0%)
Date of stool sample 1	
1	17 (100.0%)
Date of stool sample 2	

	Overall (N=17)
1	17 (100.0%)
Health District No	
1	4 (23.5%)
2	2 (11.8%)
3	1 (5.9%)
4	7 (41.2%)
5	2 (11.8%)
7	1 (5.9%)