

Homework 3b

Cameron Lane

CJL3282

1)

$$JD_{UT1} = 2459629.74930427$$

$$\theta_{ERA} = 5.094194831139650e+04 \text{ radians}$$

W Matrix: W =

$$\begin{bmatrix} 0.999999999999997 & -5.02960587195337e-11 & -8.12935581357585e-08 \\ 5.04369830566906e-11 & 0.999999999998497 & 1.73352403885434e-06 \\ 8.12935580484469e-08 & -1.73352403885844e-06 & 0.999999999998494 \end{bmatrix}$$

R Matrix: R =

$$\begin{bmatrix} -0.520437212092403 & 0.853899940431833 & 0 \\ -0.853899940431833 & -0.520437212092403 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

PN Matrix: PN =

$$\begin{bmatrix} 0.999999999773384 & -5.53896235428886e-08 & 0.00212557524464342 \\ 1.01377935810267e-08 & 0.999999999773385 & 2.12891916943869e-05 \\ -0.00212557524534093 & -2.1289122052659e-05 & 0.999997740735773 \end{bmatrix}$$

$Q_{GCRF}^{ITRF}$  Matrix: Q =

$$\begin{bmatrix} -0.520437164461401 & 0.853899965405303 & 0.00212709780895228 \\ -0.853899945538927 & -0.520437203310978 & 2.04564178565521e-05 \\ 0.00112448852811387 & -0.00180568244422723 & 0.999997737515671 \end{bmatrix}$$

2)

$$\theta_{GMST} = 4.16991131806314$$

$r_{GCRF}$  for  $Q = R_3(-\theta_{GMST})$ :

$$\begin{bmatrix} -4295.39724275466 \\ 3456.64811985493 \\ 3196.066 \end{bmatrix}$$

$r_{GCRF}$  for  $Q = PN R W$ :

$$\begin{bmatrix} -4272.28252523703 \end{bmatrix}$$

3477.59348022302  
3204.3259593577

Error = 32.2680215609783

The GMST transformation is accurate for the first component of  $r$ . The difference in the second and third components is significantly larger. GMST is easier and quicker to use whereas CIO is much more accurate.