

Worksheet 9

December 2, 2022

1. Prove that for $1 \leq k \leq n$,

$$(i) \langle a_0, a_1, \dots, a_n \rangle = \langle a_0, a_1, \dots, a_{k-1}, \langle a_k, a_{k+1}, \dots, a_n \rangle \rangle,$$

Hint : Use Induction on the number of terms in the innermost continued fraction on the right hand side $\langle a_k, a_{k+1}, \dots, a_n \rangle$.

(ii) Use part(i) to prove that:

$$\langle a_0, a_1, \dots, a_n \rangle = a_0 + \frac{1}{\langle a_1, \dots, a_n \rangle}$$

2. Convert each of the following into finite simple continued fractions

(i) 0.23

(ii) $\frac{233}{177}$

3. Find $\frac{p}{q}$ if $\frac{p}{q} = [3, 7, 15, 1]$. Convert $\frac{p}{q}$ to a decimal and compare with the value of π .

4. (i) Writing the simple continued fraction of proper fractions. Ex. $\frac{29}{67}$.

Ex. $\frac{29}{67}$;

$$\frac{29}{67} = [0, 2, 3, 4, 2]$$

$$= [a_1, a_2, a_3, a_4, a_5]$$

$$0 + \frac{1}{2 + \frac{1}{3 + \frac{1}{4 + \frac{1}{2}}}}$$

$$67 \overline{) 29} (0 = a_1$$

$$\begin{array}{r} 0 \\ 29 \overline{) 67} (2 = a_2 \\ \underline{58} \end{array}$$

$$\begin{array}{r} 58 \\ 9 \overline{) 29} (3 = a_3 \\ \underline{27} \end{array}$$

$$\begin{array}{r} 27 \\ 2 \overline{) 9} (4 = a_4 \\ \underline{8} \end{array}$$

$$\begin{array}{r} 8 \\ 1 \overline{) 2} (2 = a_5 \\ \underline{2} \\ 0 \end{array}$$

(ii) Write the continued fraction expansion for $\frac{67}{29}$.

(iii) Compare (i) & (ii) above to conclude a general result for the relation between the continued fraction of

$$\frac{p}{q} \& \frac{q}{p}, \text{ where } p > q.$$