

Leren homework # 2**Date:** November 6, 2014**Name (Student Number):** M. Pfundstein (10452397), T.M. Meijers (10647023)**Question 1**

Hypothesis function:

$$h_{\theta}(x^{(i)}) = \theta_0 + \theta_1 x_1^{(i)} + \theta_2 x_2^{(i)}$$

Cost function:

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Update function:

$$\theta'_i = \theta_i - \alpha \frac{\delta}{\delta \theta_i} J(\theta)$$

Cost after 0 iterations:

With $\theta_0 = 0.5, \theta_1 = 0.5, \theta_2 = 0.5$

$$\begin{aligned} J(\theta) &= \frac{1}{6}((0.5 + 1 + 1.5 - 6)^2 + (0.5 + 2 + 2.5 - 6)^2 + (0.5 + 2 + 1.5 - 10)^2) = \\ &= \frac{1}{6}((-3)^2 + (-1)^2 + (-6)^2) = \\ &= \frac{46}{6} = \frac{23}{3} \end{aligned}$$

Update parameters (1 iteration):

$$\begin{aligned} \theta'_0 &= 0.5 - \frac{0.1}{3}(-3 - 1 - 6) = \\ 0.5 - \frac{-1}{3} &= \frac{5}{6} \end{aligned}$$

$$\begin{aligned} \theta'_1 &= 0.5 - \frac{0.1}{3}(2 \cdot -3 + 4 \cdot -1 + 4 \cdot -6) = \\ 0.5 - \frac{0.1}{3}(-34) &= \\ 0.5 + \frac{3.4}{3} &= \frac{49}{30} \end{aligned}$$

$$\begin{aligned} \theta'_2 &= 0.5 - \frac{0.1}{3}(3 \cdot -3 + 5 \cdot -1 + 3 \cdot -6) = \\ 0.5 - \frac{0.1}{3}(-32) &= \\ 0.5 + \frac{3.2}{3} &= \frac{47}{30} \end{aligned}$$

Cost after 1 iteration:

With $\theta_0 = \frac{5}{6}, \theta_1 = \frac{49}{30}, \theta_2 = \frac{47}{30}$

$$J(\theta) = \frac{1}{6}((\frac{5}{6} + \frac{49}{30} \cdot 2 + \frac{47}{30} \cdot 3 - 6)^2 + (\frac{5}{6} + \frac{49}{30} \cdot 4 + \frac{47}{30} \cdot 5 - 6)^2 + (\frac{5}{6} + \frac{49}{30} \cdot 4 + \frac{47}{30} \cdot 3 - 10)^2) \approx 16.12518$$

Question 2

Positive and negative aspects per terminating condition (gradient descent algorithm):

- (a) Iteration:
 - + One can directly control the running time of the algorithm.
 - + The algorithm will terminate guaranteed.
 - If the number of iterations is too high, the algorithm will do redundant work.
 - If the number of iterations is too low, you can obtain parameters far from optimal.
- (b) Change in parameters:
 - + A very small change can be chosen which guarantees nearly optimal parameters.
 - + No need to calculate cost in between updating.
 - If chosen learning rate is too high, algorithm will never terminate.
 - Change in parameters does not necessarily mean a change in cost.
- (c) Change in cost:
 - + A very small change can be chosen which guarantees nearly optimal parameters.
 - + Cost is the result one wants to minimize.
 - Algorithm also needs to calculate cost, opposed to using latest change in parameters.

Question 3

Hypothesis function:

$$h_{\theta}(x^{(i)}) = \frac{1}{1+e^{-\theta^T x^{(i)}}}$$

Cost function:

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \begin{cases} -\log(h_{\theta}(x^{(i)})) & \text{if } y^{(i)} = 1, \\ -\log(1 - h_{\theta}(x^{(i)})) & \text{if } y^{(i)} = 0 \end{cases}$$

Simplified cost function:

$$J(\theta) = -\frac{1}{m} \sum_{i=1}^m y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)}))$$

Update function (using simplified $J(\theta)$):

$$\theta'_j = \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

Calculate hypothesethes:

$$h_{\theta}(x^{(1)}) = \frac{1}{1+e^{-9/2}} \approx 0.9890$$

$$h_{\theta}(x^{(2)}) = \frac{1}{1+e^{-11/2}} \approx 0.9959$$

$$h_{\theta}(x^{(3)}) = \frac{1}{1+e^{-7/2}} \approx 0.9707$$

$$h_{\theta}(x^{(4)}) = \frac{1}{1+e^{-7/2}} \approx 0.9707$$

Cost after 0 iterations (Normal cost function):

With $\theta_0 = 0.5, \theta_0 = 0.5, \theta_0 = 0.5$

$$J(\theta) = -\frac{1}{4}(\log(1 - 0.9890) + \log(1 - 0.9959) + \log(0.9707) + \log(0.9707)) \approx 2.5187$$

Update parameters (1 iteration):

$$\begin{aligned} \theta'_0 &= 0.5 - 0.1 \frac{1}{4} \sum_{i=1}^4 (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)} \\ &= 0.5 - \frac{0.1}{4} [((0.9890 - 0) * 1 + (0.9959 - 0) * 1 + (0.9707 - 1) * 1 + (0.9707 - 1) * 1)] \approx 0.4518 \end{aligned}$$

$$\begin{aligned} \theta'_1 &= 0.5 - 0.1 \frac{1}{4} \sum_{i=1}^4 (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)} \\ &= 0.5 - \frac{0.1}{4} [((0.9890 - 0) * 5 + (0.9959 - 0) * 5 + (0.9707 - 1) * 3 + (0.9707 - 1) * 2)] \approx 0.2555 \end{aligned}$$

$$\begin{aligned} \theta'_2 &= 0.5 - 0.1 \frac{1}{4} \sum_{i=1}^4 (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)} \\ &= 0.5 - \frac{0.1}{4} [((0.9890 - 0) * 3 + (0.9959 - 0) * 5 + (0.9707 - 1) * 3 + (0.9707 - 1) * 4)] \approx 0.3065 \end{aligned}$$

Cost after 1 iteration (Normal cost function):

With $\theta'_0 = 0.4518, \theta'_1 = 0.2555, \theta'_2 = 0.3065$

Calculate hypothesethes:

$$h_{\theta'}(x^{(1)}) = \frac{1}{1+e^{-9/2}} \approx 0.9339$$

$$h_{\theta'}(x^{(2)}) = \frac{1}{1+e^{-11/2}} \approx 0.9631$$

$$h_{\theta'}(x^{(3)}) = \frac{1}{1+e^{-7/2}} \approx 0.8945$$

$$h_{\theta'}(x^{(4)}) = \frac{1}{1+e^{-7/2}} \approx 0.8992$$

$$J(\theta') = -\frac{1}{4}(\log(1 - 0.9339) + \log(1 - 0.9631) + \log(0.8945) + \log(0.8992)) \approx 1.5586$$

Plotting

$$\theta_0 = \begin{bmatrix} 0.5 \\ 0.5 \\ 0.5 \end{bmatrix}$$

$$h_{\theta_0}(x_1, x_2) = 0.5 = 0.5 + 0.5x_1 + 0.5x_2$$


$$x_2 = -x_1$$

$$\theta_1 = \begin{bmatrix} 0.4518 \\ 0.2555 \\ 0.3065 \end{bmatrix}$$

$$h_{\theta_1}(x_1, x_2) = 0.5 \rightarrow 0.5 = 0.4518 + 0.2555x_1 + 0.3065x_2$$

$$\rightarrow x_2 = 0.157259 - 0.833605x_1$$

Figure 1: Class boundaries plotted for both sets of Theta's



Question 4

The cost method (accuracy or Andrew's method) used makes a difference when one has a dataset where all the data points are very apart, i.e. they're all far away from the class boundary. When one has such a dataset the accuracy method will report 100% accuracy whereas Andrew's method will return a very large cost.

Figure 2: A small dataset with significant difference between cost methods

