

Introduction

Time series analysis is important in any areas such as finical, astronomy, geophysics and so on. In the geophysics, time series are analyzed to extract the signals caused by geophysics phenomenon. This software is designed to analyze the GNSS coordinates time series and estimated the signals, such as linear, periodical signals. Also, some useful tools are provided for analysis convenience.

Requirements and Installation

TSAalyzer V2.0 is based on Python, it supports Python 2 and Python 3, so some requirements should be installed before it launched.

Requirements

- qtpy
- Matplotlib
- Numpy
- Pandas
- cvxpy (For offset detection, the solver [MOSEK](#) is better)
- cvxopt

Installation


Before installing this package, the [Anaconda](#) is recommended to be installed. *TSAalyzer* can support both Python2 and Python3.

- Download this repo, and `python setup.py install` on common line prompt.
- `pip install tsanalyzer`

After that, the package creates a command `tsanalyzer` in your path.

GUI Instructions


As TSAalyzer 2.0 was redesigned and refactoring, we discard some functions, for example, spectrum analysis, sigseg, however the discontinuity detection and interaction ability was enhanced. Also the batch functions were made easy to use.



Tool Bar

- Home plot view (alt+h)
- Previous plot view
- Next plot view
- Pan the figure (alt+p)
- Zoom the figure (alt+z)
- Subplot Configuration
- Figure Settings (lines, markers, colors)
- Offset (alt+1)
- Trend Change (alt+2)
- Earthquake Exponential Relaxation (alt+3)
- Earthquake Logarithmic Relaxation (alt+4)
- Save the figure (alt+s)

Time-Series Files Dock



The files dock support the context menu. Right click the dock, and the file menu will show up. Double click the filename in the dock, this file will be loaded. The `Export all time-series filename` is to save the files name in the dock into a text file.


Discontinuities Dock



The dock provide edit, import and export functions. The discontinuities dock only display the loaded file's, but the others time series files' discontinuities are also recorded in the memory.

Time-series Figure

With the tool bar provided, the user can do interaction with the figure. Discontinuities will be added by **left click**, once the user click the figure when the discontinuities actions (offset, trend change, exp decay or log decay) were triggered. When the discontinuity line are hovered, you can **drag** the position to specific date with **middle click** of the mouse, **right click** is to delete the discontinuity.




Analysis Dock

Discontinuity Detection

The mathematical models for L1 Extensive Model is:

$$\text{minimize} \quad \frac{1}{2} \|\mathbf{y} - \mathbf{x} - \mathbf{s} - \mathbf{w}\|_2^2 + \lambda \|D^2 \mathbf{x}\|_1 + \rho \|D^1 \mathbf{w}\|_1$$

where \mathbf{y} is the observation, \mathbf{x} is piecewise trends, $\mathbf{s} = \sum_i (a_i \sin \omega_i t_i + b_i \cos \omega_i t_i)$ is seasonal term, and \mathbf{w} are the level shifts. Here, D^n stands for n-th difference matrix. We can use the piecewise trends \mathbf{x} to find trend change points and \mathbf{w} to get potential offset points. Then F-test is used to validate. So the parameters in the groupbox are stands for these parameters. The offsets and trend change check means give the result of offset or trend change result or not.



The batch button is used to detect discontinuities in the Time-series Dock files. Once the procedure done, click the `import` button to import discontinuities to the `Discontinuity Dock`

Outliers

Big Sigma Criterion

This criterion is according the sigma (userinput, accepted one or three numbers) to mask out the data whose sigma is larger than sigma criterion to get the cleaner data. One number input means that three sigma criterion for all components are the same. If you want to specific for individual component, please use space between numbers. If sigma is not set, this criterion will not be adopted.

IQR

By using the least squares model, one could get the residuals \mathbf{v} , then the software use the window (user input) to get the percentile of 75%, median and 25%.

$$|\mathbf{v}_i - \mathbf{v}_{median}| > \text{factor}(\mathbf{v}_{75} - \mathbf{v}_{25})$$


Least Squares Analysis

$$y(t_i) = a + \sum_{i=1}^n b_i (t_i - t_0)^i + \sum_{i=1}^{n_p} (c_i \sin(2\pi t_i / p_i) + d_i \cos(2\pi t_i / p_i)) - \sum_{i=1}^{n_g} (g_i H(t_i - T_{g_j})) + \sum_{i=1}^{n_h} (h_i (t_i - T_{k_i}) H(t_i - T_{k_i})) + \sum_{i=1}^{n_l} \left(C_{l_i} + l_i H \ln(1 + \frac{t_i - t_{eq}}{\tau}) \right) + \sum_{i=1}^{n_e} (C_{e_i} + e_i H(1 - e^{-(1-(t-t_{eq})/\tau)}))$$

* if the math equations cannot be rendered, please refer to the pdf file

In the equation, b_i is polynomials, if $i = 1$ for b_i , it is the linear trend, c_i and d_i are harmonic components, the H is a step function and g_i can be used to explain the sudden change event caused the equipments or co-seismic, C_{l_i} and C_{e_i} terms are stand for logarithmic or exponential function to be fit after an Earthquake. These terms are the accumulated post-seismic motion, C is the co-seismic offset, and τ is the decay time.

This model can be used to estimated polynomials (max order is 10), harmonics (users can define the period) as well as step function (including sudden change, post-seismic log or relaxation) at specific times (users pick interactively or input offsets file), detail information is displayed in the picture.



The batch button is also provide the function to analysis all the files in Time-series files dock. The results will be saved in the directory with data and log separately.

Tools

Date Converter

In the GNSS coordinate time series analysis, there are some formats of time are given, so TSAAnalyzer provides this tool for convenience.

TSAAnalyzer - Date Tool

Date Conversion Tools

☒ Date 2018-04-17

☐ MJD 58226.0

☐ Julday 2458226

☐ Demical Year 2018.2930

☐ Year Doy 2018-107

☐ GPSWeek 19972

Convert

Header Convert Tool

TSAAnalyzer - Header Tool

Add Files **File Preview**

E:\Program\TSAAnalyzer\Example\AHAQios.gar

#	YYYY	DECM	YYYY	DOY	N(m)	E(m)	U(m)	nn(m)	ee(m)	uu(m)
2010.6260	2010	229	0.00000	0.00000	0.00000	0.00199	0.00200	0.0		
2010.6288	2010	230	-0.00176	0.00197	0.00798	0.00192	0.00186	0.0		
2010.6315	2010	231	-0.00042	0.00219	0.00470	0.00183	0.00181	0.0		
2010.6342	2010	232	0.00156	0.00082	0.00681	0.00159	0.00156	0.0		
2010.6370	2010	233	-0.00070	0.00209	0.00030	0.00163	0.00161	0.0		
2010.6397	2010	234	-0.00110	0.00000	0.00311	0.00189	0.00182	0.0		
2010.6425	2010	235	-0.00090	-0.00010	-0.00262	0.00167	0.00168	0.0		

Header Comment

Unit mm Time Unit years Scale 1000

Column indexes

1 2 3 4 5 6 7 8

Column names

year doy north east up north_sigma east_sigma up_sigma

Index columns

year doy

Directory E:\Program\TSAAnalyzer\Example\header

Delete **Write**

From the Header Comment Tool Figure, there are some parameters should be input.

- Unit, mm abbreviation for millimeter, although the original file's unit is meter
- Scale, we use scale factor 1000 to convert meter to millimeter.
- Time Unit, years and days supports at present.

- Column indexes and column names, starting from 0 in this example, we use columns year (1), doy (2), north (3), east (4), up (5), north_sigma (6), east_sigma (7) and up_sigma (8).

Index columns supports the following key words: year, month, day, hour, minute, seconds, doy, ymd, hms, mjd

- doy, day of year
 - ymd, year month day, for example 2010101
 - hms, hour minute seconds, for example 120000
 - mjd, Modified Julian Date
- Directory, save new files after adding header comments.

After adding header comments, we could use TSAAnalyzer to load these new files.

End

If this tool can help you, please consider to cite our paper: Wu, D. and H. Yan, et al. (2017). "TSAAnalyzer, a GNSS time series analysis software." GPS Solutions 21(3): 1389-1394.

For bugs and suggestion, please open the issues. Thanks.

Reference

- Anaconda. <https://anaconda.org/>
- Cvxpy. <http://cvxpy.org>
- Goudarzi, M. A. and M. Cocard, et al. (2013). "GPS interactive time series analysis software." GPS Solutions 17 (4): 595-603
- Herring, T. (2003). "MATLAB Tools for viewing GPS velocities and time series." GPS Solutions 7 (3): 194-199
- Kim, S. and K. Koh, et al. (2009). " ℓ_1 Trend Filtering." SIAM Review 51(2): 339-360.
- Little, M. A. and N. S. Jones (2010). Sparse Bayesian step-filtering for high-throughput analysis of molecular machine dynamics. Acoustics Speech and Signal Processing (ICASSP), 2010 IEEE International Conference on, IEEE.
- Matplotlib. <http://matplotlib.org/>
- MOSEK. <https://www.mosek.com/>
- Nikolaidis (2002). Observation of geodetic and seismic deformation.
- Numpy. <http://numpy.org/>
- Pandas. <http://pandas.pydata.org/>
- PyQt. <https://www.riverbankcomputing.com/>
- qtpy. <https://github.com/spyder-ide/qtpy>
- Scipy. <http://scipy.org/>

- Tian, Y. (2011). "iGPS: IDL tool package for GPS position time series analysis." GPS Solutions 15 (3): 299-303

