

INTRODUCTION TO SHIMURA VARIETIES AND THEIR COHOMOLOGY

LECTURER: OLIVIER TAIBI, TYPESETTER: MICHAŁ MRUGAŁA

Plan:

- I) Siegel modular varieties
 - II) General Shimura varieties
 - III) (Étale) Cohomology: Kottwitz conjecture
- (The cohomology gives a realization of the global Langlands correspondence for certain number fields.)

The main reference are Sophie Morel's notes from the IHES '22 summer school.

§1. SIEGEL MODULAR VARIETIES

Analytically: we want to parametrize complex tori. This is fairly simple. Let V be a \mathbf{C} -vector space of dimension $m \geq 1$, $\Lambda \subset V$ a lattice (a discrete subgroup such that V/Λ is compact), then $X = V/\Lambda$ is a complex Lie group, which is a complex torus.

Exercise. — A morphism $f : X = V/\Lambda \rightarrow X' = V'/\Lambda'$ of complex Lie groups is given by a \mathbf{C} -linear map $V \rightarrow V'$ mapping Λ to Λ' .

Question: Which complex tori are algebraizable, i.e. $X \hookrightarrow \mathbb{P}^n(\mathbf{C})$ (equivalent to $X \simeq \underline{X}^{\text{an}}$ for some projective \underline{X} by Chow). Can we find a parametrization?

Example 1.0.1. — Let $n = 1$ complex tori are always algebraic. There is the Weierstrass \wp -function

$$\wp : V/\Lambda \longrightarrow \mathbb{P}^1(\mathbf{C})$$

$$v \longmapsto \wp(v) = \frac{1}{v^2} + \sum_{\lambda \in \Lambda - 0} \frac{1}{(\lambda + v)^2} - \frac{1}{\lambda^2}.$$

It embeds V/Λ in $\mathbb{P}^2(\mathbf{C})$ via $[\wp : \wp' : 1]$ with image $y^2 = P_\Lambda(x)$ where $P_\Lambda \in \mathbf{C}[X]$ has degree 3. The coefficients of P_Λ are Eisenstein series (modular forms).

For $n > 1$, X is “almost never” algebraic.

Recall that X is algebraizable if and only if there exists $\mathcal{L} \in \text{Pic}(X)$ which is ample (see Mumford's Abelian Varieties). Recall that $\text{Pic}(X) \simeq H^1(X, \mathcal{O}_X^\times)$. There is a short exact sequence

$$(1) \quad 0 \longrightarrow \underline{\mathbb{Z}}_X \longrightarrow \mathcal{O}_X \xrightarrow{\exp(2\pi i -)} \mathcal{O}_X^\times \longrightarrow 0$$

so we get a long exact sequence. The map

$$H^0(X, \mathcal{O}_X) \simeq \mathbf{C} \longrightarrow \mathbf{C}^\times \simeq H^0(X, \mathcal{O}_X^\times)$$

is surjective so we get

$$(2) \quad \begin{array}{ccccccc} H^1(X, \mathbb{Z}) & \hookrightarrow & H^1(X, \mathcal{O}_X) & \longrightarrow & H^1(X, \mathcal{O}_X) & \xrightarrow{\delta} & \ker(H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathcal{O}_X)) \\ \downarrow \simeq & & \downarrow \simeq \text{Dolbeault} & & \downarrow \simeq & & \downarrow \simeq \\ H^1(\Lambda, \mathbb{Z}) & & \bar{T} & & H^1(\Lambda, \mathcal{O}(X)^\times) & & \text{Hom}(\wedge^2 \Lambda, \mathbb{Z}) \\ \parallel & & \uparrow \text{pr}_2 & & & & \\ \text{Hom}(\Lambda, \mathbb{Z}) & & T \oplus \bar{T} & & & & \\ & & \uparrow \simeq & & & & \\ & & \text{Hom}_{\mathbf{R}}(V, \mathbf{C}) & & & & \end{array}$$

We have $H^i(V, \mathbb{Z}) = 0$ for all $i > 0$ and $H^i(V, \mathcal{O}_V) = 0$ for all $i > 0$ so $\text{Pic}(V) = 0$. \bar{T} are the antilinear maps $V \rightarrow \mathbb{C}$ and $T = \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$. Observe that

$$\text{Pic}^0(X) = \ker \delta \simeq \frac{\bar{T}}{\text{pr}_2(\text{Hom}(\Lambda, \mathbb{Z}))}$$

is also a complex torus. The last term is the Neron-Severi group

$$\begin{aligned} NS(X) &\simeq \{E : \Lambda \times \Lambda \longrightarrow \mathbb{Z} \text{ alt.} : E(iv_1, iv_2) = E(v_1, v_2), \forall v_1, v_2 \in V\} \\ &= \{\Im H : H : V \times V \longrightarrow \mathbb{C} \text{ Hermitian such that } (\Im H)(\Lambda \times \Lambda) \subset \mathbb{Z}\}. \end{aligned}$$

The Appel-Humbert theorem completely describes $\text{Pic}(X)$ as $\{L(H, \alpha)\}$ with H as above and α an extra datum.

Theorem 1.0.1 (Lefschetz). — *The following are equivalent:*

- 1) H is positive definite.
- 2) $L(H, \alpha)$ is ample (in fact, $L(H, \alpha)^{\otimes 3}$ is enough to embed X).

Let $L \in \text{Pic}(X)$ then

$$\begin{aligned} \phi_L : X &\longrightarrow \text{Pic}^0(X) = \hat{X} \\ x &\longmapsto T_x^* L \otimes L^{-1} \end{aligned}$$

is a morphism of Lie groups (here T_x is translation by x).

Theorem 1.0.2. — *The following are equivalent:*

- L is ample.
- $\ker \phi_L$ is finite.
- ϕ_L is surjective (i.e. an isogeny).

Exercise. — Check that ϕ_L is an isomorphism if and only if $E(\cdot, \cdot)$ is perfect ($\Lambda \simeq \text{Hom}(\Lambda, \mathbb{Z})$).

Definition. — Say that such ϕ_L is a **polarization**. If ϕ_L is an isomorphism, then it is called a **principal polarization**.

Remark. — Not every algebraic X admits a principal polarization, but is isogenous to one that does.

We can define the moduli space

$$\mathcal{A}_n(\mathbb{C}) = \left\{ (X, \phi) : X = V/\Lambda \text{ of dimension } n, \phi : X \longrightarrow \hat{X} \text{ a principal polarization} \right\}.$$

Let (V, Λ, H) be a principally polarized complex torus. Choose a symplectic basis (e_1, \dots, e_{2n}) of Λ , i.e.

$$(E(e_i, e_j))_{i,j} = J_n = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

Exercise. — $L = L(H, \alpha)$ is ample if and only if e_{n+1}, \dots, e_{2n} is a basis of V over \mathbb{C} such that

$$\tau = \text{Mat}_{e_{n+1}, \dots, e_{2n}}(e_1, \dots, e_n)$$

satisfies $\tau = {}^t \tau$ and $\Im(\tau)$ is positive definite.

Definition. — \mathcal{H}_n^+ is the set of such $\tau \in M_n(\mathbb{C})$. There is an algebraic group

$$\mathbf{Sp}_{2n, \mathbb{Z}} : R \longmapsto \{g \in M_{2n}(R) : {}^t g J_n g = J_n\}.$$

There is an action of $\mathbf{Sp}_{2n}(\mathbb{Z})$ on \mathcal{H}_n^+ such that given

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

we have

$$\tau \gamma = (\tau b + d)^{-1}(\tau a + c)$$

(this corresponds to replacing $\underline{e} = (e_1, \dots, e_{2n})$ by $\underline{e}\gamma$).

We prefer left actions: let ${}^t \gamma$ act so that $\gamma \tau = \tau * {}^t \gamma$, i.e.

$$(\tau {}^t c + d)^{-1}(\tau {}^t a + {}^t b) = (a\tau + b)(c\tau + d).$$

This extends to an action of $\mathbf{Sp}_{2n}(\mathbf{R})$ on \mathcal{H}_n^+ . This action is transitive and

$$\begin{aligned} \mathrm{Stab}_{\mathbf{Sp}_{2n}(\mathbf{R})}(iI_n) &\longrightarrow U(n) \\ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} &\longmapsto a + ib \end{aligned}$$

is an isomorphism (this is a maximal compact subgroup).

So $\mathcal{A}_n(\mathbf{C}) \simeq \Gamma_n \backslash \mathcal{H}_n^+$ where $\Gamma_n \simeq \mathbf{Sp}_{2n}(\mathbf{Z})$.

Remark. — There exists $\gamma \in \Gamma_n \setminus \{\pm 1\}$ and $\tau \in \mathcal{H}_n^+$ such that $\gamma\tau = \tau$.

There is a universal object

$$\mathcal{X}_n(\mathbf{C}) = \mathbb{Z}^{2n} \rtimes \Gamma_n \backslash \mathbf{C}^n \times \mathcal{H}_n^+$$

where

$$\gamma(v, \tau) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} (v, \tau) = ((\tau^t c + {}^t d)^{-1} v, \gamma\tau)$$

and

$$(\lambda_1, \lambda_2)(v, \tau) = (v + \lambda_1 + \tau\lambda_2, \tau)$$

for $\lambda_i \in \mathbb{Z}^n$.

There is a morphism $\pi : \mathcal{X}(\mathbf{C}) \rightarrow \mathcal{A}_n(\mathbf{C})$ which admits a section e . The fiber of τ is $[\tau] \simeq \mathbf{C}^n / \Lambda_\tau$ where $\Lambda_\tau = \mathbb{Z}^n \oplus \tau\mathbb{Z}^n$. We get the **Hodge bundle**: take $\Omega^1(V/\Lambda)$ which are translation invariant 1-forms, which is isomorphic to V^* via e^* , then the Hodge bundle is

$$\mathcal{E}_n = e^* \Omega^1_{\mathcal{X}(\mathbf{C})/\mathcal{A}_n(\mathbf{C})} \simeq \Gamma_n \backslash \mathbf{C}^n \times \mathcal{H}_n^+$$

for action (??).

We can apply Schur functors (use the action of \mathfrak{S}_k on $\mathcal{E}_n^{\otimes k}$ to act on subbundles, e.g. $\bigwedge^k \mathcal{E}_n$ for $0 \leq k \leq n$). (Equivalently see \mathcal{E}_n as a $\mathbf{GL}_n(\mathbf{C})$ -bundle on $\mathcal{A}_n(\mathbf{C})$ and apply a holomorphic representation $\rho : (\mathbf{GL}_n(\mathbf{C}) \rightarrow \mathbf{GL}(W))$.) Sections of such vector bundles on $\mathcal{A}_k(\mathbf{C})$ are (level Γ_n , weight ρ) Siegel modular forms on $\mathcal{A}_n(\mathbf{C})$.

Notation: Write

$$M_\rho(\Gamma_n) = \{f \in \Gamma(A_n(\mathbf{C}), \rho(\mathcal{E}_n)) : f \text{ is holomorphic at } \infty\}$$

(the last condition is automatic if $n > 1$). We write

$$S_\rho(\Gamma_n) = \{f : \text{vanish at } \infty\} \subset M_\rho(\Gamma_n)$$

for the set of **cuspidal forms**.

We want a group theoretic description of the complex structure on $\mathcal{A}_n(\mathbf{C})$ and these vector bundles on $\mathcal{A}_n(\mathbf{C})$.

We have $Z(U(n)) \simeq U(1)$ and its centralizer in $\mathbf{Sp}_{2n}(\mathbf{R})$ is $U(n) = K(\mathbf{R})$ where $K \hookrightarrow \mathbf{Sp}_{2n, \mathbf{R}}$ is an algebraic subgroup.

Over \mathbf{C} we have

$$Z(K_{\mathbf{C}}) \simeq \mathbf{GL}_{1, \mathbf{C}} \hookrightarrow \mathbf{Sp}_{2n, \mathbf{C}}.$$

This determines two opposite parabolic subgroups $Q_+ = K_{\mathbf{C}}N_+$, $Q_- = K_{\mathbf{C}}N_-$.

1.1. Siegel modular forms as automorphic forms Let $\rho : \mathbf{GL}_n(\mathbf{C}) \rightarrow \mathbf{GL}(W)$ be a holomorphic (equivalently algebraic) representation. **Siegel modular forms** are

$$M_\rho(\Gamma_n) = \left\{ \begin{array}{l} f : \mathcal{H}_n^+ \rightarrow W \\ \text{holomorphic} \end{array} \middle| \begin{array}{l} \forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_n, \forall \tau \in \mathcal{H}_n^+, f(\gamma\tau) = \rho(c\tau + d)f(\tau) \\ \text{and } f \text{ holomorphic at } \infty \end{array} \right\} \\ \subset H^0(\mathcal{A}_n(\mathbf{C}), {}^\rho \mathcal{E}_n).$$

${}^\rho \mathcal{E}_m$ comes from a $\mathbf{Sp}_{2n}(\mathbf{R})$ -equivariant vector bundle on

$$\begin{array}{ccc} \mathcal{H}_n^+ & \hookrightarrow & \mathbf{Sp}_{2n}(\mathbf{C})/Q_-(\mathbf{C}) \\ \simeq \uparrow & & \\ \mathbf{Sp}_{2n}(\mathbf{R}) & \longrightarrow & \mathbf{Sp}_{2n}(\mathbf{R})/U(n) \end{array}$$

Define

$$j : \mathbf{Sp}_{2n}(\mathbf{R}) \times \mathcal{H}_n^+ \longrightarrow \mathbf{GL}_n(\mathbf{C}) \\ \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \tau \longmapsto c\tau + d.$$

This is a cocycle

$$j(gg', \tau) = j(g, g'\tau)j(g', \tau)$$

(so $j(-, i)|_{U(n)} : U(n) \rightarrow \mathbf{GL}_n(\mathbf{C})$ is a morphism). To $f \in M_\rho(\Gamma_n)$ associate

$$\begin{aligned} \phi_f : \Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R}) &\longrightarrow W \\ g &\longmapsto \phi_f(g) = \rho(j(g, i))^{-1}f(gi) \end{aligned}$$

a smooth function. Let $g \in \mathbf{Sp}_{2n}(\mathbf{R})$ and $k \in U(n)$, then

$$\phi_f(gk) = \rho(j(k, i))^{-1}f(gi).$$

Assume $W = \mathbf{C}$ for simplicity, e.g. ${}^\rho \mathcal{E}_n = (\wedge^n \mathcal{E}_n)^{\otimes k}$. Then

$$\phi_f \in \mathcal{A}(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R})) \subset C^\infty(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R}, \mathbf{C}))$$

(? details). This space has actions by \mathfrak{g} and $U(n)$. By the Cauchy-Riemann equations f is holomorphic if and only if ϕ_f is killed by $\text{Lie } N_- \subset \mathfrak{g} = \mathbf{C} \otimes_{\mathbf{R}} \text{Lie } \mathbf{Sp}_{2n}(\mathbf{R})$. Note that $\text{Lie}(\mathbf{Sp}_{2n}(\mathbf{R}))$ acts on $C^\infty(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R}))$ by

$$(X \cdot \phi)(g) = \left. \frac{d}{dt} \right|_{t=0} \phi(ge^{tX}).$$

ϕ_f lies in some generalized Verma module in $\mathcal{A}(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R}))$.

If $f \in S_\rho(\Gamma_m)$ (vanishes at ∞) then

$$\phi_f \in \mathcal{A}_{\text{cusp}}(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R})) \subset \mathcal{A}^2(-) \subset \mathcal{A}(-)$$

and $\mathcal{A}^2(-)$ decomposes inside $L^2(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R}))$ with the action of $\mathbf{Sp}_{2n}(\mathbf{R})$. This means that cusp forms have fast decay at cusps.

As a $(\mathfrak{g}, U(n))$ -module,

$$\mathcal{A}_{\text{cusp}} \subset \mathcal{A}^2(\Gamma_n \backslash \mathbf{Sp}_{2n}(\mathbf{R})) \simeq \bigoplus_{\substack{\pi \text{ irr} \\ (\mathfrak{g}, U(n))\text{-mod}}} \pi^{\oplus m(\pi)}.$$

Siegel cusp forms correspond to special vectors in some of these π s ($U(n)$ -equivariant and killed by $\text{Lie } N$).

1.2. Level structures Let $X = V/\Lambda$ be a complex torus with a principal polarization $\phi : X \xrightarrow{\sim} \widehat{X}$. For $M \geq 1$

$$X[M] := \ker \left(X \xrightarrow{\times M} X \right) = \frac{1}{M} \Lambda / \Lambda \simeq (\mathbb{Z}/M)^{2n}.$$

The map $[M]_X : X \rightarrow X$ is an isogeny (i.e. surjective with finite kernel). For all isogenies $f : X \rightarrow Y$ inducing $\widehat{f} = f^* : \widehat{Y} \rightarrow \widehat{X}$, also an isogeny. We get the Weil pairing

$$\begin{aligned} \ker f \times \ker \widehat{f} &\longrightarrow \mathbb{C}^\times \\ (x, [L]) &\longmapsto \langle x, [L] \rangle. \end{aligned}$$

Choose $t : f^* L \xrightarrow{\sim} \mathcal{O}_X$ we have

$$\begin{array}{ccc} T_x^* f^* L & \xrightarrow{T_x^*(t)} & T_x^* \mathcal{O}_X \\ \downarrow \simeq & & \downarrow \simeq \\ f^* L & \xrightarrow{t \times \langle x, [L] \rangle} & \mathcal{O}_X. \end{array}$$

$f = [M]_X$ is a special case, then we get $X[M] \times \widehat{X}[M] \rightarrow \mu_M(\mathbb{C})$ and using a polarization we get

$$\langle \cdot, \cdot \rangle_\phi : X[M] \times X[M] \longrightarrow \mu_M(\mathbb{C}).$$

Proposition 1.2.1. — $\langle \cdot, \cdot \rangle_\phi$ is alternating and non-degenerate.

Proof. Recall that $\phi = \phi_L$ for some $L = L(H, \alpha)$, let $E = \Im H : \Lambda \times \Lambda \rightarrow \mathbb{Z}$. Then

$$\begin{array}{ccc} X[M] \times X[M] & \xrightarrow{\langle \cdot, \cdot \rangle_\phi} & \mu_M(\mathbb{C}) \\ \downarrow \simeq & & \uparrow \exp(2\pi i -) \\ \left(\frac{1}{M} \Lambda / \Lambda \right)^2 & \xrightarrow{ME(\cdot, \cdot)} & \frac{1}{M} \mathbb{Z} / \mathbb{Z} \end{array}$$

□

Definition. — Temporarily we define a level structure on (X, ϕ) to be

$$(\mathbb{Z}/M)^{2n} \xrightarrow[\eta]{\sim} X[M]$$

such that $\eta^* \langle \cdot, \cdot \rangle_\phi$ is the standard pairing for metric J_n .

Fact. — By strong approximation $\mathbf{Sp}_{2n}(\mathbb{Z}) \twoheadrightarrow \mathbf{Sp}_{2n}(\mathbb{Z}/M\mathbb{Z})$. Define $\Gamma_m(M)$ to be the kernel.

Corollary 1.2.1. — There is a bijection

$$\{(X, \phi, \eta) | \text{PPAV with a level } M \text{ structure}\} / \sim \simeq \Gamma_n(M) \backslash \mathcal{H}_n^+ = \mathcal{A}'_n(M)(\mathbb{C}).$$

Exercise. — For $M \geq 3$, for all $\tau \in \mathcal{H}_n^+$ show that $\text{Stab}_{\Gamma_n(M)}(\tau) = \{1\}$. (?)

We get a tower $(\mathcal{A}'_n(M)(\mathbb{C}))_{M \geq 1}$ ordered by divisibility. For $M \mid M'$ we get $\mathcal{A}'_n(M')(\mathbb{C}) \rightarrow \mathcal{A}'_n(M)(\mathbb{C})$.

Given (X, ϕ)

$$\{\text{level } M \text{ structures on } (X, \phi)\}$$

is a right $\mathbf{Sp}_{2n}(\mathbb{Z}/M\mathbb{Z})$ -torsor which gives us an action of

$$\mathbf{Sp}_{2n}(\widehat{\mathbb{Z}}) = \varprojlim_M \mathbf{Sp}_{2n}(\mathbb{Z}/M)$$

on this tower.

Also

$$\mathcal{A}'_n(M)(\mathbb{C}) \simeq \mathcal{A}'_n(M')(\mathbb{C}) / (K(M)/K(M'))$$

where

$$K(M) = \ker \left(\mathbf{Sp}_{2n}(\widehat{\mathbb{Z}}) \longrightarrow \mathbf{Sp}_{2n}(\mathbb{Z}/M) \right).$$

The quotient $K(M)/K(M')$ is a finite group.

1.3. Hecke operators (adelically) The goal is to define more natural maps between $\mathcal{A}_n'(M)(\mathbf{C})$. The basic idea is that given (X, ϕ, η) , we should also consider isogeneous complex tori (i.e. quotients of X by finite subgroups). But there are some problems: this is not strictly compatible with principal polarizations. Let $f : X \rightarrow Y$ be an isogeny, ϕ be a principal polarization for Y , then $f^*\phi := \hat{f} \circ \phi \circ f$ has degree $(\deg f)^2$, so it is not principal unless f is an isomorphism.

There are two solutions:

- 1) Rescale polarizations.
- 2) Consider quasi-isogenies

$f \in \mathbf{Q} \otimes \text{Hom}(X, Y)$ such that $\exists M \geq 1$ with $Mf \in \text{Hom}(X, Y)$ an isogeny.

Let's do both.

Recall the ring of adeles $\mathbb{A} = \mathbf{R} \times \mathbb{A}_f$ where

$$\mathbb{A}_f = \prod_p' (\mathbf{Q}_p, \mathbb{Z}_p) = \left\{ (x_p)_p \text{ prime} \left| \begin{array}{l} x_p \in \mathbf{Q}_p \\ \exists \text{ finite } S \text{ such that } \forall p \notin S, x_p \in \mathbb{Z}_p \end{array} \right. \right\}.$$

Recall that

$$\begin{aligned} \text{lattices in } \mathbf{Q}\Lambda &\leftrightarrow \left\{ (\Lambda'_p)_p \left| \begin{array}{l} \Lambda'_p \subset \mathbf{Q}_p \otimes_{\mathbf{Z}} \Lambda \text{ is a } \mathbb{Z}_p\text{-lattice} \\ \exists \text{ finite } S \text{ such that } \forall p \notin S, \Lambda'_p = \mathbb{Z}_p \Lambda \end{array} \right. \right\} \\ &\leftrightarrow \mathbf{GL}(\mathbb{A}_f \otimes \Lambda) / \mathbf{GL}(\hat{\mathbf{Z}} \otimes \Lambda). \end{aligned}$$

Proof. Reduce to the case where

$$M\Lambda \subset \Lambda' \subset \frac{1}{M}\Lambda$$

and use the chinese remainder theorem. \square

Proposition 1.3.1. — *Let (X, ϕ) be a principally polarized abelian variety.*

- (a) *Let L be the set of principally polarized abelian varieties (X', ϕ') quasi-isogeneous to (X, ϕ) , i.e. there exists a quasi-isogeny $f : X' \dashrightarrow X$ such that $f^*\phi = c\phi'$, where $c \in \mathbf{Q}_{>0}$.*
- (b) *Let R be the set of $(\Lambda'_p)_p$ such that*

$$\Lambda'_p \subset V_p X := \mathbf{Q}_p \otimes_{\mathbf{Z}_p} T_p X$$

is a \mathbb{Z}_p -lattice such that there exists k_p making $p^{k_p} \langle \cdot, \cdot \rangle|_{\Lambda'_p \times \Lambda'_p}$ take values in $\mathbb{Z}_p(1) := \varprojlim_k \mu_{p^k}(\mathbf{C})$ and is perfect, as well as there is a finite S such that for all $p \notin S$

$$\Lambda'_p = T_p X := \varprojlim_k X[p^k].$$

Then

$$L / \sim \simeq R.$$

This is also isomorphic to the set of $\mathbf{GSp}_{2n}(\hat{\mathbf{Z}})$ -orbits of symplectic trivializations

$$(\mathbb{A}_f^{2n}, \text{standard } \langle \cdot, \cdot \rangle) \xrightarrow{\sim} \left(\mathbf{Q} \otimes \prod_p T_p X, \langle \cdot, \cdot \rangle_{\phi} \right).$$

Here \mathbf{GSp}_{2n} is the \mathbf{Z} -group scheme

$$\mathbf{GSp}_{2n}(R) = \{ (g, c) | g \in M_{2n}(R), c \in R^{\times}, {}^t g J_n g = c J_n \}.$$

Definition. — A **level structure** for (X, ϕ) is an isomorphism $(\mathbf{Z}/M)^{2n} \xrightarrow{\sim} X[M]$. $\mathbf{Z}/M \xrightarrow{\sim} \mu_M(\mathbf{C})$ such that the obvious diagram commutes.

We have

$$\begin{aligned} \mathcal{A}_n(M)(\mathbf{C}) &\simeq \{ (X, \phi, \eta) | \text{PPAV with level } M \text{ structure} \} / \sim \\ &\simeq \left\{ (X', \phi') \left| K(M)\text{-orbit of trivialization of } \mathbf{Q} \otimes \prod_p T_p X' \right. \right\} / \text{quasi-isogeny} \\ &\simeq \mathbf{GSp}_{2n}(\mathbf{Q}) \setminus \left(\mathcal{H}_n^{\pm} \times \mathbf{GSp}_{2n}(\mathbb{A}_f) / K(M) \right) \end{aligned}$$

where

$$\mathcal{H}_n^\pm = \mathcal{H}_n^+ \coprod \mathcal{H}_n^-$$

has an action of $\mathbf{GSp}_{2n}(\mathbf{R})$ and

$$K(M) := \ker \left(\mathbf{GSp}_{2n}(\hat{\mathbb{Z}}) \longrightarrow \mathbf{GSp}_{2n}(\mathbb{Z}/M) \right).$$