

高级算法

Advanced Topics in Algorithms

陈旭

数据科学与计算机学院



中山大學
SUN YAT-SEN UNIVERSITY

Chapter 1:

Simple Decisions Models

Some definitions

Definition

argmax is defined by the following equivalence:

$$x^* \in \operatorname{argmax}_{x \in X} f(x) \iff f(x^*) = \max_{x \in X} f(x)$$

Definition

A choice of behavior in a single-decision problem is called an *action*. The set of alternative actions available will be denoted as **A**. This will either be discrete set, e.g., $\{a_1, a_2, \dots, \}$, or a continuous set, .e.g., the unit interval $[0, 1]$.

More...

Definition

A *payoff* is a function $\pi : \mathbf{A} \rightarrow R$ that associates a numerical value with every action $a \in \mathbf{A}$.

Definition

An action a^* is an *optimal action* if

$$\pi(a^*) \geq \pi(a) \quad \forall a \in \mathbf{A}.$$

or equivalently, $a^* \in \operatorname{argmax}_{a \in \mathbf{A}} \pi(a)$.

Definition

An *affine transformation* changes payoff $\pi(a)$ into $\pi'(a)$ as

$$\pi'(a) = \alpha\pi(a) + \beta$$

where α, β are constants independent of a and $\alpha > 0$.

Theorem

The optimal action is unchanged if payoffs are altered by an affine transformation.

Proof

because $\alpha > 0$, we have

$$\begin{aligned} \operatorname{argmax}_{a \in \mathbf{A}} \pi'(a) &= \operatorname{argmax}_{a \in \mathbf{A}} [\alpha\pi(a) + \beta] \\ &= \operatorname{argmax}_{a \in \mathbf{A}} \pi(a). \end{aligned}$$

Example

The Convent Fields Soup Company needs to determine the price p . The demand function is:

$$Q(p) = \begin{cases} Q_0 \left(1 - \frac{p}{p_0}\right) & \text{if } p < p_0, \\ 0 & \text{if } p \geq p_0. \end{cases}$$

The payoff is $\pi(p) = (p - c)Q(p)$ where c is the unit production cost.

- Solving, we have $p^* = \frac{1}{2}(p_0 + c)$.
- Now, let say we need to consider a fixed cost to build the factory, the payoff function is $\pi(p) = (p - c)Q(p) - B$, where B is a constant. What is p^* ?

Uncertainty

Modeling uncertainty

- If uncertainty exists, we compare the expected outcome for each action.
- Let X be the set of states with $P(X = x)$.
- Payoff for adopting action a is:

$$\pi(a) = \sum_{x \in X} \pi(a|x)P(X = x)$$

- An optimal action is

$$a^* \in \operatorname{argmax}_{a \in A} \sum_{x \in X} \pi(a|x)P(X = x).$$

Example

- An investor has \$1000 to invest in one year. The available actions (1) put the money in the bank with 7% interest per year; (2) invest in stock which returns \$1500 if the stock market is good or returns \$600 if the stock market is bad.
- $P(\text{Good}) = P(\text{Bad}) = 0.5$.
- Expected payoff:
 - ① $\pi(a_1) = \$1070$;
 - ② $\pi(a_2) = 1500/2 + 600/2 = \1050 .
- So a_1^* and we should put the money in the bank.

Definition

Let $\Omega = \{\omega_1, \omega_2, \dots\}$ be the set of possible outcomes.

- We say $\omega_i \succ \omega_j$ if an individual *strictly prefers* outcome ω_i over ω_j .
- If the individual is indifferent: $\omega_i \sim \omega_j$.
- Either prefer or indifferent: $\omega_i \succeq \omega_j$.

Definition

An individual will be called **rational under certainty** if his preference for outcomes satisfy the following conditions:

- (Completeness) Either $\omega_i \succeq \omega_j$ or $\omega_j \succeq \omega_i$.
- (Transitivity) If $\omega_i \succeq \omega_j$ and $\omega_j \succeq \omega_k$, then $\omega_i \succeq \omega_k$.

Definition

A **utility function** is a function $u : \Omega \rightarrow \mathbf{R}$ such that

$$u(\omega_i) > u(\omega_j) \iff \omega_i \succ \omega_j$$

$$u(\omega_i) = u(\omega_j) \iff \omega_i \sim \omega_j$$

The immediate consequence of this definition is an individual who is rational under certainty should seek to maximize his utility.

What happens when an action does not produce a definite outcome and instead, we allow each outcome to occur with a known probability?

Definition

A **simple lottery**, λ , is a set of probabilities for the occurrence of every $\omega \in \Omega$. The probability that outcome ω occurs is $p(\omega|\lambda)$. The set of all possible lotteries is denoted as **A**.

Theorem

Expected Utility Theorem: *If an individual is rational, then we can define a utility function $u : \Omega \rightarrow \mathbf{R}$ and the individual will maximize the payoff function $\pi(a)$ (or the expected utility) given by*

$$\pi(a) = \sum_{\omega \in \Omega} p(w|\lambda(a))u(\omega)$$

Definition

An individual whose utility function satisfies

- $E(u(w)) < u(E(w))$, it is said to be **risk averse**,
- $E(u(w)) > u(E(w))$, it is said to be **risk prone**,
- $E(u(w)) = u(E(w))$, it is said to be **risk neutral**.

Example

Someone flip a coin. If it is head (tail), you get \$1 (\$1M). Your utility function can be:


- $u(x) = x$,
- $u(x) = x^2$,

Classify the above as risk averse, risk prone and risk neutral utility function.


https://en.wikipedia.org/wiki/Risk_aversion

https://en.wikipedia.org/wiki/Risk_aversion

Concave




Convex



Linear

Risk Neutral



Up to now, we assume finding an optimal action a^* from a given set \mathbf{A} . But the selection can be *randomized*. Does this allow one to achieve a higher payoff?

Definition

We specify a **general behavior** β by giving a list of probabilities with which each available action is chosen. We denote the probability that action a is chosen by $p(a)$ and $\sum_{a \in \mathbf{A}} p(a) = 1$. The set of all randomizing behavior is denoted by \mathbf{B} . The payoff of using behavior β is

$$\pi(\beta) = \sum_{a \in \mathbf{A}} p(a)\pi(a).$$

An **optimal behavior** β^* is one for which

$$\pi(\beta^*) \geq \pi(\beta) \quad \forall \beta \in \mathbf{B}.$$

or $\beta^* \in \operatorname{argmax}_{\beta \in \mathbf{B}} \pi(\beta)$.

Definition

The support of a behavior β is the set $\mathbf{A}(\beta) \subseteq \mathbf{A}$ of all the actions for which β specifies $p(a) > 0$.

Theorem

Let β^ be an optimal behavior with support \mathbf{A}^* . Then*

$$\pi(a) = \pi(\beta^*) \quad \forall a \in \mathbf{A}^*.$$

The consequence of the above theorem is that if a randomized behavior is optimal, then two or more actions are optimal as well. So randomization is not necessary but it may be used to break a tie.

Example

A firm may make one of the marketing actions $\{a_1, a_2, a_3\}$. The profit for each action depends on the state of the economy $\mathbf{X} = \{x_1, x_2, x_3\}$:

	x_1	x_2	x_3
a_1	6	5	3
a_2	3	5	4
a_3	5	9	1

If $P(X = x_1) = 1/2$, $P(X = x_2) = P(X = x_3) = 1/4$. What *are* the optimal behaviors?

Answer

Because $\pi(a_1) = \pi(a_3) = 5$ and $\pi(a_2) = 3.75$, optimal randomizing behaviors have support $\mathbf{A}^* = \{a_1, a_3\}$ with $p(a_1) = p$ and $p(a_3) = 1 - p$ ($0 < p < 1$). Using either a_1 or a_3 with probability 1 is also an optimal behavior.

Reading

- Chapter 1 Simple Decision Models
- Appendix A Constrained Optimisation