

Contents

1	Scientific, Technical, and Management Section	2
1.1	Science Objectives	2
1.1.1	The Inflationary Gravitational Wave Background	2
1.1.2	Neutrinos and Light Relics	3
1.1.3	CMB spectral distortion science	3
1.2	The Challenges: Foregrounds, systematics	4
1.3	Current and Forthcoming Efforts and the CMB Probe	4
1.4	State of Technologies	4
1.5	Mission Study, and Management Plan	4
2	Curriculum Vitae	10
3	Summary of Work Effort	19
4	Current and Pending Support	19
5	Letters of Support	33
6	Budget Details - Narrative	37
6.1	Team, and Work Effort	37
6.1.1	Funded Team Members	37
6.1.2	Non-Funded Team Members	37
6.2	Costing Principles	37
6.3	University of Minnesota Budget	37
6.3.1	Direct Labor	37
6.3.2	Supplies	37
6.3.3	Travel	37
6.3.4	Other Direct Costs	37
6.3.5	Facilities and Administrative Costs	37
7	Budget Sheets	38

1 Scientific, Technical, and Management Section

0.5 pg. executive summary goes here: the sky is sunny

1.1 Science Objectives

5 pages for all science goals including the (temporary) two sections below.

1.1.1 The Inflationary Gravitational Wave Background

The verbiage below is taken from another proposal. Here we need to explain what are the science objectives of the CMBProbe, how the science objectives relate to the current state of knowledge, and to NASA's goals

The paradigm of inflation [1, 2, 3, 4, 5] makes several predictions that are consistent with all current astrophysical measurements [6, 7, 8, 9]. A robust prediction of inflation is the existence of a stochastic background of gravitational radiation with an amplitude depending on the mechanism driving the accelerated expansion [10, 11, 12, 13, 14]. In most scenarios, this ‘inflationary gravitational wave background’ (IGB) is predicted to have a spatial power spectrum whose amplitude is proportional to the energy scale of inflation $V^{1/4}$ via $V^{1/4} = 3.7 \times 10^{16} r^{1/4}$ GeV, where V is the inflaton potential and r is the ratio of the temperature quadrupoles produced by gravitational waves and by density perturbations. There are theoretical reasons $V^{1/4}$ may be close to the Grand Unification scale of 10^{16} GeV, suggesting detectable r values between ~ 0.001 and ~ 0.1 . In addition to determining the energy scale of inflation, measurements of the IGB probe the scalar field potential at or above the Planck scale, which is particularly relevant for inflation models motivated by string theory [15]. Measurements of the IGB thus probe fundamental physics at the highest possible energy scales.



Figure 1: Sample Figure of Sunny Skies

The most promising way to search for the IGB is through its signature on the CMB polarization [16, 17]. Primordial energy density perturbations produce only a curl-free, or ‘E-mode’, pattern of polarization. Gravitational waves also produce a curl, or ‘B-mode’, pattern of polarization that density perturbations cannot produce [18, 19]. The amplitude of the B mode is related to the energy scale of inflation by $V^{1/4} = 2 \times 10^{16} (B_{peak}/0.1 \mu\text{K})^{1/2}$ GeV, where B_{peak} is the amplitude of the power spectrum of the B mode in μK at $\ell = 80$; see Fig. 1. In its recent report New Worlds New Horizons (NWNH), the decadal survey committee strongly endorsed sub-orbital searches for the B-mode signal from inflation saying that “The convincing detection of B-mode polarization in the CMB produced in the epoch of reionization would represent a watershed discovery.” [20]

B-mode signatures near the expected IGB peak at $\ell = 80$ have recently been detected by BICEP2 [21]. However, the combination of Planck data with those from the BICEP2 and Keck Array collaborations have demonstrated that the B-mode signal measured is entirely consistent with contributions from polarized emission of Galactic dust and the signal from the gravitational lensing of CMB photons by the large scale structure of the Universe (see Section ??) [22, 23?]. These data




Figure 2: [Placeholders] *Left*: Neutrino mass constraints as a function of the prior on τ . *Right* N_{eff} Forecasts as a function of resolution in arc-min and temperature noise in $\mu\text{K-arcmin}$ assuming $f_{\text{sky}} = 0.7$.

give an upper limit of $r < 0.09$ at 95% confidence level. Most importantly, the constraint is largely limited by Planck’s noisy measurement of the dust properties in the 353 GHz band; a noiseless dust map could shrink the constraint by a factor of two [22]. Further progress — detections or improved limits — requires instruments with higher sensitivity at *both* the dust and CMB frequency bands so that this Galactic foreground can be properly identified and removed.

1.1.2 Neutrinos and Light Relics

Much of the information about our thermal history and the particle content of the universe is encoded in the T and E power spectra. A high-precision measurement of these spectra over the full sky is expected to significantly improve our understanding of the post-inflationary universe. This is particularly true in E -mode polarization where far fewer modes have been measured at the level of cosmic variance.

1.1.3 CMB spectral distortion science

In addition to the CMB temperature and polarization anisotropies targeted by CMB imagers, *unique* new information about early-universe physics can be gained by studying the energy spectrum of the CMB [24, 25, 26, 27]. The measurements of COBE/FIRAS have shown that the average CMB spectrum is extremely close to that of a blackbody at a temperature $T_0 = (2.726 \pm 0.001) \text{ K}$ [28, 29]. However, several standard processes are expected to distort the CMB spectrum [e.g., 30] at a level that is within reach of present-day technology [31, 32]. The classical distortion shapes are known as Compton- y and chemical potential (μ -type) distortions [33, 34] and are caused by energy exchange of CMB photons with free electrons. A μ -distortion can only be produced in a hot and dense environment present at redshifts $z \gtrsim 5 \times 10^4$, while y -type distortions appear at lower redshifts. This makes μ -distortions a unique messenger from the early Universe.

Future studies of the CMB spectrum will open a new unexplored window to early phases before the recombination era ($z > 10^3$), which cannot be probed in any other way. This will not only allow us to test the standard cosmological paradigm (e.g., inflation and reionization) but also opens up a huge discovery space to non-standard physics (e.g., decaying/annihilating particles) and new tests of the nature of dark matter. This immense potential and complementarity with CMB anisotropy

studies has made CMB spectral distortions an important future target and identifying experimental routes towards measuring these tiny signals from the early Universe will be one objective of the proposed mission study.

The largest guaranteed distortion is caused by the late-time energy release of forming structures and from reionization [35, 36, 37, 38, 39], creating a y -type distortion with $y \simeq 2 \times 10^{-6}$ [e.g., 39, 40]. This signal is only one order of magnitude below the current limit from COBE/FIRAS and, even with most pessimistic assumptions about foregrounds, should be clearly seen with next-generation spectrometers, telling us about the total energy output of first stars, AGN and galaxy clusters. In particular, group-size clusters ($M \simeq 10^{13} M_{\odot}$) contribute to this distortion. These are still sufficiently hot (temperature $kT_e \simeq 1$ keV) to create a visible relativistic temperature correction to this large y -distortion, which could be used to constrain cluster feedback models [40]. These two inevitable signals probe the low-redshift Universe and provide clear targets for future spectral distortions measurements and their requirements in the presence of foregrounds.

Next generation CMB spectrometers are also expected to improve the μ -distortion limits. This will allow us to place stringent bounds on the presence of long-lived decaying particles [41, 42, 43, 44] and other new physics [e.g., 45, 46, 47, 48, 49], but a clear target is predicted from the dissipation of small-scale perturbation....

1.2 The Challenges: Foregrounds, systematics

3 pages. Discuss the challenge of Foregrounds and Systematics.

1.3 Current and Forthcoming Efforts and the CMB Probe

2.5 pages. S3 experiments, forthcoming S4, Baselines CMB Probe options and their complementarity with S4.

1.4 State of Technologies

2 pages. Discuss the technologies, their TRL, and what will be studied

1.5 Mission Study, and Management Plan

1.5 pages; Describe what we want to do, who is doing what, what we are funding

References

- [1] A. H. Guth. Inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D.*, 23:347–356, January 1981.
- [2] A. D. Linde. A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems. *Phys. Lett.*, B108:389–393, 1982.
- [3] A. Albrecht and P. J. Steinhardt. Cosmology for grand unified theories with radiatively induced symmetry breaking. *Phys. Rev. Lett.*, 48:1220–1223, 1982.
- [4] K. Sato. First-order phase transition of a vacuum and the expansion of the Universe. *MNRAS*, 195:467–479, May 1981.
- [5] E. W. Kolb and M. S. Turner. *The Early Universe*. Addison-Wesley, Redwood City, CA, 2nd. edition, 1994.
- [6] D. N. Spergel, R. Bean, O. Doré, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page, H. V. Peiris, L. Verde, M. Halpern, R. S. Hill, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright. Three-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Implications for Cosmology, June 2007.
- [7] M. Tegmark, D. Eisenstein, M. Strauss, D. Weinberg, M. Blanton, J. Frieman, M. Fukugita, J. Gunn, A. Hamilton, G. Knapp, R. Nichol, J. Ostriker, N. Padmanabhan, W. Percival, D. Schlegel, D. Schneider, R. Scoccimarro, U. Seljak, H. Seo, M. Swanson, A. Szalay, M. Vogeley, J. Yoo, I. Zehavi, K. Abazajian, S. Anderson, J. Annis, N. Bahcall, B. Bassett, A. Berlind, J. Brinkmann, T. Budavari, F. Castander, A. Connolly, I. Csabai, M. Doi, D. Finkbeiner, B. Gillespie, K. Glazebrook, G. Hennessy, D. Hogg, Z. Ivezic, B. Jain, D. Johnston, S. Kent, D. Lamb, B. Lee, H. Lin, J. Loveday, R. Lupton, J. Munn, K. Pan, C. Park, J. Peoples, J. Pier, A. Pope, M. Richmond, C. Rockosi, R. Scranton, R. Sheth, A. Stebbins, C. Stoughton, I. Szapudi, D. Tucker, D. Vanden Berk, B. Yanny, and D. York. Cosmological Constraints from the SDSS Luminous Red Galaxies. *Phys. Rev. D.*, 74:123507, 2006.
- [8] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al. Planck 2015 results. XIII. Cosmological parameters. *ArXiv e-prints*, February 2015.
- [9] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, F. Arroja, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, and et al. Planck 2015. XX. Constraints on inflation. *ArXiv e-prints*, February 2015.
- [10] A. A. Starobinsky. Dynamic of Phase Transition in the New Inflationary Universe Scenario and Generation of Perturbationss. *Phys. Lett.*, B117:175–178, 1982.
- [11] A. A. Starobinskii. The Perturbation Spectrum Evolving from a Nonsingular Initially De-Sitter Cosmology and the Microwave Background Anisotropy. *Soviet Astronomy Letters*, 9:302–+, June 1983.

- [12] V. A. Rubakov, M. V. Sazhin, and A. V. Veryaskin. Graviton creation in the inflationary universe and the grand unification scale. *Phys. Lett. B.*, 115:189–192, September 1982.
- [13] L. P. Grishchuk. Amplification of gravitational waves in an isotropic universe. *Sov. Phys. JETP*, 40:409–415, 1975.
- [14] L. F. Abbott and M. B. Wise. Constraints on generalized inflationary cosmologies. *Nuclear Physics B*, 244:541–548, October 1984.
- [15] K. N. Abazajian, K. Arnold, J. Austermann, B. A. Benson, C. Bischoff, J. Bock, J. R. Bond, J. Borrill, I. Buder, D. L. Burke, E. Calabrese, J. E. Carlstrom, C. S. Carvalho, C. L. Chang, H. C. Chiang, S. Church, A. Cooray, T. M. Crawford, B. P. Crill, K. S. Dawson, S. Das, M. J. Devlin, M. Dobbs, S. Dodelson, O. Doré, J. Dunkley, J. L. Feng, A. Fraisse, J. Gallicchio, S. B. Giddings, D. Green, N. W. Halverson, S. Hanany, D. Hanson, S. R. Hildebrandt, A. Hincks, R. Hlozek, G. Holder, W. L. Holzapfel, K. Honscheid, G. Horowitz, W. Hu, J. Hubmayr, K. Irwin, M. Jackson, W. C. Jones, R. Kallosh, M. Kamionkowski, B. Keating, R. Keisler, W. Kinney, L. Knox, E. Komatsu, J. Kovac, C.-L. Kuo, A. Kusaka, C. Lawrence, A. T. Lee, E. Leitch, A. Linde, E. Linder, P. Lubin, J. Maldacena, E. Martinec, J. McMahon, A. Miller, V. Mukhanov, L. Newburgh, M. D. Niemack, H. Nguyen, H. T. Nguyen, L. Page, C. Pryke, C. L. Reichardt, J. E. Ruhl, N. Sehgal, U. Seljak, L. Senatore, J. Sievers, E. Silverstein, A. Slosar, K. M. Smith, D. Spergel, S. T. Staggs, A. Stark, R. Stompor, A. G. Vieregg, G. Wang, S. Watson, E. J. Wollack, W. L. K. Wu, K. W. Yoon, O. Zahn, and M. Zaldarriaga. Inflation Physics from the Cosmic Microwave Background and Large Scale Structure. *ArXiv e-prints*, September 2013.
- [16] M. Kamionkowski, A. Kosowsky, and A. Stebbins. Statistics of Cosmic Microwave Background Polarization. *Phys. Rev. D.*, 55:7368–7388, June 1997.
- [17] U. Seljak and M. Zaldarriaga. Signature of Gravity Waves in the Polarization of the Microwave Background. *Phys. Rev. Lett.*, 78:2054–2057, March 1997. astro-ph/9609169.
- [18] M. Kamionkowski, A. Kosowsky, and A. Stebbins. A Probe of Primordial Gravity Waves and Vorticity. *Phys. Rev. Lett.*, 78:2058–2061, March 1997. astro-ph/9609132.
- [19] M. Zaldarriaga and U. Seljak. All-sky analysis of polarization in the microwave background. *Phys. Rev. D.*, 55:1830–1840, 1997.
- [20] Committee for a Decadal Survey of Astronomy and Astrophysics. *New Worlds, New Horizons in Astronomy and Astrophysics*. National Academy Press, 2010.
- [21] P. A. R. Ade, R. W. Aikin, D. Barkats, S. J. Benton, C. A. Bischoff, J. J. Bock, J. A. Brevik, I. Buder, E. Bullock, C. D. Dowell, L. Duband, J. P. Filippini, S. Fliescher, S. R. Golwala, M. Halpern, M. Hasselfield, S. R. Hildebrandt, G. C. Hilton, V. V. Hristov, K. D. Irwin, K. S. Karkare, J. P. Kaufman, B. G. Keating, S. A. Kernasovskiy, J. M. Kovac, C. L. Kuo, E. M. Leitch, M. Lueker, P. Mason, C. B. Netterfield, H. T. Nguyen, R. O’Brien, R. W. Ogburn, A. Orlando, C. Pryke, C. D. Reintsema, S. Richter, R. Schwarz, C. D. Sheehy, Z. K. Staniszewski, R. V. Sudiwala, G. P. Teply, J. E. Tolan, A. D. Turner, A. G. Vieregg, C. L. Wong, K. W. Yoon, and Bicep2 Collaboration. Detection of B-Mode Polarization at Degree Angular Scales by BICEP2. *Physical Review Letters*, 112(24):241101, June 2014.

- [22] BICEP2/Keck, Planck Collaborations, :, P. A. R. Ade, N. Aghanim, Z. Ahmed, R. W. Aikin, K. D. Alexander, M. Arnaud, J. Aumont, and et al. A Joint Analysis of BICEP2/Keck Array and Planck Data. *ArXiv e-prints*, February 2015.
- [23] Planck Collaboration, R. Adam, P. A. R. Ade, N. Aghanim, M. Arnaud, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al. Planck intermediate results. XXX. The angular power spectrum of polarized dust emission at intermediate and high Galactic latitudes. *ArXiv e-prints*, September 2014.
- [24] R. A. Sunyaev and Y. B. Zeldovich. The Spectrum of Primordial Radiation, its Distortions and their Significance. *Comments on Astrophysics and Space Physics*, 2:66, March 1970.
- [25] C. Burigana. Distortions of the CMB Spectrum by Continuous Heating. In G. L. Chincarini, A. Iovino, T. Maccacaro, & D. Maccagni, editor, *Observational Cosmology*, volume 51 of *Astronomical Society of the Pacific Conference Series*, pages 554–+, January 1993.
- [26] W. Hu and J. Silk. Thermalization and spectral distortions of the cosmic background radiation. *Phys. Rev. D.*, 48:485–502, July 1993.
- [27] J. Chluba and R. A. Sunyaev. The evolution of CMB spectral distortions in the early Universe. *MNRAS*, 419:1294–1314, January 2012.
- [28] J. C. Mather, E. S. Cheng, D. A. Cottingham, R. E. Eplee, Jr., D. J. Fixsen, T. Hewagama, R. B. Isaacman, K. A. Jensen, S. S. Meyer, P. D. Noerdlinger, S. M. Read, and L. P. Rosen. Measurement of the cosmic microwave background spectrum by the COBE FIRAS instrument. *Ap. J.*, 420:439–444, January 1994.
- [29] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright. The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set. *Ap. J.*, 473:576–+, December 1996.
- [30] J. Chluba. Which spectral distortions does Λ CDM actually predict? *MNRAS*, 460:227–239, July 2016.
- [31] A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M. Meyer, S. H. Moseley, M. D. Seiffert, D. N. Spergel, and E. J. Wollack. The Primordial Inflation Explorer (PIXIE): a nulling polarimeter for cosmic microwave background observations. *JCAP*, 7:25–+, July 2011.
- [32] P. André, C. Baccigalupi, A. Banday, D. Barbosa, B. Barreiro, J. Bartlett, N. Bartolo, E. Battistelli, R. Battye, G. Bendo, A. Beno#523t, J.-P. Bernard, M. Bersanelli, M. Béthermin, P. Bielewicz, A. Bonaldi, F. Bouchet, F. Boulanger, J. Brand, M. Bucher, C. Burigana, Z.-Y. Cai, P. Camus, F. Casas, V. Casasola, G. Castex, A. Challinor, J. Chluba, G. Chon, S. Colafrancesco, B. Comis, F. Cuttaia, G. D’Alessandro, A. Da Silva, R. Davis, M. de Avillez, P. de Bernardis, M. de Petris, A. de Rosa, G. de Zotti, J. Delabrouille, F.-X. Désert, C. Dickinson, J. M. Diego, J. Dunkley, T. Enßlin, J. Errard, E. Falgarone, P. Ferreira, K. Ferrière, F. Finelli, A. Fletcher, P. Fosalba, G. Fuller, S. Galli, K. Ganga, J. García-Bellido, A. Ghribi, M. Giard, Y. Giraud-Héraud, J. Gonzalez-Nuevo, K. Grainge, A. Gruppuso, A. Hall, J.-C. Hamilton, M. Haverkorn, C. Hernandez-Monteagudo, D. Herranz, M. Jackson, A. Jaffe,

- R. Khatri, M. Kunz, L. Lamagna, M. Lattanzi, P. Leahy, J. Lesgourgues, M. Liguori, E. Liguori, M. Lopez-Caniego, J. Macias-Perez, B. Maffei, D. Maino, A. Mangilli, E. Martinez-Gonzalez, C. Martins, S. Masi, M. Massardi, S. Matarrese, A. Melchiorri, J.-B. Melin, A. Mennella, A. Mignano, M.-A. Miville-Deschênes, A. Monfardini, A. Murphy, P. Naselsky, F. Nati, P. Natoli, M. Negrello, F. Noviello, C. O’Sullivan, F. Paci, L. Pagano, R. Paladino, N. Palanque-Delabrouille, D. Paoletti, H. Peiris, F. Perrotta, F. Piacentini, M. Piat, L. Piccirillo, G. Pisano, G. Polenta, A. Pollo, N. Ponthieu, M. Remazeilles, S. Ricciardi, M. Roman, C. Rosset, J.-A. Rubino-Martin, M. Salatino, A. Schillaci, P. Shellard, J. Silk, A. Starobinsky, R. Stompor, R. Sunyaev, A. Tartari, L. Terenzi, L. Toffolatti, M. Tomasi, N. Trappe, M. Tristram, T. Trombetti, M. Tucci, R. Van de Weijgaert, B. Van Tent, L. Verde, P. Vielva, B. Wandelt, R. Watson, and S. Withington. PRISM (Polarized Radiation Imaging and Spectroscopy Mission): an extended white paper. *JCAP*, 2:6, February 2014.
- [33] Y. B. Zeldovich and R. A. Sunyaev. The Interaction of Matter and Radiation in a Hot-Model Universe. *ApSS*, 4:301–316, July 1969.
- [34] R. A. Sunyaev and Y. B. Zeldovich. The interaction of matter and radiation in the hot model of the Universe, II. *ApSS*, 7:20–30, April 1970.
- [35] R. A. Sunyaev and Y. B. Zeldovich. Formation of Clusters of Galaxies; Protocluster Fragmentation and Intergalactic Gas Heating. *Astron. Astrophys.*, 20:189–+, August 1972.
- [36] W. Hu, D. Scott, and J. Silk. Reionization and cosmic microwave background distortions: A complete treatment of second-order Compton scattering. *Phys. Rev. D.*, 49:648–670, January 1994.
- [37] S. P. Oh, A. Cooray, and M. Kamionkowski. Sunyaev-Zeldovich fluctuations from the first stars? *MNRAS*, 342:L20–L24, June 2003.
- [38] R. Cen and J. P. Ostriker. Where Are the Baryons? *Ap. J.*, 514:1–6, March 1999.
- [39] A. Refregier, E. Komatsu, D. N. Spergel, and U.-L. Pen. Power spectrum of the Sunyaev-Zel’dovich effect. *Phys. Rev. D.*, 61(12):123001, June 2000.
- [40] J. C. Hill, N. Battaglia, J. Chluba, S. Ferraro, E. Schaan, and D. N. Spergel. Taking the Universe’s Temperature with Spectral Distortions of the Cosmic Microwave Background. *Physical Review Letters*, 115(26):261301, December 2015.
- [41] W. Hu and J. Silk. Thermalization constraints and spectral distortions for massive unstable relic particles. *Physical Review Letters*, 70:2661–2664, May 1993.
- [42] J. Chluba. Distinguishing different scenarios of early energy release with spectral distortions of the cosmic microwave background. *MNRAS*, 436:2232–2243, December 2013.
- [43] J. Chluba and D. Jeong. Teasing bits of information out of the CMB energy spectrum. *MNRAS*, 438:2065–2082, March 2014.
- [44] E. Dimastrogiovanni, L. M. Krauss, and J. Chluba. Constraints on gravitino decay and the scale of inflation using CMB spectral distortions. *Phys. Rev. D.*, 94(2):023518, July 2016.

- [45] K. Jedamzik, V. Katalinić, and A. V. Olinto. Limit on Primordial Small-Scale Magnetic Fields from Cosmic Microwave Background Distortions. *PRL*, 85:700–703, July 2000.
- [46] H. Tashiro, E. Sabancilar, and T. Vachaspati. CMB distortions from superconducting cosmic strings. *Phys. Rev. D.*, 85(10):103522, May 2012.
- [47] A. D. Dolgov and D. Ejlli. Resonant high energy graviton to photon conversion at the post-recombination epoch. *Phys. Rev. D.*, 87(10):104007, May 2013.
- [48] H. Tashiro, J. Silk, and D. J. E. Marsh. Constraints on primordial magnetic fields from CMB distortions in the axiverse. *Phys. Rev. D.*, 88(12):125024, December 2013.
- [49] R. R. Caldwell and N. A. Maksimova. Spectral distortion in a radially inhomogeneous cosmology. *Phys. Rev. D.*, 88(10):103502, November 2013.

2 Curriculum Vitae

3 Summary of Work Effort

4 Current and Pending Support

5 Letters of Support

6 Budget Details - Narrative

6.1 Team, and Work Effort

6.1.1 Funded Team Members

6.1.2 Non-Funded Team Members

6.2 Costing Principles

- **Summer Salaries:**
 - **Workshop:**

6.3 University of Minnesota Budget

6.3.1 Direct Labor

6.3.2 Supplies

6.3.3 Travel

6.3.4 Other Direct Costs

Publications and Teleconferencing

Other Subcontracts

6.3.5 Facilities and Administrative Costs

7 Budget Sheets