Contents

1   Scientific, Technical, and Management Section

0.5 pg. executive summary goes here: the sky is sunny

1.1   Science Objectives

5 pages for all science goals including the (temporary) two sections below.

1.1.1   The Inflationary Gravitional Wave Background

The verbiage below is taken from another proposal. Here we need to explain what are the science objectives of the CMBProbe, how the science objectives relate to the current state of knowledge, and to NASA's goals

The paradigm of inflation [1, 2, 3, 4, 5] makes several predictions that are consistent with all current astrophysical measurements [6, 7, 8, 9]. A robust prediction of inflation is the existence of a stochastic background of gravitational radiation with an amplitude depending on the mechanism driving the accelerated expansion [10, 11, 12, 13, 14]. In most scenarios, this 'inflationary gravitional wave background' (IGB) is predicted to have a spatial power spectrum whose amplitude is proportional to the energy scale of inflation $V^{1/4}$ via $V^{1/4} = 3.7 \times 10^{16} \, r^{1/4}$ GeV, where V is the inflaton potential and r is the ratio of the temperature quadrupoles produced by gravitational waves and by density perturbations. There are theoretical reasons $V^{1/4}$ may be close to the Grand Unification scale of 10^{16} GeV, suggesting detectable r values between ~ 0.001 and ~ 0.1. In addition to determining the energy scale of inflation, measurements of the IGB probe the scalar field potential at or above the Planck scale, which is particularly relevant for inflation models motivated by string theory [15]. Measurements of the IGB thus probe fundamental physics at the highest possible energy scales.

Figure 1:   Sample Figure of Sunny Skies

The most promising way to search for the IGB is through its signature on the CMB polarization [16, 17]. Primordial energy density perturbations produce only a curl-free, or 'E-mode', pattern of polarization. Gravitational waves also produce a curl, or 'B-mode', pattern of polarization that density perturbations cannot produce [18, 19]. The amplitude of the B mode is related to the energy scale of inflation by $V^{1/4} = 2 \times 10^{16} \, (B_{peak}/0.1 \, \mu\text{K})^{1/2}$ GeV, where B_{peak} is the amplitude of the power spectrum of the B mode in μK at $\ell = 80$; see Fig. 1. In its recent report New Worlds New Horizons (NWNH), the decadal survey committee strongly endorsed sub-orbital searches for the B-mode signal from inflation saying that "The convincing detection of B-mode polarization in the CMB produced in the epoch of reionization would represent a watershed discovery." [20]

B-mode signatures near the expected IGB peak at $\ell = 80$ have recently been detected by BI-CEP2 [21]. However, the combination of Planck data with those from the BICEP2 and Keck Array collaborations have demonstrated that the B-mode signal measured is entirely consistent with contributions from polarized emission of Galactic dust and the signal from the gravitational lensing of CMB photons by the large scale structure of the Universe (see Section **??**) [22, 23**?**]. These data

Figure 2: [Placeholders] *Left:* Neutrino mass constraints as a function of the prior on τ. *Right* N_{eff} Forecasts as a function of resolution in arc-min and temperature noise in μK-arcmin assuming $f_{\mathrm{sky}} = 0.7$.

give an upper limit of $r < 0.09$ at 95% confidence level. Most importantly, the constraint is largely limited by Planck's noisy measurement of the dust properties in the 353 GHz band; a noiseless dust map could shrink the constraint by a factor of two [22]. Further progress — detections or improved limits — requires instruments with higher sensitivity at *both* the dust and CMB frequency bands so that this Galactic foreground can be properly identified and removed.

1.1.2 Neutrinos and Light Relics

Much of the information about our thermal history and the particle content of the universe is encoded in the T and E power spectra. A high-precision measurement of these spectra over the full sky is expected to significantly improve our understanding of the post-inflationary universe. This is particularly true in E-mode polarization where far fewer modes have be measured at the level of cosmic variance.

1.1.3 CMB spectral distortion science

In addition to the CMB temperature and polarization anisotropies targeted by CMB imagers, *unique* new information about early-universe physics can be gained by studying the energy spectrum of the CMB [24, 25, 26, 27]. The measurements of COBE/FIRAS have shown that the average CMB spectrum is extremely close to that of a blackbody at a temperature $T_0 = (2.726 \pm 0.001)\,\mathrm{K}$ [28, 29]. However, several standard processes are expected to *distort* the CMB spectrum [e.g., 30] at a level that is within reach of present-day technology [31, 32]. The classical distortion shapes are known as Compton-y and chemical potential (μ-type) distortions [33, 34] and are caused by energy exchange of CMB photons with free electrons. A μ-distortion can only be produced in a hot and dense environment present at redshifts $z \gtrsim 5 \times 10^4$, while y-type distortions appear at lower redshifts. This makes μ-distortions a unique messenger from the early Universe.

The largest guaranteed distortion is caused by the late-time energy release of forming structures and from reionization [35, 36, 37, 38, 39], imprinting a y-type distortion with $y \simeq 2 \times 10^{-6}$ [e.g., 39, 40]. This distortion is only one order of magnitude below the current limit from COBE/FIRAS and, even with most pessimistic assumptions about foregrounds, should be clearly seen with next-generation spectrometers, telling us about the total energy output of first stars, AGN and galaxy

clusters. In particular, group-size clusters ($M \simeq 10^{13}\,M_\odot$) contribute significantly to the signal. These are still sufficiently hot (temperature $kT_e \simeq 1\,\mathrm{keV}$) to create a visible relativistic temperature correction to this large y-distortion, which could be used to constrain cluster feedback models [40]. These two inevitable signals probe the low-redshift Universe and provide clear targets for future spectral distortions measurements and their requirements in the presence of foregrounds.

Next generation CMB spectrometers are also expected to greatly improve the μ-distortion limits of COBE/FIRAS [31]. This will allow us to place stringent bounds on the presence of long-lived decaying particles [41, 42, 43, 44] and other new physics [e.g., 45, 46, 47, 48, 49, 50], but a clear target is predicted by the dissipation of small-scale perturbation through Silk-damping [51, 52, 53, 54]. This process allows us to place stringent constraints on the amplitude of the small-scale curvature power spectrum, present at scales (wavelength $0.1\,\mathrm{kpc} \lesssim \lambda \lesssim 1\,\mathrm{Mpc}$) and epochs ($10^4 \lesssim z \lesssim 10^6$) inaccessible through any other observation. This delivers a complementary test for the inflation paradigm [43, 55, 56, 57, 58], with $\mu = (2.0 \pm 0.14) \times 10^{-8}$ expected in ΛCDM [30]. Precise measurements of signals at this level will be extremely challenging and requires unprecedented control of systematics and modeling of foregrounds. It would also bring us to the sensitivity level required to detect the cosmological recombination radiation [59, 60] imprinted by the recombination of hydrogen and helium at redshift $z \simeq 10^3 - 10^4$. Optimizing next-generation CMB spectrometers for these purposes requires extensive studies.

Aside from the average CMB distortion signals, the CMB spectrum can also vary across the sky. One source of anisotropic distortions is related to clusters of galaxies and has already been measured [61]. A combination of precise CMB imaging and spectroscopic measurements might allow observing the relativistic temperature correction [62, 63, 64] of individual SZ clusters. This could allow us to calibrate cluster scaling relations and learn about the dynamical state of the cluster atmosphere. Anisotropies in the μ-distortion can be created through ultra-squeezed limit non-Gaussianity [65, 66] and could be used to probe scale-dependent non-Gaussianity [67, 68]. Finally, resonant scattering signals in the recombination [69, 70, 71] and post-recombination eras [72, 73] can lead to spectral-spatial CMB signals that can be used to constrain the presence of metals in the dark ages and the physics of recombination. For all these applications, instrumental synergies between CMB imaging and spectroscopy need to be studied in detail.

In summary, future studies of the CMB spectrum will open a new *unexplored* window to early phases of the Universe ($z > 10^3$), which cannot be probed in any other way. This will not only allow us to test the standard cosmological paradigm (e.g., inflation and reionization) but also opens up a huge discovery space to non-standard physics (e.g., decaying/annihilating particles). This immense potential and complementarity with CMB anisotropy studies makes CMB spectral distortions an important future target and identifying experimental routes towards extracting these tiny signals from the early Universe will be one main objective of the proposed mission study.

1.2 The Challenges: Foregrounds, systematics

3 pages. Discuss the challenge of Foregrounds and Systematics.

1.3 Current and Forthcoming Efforts and the CMB Probe

2.5 pages. S3 experiments, forthcoming S4, Baselines CMB Probe options and their complementarity with S4.

1.4 State of Technologies

2 pages. Discuss the technologies, their TRL, and what will be studied

1.5 Mission Study, and Management Plan

1.5 pages; Describe what we want to do, who is doing what, what we are funding

References

[1] A. H. Guth. Inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D.*, 23:347–356, January 1981.

[2] A. D. Linde. A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems. *Phys. Lett.*, B108:389–393, 1982.

[3] A. Albrecht and P. J. Steinhardt. Cosmology for grand unified theories with radiatively induced symmetry breaking. *Phys. Rev. Lett.*, 48:1220–1223, 1982.

[4] K. Sato. First-order phase transition of a vacuum and the expansion of the Universe. *MNRAS*, 195:467–479, May 1981.

[5] E. W. Kolb and M. S. Turner. *The Early Universe*. Addison-Wesley, Redwood City, CA, 2nd. edition, 1994.

[6] D. N. Spergel, R. Bean, O. Doré, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page, H. V. Peiris, L. Verde, M. Halpern, R. S. Hill, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright. Three-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Implications for Cosmology, June 2007.

[7] M. Tegmark, D. Eisenstein, M. Strauss, D. Weinberg, M. Blanton, J. Frieman, M. Fukugita, J. Gunn, A. Hamilton, G. Knapp, R. Nichol, J. Ostriker, N. Padmanabhan, W. Percival, D. Schlegel, D. Schneider, R. Scoccimarro, U. Seljak, H. Seo, M. Swanson, A. Szalay, M. Vogeley, J. Yoo, I. Zehavi, K. Abazajian, S. Anderson, J. Annis, N. Bahcall, B. Bassett, A. Berlind, J. Brinkmann, T. Budavari, F. Castander, A. Connolly, I. Csabai, M. Doi, D. Finkbeiner, B. Gillespie, K. Glazebrook, G. Hennessy, D. Hogg, Z. Ivezic, B. Jain, D. Johnston, S. Kent, D. Lamb, B. Lee, H. Lin, J. Loveday, R. Lupton, J. Munn, K. Pan, C. Park, J. Peoples, J. Pier, A. Pope, M. Richmond, C. Rockosi, R. Scranton, R. Sheth, A. Stebbins, C. Stoughton, I. Szapudi, D. Tucker, D. Vanden Berk, B. Yanny, and D. York. Cosmological Constraints from the SDSS Luminous Red Galaxies. *Phys. Rev. D.*, 74:123507, 2006.

[8] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al. Planck 2015 results. XIII. Cosmological parameters. *ArXiv e-prints*, February 2015.

[9] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, F. Arroja, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, and et al. Planck 2015. XX. Constraints on inflation. *ArXiv e-prints*, February 2015.

[10] A. A. Starobinsky. Dynamic of Phase Transition in the New Inflationary Universe Scenario and Generation of Perturbationss. *Phys. Lett.*, B117:175–178, 1982.

[11] A. A. Starobinskii. The Perturbation Spectrum Evolving from a Nonsingular Initially De-Sitter Cosmology and the Microwave Background Anisotropy. *Soviet Astronomy Letters*, 9:302–+, June 1983.

[12] V. A. Rubakov, M. V. Sazhin, and A. V. Veryaskin. Graviton creation in the inflationary universe and the grand unification scale. *Phys. Lett. B.*, 115:189–192, September 1982.

[13] L. P. Grishchuk. Amplification of gravitational waves in an istropic universe. *Sov. Phys. JETP*, 40:409–415, 1975.

[14] L. F. Abbott and M. B. Wise. Constraints on generalized inflationary cosmologies. *Nuclear Physics B*, 244:541–548, October 1984.

[15] K. N. Abazajian, K. Arnold, J. Austermann, B. A. Benson, C. Bischoff, J. Bock, J. R. Bond, J. Borrill, I. Buder, D. L. Burke, E. Calabrese, J. E. Carlstrom, C. S. Carvalho, C. L. Chang, H. C. Chiang, S. Church, A. Cooray, T. M. Crawford, B. P. Crill, K. S. Dawson, S. Das, M. J. Devlin, M. Dobbs, S. Dodelson, O. Doré, J. Dunkley, J. L. Feng, A. Fraisse, J. Gallicchio, S. B. Giddings, D. Green, N. W. Halverson, S. Hanany, D. Hanson, S. R. Hildebrandt, A. Hincks, R. Hlozek, G. Holder, W. L. Holzapfel, K. Honscheid, G. Horowitz, W. Hu, J. Hubmayr, K. Irwin, M. Jackson, W. C. Jones, R. Kallosh, M. Kamionkowski, B. Keating, R. Keisler, W. Kinney, L. Knox, E. Komatsu, J. Kovac, C.-L. Kuo, A. Kusaka, C. Lawrence, A. T. Lee, E. Leitch, A. Linde, E. Linder, P. Lubin, J. Maldacena, E. Martinec, J. McMahon, A. Miller, V. Mukhanov, L. Newburgh, M. D. Niemack, H. Nguyen, H. T. Nguyen, L. Page, C. Pryke, C. L. Reichardt, J. E. Ruhl, N. Sehgal, U. Seljak, L. Senatore, J. Sievers, E. Silverstein, A. Slosar, K. M. Smith, D. Spergel, S. T. Staggs, A. Stark, R. Stompor, A. G. Vieregg, G. Wang, S. Watson, E. J. Wollack, W. L. K. Wu, K. W. Yoon, O. Zahn, and M. Zaldarriaga. Inflation Physics from the Cosmic Microwave Background and Large Scale Structure. *ArXiv e-prints*, September 2013.

[16] M. Kamionkowski, A. Kosowsky, and A. Stebbins. Statistics of Cosmic Microwave Background Polarization. *Phys. Rev. D.*, 55:7368–7388, June 1997.

[17] U. Seljak and M. Zaldarriaga. Signature of Gravity Waves in the Polarization of the Microwave Background. *Phys. Rev. Lett.*, 78:2054–2057, March 1997. astro-ph/9609169.

[18] M. Kamionkowski, A. Kosowsky, and A. Stebbins. A Probe of Primordial Gravity Waves and Vorticity. *Phys. Rev. Lett.*, 78:2058–2061, March 1997. astro-ph/9609132.

[19] M. Zaldarriaga and U. Seljak. All-sky analysis of polarization in the microwave background. *Phys. Rev. D.*, 55:1830–1840, 1997.

[20] Committee for a Decadal Survey of Astronomy and Astrophysics. *New Worlds, New Horizons in Astronomy and Astrophysics*. National Academy Press, 2010.

[21] P. A. R. Ade, R. W. Aikin, D. Barkats, S. J. Benton, C. A. Bischoff, J. J. Bock, J. A. Brevik, I. Buder, E. Bullock, C. D. Dowell, L. Duband, J. P. Filippini, S. Fliescher, S. R. Golwala, M. Halpern, M. Hasselfield, S. R. Hildebrandt, G. C. Hilton, V. V. Hristov, K. D. Irwin, K. S. Karkare, J. P. Kaufman, B. G. Keating, S. A. Kernasovskiy, J. M. Kovac, C. L. Kuo, E. M. Leitch, M. Lueker, P. Mason, C. B. Netterfield, H. T. Nguyen, R. O'Brient, R. W. Ogburn, A. Orlando, C. Pryke, C. D. Reintsema, S. Richter, R. Schwarz, C. D. Sheehy, Z. K. Staniszewski, R. V. Sudiwala, G. P. Teply, J. E. Tolan, A. D. Turner, A. G. Vieregg, C. L. Wong, K. W. Yoon, and Bicep2 Collaboration. Detection of B-Mode Polarization at Degree Angular Scales by BICEP2. *Physical Review Letters*, 112(24):241101, June 2014.

[22] BICEP2/Keck, Planck Collaborations, :, P. A. R. Ade, N. Aghanim, Z. Ahmed, R. W. Aikin, K. D. Alexander, M. Arnaud, J. Aumont, and et al. A Joint Analysis of BICEP2/Keck Array and Planck Data. *ArXiv e-prints*, February 2015.

[23] Planck Collaboration, R. Adam, P. A. R. Ade, N. Aghanim, M. Arnaud, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al. Planck intermediate results. XXX. The angular power spectrum of polarized dust emission at intermediate and high Galactic latitudes. *ArXiv e-prints*, September 2014.

[24] R. A. Sunyaev and Y. B. Zeldovich. The Spectrum of Primordial Radiation, its Distortions and their Significance. *Comments on Astrophysics and Space Physics*, 2:66, March 1970.

[25] C. Burigana. Distortions of the CMB Spectrum by Continuous Heating. In G. L. Chincarini, A. Iovino, T. Maccacaro, & D. Maccagni, editor, *Observational Cosmology*, volume 51 of *Astronomical Society of the Pacific Conference Series*, pages 554–+, January 1993.

[26] W. Hu and J. Silk. Thermalization and spectral distortions of the cosmic background radiation. *Phys. Rev. D.*, 48:485–502, July 1993.

[27] J. Chluba and R. A. Sunyaev. The evolution of CMB spectral distortions in the early Universe. *MNRAS*, 419:1294–1314, January 2012.

[28] J. C. Mather, E. S. Cheng, D. A. Cottingham, R. E. Eplee, Jr., D. J. Fixsen, T. Hewagama, R. B. Isaacman, K. A. Jensen, S. S. Meyer, P. D. Noerdlinger, S. M. Read, and L. P. Rosen. Measurement of the cosmic microwave background spectrum by the COBE FIRAS instrument. *Ap. J.*, 420:439–444, January 1994.

[29] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright. The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set. *Ap. J.*, 473:576–+, December 1996.

[30] J. Chluba. Which spectral distortions does ΛCDM actually predict? *MNRAS*, 460:227–239, July 2016.

[31] A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M. Meyer, S. H. Moseley, M. D. Seiffert, D. N. Spergel, and E. J. Wollack. The Primordial Inflation Explorer (PIXIE): a nulling polarimeter for cosmic microwave background observations. *JCAP*, 7:25–+, July 2011.

[32] P. André, C. Baccigalupi, A. Banday, D. Barbosa, B. Barreiro, J. Bartlett, N. Bartolo, E. Battistelli, R. Battye, G. Bendo, A. Beno#523t, J.-P. Bernard, M. Bersanelli, M. Béthermin, P. Bielewicz, A. Bonaldi, F. Bouchet, F. Boulanger, J. Brand, M. Bucher, C. Burigana, Z.-Y. Cai, P. Camus, F. Casas, V. Casasola, G. Castex, A. Challinor, J. Chluba, G. Chon, S. Colafrancesco, B. Comis, F. Cuttaia, G. D'Alessandro, A. Da Silva, R. Davis, M. de Avillez, P. de Bernardis, M. de Petris, A. de Rosa, G. de Zotti, J. Delabrouille, F.-X. Désert, C. Dickinson, J. M. Diego, J. Dunkley, T. Enßlin, J. Errard, E. Falgarone, P. Ferreira, K. Ferrière, F. Finelli, A. Fletcher, P. Fosalba, G. Fuller, S. Galli, K. Ganga, J. García-Bellido, A. Ghribi, M. Giard, Y. Giraud-Héraud, J. Gonzalez-Nuevo, K. Grainge, A. Gruppuso, A. Hall, J.-C. Hamilton, M. Haverkorn, C. Hernandez-Monteagudo, D. Herranz, M. Jackson, A. Jaffe,

R. Khatri, M. Kunz, L. Lamagna, M. Lattanzi, P. Leahy, J. Lesgourgues, M. Liguori, E. Liuzzo, M. Lopez-Caniego, J. Macias-Perez, B. Maffei, D. Maino, A. Mangilli, E. Martinez-Gonzalez, C. Martins, S. Masi, M. Massardi, S. Matarrese, A. Melchiorri, J.-B. Melin, A. Mennella, A. Mignano, M.-A. Miville-Deschênes, A. Monfardini, A. Murphy, P. Naselsky, F. Nati, P. Natoli, M. Negrello, F. Noviello, C. O'Sullivan, F. Paci, L. Pagano, R. Paladino, N. Palanque-Delabrouille, D. Paoletti, H. Peiris, F. Perrotta, F. Piacentini, M. Piat, L. Piccirillo, G. Pisano, G. Polenta, A. Pollo, N. Ponthieu, M. Remazeilles, S. Ricciardi, M. Roman, C. Rosset, J.-A. Rubino-Martin, M. Salatino, A. Schillaci, P. Shellard, J. Silk, A. Starobinsky, R. Stompor, R. Sunyaev, A. Tartari, L. Terenzi, L. Toffolatti, M. Tomasi, N. Trappe, M. Tristram, T. Trombetti, M. Tucci, R. Van de Weijgaert, B. Van Tent, L. Verde, P. Vielva, B. Wandelt, R. Watson, and S. Withington. PRISM (Polarized Radiation Imaging and Spectroscopy Mission): an extended white paper. *JCAP*, 2:6, February 2014.

[33] Y. B. Zeldovich and R. A. Sunyaev. The Interaction of Matter and Radiation in a Hot-Model Universe. *ApSS*, 4:301–316, July 1969.

[34] R. A. Sunyaev and Y. B. Zeldovich. The interaction of matter and radiation in the hot model of the Universe, II. *ApSS*, 7:20–30, April 1970.

[35] R. A. Sunyaev and Y. B. Zeldovich. Formation of Clusters of Galaxies; Protocluster Fragmentation and Intergalactic Gas Heating. *Astron. Astrophys.*, 20:189–+, August 1972.

[36] W. Hu, D. Scott, and J. Silk. Reionization and cosmic microwave background distortions: A complete treatment of second-order Compton scattering. *Phys. Rev. D.*, 49:648–670, January 1994.

[37] S. P. Oh, A. Cooray, and M. Kamionkowski. Sunyaev-Zeldovich fluctuations from the first stars? *MNRAS*, 342:L20–L24, June 2003.

[38] R. Cen and J. P. Ostriker. Where Are the Baryons? *Ap. J.*, 514:1–6, March 1999.

[39] A. Refregier, E. Komatsu, D. N. Spergel, and U.-L. Pen. Power spectrum of the Sunyaev-Zel'dovich effect. *Phys. Rev. D.*, 61(12):123001, June 2000.

[40] J. C. Hill, N. Battaglia, J. Chluba, S. Ferraro, E. Schaan, and D. N. Spergel. Taking the Universe's Temperature with Spectral Distortions of the Cosmic Microwave Background. *Physical Review Letters*, 115(26):261301, December 2015.

[41] W. Hu and J. Silk. Thermalization constraints and spectral distortions for massive unstable relic particles. *Physical Review Letters*, 70:2661–2664, May 1993.

[42] J. Chluba. Distinguishing different scenarios of early energy release with spectral distortions of the cosmic microwave background. *MNRAS*, 436:2232–2243, December 2013.

[43] J. Chluba and D. Jeong. Teasing bits of information out of the CMB energy spectrum. *MNRAS*, 438:2065–2082, March 2014.

[44] E. Dimastrogiovanni, L. M. Krauss, and J. Chluba. Constraints on gravitino decay and the scale of inflation using CMB spectral distortions. *Phys. Rev. D.*, 94(2):023518, July 2016.

[45] K. Jedamzik, V. Katalinić, and A. V. Olinto. Limit on Primordial Small-Scale Magnetic Fields from Cosmic Microwave Background Distortions. *PRL*, 85:700–703, July 2000.

[46] H. Tashiro, E. Sabancilar, and T. Vachaspati. CMB distortions from superconducting cosmic strings. *Phys. Rev. D.*, 85(10):103522, May 2012.

[47] A. D. Dolgov and D. Ejlli. Resonant high energy graviton to photon conversion at the post-recombination epoch. *Phys. Rev. D.*, 87(10):104007, May 2013.

[48] H. Tashiro, J. Silk, and D. J. E. Marsh. Constraints on primordial magnetic fields from CMB distortions in the axiverse. *Phys. Rev. D.*, 88(12):125024, December 2013.

[49] R. R. Caldwell and N. A. Maksimova. Spectral distortion in a radially inhomogeneous cosmology. *Phys. Rev. D.*, 88(10):103502, November 2013.

[50] Y. Ali-Haïmoud, J. Chluba, and M. Kamionkowski. Constraints on Dark Matter Interactions with Standard Model Particles from Cosmic Microwave Background Spectral Distortions. *Physical Review Letters*, 115(7):071304, August 2015.

[51] R. A. Sunyaev and Y. B. Zeldovich. Small scale entropy and adiabatic density perturbations - Antimatter in the Universe. *ApSS*, 9:368–382, December 1970.

[52] R. A. Daly. Spectral distortions of the microwave background radiation resulting from the damping of pressure waves. *Ap. J.*, 371:14–28, April 1991.

[53] W. Hu, D. Scott, and J. Silk. Power spectrum constraints from spectral distortions in the cosmic microwave background. *Ap. J. Lett.*, 430:L5–L8, July 1994.

[54] J. Chluba, R. Khatri, and R. A. Sunyaev. CMB at 2 x 2 order: the dissipation of primordial acoustic waves and the observable part of the associated energy release. *MNRAS*, 425:1129–1169, September 2012.

[55] J. Chluba, A. L. Erickcek, and I. Ben-Dayan. Probing the Inflaton: Small-scale Power Spectrum Constraints from Measurements of the Cosmic Microwave Background Energy Spectrum. *Ap. J.*, 758:76, October 2012.

[56] J. B. Dent, D. A. Easson, and H. Tashiro. Cosmological constraints from CMB distortion. *Phys. Rev. D.*, 86(2):023514, July 2012.

[57] S. Clesse, B. Garbrecht, and Y. Zhu. Testing inflation and curvaton scenarios with CMB distortions. *JCAP*, 10:046, October 2014.

[58] G. Cabass, A. Melchiorri, and E. Pajer. μ distortions or running: A guaranteed discovery from CMB spectrometry. *Phys. Rev. D.*, 93(8):083515, April 2016.

[59] R. A. Sunyaev and J. Chluba. Signals from the epoch of cosmological recombination (Karl Schwarzschild Award Lecture 2008). *Astronomische Nachrichten*, 330:657–+, 2009.

[60] J. Chluba and Y. Ali-Haïmoud. COSMOSPEC: fast and detailed computation of the cosmological recombination radiation from hydrogen and helium. *MNRAS*, 456:3494–3508, March 2016.

[61] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al. Planck 2013 results. XX. Cosmology from Sunyaev-Zeldovich cluster counts. *Astron. Astrophys.*, 571:A20, November 2014.

[62] S. Y. Sazonov and R. A. Sunyaev. Cosmic Microwave Background Radiation in the Direction of a Moving Cluster of Galaxies with Hot Gas: Relativistic Corrections. *Ap. J.*, 508:1–5, November 1998.

[63] N. Itoh, Y. Kohyama, and S. Nozawa. Relativistic corrections to the sunyaev-zeldovich effect for clusters of galaxies. *Ap. J.*, 502:7, July 1998.

[64] A. Challinor and A. Lasenby. Relativistic corrections to the sunyaev-zeldovich effect. *Ap. J.*, 499:1, May 1998.

[65] E. Pajer and M. Zaldarriaga. New Window on Primordial Non-Gaussianity. *Physical Review Letters*, 109(2):021302, July 2012.

[66] J. Ganc and E. Komatsu. Scale-dependent bias of galaxies and μ-type distortion of the cosmic microwave background spectrum from single-field inflation with a modified initial state. *Phys. Rev. D.*, 86(2):023518, July 2012.

[67] M. Biagetti, H. Perrier, A. Riotto, and V. Desjacques. Testing the running of non-Gaussianity through the CMB μ-distortion and the halo bias. *Phys. Rev. D.*, 87(6):063521, March 2013.

[68] R. Emami, E. Dimastrogiovanni, J. Chluba, and M. Kamionkowski. Probing the scale dependence of non-Gaussianity with spectral distortions of the cosmic microwave background. *Phys. Rev. D.*, 91(12):123531, June 2015.

[69] J. A. Rubiño-Martín, C. Hernández-Monteagudo, and R. A. Sunyaev. The imprint of cosmological hydrogen recombination lines on the power spectrum of the CMB. *Astron. Astrophys.*, 438:461–473, August 2005.

[70] C. Hernández-Monteagudo, J. A. Rubiño-Martín, and R. A. Sunyaev. On the influence of resonant scattering on cosmic microwave background polarization anisotropies. *MNRAS*, 380:1656–1668, October 2007.

[71] A. Lewis. Rayleigh scattering: blue sky thinking for future CMB observations. *JCAP*, 8:053, August 2013.

[72] K. Basu, C. Hernández-Monteagudo, and R. A. Sunyaev. CMB observations and the production of chemical elements at the end of the dark ages. *Astron. Astrophys.*, 416:447–466, March 2004.

[73] D. R. G. Schleicher, D. Galli, F. Palla, M. Camenzind, R. S. Klessen, M. Bartelmann, and S. C. O. Glover. Effects of primordial chemistry on the cosmic microwave background. *Astron. Astrophys.*, 490:521–535, November 2008.

2 Curriculum Vitae

3   Summary of Work Effort

4   Current and Pending Support

5 Letters of Support

6 Budget Details - Narrative

6.1 Team, and Work Effort

6.1.1 Funded Team Members

6.1.2 Non-Funded Team Members

6.2 Costing Principles

- **Summer Salaries:**
  - **Workshop:**

6.3 University of Minnesota Budget

6.3.1 Direct Labor

6.3.2 Supplies

6.3.3 Travel

6.3.4 Other Direct Costs

Publications and Teleconferencing
   Other Subcontracts

6.3.5 Facilities and Administrative Costs

7 Budget Sheets