

Characteristics of the Mississippi Sound Estuary and Watershed

Mississippi Sound Estuary Program

August 7, 2025

Table of contents

Preface	3
I Geographic Setting	4
1 Watersheds	6
1.1 HUCs	6
1.2 MDEQ Basin Boundaries	6
About	6
2 EPA Ecoregions	7
2.1 Level 3 Ecoregions	7
2.2 Level 4 Ecoregions	9
About	9
II Climatology	10
3 Precipitation	12
3.1 Annual Totals	12
3.2 Monthly Totals (30-year normal)	13
About	13
4 Temperature	14
4.1 Annual Mean Temperature	14
4.2 Monthly Mean Temperature	15
4.2.1 Cool Months	15
4.2.2 Warm Months	16
About	16
III Hydrology	17
5 Streams and Waterbodies	19
5.1 Amount of water	19
5.1.1 Stream Lengths	19

5.1.2	Waterbody Areas	19
5.2	Designated Uses	20
5.3	Impairments	21
	About	21
6	Freshwater Inflows	22
	About	22
7	Salinity	23
	About	23
IV	People and Land Use	24
8	Population	26
8.1	Entire watershed	26
8.2	Watershed in the state of MS	26
	About	26
9	Land Use / Land Cover	27
	About	27
	Summary	28
	Appendices	29
A	Data Processing	29
A.1	Geographic Setting	29
A.1.1	Watersheds	29
A.1.2	EPA Ecoregions	29
A.2	Climatology	30
A.2.1	Precipitation	30
A.2.2	Temperature	30
A.3	Hydrology	31
A.3.1	Streams and Waterbodies	31
A.3.2	Freshwater Inflows	32
A.3.3	Salinity	32
A.4	People and Land Use	32
A.4.1	Population	32
A.4.2	Land Use / Land Cover	33
	About	33

Preface

This website is a collection of documents characterizing the Mississippi Sound Watershed, primarily through use of geospatial data. The outputs are compiled as a ‘[quarto book](#)’. Each chapter is produced by a separate `.qmd` file.

The purpose of the book format is to make the outputs readily findable and accessible. It may also serve as an outline for future undertakings like the State of the Sound report.

Part I

Geographic Setting

Some info here about different ways of describing where we are and what comprises the watershed: USGS HUCs, DEQ's division into different basins, and EPA EcoRegions.

Possibly also the definitions from the CCMP of MS Sound, MS Sound Estuary, and MS Sound Watershed.

1 Watersheds

1.1 HUCs

Nice explanation of HUCs on [this USGS page](#).

The HUCs that make up the Mississippi Sound Watershed are 0317 (Pascagoula) and 0318 (Pearl).

Show some mappy breakdowns.

1.2 MDEQ Basin Boundaries

There are several waterways that don't actually end up in either the Pearl or Pascagoula Rivers. So the Mississippi Department of Environmental Quality uses a slightly different division.

NEED TO CONFIRM, but I think: HUC12s are the same between MDEQ's classification and the USGS HUCs. They're just grouped differently, and the streams that empty into bays or straight into the Sound are the 'Coastal Streams Basin'. There is also a 'Coastal Offshore' grouping, that is comprised of the Sound itself, the islands, and the 3-mile zone beyond the islands.

About

The `.qmd` file that generated this section was: `watersheds.qmd`.

For information on data processing, see Appendix [A.1.1](#).

2 EPA Ecoregions

For some geological classification that isn't based on watershed boundaries, we turn to the [EPA's Ecoregions](#).

2.1 Level 3 Ecoregions




Figure 2.1: Level 3 Ecoregions in Louisiana, Mississippi, and Alabama; the Mississippi Sound Watershed is outlined in black.

The Mississippi Sound Watershed contains 4 level-3 ecoregions:

- 65 Southeastern Plains
- 73 Mississippi Alluvial Plain (in the very southwest corner)

- 74 Mississippi Valley Loess Plains
- 75 Southern Coastal Plain




Figure 2.2: Level 3 Ecoregions in the Mississippi Sound Watershed only.

2.2 Level 4 Ecoregions




Figure 2.3: Level 4 Ecoregions in the Mississippi Sound Watershed.

About

The `.qmd` file that generated this section was: **EPA__Ecoregions.qmd**.

For information on data processing, see Appendix [A.1.2](#).

Part II

Climatology

Pieces of this part will be things like precipitation, temperature, etc.

3 Precipitation

The maps and information below were generated from gridded (spatial) climate normals: values averaged over 15, 30, or 100 years. Data were obtained from the National Centers for Environmental Information (NCEI).

3.1 Annual Totals





Figure 3.1: Annual Precipitation averaged over 100, 30, and 15 years.

3.2 Monthly Totals (30-year normal)



Data source: National Centers for Environmental Information

Figure 3.2: Monthly Precipitation, averaged over 30 years.

About

The `.qmd` file that generated this section was: `precipitation.qmd`.

For information on data processing, see Appendix [A.2.1](#).

4 Temperature

The maps and information below were generated from gridded (spatial) climate normals: values averaged over 15, 30, or 100 years. Data were obtained from the National Centers for Environmental Information (NCEI).

4.1 Annual Mean Temperature





Figure 4.1: Annual Mean Temperature averaged over 100, 30, and 15 years.

4.2 Monthly Mean Temperature

4.2.1 Cool Months



Data source: National Centers for Environmental Information

Figure 4.2: Comparison of long-term winter temperature averages across a 100-year baseline, and 30- and 15-year normals.

4.2.2 Warm Months


100-year baseline



30-year normal



15-year normal



Data source: National Centers for Environmental Information

Figure 4.3: Comparison of long-term spring and summer temperature averages across a 100-year baseline, and 30- and 15-year normals.

About

The `.qmd` file that generated this section was: `temperature.qmd`.

For information on data processing, see Appendix [A.2.2](#).

Part III

Hydrology

Freshwater inflows from the different sources, salinity characteristics, etc.

5 Streams and Waterbodies

5.1 Amount of water

e.g. how many miles of streams, how many acres of lakes/ponds/etc. in the watershed

TO-DO: Calculate this by basin (MDEQ definitions, not HUCs)

5.1.1 Stream Lengths

Table 5.1: Length (mi) of stream types in the MSEP watershed

Type	total_miles
Artificial Path	1,496.8
Canal/Ditch	145.3
Coastline	308.7
Connector	26.6
Stream/River	5.2
Stream/River: Intermittent	20,350.3
Stream/River: Perennial	11,730.7

5.1.2 Waterbody Areas

Table 5.2: Total area (acres) of water body types in the MSEP watershed

Type	total_acres
Lake/Pond: Intermittent	70.6
Lake/Pond: Perennial	67,602.0
Lake/Pond: Perennial; Average Water Elevation	27,726.5
Lake/Pond: Perennial; Normal Pool	9.5
Reservoir	48.2
Reservoir: Aquaculture	471.6
Reservoir; Treatment	954.8

Table 5.2: Total area (acres) of water body types in the MSEP watershed

Type	total_acres
Swamp/Marsh	251,495.2

5.2 Designated Uses

The Mississippi Department of Environmental Quality (MDEQ), per the federal Clean Water Act of 1972, is responsible for classifying surface waters in Mississippi into ‘designated uses’ (e.g. fish and wildlife, recreation, shellfish harvesting, public water supply) and creating water quality standards to ensure designated uses can be met. More detail on designated uses and water quality standards can be found on [MDEQ’s website](#).

A 305(b) report (the most recent as of this writing is from 2024; [pdf available here](#)) is generated every other year, and quantifies how many and which water bodies are / are not meeting the criteria for their designated use.

Here, we are only summarizing designated uses within the Mississippi Sound Watershed - not assessment of whether criteria are met. See the 305(b) report linked above for more detailed information.

The tables below do not include the general ‘Fish and Wildlife’ designation; that designation applies to most waters of the state.

Table 5.3: Miles of streams with each designated use in the MS Sound Watershed

Code	Description	Miles
REC	Recreation	1,011.6
PWS	Public Water Supply	26.3
SHF-REC	Shellfish Harvesting & Recreation	25.3
FW1	Fish and Wildlife	10.4
EPH	Ephemeral	1.8

Table 5.4: Area of non-linear waterbodies with each designated use in the MS Sound Watershed

Code	Description	Square Miles	Acres
REC	Recreation	742.2	475,019.3
PWS	Public Water Supply	34.8	22,259.2
SHF-REC	Shellfish Harvesting & Recreation	22.0	14,100.5
PWS-REC	Public Water Supply & Recreation	0.2	119.8




Figure 5.1: Designated uses of streams and waterbodies of the Mississippi Sound Watershed (navy outline on map), within the state of MS, as defined by MDEQ in the 2015 Water Quality Standards dataset. Data provided by MDEQ.

5.3 Impairments

303d, TMDLs - How many, where are they, what are the TMDLs for. What problems are we dealing with in the watershed.

About

The `.qmd` file that generated this section was: `streams_waterbodies.qmd`.

For information on data processing, see Appendix [A.3.1](#).

6 Freshwater Inflows

Pearl River, Pascagoula River, bays. Do some summarizing based on USGS gage data?

About

The `.qmd` file that generated this section was: `freshwater__inflows.qmd`.

For information on data processing, see Appendix [A.3.2](#).

7 Salinity

Summarize the info from the Gulf Data Atlas (e.g. the maps that went into the CCMP). Would be great to summarize the USGS salinity data and DMR's sampling data as well - some seasonal and annual details would be great.

About

The `.qmd` file that generated this section was: `salinity.qmd`.

For information on data processing, see Appendix [A.3.3](#).

Part IV

People and Land Use

Here's where population calculations and land use/land cover summaries can be generated.

8 Population

The values calculated below are based on 2020 US Census data. The population estimate for the entire state of Mississippi was ~2.96 million in 2020.

8.1 Entire watershed

This estimate includes the portions of the watershed in Louisiana and Alabama:

1,718,504

8.2 Watershed in the state of MS

This estimate includes the state of Mississippi only:

1,539,090

About

The `.qmd` file that generated this section was: **population.qmd**.

For information on data processing, see Appendix [A.4.1](#).

9 Land Use / Land Cover

Summarize LULC info by MDEQ basin here

About

The `.qmd` file that generated this section was: `landuse_landcover.qmd`.

For information on data processing, see Appendix [A.4.2](#).

Summary

Unsure what would go in a summary chapter

A Data Processing

In general, links to original data sources are provided in each section below. Because data files are routinely updated, the files as downloaded and used in this document are also available online. These files are provided for reproducibility and transparency; for your own analyses, we encourage you to seek out data from the original linked sources.

A.1 Geographic Setting

A.1.1 Watersheds

Jump to [Chapter 1](#)

A.1.1.1 HUCs

stuff

A.1.1.2 MDEQ Basins

Shapefiles were provided by the Mississippi Department of Environmental Quality in August 2025. The zipped shapefiles are available in the [DEQ_basins](#) folder of the [msepCharacterization data files](#) directory. **However, we encourage you to contact MDEQ to get the most up-to-date files for your own analyses.**

A.1.2 EPA Ecoregions

Jump to [Chapter 2](#)

Level 3 and 4 Ecoregion shapefiles were downloaded from the [EPA's website](#) on 16 June 2025 for the states of MS, AL, and LA.

The original downloaded files along with their metadata files are available in the [EPA EcoRegions](#) folder of the [msepCharacterization data files](#) directory.

Those files were read in, combined, and clipped to the Mississippi Sound Watershed boundaries (using the `outline_full` object from the [{msepBoundaries}](#) package) using the script `EPAEcoRegion_subsetting.R` ([on github](#)) in the `R_preprocessing` folder. Three objects resulted: `level3_lamsal`, level 3 ecoregions for the 3 states combined; `level3_mssoundwatershed`, level 3 ecoregions trimmed to only the Mississippi Sound Watershed area; and `level4_mssoundwatershed`, level 4 ecoregions trimmed to only the Mississippi Sound Watershed area. The level 4 ecoregions for the entire three states were not retained because there were so many categories; it really only makes sense to look at these at the smaller level.

The resulting objects were saved as layers of a geopackage, `EPA_EcoRegion.gpkg` in the `data/processed` folder. These files are available in the [processed](#) folder of the [msepCharacterization data files](#) directory.

A.2 Climatology

A.2.1 Precipitation

Jump to Chapter 3

Precipitation normals (over 15 and 30 years, and a 100-year baseline: 2006-2020, 1991-2020, and 1901-2000 respectively) were downloaded as “gridded normals” from the [National Centers for Environmental Information Climate Normals website](#) as netCDF (.nc) files. The downloaded files as used are available in the [Precipitation_NCEI](#) folder of the [msepCharacterization data files](#) directory.

These were subsetted to only monthly and annual value layers (removing layers with sd, min, max, and flags) and trimmed geographically to the states of MS, LA, and AL in the file `Precipitation_subsetting.R` ([on github](#)) in the `R_preprocessing` folder.

The resulting files were saved as netCDF files `prcp_15yearNormals.nc`, `prcp_30yearNormals.nc`, and `prcp_100yearBaseline.nc` in the `data/processed` folder. These files are available in the [processed](#) folder of the [msepCharacterization data files](#) directory.

A.2.2 Temperature

Jump to Chapter 4

Average Temperature normals (over 15 and 30 years, and a 100-year baseline: 2006-2020, 1991-2020, and 1901-2000 respectively) were downloaded as “gridded normals” from the [National Centers for Environmental Information Climate Normals website](#) as netCDF (.nc) files. The downloaded files as used are available in the [Temperature_NCEI](#) folder of the [msepCharacterization data files](#) directory.

These were subsetted to only monthly and annual value layers (removing layers with sd, min, max, and flags) and trimmed geographically to the states of MS, LA, and AL in the file `Temperature_subsetting.R` ([on github](#)) in the `R_preprocessing` folder.

The resulting files were saved as netCDF files `tempAvg_15yearNormals.nc`, `tempAvg_30yearNormals.nc`, and `tempAvg_100yearBaseline.nc` in the `data/processed` folder. These files are available in the [processed](#) folder of the [msepCharacterization data files](#) directory.

A.3 Hydrology

A.3.1 Streams and Waterbodies

Jump to Chapter 5

A.3.1.1 Stream Lengths and Waterbody Areas

Stream Lengths and Waterbody Areas were calculated from the [EPA's NHDPlus](#) dataset. Files for the [South Atlantic \(03f\)](#) and [Lower Mississippi River \(03g\)](#) regions were downloaded on 12 June 2025. The original downloaded files are available in the [EPA NHDPlus](#) folder of the [msepCharacterization data files](#) directory.

Those files were read in, combined, and clipped to only the Mississippi Sound Watershed boundaries (using the `outline_full` object from the [{msepBoundaries}](#) package) using the script `EPANHD_subsetting.R` ([on github](#)) in the `R_preprocessing` folder. The resulting files were saved as geopackages `EPA_NHDplus_flowline_MSEP.gpkg` and `EPA_NHDplus_waterbody_MSEP.gpkg` in the `data/processed` folder. These files are available in the [processed](#) folder of the [msepCharacterization data files](#) directory.

A.3.1.2 Designated Uses

Shapefiles of designated uses, already limited to the portion of the Mississippi Sound Watershed that lies within the state of MS, were provided by the Mississippi Department of Environmental Quality (MDEQ) in April 2025. These files provide the designated uses for each waterbody from the 2015 Water Quality Standards dataset. The Mississippi Sound Watershed is made up of MDEQ's Pearl River, Pascagoula River, and Coastal Streams basins. Per MDEQ, "all waters in those 3 basins not specifically listed in the data provided are classified as Fish and Wildlife." See [their website](#) for more information.

The zipped shapefiles are available in the [DEQ_designated_uses](#) folder of the [msepCharacterization data files](#) directory. **However, we encourage you to contact MDEQ to get the most up-to-date files for your own analyses.**

A.3.1.3 Impairments

Not done yet but see MDEQ's online [TMDL Tool](#).

A.3.2 Freshwater Inflows

Jump to Chapter 6

Stuff

A.3.3 Salinity

Jump to Chapter 7

Stuff - Gulf Data Atlas especially

A.4 People and Land Use

A.4.1 Population

Jump to Chapter 8

A.4.1.1 Dasymetric Population Estimates

The [EPA EnviroAtlas](#)'s dasymetric population datasets “intelligently reallocate ... population from census blocks to 30 meter pixels based on land cover and slope”. The national datasets for 2010 and 2020 were downloaded as raster files on 24 July 2025. Only the 2020 dataset has been used here.

The original downloaded file (zipped; ~3.6 GB) is available in the [Population](#) folder of the [msepCharacterization data files](#) directory.

The dataset was trimmed to Mississippi Sound Watershed boundaries (using the `outline_full` object from the [{msepBoundaries}](#) package) using the script `Population_2020_subsetting.R` ([on github](#)) in the `R_preprocessing` folder.

The resulting file, `Population_Dasymetric_2020.tif`, is available in the [processed](#) folder of the [msepCharacterization data files](#) directory.

A.4.2 Land Use / Land Cover

Jump to Chapter [9](#)

Stuff

About

The `.qmd` file that generated this section was: `data_processing.qmd`.