


# Salem Sets Avoiding Patterns

Jacob Denson

November 27, 2020

# General Research Question



- ▶ Phenomenon: Structure appears in suitably large objects.
- ▶ Like Ramsey Theory, but with a more analytical foundation, e.g. Geometric Measure Theory / Harmonic Analysis.

# Examples

# Examples

- ▶ How large can a subset  $X$  of  $\mathbf{R}^d$  be such that no right angle is formed by any three points in  $X$ .

# Examples

- ▶ How large can a subset  $X$  of  $\mathbf{R}^d$  be such that no right angle is formed by any three points in  $X$ .
- ▶ How large can an additive subgroup  $G$  of  $\mathbf{R}$  be, such that  $G \cap \mathbf{Q} = \{0\}$ .

# Examples

- ▶ How large can a subset  $X$  of  $\mathbf{R}^d$  be such that no right angle is formed by any three points in  $X$ .
- ▶ How large can an additive subgroup  $G$  of  $\mathbf{R}$  be, such that  $G \cap \mathbf{Q} = \{0\}$ .
- ▶ Our problem isn't well specified: No subset of  $\mathbf{R}^d$  with positive measure can satisfy the constraints of these problems, but we can find discrete sets of arbitrarily large cardinality which do satisfy these constraints.

# Examples

- ▶ How large can a subset  $X$  of  $\mathbf{R}^d$  be such that no right angle is formed by any three points in  $X$ .
- ▶ How large can an additive subgroup  $G$  of  $\mathbf{R}$  be, such that  $G \cap \mathbf{Q} = \{0\}$ .
- ▶ Our problem isn't well specified: No subset of  $\mathbf{R}^d$  with positive measure can satisfy the constraints of these problems, but we can find discrete sets of arbitrarily large cardinality which do satisfy these constraints.
- ▶ What does 'largeness' mean?

# Dimension Theory

- ▶ We use *fractal dimension* to measure largeness / thickness.



# Dimension Theory

- ▶ We use *fractal dimension* to measure largeness / thickness.
  - ▶ It takes  $N^1$  sidelength  $1/N$  intervals to cover  $[0, 1]$ .
  - ▶ It takes  $N^2$  sidelength  $1/N$  squares to cover  $[0, 1]^2$ .
  - ▶ It takes  $N^3$  sidelength  $1/N$  cubes to cover  $[0, 1]^3$ .

# Dimension Theory

- ▶ We use *fractal dimension* to measure largeness / thickness.
  - ▶ It takes  $N^1$  sidelength  $1/N$  intervals to cover  $[0, 1]$ .
  - ▶ It takes  $N^2$  sidelength  $1/N$  squares to cover  $[0, 1]^2$ .
  - ▶ It takes  $N^3$  sidelength  $1/N$  cubes to cover  $[0, 1]^3$ .
- ▶ A set  $X \subset \mathbf{R}^d$  has *Minkowski dimension* at least  $s$  if it takes at least  $\Omega(N^s)$  radius  $1/N$  balls to cover  $X$ .

# Dimension Theory

- ▶ We use *fractal dimension* to measure largeness / thickness.
  - ▶ It takes  $N^1$  sidelength  $1/N$  intervals to cover  $[0, 1]$ .
  - ▶ It takes  $N^2$  sidelength  $1/N$  squares to cover  $[0, 1]^2$ .
  - ▶ It takes  $N^3$  sidelength  $1/N$  cubes to cover  $[0, 1]^3$ .
- ▶ A set  $X \subset \mathbf{R}^d$  has *Minkowski dimension* at least  $s$  if it takes at least  $\Omega(N^s)$  radius  $1/N$  balls to cover  $X$ .
- ▶ *Hausdorff dimension*  $\approx$  Minkowski dimension for compact  $X$ .

# The Generic Problem

- **Avoidance Problem:** Given  $Z \subset \mathbb{R}^{nd}$ , find  $X \subset \mathbb{R}^d$  with large dimension such that for distinct points  $x_1, \dots, x_n \in X$ ,  $(x_1, \dots, x_n) \notin Z$ . We say  $X$  *avoids*  $Z$ .

# The Generic Problem

- ▶ **Avoidance Problem:** Given  $Z \subset \mathbf{R}^{nd}$ , find  $X \subset \mathbf{R}^d$  with large dimension such that for distinct points  $x_1, \dots, x_n \in X$ ,  $(x_1, \dots, x_n) \notin Z$ . We say  $X$  *avoids*  $Z$ .
- ▶ Let  $Z = \{(x, y, z) \in (\mathbf{R}^d)^3 : (x - z) \cdot (y - z) = 0\}$ .
  - ▶  $X \subset \mathbf{R}^d$  avoids  $Z$  iff  $X$  does not contain any right angles.

# The Generic Problem

- ▶ **Avoidance Problem:** Given  $Z \subset \mathbf{R}^{nd}$ , find  $X \subset \mathbf{R}^d$  with large dimension such that for distinct points  $x_1, \dots, x_n \in X$ ,  $(x_1, \dots, x_n) \notin Z$ . We say  $X$  *avoids*  $Z$ .
- ▶ Let  $Z = \{(x, y, z) \in (\mathbf{R}^d)^3 : (x - z) \cdot (y - z) = 0\}$ .
  - ▶  $X \subset \mathbf{R}^d$  avoids  $Z$  iff  $X$  does not contain any right angles.
- ▶ How does the geometry of  $Z$  help us?

# The Generic Problem

- ▶ **Avoidance Problem:** Given  $Z \subset \mathbf{R}^{nd}$ , find  $X \subset \mathbf{R}^d$  with large dimension such that for distinct points  $x_1, \dots, x_n \in X$ ,  $(x_1, \dots, x_n) \notin Z$ . We say  $X$  *avoids*  $Z$ .
- ▶ Let  $Z = \{(x, y, z) \in (\mathbf{R}^d)^3 : (x - z) \cdot (y - z) = 0\}$ .
  - ▶  $X \subset \mathbf{R}^d$  avoids  $Z$  iff  $X$  does not contain any right angles.
- ▶ How does the geometry of  $Z$  help us?
- ▶ e.g.  $Z$  is a degree 2 algebraic hypersurface in last example.

# Results in Literature



## Results in Literature

- ▶ Mathé (2012): If  $Z \subset \mathbb{R}^{nd}$  is an algebraic hypersurfaces specified by a rational coefficient polynomial with degree at most  $r$ , then we can  $X \subset \mathbb{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{d}{r}.$$

# Results in Literature

- ▶ Mathé (2012): If  $Z \subset \mathbf{R}^{nd}$  is an algebraic hypersurfaces specified by a rational coefficient polynomial with degree at most  $r$ , then we can  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{d}{r}.$$

- ▶ Fraser and Pramanik (2016): If  $Z \subset \mathbf{R}^{nd}$  is a smooth hypersurface with dimension at most  $m$ , we can find  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{nd - m}{n - 1}.$$

# Results in Literature

- ▶ Mathé (2012): If  $Z \subset \mathbf{R}^{nd}$  is an algebraic hypersurfaces specified by a rational coefficient polynomial with degree at most  $r$ , then we can  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{d}{r}.$$

- ▶ Fraser and Pramanik (2016): If  $Z \subset \mathbf{R}^{nd}$  is a smooth hypersurface with dimension at most  $m$ , we can find  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{nd - m}{n - 1}.$$

- ▶ What if we use less rigid geometric information, i.e. the fractal dimension of the set  $Z$ ?

# Our Results

- Denson, Pramanik, and Zahl (2019): If  $Z \subset \mathbf{R}^{nd}$  is a set with Minkowski dimension bounded by  $s$ , we can find  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{nd - s}{n - 1}.$$

# Our Results

- ▶ Denson, Pramanik, and Zahl (2019): If  $Z \subset \mathbf{R}^{nd}$  is a set with Minkowski dimension bounded by  $s$ , we can find  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{nd - s}{n - 1}.$$

- ▶ Denson (2019): If  $Z \subset \mathbf{R}^n$ , and  $\pi : \mathbf{R}^n \rightarrow \mathbf{R}^m$  is a rational coefficient projection map such that  $\pi(Z)$  has Minkowski dimension bounded by  $s$ , then we can find  $X \subset \mathbf{R}$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{m - s}{m}.$$

# Our Results

- ▶ Denson, Pramanik, and Zahl (2019): If  $Z \subset \mathbf{R}^{nd}$  is a set with Minkowski dimension bounded by  $s$ , we can find  $X \subset \mathbf{R}^d$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{nd - s}{n - 1}.$$

- ▶ Denson (2019): If  $Z \subset \mathbf{R}^n$ , and  $\pi : \mathbf{R}^n \rightarrow \mathbf{R}^m$  is a rational coefficient projection map such that  $\pi(Z)$  has Minkowski dimension bounded by  $s$ , then we can find  $X \subset \mathbf{R}^m$  avoiding  $Z$  with

$$\dim_{\mathbf{H}}(X) = \frac{m - s}{m}.$$

- ▶ (Proved in Msc Thesis, but want to find higher dimensional result before full publication).

# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?

# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?
- ▶ **Theorem:** Let  $H \subset \mathbf{R}$  be a set with Minkowski dimension  $s$ . Then we can find an additive subgroup  $G \subset \mathbf{R}$  such that  $G \cap H \subset \{0\}$  and such that  $\dim_{\mathbf{H}}(G) = 1 - s$ .



# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?
- ▶ **Theorem:** Let  $H \subset \mathbf{R}$  be a set with Minkowski dimension  $s$ . Then we can find an additive subgroup  $G \subset \mathbf{R}$  such that  $G \cap H \subset \{0\}$  and such that  $\dim_{\mathbf{H}}(G) = 1 - s$ .
  - ▶ For each  $n > 0$  and  $m \in \mathbf{Z}^n$ , let

$$Z_m = \{(x_1, \dots, x_n) \in \mathbf{R}^{nd} : m_1 x_1 + \dots + m_n x_n \in H\}.$$

If  $\pi_m(x_1, \dots, x_n) = m_1 x_1 + \dots + m_n x_n$ , then  $\pi_m(Z_m) = H$  has Minkowski dimension  $s$ .

# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?
- ▶ **Theorem:** Let  $H \subset \mathbf{R}$  be a set with Minkowski dimension  $s$ . Then we can find an additive subgroup  $G \subset \mathbf{R}$  such that  $G \cap H \subset \{0\}$  and such that  $\dim_{\mathbf{H}}(G) = 1 - s$ .
  - ▶ For each  $n > 0$  and  $m \in \mathbf{Z}^n$ , let

$$Z_m = \{(x_1, \dots, x_n) \in \mathbf{R}^{nd} : m_1x_1 + \dots + m_nx_n \in H\}.$$

If  $\pi_m(x_1, \dots, x_n) = m_1x_1 + \dots + m_nx_n$ , then  $\pi_m(Z_m) = H$  has Minkowski dimension  $s$ .

- ▶ Thus we can apply the theorem to find  $X$  avoiding  $Z_m$  with Hausdorff dimension  $1 - s$ .

# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?
- ▶ **Theorem:** Let  $H \subset \mathbf{R}$  be a set with Minkowski dimension  $s$ . Then we can find an additive subgroup  $G \subset \mathbf{R}$  such that  $G \cap H \subset \{0\}$  and such that  $\dim_{\mathbf{H}}(G) = 1 - s$ .
  - ▶ For each  $n > 0$  and  $m \in \mathbf{Z}^n$ , let

$$Z_m = \{(x_1, \dots, x_n) \in \mathbf{R}^{nd} : m_1x_1 + \dots + m_nx_n \in H\}.$$

If  $\pi_m(x_1, \dots, x_n) = m_1x_1 + \dots + m_nx_n$ , then  $\pi_m(Z_m) = H$  has Minkowski dimension  $s$ .

- ▶ Thus we can apply the theorem to find  $X$  avoiding  $Z_m$  with Hausdorff dimension  $1 - s$ .
- ▶ Repeat countably many times to find  $X$  avoiding  $Z_m$  for all  $m$  with Hausdorff dimension  $1 - s$ .

# Applications

- ▶ Given a subgroup  $H \subset \mathbf{R}$ , is it possible to find  $G \subset \mathbf{R}$  such that  $G + H = \mathbf{R}$ ?
- ▶ **Theorem:** Let  $H \subset \mathbf{R}$  be a set with Minkowski dimension  $s$ . Then we can find an additive subgroup  $G \subset \mathbf{R}$  such that  $G \cap H \subset \{0\}$  and such that  $\dim_{\mathbf{H}}(G) = 1 - s$ .
  - ▶ For each  $n > 0$  and  $m \in \mathbf{Z}^n$ , let

$$Z_m = \{(x_1, \dots, x_n) \in \mathbf{R}^{nd} : m_1 x_1 + \dots + m_n x_n \in H\}.$$

If  $\pi_m(x_1, \dots, x_n) = m_1 x_1 + \dots + m_n x_n$ , then  $\pi_m(Z_m) = H$  has Minkowski dimension  $s$ .

- ▶ Thus we can apply the theorem to find  $X$  avoiding  $Z_m$  with Hausdorff dimension  $1 - s$ .
- ▶ Repeat countably many times to find  $X$  avoiding  $Z_m$  for all  $m$  with Hausdorff dimension  $1 - s$ .
- ▶ We would hope that whatever higher dimensional generalization would construct  $G \subset \mathbf{R}^d$  with Hausdorff dimension  $d - s$  for any  $H$  of Minkowski dimension  $s$ .

# Applications

- ▶ Since we use 'rough' geometric information about  $Z$ , our method can even consider 'avoidance problems on fractals'.

# Applications

- ▶ Since we use 'rough' geometric information about  $Z$ , our method can even consider 'avoidance problems on fractals'.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .

# Applications

- ▶ Since we use ‘rough’ geometric information about  $Z$ , our method can even consider ‘avoidance problems on fractals’.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .
  - ▶ If  $\pi : E \rightarrow \mathbf{R}$  is a projection onto the line at a  $45^\circ$  angle,  $\pi(E)$  is an interval.

# Applications

- ▶ Since we use 'rough' geometric information about  $Z$ , our method can even consider 'avoidance problems on fractals'.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .
  - ▶ If  $\pi : E \rightarrow \mathbf{R}$  is a projection onto the line at a  $45^\circ$  angle,  $\pi(E)$  is an interval.
  - ▶ Let

$$Z = \left\{ (x, y, z) \in \pi(E)^3 : \begin{array}{l} \text{there is } x_0, y_0, z_0 \in E \\ \text{s.t. } (x_0 - z_0) \cdot (y_0 - z_0) = 0 \end{array} \right\}.$$



# Applications

- ▶ Since we use ‘rough’ geometric information about  $Z$ , our method can even consider ‘avoidance problems on fractals’.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .
  - ▶ If  $\pi : E \rightarrow \mathbf{R}$  is a projection onto the line at a  $45^\circ$  angle,  $\pi(E)$  is an interval.
  - ▶ Let

$$Z = \left\{ (x, y, z) \in \pi(E)^3 : \begin{array}{l} \text{there is } x_0, y_0, z_0 \in E \\ \text{s.t. } (x_0 - z_0) \cdot (y_0 - z_0) = 0 \end{array} \right\}.$$

- ▶ Basic considerations suggest that  $\dim_{\mathbf{M}}(Z) = 2$ , so that we can find  $X \subset \pi(E)$  avoiding  $Z$  with Hausdorff dimension  $1/2$ .

# Applications

- ▶ Since we use 'rough' geometric information about  $Z$ , our method can even consider 'avoidance problems on fractals'.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .
  - ▶ If  $\pi : E \rightarrow \mathbf{R}$  is a projection onto the line at a  $45^\circ$  angle,  $\pi(E)$  is an interval.
  - ▶ Let

$$Z = \left\{ (x, y, z) \in \pi(E)^3 : \begin{array}{l} \text{there is } x_0, y_0, z_0 \in E \\ \text{s.t. } (x_0 - z_0) \cdot (y_0 - z_0) = 0 \end{array} \right\}.$$


- ▶ Basic considerations suggest that  $\dim_{\mathbf{M}}(Z) = 2$ , so that we can find  $X \subset \pi(E)$  avoiding  $Z$  with Hausdorff dimension  $1/2$ .
- ▶ Then  $\pi^{-1}(X)$  avoids right angles and  $\dim_{\mathbf{H}}(\pi^{-1}(X)) \geq 1/2$ .

# Applications

- ▶ Since we use 'rough' geometric information about  $Z$ , our method can even consider 'avoidance problems on fractals'.
  - ▶ Consider the Cantor dust  $E$ , with  $\dim_{\mathbf{M}}(E) = 1$ .
  - ▶ If  $\pi : E \rightarrow \mathbf{R}$  is a projection onto the line at a  $45^\circ$  angle,  $\pi(E)$  is an interval.
  - ▶ Let

$$Z = \left\{ (x, y, z) \in \pi(E)^3 : \begin{array}{l} \text{there is } x_0, y_0, z_0 \in E \\ \text{s.t. } (x_0 - z_0) \cdot (y_0 - z_0) = 0 \end{array} \right\}.$$

- ▶ Basic considerations suggest that  $\dim_{\mathbf{M}}(Z) = 2$ , so that we can find  $X \subset \pi(E)$  avoiding  $Z$  with Hausdorff dimension  $1/2$ .
  - ▶ Then  $\pi^{-1}(X)$  avoids right angles and  $\dim_{\mathbf{H}}(\pi^{-1}(X)) \geq 1/2$ .
- ▶ We have also used this technique to bound the existence of isosceles triangles on Lipschitz curves.



# Fourier Dimension

# Fourier Dimension

- ▶ A set  $X \subset \mathbf{R}^d$  has *Fourier dimension* at least  $s$  if there exists a finite measure  $\mu$  supported on  $X$  such that  $|\hat{\mu}(\xi)| \lesssim |\xi|^{-s/2}$ .

# Fourier Dimension

- ▶ A set  $X \subset \mathbf{R}^d$  has *Fourier dimension* at least  $s$  if there exists a finite measure  $\mu$  supported on  $X$  such that  $|\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}$ .
- ▶ Often gives much more structural information about a set than Minkowski dimension does.





# Fourier Dimension

- ▶ A set  $X \subset \mathbf{R}^d$  has *Fourier dimension* at least  $s$  if there exists a finite measure  $\mu$  supported on  $X$  such that  $|\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}$ .
- ▶ Often gives much more structural information about a set than Minkowski dimension does.
- ▶ (Keleti, 1998) There exist an 'independent' set  $X$  with full Hausdorff dimension such that there exists no nontrivial solutions to  $m_1x_1 + \dots + m_nx_n = 0$  for any  $m \in \mathbf{Z}^n$  and  $x_1, \dots, x_n \in X$ .



# Fourier Dimension

- ▶ A set  $X \subset \mathbf{R}^d$  has *Fourier dimension* at least  $s$  if there exists a finite measure  $\mu$  supported on  $X$  such that  $|\hat{\mu}(\xi)| \lesssim |\xi|^{-s/2}$ .
- ▶ Often gives much more structural information about a set than Minkowski dimension does.
- ▶ (Keleti, 1998) There exist an 'independent' set  $X$  with full Hausdorff dimension such that there exists no nontrivial solutions to  $m_1x_1 + \dots + m_nx_n = 0$  for any  $m \in \mathbf{Z}^n$  and  $x_1, \dots, x_n \in X$ .
- ▶ (Rudin, 1960) If  $X$  has Fourier dimension greater than  $1/n$ , then there exists some  $m \in \mathbf{Z}^n$  and some  $x_1, \dots, x_n \in X$  such that  $m_1x_1 + \dots + m_nx_n = 0$ .



# Getting Fourier Dimension Bounds Is Hard

- ▶ For any set  $X$ ,  $\dim_{\mathbf{F}}(X) \leq \dim_{\mathbf{H}}(X) \leq \dim_{\mathbf{M}}(X)$ .

# Getting Fourier Dimension Bounds Is Hard


- ▶ For any set  $X$ ,  $\dim_{\mathbf{F}}(X) \leq \dim_{\mathbf{H}}(X) \leq \dim_{\mathbf{M}}(X)$ .
- ▶  $X$  is a *Salem Set* if  $\dim_{\mathbf{F}}(X) = \dim_{\mathbf{H}}(X)$ .

# Getting Fourier Dimension Bounds Is Hard

- ▶ For any set  $X$ ,  $\dim_{\mathbf{F}}(X) \leq \dim_{\mathbf{H}}(X) \leq \dim_{\mathbf{M}}(X)$ .
- ▶  $X$  is a *Salem Set* if  $\dim_{\mathbf{F}}(X) = \dim_{\mathbf{H}}(X)$ .
- ▶ All sets are Salem 'generically', but for most explicit constructions, the Fourier dimension is equal to zero.


# Getting Fourier Dimension Bounds Is Hard

- ▶ For any set  $X$ ,  $\dim_{\mathbf{F}}(X) \leq \dim_{\mathbf{H}}(X) \leq \dim_{\mathbf{M}}(X)$ .
- ▶  $X$  is a *Salem Set* if  $\dim_{\mathbf{F}}(X) = \dim_{\mathbf{H}}(X)$ .
- ▶ All sets are Salem 'generically', but for most explicit constructions, the Fourier dimension is equal to zero.
- ▶ If  $X$  is the Cantor set, then  $\dim_{\mathbf{H}}(X) = \dim_{\mathbf{M}}(X) = \log_3(2)$ , but  $\dim_{\mathbf{F}}(X) = 0$ .



# Getting Fourier Dimension Bounds Is Hard

- ▶ For any set  $X$ ,  $\dim_{\mathbf{F}}(X) \leq \dim_{\mathbf{H}}(X) \leq \dim_{\mathbf{M}}(X)$ .
- ▶  $X$  is a *Salem Set* if  $\dim_{\mathbf{F}}(X) = \dim_{\mathbf{H}}(X)$ .
- ▶ All sets are Salem 'generically', but for most explicit constructions, the Fourier dimension is equal to zero.
- ▶ If  $X$  is the Cantor set, then  $\dim_{\mathbf{H}}(X) = \dim_{\mathbf{M}}(X) = \log_3(2)$ , but  $\dim_{\mathbf{F}}(X) = 0$ .



- ▶ Heuristic: Typically need 'square root cancellation' to obtain optimal Fourier decay, e.g. by using randomness.

# Salem Set Result



# Salem Set Result

## Theorem (2020, Denson)

*If  $Z$  has lower Minkowski dimension bounded by  $s$ , we can find  $X$  avoiding  $Z$  with*

$$\dim_{\mathrm{H}}(X) = \dim_{\mathrm{F}}(X) = \frac{nd - s}{n - 1/2}.$$

# Salem Set Result

## Theorem (2020, Denson)

*If  $Z$  has lower Minkowski dimension bounded by  $s$ , we can find  $X$  avoiding  $Z$  with*

$$\dim_{\mathrm{H}}(X) = \dim_{\mathrm{F}}(X) = \frac{nd - s}{n - 1/2}.$$

*Conjecture: If  $Z$  is 'suitably smooth', then we can find  $X$  with  $\dim_{\mathrm{H}}(X) = \dim_{\mathrm{F}}(X) = (nd - s)/(n - 1)$ .*

Thanks for listening!