

Algorithmic Aspects of the Brascamp Lieb Inequality

Jacob Denson (Based on Gurvits, 2004)

University of Wisconsin Madison

September 28, 2021

Lieb's Theorem

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

Lieb's Theorem

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- (Lieb, 1990) To calculate $\text{BL}(B, p)$, plug in Gaussians.

Lieb's Theorem

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Lieb, 1990) To calculate $\text{BL}(B, p)$, plug in Gaussians.
- ▶ If $f_i(x) = e^{-\pi(A_i x) \cdot x}$ for $A_i \succ 0$, then

Lieb's Theorem

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Lieb, 1990) To calculate $\text{BL}(B, p)$, plug in Gaussians.
- ▶ If $f_i(x) = e^{-\pi(A_i x) \cdot x}$ for $A_i \succ 0$, then

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx = (\sum p_i B_i^* A_i B_i)^{-1/2}$$

Lieb's Theorem

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Lieb, 1990) To calculate $\text{BL}(B, p)$, plug in Gaussians.
- ▶ If $f_i(x) = e^{-\pi(A_i x) \cdot x}$ for $A_i \succ 0$, then

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx = (\sum p_i B_i^* A_i B_i)^{-1/2}$$

$$\prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i} = (\prod_{i=1}^m \det(A_i)^{p_i})^{-1/2}.$$

Thus

$$(\sum p_i B_i^* A_i B_i)^{-1/2} \leq \text{BL}(B, p) \cdot (\prod_{i=1}^m \det(A_i)^{p_i})^{-1/2}$$

Lieb's Theorem

- Rearranging gives

$$\mathrm{BL}(B, p) = \left(\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i B_i^* A_i B_i)} \right)^{1/2}.$$

Lieb's Theorem

- ▶ Rearranging gives

$$\text{BL}(B, p) = \left(\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i B_i^* A_i B_i)} \right)^{1/2}.$$

- ▶ Non convex objective, so tricky to optimize (NP hard).

Lieb's Theorem

- ▶ Rearranging gives

$$\text{BL}(B, p) = \left(\sup_{A_1, \dots, A_m \succcurlyeq 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i B_i^* A_i B_i)} \right)^{1/2}.$$

- ▶ Non convex objective, so tricky to optimize (NP hard).
- ▶ Checking Finiteness is also non-obvious.

Lieb's Theore

- ▶ Rearranging gives

$$\text{BL}(B, p) = \left(\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i B_i^* A_i B_i)} \right)^{1/2}.$$

- ▶ Non convex objective, so tricky to optimize (NP hard).
- ▶ Checking Finiteness is also non-obvious.
- ▶ BCCT gives a family of conditions

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

Since $0 \leq \dim(V) \leq n$ and $0 \leq \dim(B_i V) \leq n_i$, only finitely many linear conditions on the p_i .

Lieb's Theore

- ▶ Rearranging gives

$$\text{BL}(B, p) = \left(\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i B_i^* A_i B_i)} \right)^{1/2}.$$

- ▶ Non convex objective, so tricky to optimize (NP hard).
- ▶ Checking Finiteness is also non-obvious.
- ▶ BCCT gives a family of conditions

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

Since $0 \leq \dim(V) \leq n$ and $0 \leq \dim(B_i V) \leq n_i$, only finitely many linear conditions on the p_i .

- ▶ But there can be exponentially many, so still tricky to compute in practice.

Geometric Brascamp Lieb

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}$$

Geometric Brascamp Lieb

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}$$

- A Brascamp-Lieb inequality is *geometric* if each B_i is a projection ($B_i B_i^* = I$), and $\sum_i p_i B_i^* B_i = I$.

Geometric Brascamp Lieb

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}$$

- ▶ A Brascamp-Lieb inequality is *geometric* if each B_i is a projection ($B_i B_i^* = I$), and $\sum_i p_i B_i^* B_i = I$.
- ▶ Hölder's inequality and Loomis-Whitney are special cases.

Geometric Brascamp Lieb

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}$$

- ▶ A Brascamp-Lieb inequality is *geometric* if each B_i is a projection ($B_i B_i^* = I$), and $\sum_i p_i B_i^* B_i = I$.
- ▶ Hölder's inequality and Loomis-Whitney are special cases.
- ▶ Plugging in $f_i(x) = e^{-\pi|x|^2}$ gives $\text{BL}(B, p) \geq 1$.

Geometric Brascamp Lieb

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}$$

- ▶ A Brascamp-Lieb inequality is *geometric* if each B_i is a projection ($B_i B_i^* = I$), and $\sum_i p_i B_i^* B_i = I$.
- ▶ Hölder's inequality and Loomis-Whitney are special cases.
- ▶ Plugging in $f_i(x) = e^{-\pi|x|^2}$ gives $\text{BL}(B, p) \geq 1$.
- ▶ (Barthe, 1998), generalizing (Ball, 1989), showed that these functions are extremizers, i.e. $\text{BL}(B, p) = 1$.

Operator Rescaling

- ▶ What happens to $\text{BL}(B, p)$ if we ‘rescale / change coordinates’.

Operator Rescaling

- ▶ What happens to $\text{BL}(B, p)$ if we 'rescale / change coordinates'.
- ▶ Fix invertible matrices M and M_1, \dots, M_m , and consider the Brascamp-Lieb inequality with matrices $B'_i = M_i B_i M_i$,

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(M_i B_i M x)|^{p_i} dx \leq \text{BL}(B', p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

Operator Rescaling

- ▶ What happens to $\text{BL}(B, p)$ if we 'rescale / change coordinates'.
- ▶ Fix invertible matrices M and M_1, \dots, M_m , and consider the Brascamp-Lieb inequality with matrices $B'_i = M_i B_i M_i$,

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(M_i B_i M x)|^{p_i} dx \leq \text{BL}(B', p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ Then

$$\begin{aligned} \text{BL}(B', p) &= \sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum_i p_i M_i^* B_i^* M_i^* A_i M_i B_i M)} \\ &= \sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det((M_i^{-1})^* A_i M_i^{-1})^{p_i}}{\det(M^* (\sum_i p_i B_i^* A_i B_i) M)} \\ &= \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \text{BL}(B, p). \end{aligned}$$

Operator Rescaling

$$\mathrm{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \mathrm{BL}(B, p)$$

Operator Rescaling

$$\mathrm{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \mathrm{BL}(B, p)$$

- If (B', p) is geometric, $\mathrm{BL}(B', p) = 1$, so

$$\mathrm{BL}(B, p) = \det(M)^2 \prod_i \det(M_i)^{2p_i}.$$

Operator Rescaling

$$\mathrm{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \mathrm{BL}(B, p)$$

- ▶ If (B', p) is geometric, $\mathrm{BL}(B', p) = 1$, so

$$\mathrm{BL}(B, p) = \det(M)^2 \prod_i \det(M_i)^{2p_i}.$$

- ▶ (Bennett, Carbery, Christ, Tao, 2008) Geometric rescaling possible iff the Brascamp-Lieb equation has extremizers.

Operator Rescaling

$$\text{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \text{BL}(B, p)$$

- ▶ If (B', p) is geometric, $\text{BL}(B', p) = 1$, so

$$\text{BL}(B, p) = \det(M)^2 \prod_i \det(M_i)^{2p_i}.$$

- ▶ (Bennett, Carbery, Christ, Tao, 2008) Geometric rescaling possible iff the Brascamp-Lieb equation has extremizers.
- ▶ (Garg, Gurvits, Oliveira, Wigderson, 2018) If $\text{BL}(B, p) < \infty$, we can rescale to be within a ε of a geometric Brascamp-Lieb equation.

Operator Rescaling

$$\text{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \text{BL}(B, p)$$

- ▶ If (B', p) is geometric, $\text{BL}(B', p) = 1$, so

$$\text{BL}(B, p) = \det(M)^2 \prod_i \det(M_i)^{2p_i}.$$

- ▶ (Bennett, Carbery, Christ, Tao, 2008) Geometric rescaling possible iff the Brascamp-Lieb equation has extremizers.
- ▶ (Garg, Gurvits, Oliveira, Wigderson, 2018) If $\text{BL}(B, p) < \infty$, we can rescale to be within a ε of a geometric Brascamp-Lieb equation.
 - ▶ We can do this algorithmically, i.e. a computer can compute a ε -approximate geometric rescaling in $\text{Poly}(\text{Bits}(B), \log(p), 1/\varepsilon)$ computations.

Operator Rescaling

$$\text{BL}(B', p) = \det(M)^{-2} \prod_i \det(M_i)^{-2p_i} \cdot \text{BL}(B, p)$$

- ▶ If (B', p) is geometric, $\text{BL}(B', p) = 1$, so

$$\text{BL}(B, p) = \det(M)^2 \prod_i \det(M_i)^{2p_i}.$$

- ▶ (Bennett, Carbery, Christ, Tao, 2008) Geometric rescaling possible iff the Brascamp-Lieb equation has extremizers.
- ▶ (Garg, Gurvits, Oliveira, Wigderson, 2018) If $\text{BL}(B, p) < \infty$, we can rescale to be within a ε of a geometric Brascamp-Lieb equation.
 - ▶ We can do this algorithmically, i.e. a computer can compute a ε -approximate geometric rescaling in $\text{Poly}(\text{Bits}(B), \log(p), 1/\varepsilon)$ computations.
 - ▶ Conversely, we can determine if $\text{BL}(B, p) = \infty$ in $\text{Poly}(\text{Bits}(B), \log(p))$ computations.

Computing Permanents (An Analogous Problem)

- ▶ Given an $n \times n$ matrix A with non-negative entries, define

$$\text{Perm}(A) = \sum_{\sigma \in S_n} \prod_i A_{i\sigma(i)}.$$

Computing Permanents (An Analogous Problem)

- ▶ Given an $n \times n$ matrix A with non-negative entries, define

$$\text{Perm}(A) = \sum_{\sigma \in S_n} \prod_i A_{i\sigma(i)}.$$

- ▶ This is (maybe suprisingly) NP hard to compute.

Computing Permanents (An Analogous Problem)

- ▶ Given an $n \times n$ matrix A with non-negative entries, define

$$\text{Perm}(A) = \sum_{\sigma \in S_n} \prod_i A_{i\sigma(i)}.$$

- ▶ This is (maybe suprisingly) NP hard to compute.
 - ▶ If $R = \text{diag}(\lambda_1, \dots, \lambda_n)$ and $C = \text{diag}(\gamma_1, \dots, \gamma_n)$, then

$$\begin{aligned}\text{Perm}(RAC) &= (\lambda_1 \dots \lambda_n)(\gamma_1 \dots \gamma_n)\text{Perm}(A) \\ &= \det(R) \cdot \det(C) \cdot \text{Perm}(A).\end{aligned}$$

Computing Permanents (An Analogous Problem)

- ▶ Given an $n \times n$ matrix A with non-negative entries, define

$$\text{Perm}(A) = \sum_{\sigma \in S_n} \prod_i A_{i\sigma(i)}.$$

- ▶ This is (maybe suprisingly) NP hard to compute.
 - ▶ If $R = \text{diag}(\lambda_1, \dots, \lambda_n)$ and $C = \text{diag}(\gamma_1, \dots, \gamma_n)$, then

$$\begin{aligned}\text{Perm}(RAC) &= (\lambda_1 \dots \lambda_n)(\gamma_1 \dots \gamma_n)\text{Perm}(A) \\ &= \det(R) \cdot \det(C) \cdot \text{Perm}(A).\end{aligned}$$

- ▶ (Egorychev, 1981), (Falikman, 1981) If A is *doubly stochastic*, $e^{-n} \leq \text{Perm}(A) \leq 1$.

Computing Permanents (An Analogous Problem)

- ▶ Given an $n \times n$ matrix A with non-negative entries, define

$$\text{Perm}(A) = \sum_{\sigma \in S_n} \prod_i A_{i\sigma(i)}.$$

- ▶ This is (maybe suprisingly) NP hard to compute.
 - ▶ If $R = \text{diag}(\lambda_1, \dots, \lambda_n)$ and $C = \text{diag}(\gamma_1, \dots, \gamma_n)$, then

$$\begin{aligned}\text{Perm}(RAC) &= (\lambda_1 \dots \lambda_n)(\gamma_1 \dots \gamma_n)\text{Perm}(A) \\ &= \det(R) \cdot \det(C) \cdot \text{Perm}(A).\end{aligned}$$

- ▶ (Egorychev, 1981), (Falikman, 1981) If A is *doubly stochastic*, $e^{-n} \leq \text{Perm}(A) \leq 1$.
- ▶ If RAC is doubly stochastic, then

$$\text{Perm}(A) \leq \det(R)^{-1} \det(C)^{-1} \leq e^n \cdot \text{Perm}(A),$$

so $\text{Perm}(A) \approx \det(R)^{-1} \det(C)^{-1}$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.
- ▶ **Two key facts:**

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.
- ▶ **Two key facts:**
 - (1) $\text{Per}(A_i) \leq 1$ for all $i \geq 1$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.
- ▶ **Two key facts:**
 - (1) $\text{Per}(A_i) \leq 1$ for all $i \geq 1$.
 - (2) Let $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.
- ▶ **Two key facts:**
 - (1) $\text{Per}(A_i) \leq 1$ for all $i \geq 1$.
 - (2) Let $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$.
 - ▶ If $\Delta_i \leq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.
 - ▶ If $\Delta_i \geq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C)\text{Per}(A_i)$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.
- ▶ **Two key facts:**
 - (1) $\text{Per}(A_i) \leq 1$ for all $i \geq 1$.
 - (2) Let $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$.
 - ▶ If $\Delta_i \leq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.
 - ▶ If $\Delta_i \geq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C)\text{Per}(A_i)$.
- ▶ Thus $\text{Per}(A_i)$ is bounded, monotonic, converges to $P \leq 1$.

- ▶ We continue this iteration, obtaining a sequence

$$A \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- ▶ Claim: If $\text{Per}(A) > 0$, these matrices become arbitrarily close to being doubly stochastic as $i \rightarrow \infty$.
- ▶ Let $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, be the row / column sums of A_i .
 - ▶ i.e. so that $A_1 = \text{diag}(1/\gamma_0)A$, $A_2 = A_1 \text{diag}(1/\gamma_1)$, etc.

- ▶ **Two key facts:**

(1) $\text{Per}(A_i) \leq 1$ for all $i \geq 1$.

(2) Let $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$.

▶ If $\Delta_i \leq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.

▶ If $\Delta_i \geq 1$, then $\text{Per}(A_{i+1}) \geq (1 + C)\text{Per}(A_i)$.

- ▶ Thus $\text{Per}(A_i)$ is bounded, monotonic, converges to $P \leq 1$.
- ▶ If $\text{Per}(A_i) \geq P - \varepsilon$ for $\varepsilon \ll 1$, then

$$P \geq \text{Per}(A_{i+1}) \geq (1 + C \cdot \Delta_i) \cdot \text{Per}(A_i) \geq (1 + C \cdot \Delta_i)(P - \varepsilon).$$

Thus $\Delta_i \leq (C_0/P)\varepsilon$. Taking $\varepsilon \rightarrow 0$ shows $\Delta \rightarrow 0$.

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**
 - ▶ Simple inductive argument: the hypothesis is that if the rows of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$. Now expand along minors (ala the determinant).

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**
 - ▶ Simple inductive argument: the hypothesis is that if the rows of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$. Now expand along minors (ala the determinant).
- ▶ **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**
 - ▶ Simple inductive argument: the hypothesis is that if the rows of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$. Now expand along minors (ala the determinant).
- ▶ **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**
 - ▶ Write $\gamma_i = 1 + \delta_i$.

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**
 - ▶ Simple inductive argument: the hypothesis is that if the rows of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$. Now expand along minors (ala the determinant).
- ▶ **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**
 - ▶ Write $\gamma_i = 1 + \delta_i$.
 - ▶ Then $\sum \delta_i^2 = \Delta_i$, and $\sum \delta_i = 0$.

Sinkhorn Iteration

- ▶ **Proof that $\text{Per}(A_i) \leq 1$:**

- ▶ Simple inductive argument: the hypothesis is that if the rows of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$. Now expand along minors (ala the determinant).

- ▶ **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**

- ▶ Write $\gamma_i = 1 + \delta_i$.

- ▶ Then $\sum \delta_i^2 = \Delta_i$, and $\sum \delta_i = 0$.

- ▶ Since $1 + t \leq \exp(t - t^2/2 + t^3/3)$,

$$\begin{aligned}\text{Per}(A_i)/\text{Per}(A_{i+1}) &= \gamma_1 \dots \gamma_n \\ &= (1 + \delta_1) \dots (1 + \delta_n) \\ &\leq \exp\left(\sum \delta_i - \sum \delta_i^2/2 + \sum \delta_i^3/3\right) \\ &\leq \exp(0 - \Delta/2 + \Delta^{3/2}/3) \\ &= 1 - \Delta/2 + O(\Delta^{3/2}).\end{aligned}$$

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- Goal: Rescale our inputs so that

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- ▶ Goal: Rescale our inputs so that
 - ▶ (Isotropy) $\sum p_i B_i^* B_i = I$.

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- ▶ Goal: Rescale our inputs so that
 - ▶ (Isotropy) $\sum p_i B_i^* B_i = I$.
 - ▶ (Projection) $B_i B_i^* = I$ for each i .

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - ▶ (Projection Normalization)

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - ▶ (Projection Normalization)
 - ▶ Let $M_i = B_i B_i^*$.

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - ▶ (Projection Normalization)
 - ▶ Let $M_i = B_i B_i^*$.
 - ▶ Replace B_i with $B'_i = M_i^{-1/2} B_i$.

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - ▶ (Projection Normalization)
 - ▶ Let $M_i = B_i B_i^*$.
 - ▶ Replace B_i with $B'_i = M_i^{-1/2} B_i$.
 - ▶ Then $(B'_i)^* B'_i = I$ for each i .

Iteration of Operator Rescaling

- ▶ Sinkhorn: Alternately apply the following two procedures:
 - ▶ (Isotropy Normalization)
 - ▶ Let $M = \sum_i p_i B_i^* B_i$.
 - ▶ Replace B_i with $B'_i = B_i M^{-1/2}$.
 - ▶ Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - ▶ (Projection Normalization)
 - ▶ Let $M_i = B_i B_i^*$.
 - ▶ Replace B_i with $B'_i = M_i^{-1/2} B_i$.
 - ▶ Then $(B'_i)^* B'_i = I$ for each i .
- ▶ We obtain a sequence $B \rightarrow B_1 \rightarrow B_2 \rightarrow \dots$

Iteration of Operator Rescaling

- ▶ **Three Key Facts Ensuring Convergence:**

Iteration of Operator Rescaling

► **Three Key Facts Ensuring Convergence:**

(1) $\text{BL}(B_i, \rho) \geq 1$ for all $i \geq 1$.

Iteration of Operator Rescaling

► Three Key Facts Ensuring Convergence:

(1) $\text{BL}(B_i, p) \geq 1$ for all $i \geq 1$.

(2) :

- If $1 + \varepsilon \leq \text{BL}(B_i, p) \leq 2$,
 $\text{BL}(B_{i+1}, p) \leq (1 - C_1 \varepsilon^k) \text{BL}(B_i, p)$.
- If $\text{BL}(B_i, p) > 2$, then $\text{BL}(B_{i+1}, p) \leq (1 - C_2) \text{BL}(B_i, p)$.

Iteration of Operator Rescaling

► Three Key Facts Ensuring Convergence:

(1) $\text{BL}(B_i, p) \geq 1$ for all $i \geq 1$.

(2) :

► If $1 + \varepsilon \leq \text{BL}(B_i, p) \leq 2$,
 $\text{BL}(B_{i+1}, p) \leq (1 - C_1 \varepsilon^k) \text{BL}(B_i, p)$.

► If $\text{BL}(B_i, p) > 2$, then $\text{BL}(B_{i+1}, p) \leq (1 - C_2) \text{BL}(B_i, p)$.

(3) If isotropy or projection holds, and $\text{BL}(B_i, p) \leq 2$, then

$$\left\| \sum p_i B_i^* B_i - I \right\| + \sum_i \|B_i^* B_i - I\| \lesssim \log(\text{BL}(B, p)).$$

Iteration of Operator Rescaling

► Three Key Facts Ensuring Convergence:

(1) $\text{BL}(B_i, p) \geq 1$ for all $i \geq 1$.

(2) :

► If $1 + \varepsilon \leq \text{BL}(B_i, p) \leq 2$,
 $\text{BL}(B_{i+1}, p) \leq (1 - C_1 \varepsilon^k) \text{BL}(B_i, p)$.

► If $\text{BL}(B_i, p) > 2$, then $\text{BL}(B_{i+1}, p) \leq (1 - C_2) \text{BL}(B_i, p)$.

(3) If isotropy or projection holds, and $\text{BL}(B_i, p) \leq 2$, then

$$\left\| \sum p_i B_i^* B_i - I \right\| + \sum_i \|B_i^* B_i - I\| \lesssim \log(\text{BL}(B, p)).$$

► Thus convergence to the family of geometric Brascamp-Lieb datum occurs as with Sinkhorn iteration provided that $\text{BL}(B, p) < \infty$.

Iteration of Operator Rescaling

► Three Key Facts Ensuring Convergence:

(1) $\text{BL}(B_i, \rho) \geq 1$ for all $i \geq 1$.

(2) :

► If $1 + \varepsilon \leq \text{BL}(B_i, \rho) \leq 2$,
 $\text{BL}(B_{i+1}, \rho) \leq (1 - C_1 \varepsilon^k) \text{BL}(B_i, \rho)$.

► If $\text{BL}(B_i, \rho) > 2$, then $\text{BL}(B_{i+1}, \rho) \leq (1 - C_2) \text{BL}(B_i, \rho)$.

(3) If isotropy or projection holds, and $\text{BL}(B_i, \rho) \leq 2$, then

$$\left\| \sum p_i B_i^* B_i - I \right\| + \sum_i \|B_i^* B_i - I\| \lesssim \log(\text{BL}(B, \rho)).$$

► Thus convergence to the family of geometric Brascamp-Lieb datum occurs as with Sinkhorn iteration provided that $\text{BL}(B, \rho) < \infty$.

► Obtain (1), (2), and (3) by studying *positive operators*.

Another Viewpoint: Positive Operators

- ▶ For simplicity, assume that all spaces have the same ambient dimension (all B_i are square matrices).

Another Viewpoint: Positive Operators

- ▶ For simplicity, assume that all spaces have the same ambient dimension (all B_i are square matrices).
- ▶ $\text{BL}(B, p) < \infty$ can only hold if $\sum p_i = 1$.

Another Viewpoint: Positive Operators

- ▶ For simplicity, assume that all spaces have the same ambient dimension (all B_i are square matrices).
- ▶ $\text{BL}(B, p) < \infty$ can only hold if $\sum p_i = 1$.
- ▶ Also assume all A_i are equal, and let us consider optimizing the quantity

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

analogous to

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

Positive Operators

- ▶ A linear map $T : M_n \rightarrow M_m$ is *completely positive* if there are $m \times n$ matrices B_1, \dots, B_K such that

$$T(A) = \sum B_i A B_i^*.$$

For simplicity, focus on the case $n = m$.

Positive Operators

- ▶ A linear map $T : M_n \rightarrow M_m$ is *completely positive* if there are $m \times n$ matrices B_1, \dots, B_K such that

$$T(A) = \sum B_i A B_i^*.$$

For simplicity, focus on the case $n = m$.

- ▶ Important example: $T(A) = \sum p_i B_i^* A B_i$.

Positive Operators

- ▶ A linear map $T : M_n \rightarrow M_m$ is *completely positive* if there are $m \times n$ matrices B_1, \dots, B_K such that

$$T(A) = \sum B_i A B_i^*.$$

For simplicity, focus on the case $n = m$.

- ▶ Important example: $T(A) = \sum p_i B_i^* A B_i$.
- ▶ Given T , we have $T^*(A) = \sum B_i^* A B_i$.

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- The *capacity* of a positive operator $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- ▶ The *capacity* of a positive operator $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

- ▶ Capacity introduced in (Gurvits, 2004) in $n = m$ case, in general introduced in (Garg et al, 2018).

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- ▶ The *capacity* of a positive operator $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

- ▶ Capacity introduced in (Gurvits, 2004) in $n = m$ case, in general introduced in (Garg et al, 2018).
- ▶ For any Brascamp-Lieb data (B, p) , there exists a positive $T : M_n \rightarrow M_m$ such that $\text{Cap}(T) = 1/\text{BL}(B, p)^2$.

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- ▶ The *capacity* of a positive operator $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

- ▶ Capacity introduced in (Gurvits, 2004) in $n = m$ case, in general introduced in (Garg et al, 2018).
- ▶ For any Brascamp-Lieb data (B, p) , there exists a positive $T : M_n \rightarrow M_m$ such that $\text{Cap}(T) = 1/\text{BL}(B, p)^2$.
- ▶ Positive operators are well studied in the quantum information theory literature, so reduction of BL to this theory gives new insights.

Doubly Stochastic Positive Operators

- ▶ (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.

Doubly Stochastic Positive Operators

- ▶ (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - ▶ $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.

Doubly Stochastic Positive Operators

- ▶ (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - ▶ $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.
- ▶ If (B, p) is a Brascamp-Lieb datum with associated operator $T : M_n \rightarrow M_m$, then (B, p) is geometric if and only if T is *doubly stochastic*. For $n = m$ this means $T(I) = I$ and $T^*(I) = I$.

Operator Rescaling for Positive Operators

- ▶ If T is doubly stochastic, $\text{Cap}(T) = 1$.

Operator Rescaling for Positive Operators

- ▶ If T is doubly stochastic, $\text{Cap}(T) = 1$.
- ▶ We can rescale. If

$$T_{M_1 M_2}(A) = M_2^* T(M_1^* A M_1) M_2,$$

then $\text{Cap}(T_{M_1, M_2}) = \det(M_1)^2 \det(M_2)^2 \cdot \text{Cap}(T)$.

Operator Rescaling for Positive Operators

- ▶ If T is doubly stochastic, $\text{Cap}(T) = 1$.
- ▶ We can rescale. If

$$T_{M_1 M_2}(A) = M_2^* T(M_1^* A M_1) M_2,$$

then $\text{Cap}(T_{M_1 M_2}) = \det(M_1)^2 \det(M_2)^2 \cdot \text{Cap}(T)$.

- ▶ Sinkhorn iteration (alternately iterating $T \mapsto T_{I, T(I)^{-1/2}}$ and $T \mapsto T_{T^*(I)^{-1/2}, I}$) yields a method for rescaling any T with $\text{Cap}(T) > 0$ to be arbitrarily close to a doubly stochastic operator, allowing us to approximate $\text{Cap}(T)$.

Rank Decreasing Operators

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

Rank Decreasing Operators

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbf{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

Rank Decreasing Operators

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbf{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

- ▶ An operator $T : M_n \rightarrow M_n$ is *rank non-decreasing* if for any $A \succeq 0$, $\text{Rank}(TA) \geq \text{Rank}(A)$.

Rank Decreasing Operators

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbf{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

- ▶ An operator $T : M_n \rightarrow M_n$ is *rank non-decreasing* if for any $A \succeq 0$, $\text{Rank}(TA) \geq \text{Rank}(A)$.
- ▶ (Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

Rank Decreasing Operators

$$\int_{\mathbf{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbf{R}^{n_i})}^{p_i}.$$

- ▶ (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbf{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

- ▶ An operator $T : M_n \rightarrow M_n$ is *rank non-decreasing* if for any $A \succeq 0$, $\text{Rank}(TA) \geq \text{Rank}(A)$.
- ▶ (Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.
- ▶ Generalized in (Garg et al, 2018). For $T : M_n \rightarrow M_m$, $\text{Cap}(T) > 0$ if and only if T is *fractional rank non-decreasing*.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ij} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ij} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ij} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- ▶ Define $T_U = D_U \circ T$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- ▶ Define $T_U = D_U \circ T$.
- ▶ Result follows from the following two facts:

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- ▶ Define $T_U = D_U \circ T$.
- ▶ Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- ▶ Define $T_U = D_U \circ T$.
- ▶ Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .
 - (2) $\text{Cap}(T) = \inf_U \text{Cap}(T_U)$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- ▶ Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- ▶ Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- ▶ Define $T_U = D_U \circ T$.
- ▶ Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .
 - (2) $\text{Cap}(T) = \inf_U \text{Cap}(T_U)$.
- ▶ To prove (1) and (2), use a simple trick: Given $A \succeq 0$, find U diagonalizing $T(A)$. Then $T(A) = T_U(A)$.

Thanks For Listening!