

Harmonic Analysis

Jacob Denson

October 11, 2022

Table Of Contents

I	Classical Fourier Analysis	2
1	Introduction	4
1.1	Obtaining the Fourier Coefficients	5
1.2	Orthogonality	5
1.3	The Fourier Transform	7
1.4	Multidimensional Theory	8
1.5	Examples of Expansions	9
2	Fourier Series	12
2.1	Basic Properties of Fourier Series	13
2.2	Unique Representation of a Function?	15
2.3	Quantitative Bounds on Fourier Coefficients	17
2.4	Boundedness of Partial Sums	21
2.5	Asymptotic Decay of Fourier Series	23
2.6	Smoothness and Decay	24
2.7	Convolution and Kernel Methods	27
2.8	The Dirichlet Kernel	33
2.9	Countercultural Methods of Summation	35
2.10	Fejer Summation	37
2.11	Abel Summation	38
2.12	The De la Vallée Poisson Kernel	40
2.13	Pointwise Convergence	41
2.14	Gibbs Phenomenon	45
3	Applications of Fourier Series	48
3.1	Tchebychev Polynomials	48
3.2	Exponential Sums and Equidistribution	50
3.3	The Isoperimetric Inequality	51

3.4	The Poisson Equation	51
3.5	The Heat Equation on a Torus	53
4	The Fourier Transform	56
4.1	Basic Calculations	57
4.2	The Fourier Algebra	61
4.3	Basic Convergence Properties	63
4.4	Alternative Summation Methods	65
4.5	The L^2 Theory	80
4.6	The Hausdorff-Young Inequality	83
4.7	The Poisson Summation Formula	86
4.8	Convergence of Fourier Series via Poisson Summation . .	87
4.9	Radial Functions	88
4.10	Poisson Integrals	88
5	Applications of the Fourier Transform	89
5.1	Applications to Partial Differential Equations	89
5.2	Shannon-Nyquist Sampling Theorem	90
5.3	The Uncertainty Principle	93
5.3.1	The Heisenberg Uncertainty Principle	94
5.3.2	Bernstein's Inequality	97
5.4	Sums of Random Variables	100
5.5	The Wirtinger Inequality on an Interval	103
5.6	Energy Preservation in the String equation	103
5.7	Harmonic Functions	103
6	Partial Derivatives and Harmonic Functions	105
6.1	Conjugate Poisson Kernel	105
7	Finite Character Theory	106
7.1	Fourier Analysis on Cyclic Groups	107
7.2	An Arbitrary Finite Abelian Group	109
7.3	Convolutions	111
7.4	The Fast Fourier Transform	113
7.5	Dirichlet's Theorem	114

8	Complex Methods	116
8.1	Fourier Transforms of Holomorphic Functions	116
8.2	Classical Theorems by Contours	120
8.3	The Laplace Transform	122
8.4	Asymptotics via the Laplace Transform	126
II	Distributional Methods	128
9	The Theory of Distributions	129
9.1	Distributions	130
9.2	Test Functions and Distributions	132
9.3	Topologies on the Space of Distributions	146
9.4	Homogeneous Distributions	149
9.5	Localization of Distributions	152
9.6	Derivatives of Continuous Functions	159
9.7	Convolutions of Distributions	160
9.8	Schwartz Space and Tempered Distributions	167
9.9	Kernel Operators	180
9.10	Test Functions on a Manifold	185
9.11	Paley-Wiener Theorem	187
9.12	Hyperfunctions	189
10	Microlocal Analysis of Singularities	190
10.1	Oscillatory Integral Distributions	198
10.2	Singular Operations on Distributions	205
10.3	Propagation of Singularities	216
11	Distributional Methods to PDEs	219
11.1	Fundamental Solutions	223
11.2	The Cauchy Problem	242
11.2.1	First Order Operators	242
11.2.2	Calderon's Uniqueness Theorem	243
12	Symbol Classes	247
13	Pseudodifferential Operators	251
13.1	Basic Definitions	257
13.2	Compound Operators and Quantization	266

13.3	Compositions of Ψ DOs	270
13.4	Parametrices for Elliptic Operators	273
13.5	Regularity Theory	275
13.6	Pseudodifferential Operators on Manifolds	280
13.7	Ψ DOs and Microsupport	289
13.8	Vector-Valued Pseudodifferential Operators	292
13.9	The Index Theorem	295
13.10	Self-Adjoint Elliptic Pseudo-Differential Operators on a Compact Manifold	297
13.11	Distribution of Eigenvalues and the Half Wave Operator .	303
13.12	Discrete Restriction Theory	316
14	Fourier Integral Operators	323
14.1	Hyperbolic Equations	326
14.2	Local Symplectic Geometry	329
14.3	Local Theory	334
III	Calderon-Zygmund Theory	340
15	Monotone Rearrangement Invariant Norms	344
15.1	The L^p norms	345
15.2	Decreasing Rearrangements	356
15.3	Weak Norms	357
15.4	Lorentz Spaces	365
15.5	Mixed Norm Spaces	378
15.6	Orlicz Spaces	381
16	Sobolev Spaces	386
16.1	Hölder Spaces	387
17	Sobolev Spaces	390
17.1	Smoothing	392
18	Interpolation Theory	394
18.1	Interpolation of Functions	394
18.2	Complex Interpolation	395
18.3	Interpolation of Operators	398
18.4	Complex Interpolation of Operators	399

18.5	Real Interpolation of Operators	404
19	Basic Integral Operator Estimates	409
19.1	Schur's Lemma	411
19.2	Change of Variables	419
19.3	Localization In Space	419
19.4	The Christ-Kiselev Lemma	422
20	Maximal Averages	427
20.1	The Method of Maximal Functions	428
20.2	Covering Methods	430
20.3	Dyadic Methods	436
20.4	Linearization of Maximal Operators	440
20.5	TT^* Arguments	441
20.6	Lebesgue Density Theorem	442
20.7	Ergodic Averages	444
20.8	Approximations to the Identity	447
20.9	H^p Spaces	451
20.10	The Strong Maximal Function	452
20.11	The Tangential Poisson Maximal Function	452
21	Differentiability of Measurable Functions	453
21.1	Absolute Continuity	460
21.2	Differentiability of Jump Functions	467
21.3	Rectifiable Curves	470
21.4	Bounded Variation in Higher Dimensions	471
21.5	Minkowski Content	473
21.6	The Isoperimetric Inequality	476
22	Singular Integral Operators	480
22.1	The Calderon-Zygmund Decomposition	480
23	Fourier Multiplier Operators	486
23.1	Frequency Localization	489
23.2	L^p Regularity	491
23.3	Transference	507
24	Sobolev Spaces	514

25 Time Frequency Analysis	515
25.1 Localization in Time and Space	517
25.2 Convergence in L^p and the Hilbert Transform	517
25.3 A Divergent Fourier Series	521
25.4 Conjugate Fourier Series	522
26 Higher Dimensional Fourier Analysis	523
27 Oscillatory Integrals	527
27.1 One Dimensional Theory	528
27.2 Stationary Phase in Multiple Variables	548
27.3 Variable Coefficient Results	550
27.4 Surface Carried Measures	552
27.5 Oscillatory Integral Operators	558
28 Restriction Theorems	568
28.1 Stein-Tomas Theorem	574
28.2 Bochner-Riesz Multipliers	578
28.3 Restriction on the Paraboloid	580
28.4 Restriction to the Cone	582
28.5 TODO: Hardy Littlewood Majorant Conjecture	582
29 Almost Orthogonality	583
30 Weighted Estimates	589
30.1 Hardy-Littlewood Maximal Function	589
31 Bellman Function Methods	592
32 Maximal Averages Over Curves	596
32.1 Averages over a Parabola	596
IV Linear Partial Differential Equations	598
33 Green's Functions	600
34 Elliptic Equations	602

35 Dispersive Equations	603
35.1 Strichartz Estimates	606
 V Geometry of Eigenfunctions	 608
36 Self-Adjoint, Elliptic Operators on a Compact Manifold	611
37 Distribution of Eigenvalues and the Half Wave Operator	619
38 Nodal Sets	633
38.1 Nodal Sets for Open Subsets of \mathbf{R}^d	635
38.2 Nodal Sets on the Sphere	636
38.3 Brownian Motion and Nodal Sets	638
38.3.1 Probabilistic Tools	638
38.4 Nodal Sets Via Brownian Motion	640
 VI Harmonic Analysis and Fractals	 642
39 Riemann's Theory of Trigonometric Series	643
 VII Abstract Harmonic Analysis	 645
40 Topological Groups	647
40.1 Basic Results	648
40.2 Quotient Groups	650
40.3 Uniform Continuity	651
40.4 Ordered Groups	653
40.5 Topological Groups arising from Normal subgroups . . .	655
 41 The Haar Measure	 657
41.1 Fubini, Radon Nikodym, and Duality	664
41.2 Unimodularity	665
41.3 Convolution	667
41.4 The Riesz Thorin Theorem	676
41.5 Homogenous Spaces and Haar Measures	676

42 The Character Space	680
43 Banach Algebra Techniques	688
44 Vector Spaces	689
45 Interpolation of Besov and Sobolev spaces	690
45.1 Besov Spaces	693
45.2 Proof of The Projection Result	693
46 The Cap Set Problem	694
 VIII Decoupling	 697
47 The General Framework	699
47.1 Localized Estimates	705
47.2 Vinogradov Systems	706

Part I

Classical Fourier Analysis

Deep mathematical knowledge often arises hand in hand with the characterization of symmetry. Nowhere is this more clear than in the foundations of harmonic analysis, where we attempt to understand mathematical ‘signals’ by the ‘frequencies’ from which they are composed. In the mid 18th century, problems in mathematical physics led D. Bernoulli, D’Alembert, Lagrange, and Euler to consider periodic functions representable as a trigonometric series

$$f(t) = A + \sum_{m=1}^{\infty} B_m \cos(2\pi mt) + C_m \sin(2\pi mt).$$

In his book, *Théorie Analytique de la Chaleur*, published in 1811, Joseph Fourier had the audacity to announce that *all* functions were representable in this form, and used it to solve linear partial differential equations in physics. His conviction is the reason the classical theory of harmonic analysis is often named Fourier analysis, where we analyze the degree to which Fourier’s proclamation holds, as well as its paired statement on the real line, that a function f on the real line can be written as

$$f(t) = \int_{-\infty}^{\infty} A(\xi) \cos(2\pi \xi t) + B(\xi) \sin(2\pi \xi t) d\xi.$$

for some functions A and B on the line.

In the 1820s, Poisson, Cauchy, and Dirichlet all attempted to form rigorous proofs that ‘Fourier summation’ holds for all functions. Their work is responsible for most of the modern subject of analysis we know today. In particular, it is essential to utilize all the convergence techniques developed through the rigorous study of analysis. Under pointwise convergence, the representation of a function by Fourier series need not be unique. Uniform convergence is more useful, and uniform convergence holds for all smooth functions, but does not hold if we only assume a function is continuous. Thus we must introduce more subtle methods.

Chapter 1

Introduction

One fundamental family of oscillatory functions in mathematics are the trigonometric functions

$$f(t) = A \cos(st) + B \sin(st) = C \cos(st + \phi).$$

The value ϕ is the *phase* of the oscillation, C is the *amplitude*, and $s/2\pi$ is the *frequency* of the oscillation. These oscillatory functions occur in many situations; for instance, in the study of the solution of the harmonic oscillator. The main topic of Fourier analysis is to study how well one may represent a general function as an analytical combination of these trigonometric functions. In the periodic setting, we fix a function $f : \mathbf{R} \rightarrow \mathbf{C}$ such that $f(x+1) = f(x)$ for all $x \in \mathbf{R}$, and try and find coefficients $\{A_m\}$, $\{B_m\}$, and C such that

$$f(t) \sim C + \sum_{m=1}^{\infty} A_m \cos(2\pi m t) + B_m \sin(2\pi m t).$$

In the continuous setting, we fix a function $f : \mathbf{R} \rightarrow \mathbf{C}$, trying to find values $A(s)$, $B(s)$, and C such that

$$f(t) \sim C + \int_0^{\infty} A(s) \cos(2\pi s t) + B(s) \sin(2\pi s t) ds.$$

The main contribution of Fourier was a method to formally find a reliable choice of coefficients which represents f . This choice is given by the *Fourier transform* of f in the continuous case, and the *Fourier series* in the discrete case.

1.1 Obtaining the Fourier Coefficients

A *formal trigonometric series* is a formal sum of the form

$$C + \sum_{m=1}^{\infty} A_m \cos(2\pi mt) + B_m \sin(2\pi mt).$$

Our goal, given a function f , is to find a family $\{A_m\}$, $\{B_m\}$, and C which ‘represents’ the function f . In particular, we say a periodic function f *admits a trigonometric expansion* if there is a series such that for each $t \in \mathbf{R}$,

$$f(t) = C + \sum_{m=1}^{\infty} A_m \cos(2\pi mt) + B_m \sin(2\pi mt).$$

It is a *very difficult question* to characterize which functions f admit a trigonometric expansion. Nonetheless, Fourier found a way to formally associate a formal trigonometric series with any integrable periodic function. If the function is differentiable, then the trigonometric series gives a trigonometric expansion for the function. But even if this series does not give a trigonometric expansion for this function, the series itself still reflects many important properties of the function, which are of interest independent of their convergence to the function f .

1.2 Orthogonality

The key technique Fourier realized could be used to come up with a canonical trigonometric series for a function is *orthogonality*. Note that the various frequencies of sine functions are orthogonal to one another, in the sense that

$$\int_0^1 \sin(2\pi mt) \sin(2\pi nt) = \int_0^1 \cos(2\pi mt) \cos(2\pi nt) = \begin{cases} 0 & : m \neq n, \\ 1/2 & : m = n, \end{cases}$$

and for any $m, n \in \mathbf{Z}$,

$$\int_0^1 \sin(2\pi mt) \cos(2\pi nt) = 0.$$

This means that for a finite trigonometric sum

$$f(t) = C + \sum_{m=1}^N A_m \cos(2\pi mt) + B_m \sin(2\pi mt),$$

we have

$$C = \int_0^1 f(t) dt,$$

$$A_m = 2 \int_0^1 f(t) \cos(2\pi mt) dt, \quad \text{and} \quad B_m = 2 \int_{-\pi}^{\pi} f(t) \sin(2\pi mt) dt.$$

We note that these values may still be defined even if f is not a trigonometric polynomial. Thus given *any* periodic integrable function f , a reasonable candidate for the coefficients is given by the values A_m , B_m , and C above. Unlike when f is a trigonometric polynomial, we can have infinitely many non-zero coefficients.

There is an additional choice of oscillatory functions, which replaces the sine and cosine with a single family of trigonometric functions, and thus gives a more notationally convenient analysis. For $\xi, t \in \mathbf{R}$, we let $e_\xi(t) = e^{2\pi i \xi t}$. For each integer $n \in \mathbf{Z}$, e_n is periodic with period 1. Applying orthogonality again, we find

$$\int_0^1 e_n(t) \overline{e_m(t)} dt = \int_0^1 e_{n-m}(t) dt = \begin{cases} 0 & : m \neq n, \\ 1 & : m = n. \end{cases}$$

Thus we can use orthogonality to find a natural choice of an expansion

$$f(t) \sim \sum_{n \in \mathbf{Z}} C_n e^{2\pi i n t},$$

given by setting

$$C_n = \int_0^1 f(t) \overline{e_n(t)} dt = \int_0^1 f(t) e^{-2\pi i n t} dt.$$

Euler's formula $e^{nit} = \cos(nt) + i \sin(nt)$ shows this is the same as the Fourier expansion in sines and cosines. Thus the values $\{A_m, B_m, C : m \geq 0\}$ can be recovered from the values of $\{C_m : m \in \mathbf{Z}\}$. Because of its elegance,

unifying the three families of coefficients, the expansion by complex exponentials is the most standard used in Fourier analysis today.

To summarize, we have shown a periodic integrable function $f : \mathbf{R} \rightarrow \mathbf{C}$ gives rise to a formal trigonometric series

$$\sum_{m \in \mathbf{Z}} C_m e_m(t).$$

This is the *Fourier series* of f . Because we will be concentrating on the Fourier series of a function, it is worth reserving a particular notation. Given a periodic, integrable function f , and an integer $m \in \mathbf{Z}$, we set

$$\hat{f}(m) = \int_0^1 f(t) \overline{e_m(t)} dt.$$

The Fourier series representation in terms of complex exponentials will be our choice throughout the rest of these notes. No deep knowledge of the complex numbers is used here. For most basic purposes, the exponential notation is just a simple way to represent the oscillations of sines and cosines in a unified manner.

1.3 The Fourier Transform

For a general function $f : \mathbf{R} \rightarrow \mathbf{C}$, we cannot rely *just* on orthogonality, because the functions $\sin(2\pi mx)$ are not integrable on the entirety of \mathbf{R} , and therefore cannot be integrated against one another. Nonetheless, we can consider the functions $g_N : [0, 1] \rightarrow \mathbf{C}$ by setting $g_N(s) = f(N(s - 1/2))$. Then for $|t| \leq N/2$, we can apply the usual Fourier series to conclude

$$\begin{aligned} f(t) &= g_N(t/N + 1/2) \\ &\sim \sum_{m \in \mathbf{Z}} \widehat{g_N}(m) e^{2\pi m i(t/N + 1/2)} \\ &= \sum_{m \in \mathbf{Z}} (-1)^m \left(\int_0^1 f(N(s - 1/2)) e^{-2\pi m i s} ds \right) e^{2\pi(m/N)it} \\ &= \sum_{m \in \mathbf{Z}} \frac{1}{N} \left(\int_{-N/2}^{N/2} f(s) e^{-2\pi(m/N)is} ds \right) e^{(m/N)it}. \end{aligned}$$

If we take $N \rightarrow \infty$, the exterior sum operates like a Riemann sum, so we might expect

$$f(t) \sim \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} f(s) e^{-2\pi \xi i s} ds \right) e^{2\pi \xi i t} d\xi.$$

The interior integral defines the *Fourier transform* of the function f , given for each $\xi \in \mathbf{R}$ as

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(s) e^{-2\pi \xi i s} ds.$$

Thus the resultant *Fourier inversion formula* takes the form

$$f(t) \sim \int_{-\infty}^{\infty} \hat{f}(\xi) e_{\xi}(t) d\xi.$$

As the *limit* of a discrete series defined in terms of orthogonality, the Fourier transform possesses many of the same properties at the Fourier series. But the non-compactness causes issues which are not present in the case of Fourier series, and so the Fourier series theory is often a simpler theory to begin with.

1.4 Multidimensional Theory

Finally, we note that the Fourier series and Fourier transform are not relegated to a one dimensional theory. If $f : \mathbf{R}^d \rightarrow \mathbf{C}$ is periodic, in the sense that $f(x + n) = f(x)$ for each $x \in \mathbf{R}^d$ and $n \in \mathbf{Z}^d$, then we can consider the natural higher dimensional Fourier series

$$f(t) \sim \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e_n(t)$$

where for each $\xi \in \mathbf{R}^d$, $e_{\xi} : \mathbf{R}^d \rightarrow \mathbf{C}$ is the function given for each $t \in \mathbf{R}^d$ by setting $e_{\xi}(t) = e^{2\pi i \xi \cdot t}$, and

$$\hat{f}(n) = \int_{[0,1]^d} f(t) \overline{e_n(t)} dt$$

Similarly, for $f : \mathbf{R}^d \rightarrow \mathbf{C}$, we can consider the Fourier inversion formula

$$f(t) \sim \int_{\mathbf{R}^d} \hat{f}(\xi) e_{\xi}(x) d\xi$$

where for each $\xi \in \mathbf{R}^d$,

$$\hat{f}(\xi) = \int_{\mathbf{R}^d} f(t) \overline{e_{\xi}(t)} dt$$

The basic theory of Fourier series and the Fourier transform in one dimension extends naturally to higher dimensions, as do the basic theories of orthogonality. On the other hand, the theory of convergence in higher dimensions requires much greater regularity in higher dimensions and many fundamental questions about the convergence of Fourier series here more nuance than in the lower dimensional theory.

1.5 Examples of Expansions

Before we get to the real work, let's start by computing some examples of Fourier series and examples of the Fourier transform. We also illustrate the convergence properties of these series, which we shall look at in more detail later.

Example. Consider the function $f : [0, \pi] \rightarrow \mathbf{R}$ defined by $f(x) = x(\pi - x)$. Then a series of integration by parts gives that

$$\int x(\pi - x) \sin(nx) = \frac{x(\pi - x) \cos(nx)}{n} + \frac{(\pi - 2x) \sin(nx)}{n^2} - \frac{2 \cos(nx)}{n^3}.$$

Thus

$$\frac{2}{\pi} \int_0^{\pi} x(\pi - x) \sin(nx) = \frac{4(1 - \cos(n\pi))}{n^3} = \begin{cases} \frac{8}{\pi n^3} & n \text{ odd}, \\ 0 & n \text{ even}. \end{cases}$$

Thus we have a formal representation

$$f(x) \sim \sum_{n \text{ odd}} \frac{8 \sin(nx)}{\pi n^3}.$$

This sum converges absolutely and uniformly for $x \in [0, \pi]$. If we extend the domain of f to $[-\pi, \pi]$ by making f odd, then

$$\hat{f}(n) = \begin{cases} \frac{4}{\pi i n^3} & : n \text{ odd}, \\ 0 & : n \text{ even}. \end{cases}$$

In this case, we still have

$$f(x) \sim \sum_{\substack{n \text{ odd} \\ n > 0}} \frac{4}{\pi i n^3} [e_n(x) - e_n(-x)] = \sum_{n \text{ odd}} \frac{8 \sin(nx)}{\pi n^3}.$$

This sum converges absolutely and uniformly on the entire real line.

Example. The tent function

$$f(x) = \begin{cases} 1 - \frac{|x|}{\delta} & : |x| < \delta, \\ 0 & : |x| \geq \delta. \end{cases}$$

is even, and therefore has a purely real Fourier expansion

$$\hat{f}(0) = \frac{\delta}{2\pi}, \quad \hat{f}(n) = \frac{1 - \cos(n\delta)}{\delta \pi n^2}.$$

Thus we obtain an expansion

$$f(x) = \frac{\delta}{2\pi} + \sum_{n \neq 0} \frac{1 - \cos(n\delta)}{\delta \pi n^2} e_n(x) = \frac{\delta}{2\pi} + 2 \sum_{n=1}^{\infty} \frac{1 - \cos(n\delta)}{\delta \pi n^2} \cos(nx).$$

This sum also converges absolutely and uniformly on the entire real line.

Example. Consider the characteristic function

$$\chi_{(a,b)}(x) = \begin{cases} 1 & : x \in (a,b), \\ 0 & : x \notin (a,b). \end{cases}$$

Then

$$\hat{\chi}_{(a,b)}(n) = \frac{1}{2\pi} \int_a^b e_n(-x) = \frac{e_n(-a) - e_n(-b)}{2\pi i n}.$$

Hence we may write

$$\begin{aligned} \chi_{(a,b)}(x) &= \frac{b-a}{2\pi} + \sum_{n \neq 0} \frac{e_n(-a) - e_n(-b)}{2\pi i n} e_n(x) \\ &= \frac{b-a}{2\pi} + \sum_{n=1}^{\infty} \frac{\sin(nb) - \sin(na)}{\pi n} \cos(nx) + \frac{\cos(na) - \cos(nb)}{\pi n} \sin(nx). \end{aligned}$$

This sum does not converge absolutely for any value of x (except when a and b are chosen trivially). To see this, note that

$$\left| \frac{e_n(-b) - e_n(-a)}{2\pi n} \right| = \left| \frac{1 - e_n(b-a)}{2\pi n} \right| \geq \left| \frac{\sin(n(b-a))}{2\pi n} \right|,$$

so that it suffices to show $\sum |\sin(nx)|n^{-1} = \infty$ for every $x \notin \pi\mathbf{Z}$. This follows because the values of $|\sin(nx)|$ are often large, so that we may apply the divergence of $\sum n^{-1}$. First, assume $x \in (0, \pi/2)$. If

$$m\pi - x/2 < nx < m\pi + x/2$$

for some $m \in \mathbf{Z}$, then

$$m\pi + x/2 < (n+1)x < m\pi + 3x/2 < (m+1)\pi - x/2.$$

Thus if $nx \in (-x/2, x/2) + \pi\mathbf{Z}$, $(n+1)x \notin (-x/2, x/2) + \pi\mathbf{Z}$. For y outside of $(-x/2, x/2) + \pi\mathbf{Z}$, we have $|\sin(y)| > |\sin(x/2)|$, and therefore for any n ,

$$\frac{|\sin(nx)|}{n} + \frac{|\sin((n+1)x)|}{n+1} > \frac{|\sin(x/2)|}{n+1}.$$

This means

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{|\sin(nx)|}{n} &= \sum_{n=1}^{\infty} \frac{|\sin(2nx)|}{2n} + \frac{|\sin((2n+1)x)|}{2n+1} \\ &> |\sin(x/2)| \sum_{n=1}^{\infty} \frac{1}{2n+1} = \infty \end{aligned}$$

In general, we may replace x with $x - k\pi$, with no effect to the values of the sum, so we may assume $0 < x < \pi$. If $\pi/2 < x < \pi$, then

$$\sin(nx) = \sin(n(\pi - x)),$$

and $0 < \pi - x < \pi/2$, completing the proof, except when $x = \pi$, in which case

$$\sum_{n=1}^{\infty} \left| \frac{1 - e_n(\pi)}{2\pi n} \right| = \sum_{n \text{ even}} \left| \frac{1}{\pi n} \right| = \infty.$$

Thus the convergence of a Fourier series need not be absolute.

Chapter 2

Fourier Series

Let us now focus on the theory of *Fourier series* we introduced in the last chapter. We write $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, so that a function $f : \mathbf{T} \rightarrow \mathbf{C}$ is a complex-valued periodic function on the real line. We then have a metric on \mathbf{T} given by setting $d(t, s) = |t - s|$, where $|t| = \min_{n \in \mathbf{Z}} |t + n|$ for $t \in \mathbf{T}$. The Lebesgue measure on \mathbf{R} induces a natural Borel measure on \mathbf{T} , such that for any periodic function $f : \mathbf{T} \rightarrow \mathbf{C}$,

$$\int_{\mathbf{T}} f(t) dt = \int_0^1 f(t) dt.$$

It will also be of interest to consider the higher dimensional torii $\mathbf{T}^d = \mathbf{R}^d / \mathbf{Z}^d$, which naturally has the induced product metric and measure from \mathbf{T} . For each $f \in L^1(\mathbf{T}^d)$, we associate the *formal trigonometric series*

$$\sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot t}$$

where for each $n \in \mathbf{Z}^d$,

$$\hat{f}(n) = \int_{\mathbf{T}^d} f(t) e^{-2\pi i n \cdot t} dt.$$

If $\{\hat{f}(n)\}$ is an absolutely summable sequence, then one can interpret the formal trigonometric series nonformally as an infinite series, and we would then hope that for each $t \in \mathbf{T}^d$,

$$f(t) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot t}.$$

It turns out that, under the assumption of absolute summability, this equation does hold provided f is a continuous function on \mathbf{T}^d . We will eventually see that the condition that the Fourier series of f is absolutely summable under the assumption that $f \in C^\infty(\mathbf{T}^d)$. We will be able to prove these facts immediately after we prove some basic symmetry properties of the Fourier series.

2.1 Basic Properties of Fourier Series

One of the most important properties of the Fourier series is that the coefficients are controlled by reasonable transformations. A basic, but unappreciated property of the Fourier transform is *linearity*: For any two functions f and g , if $h = f + g$, then $\widehat{h} = \widehat{f} + \widehat{g}$. Linearity is *essential* to most methods in this book; many problems about nonlinear transforms remain unsolved. The Fourier series is also stable under various transformations which occur in analysis, which makes the Fourier series tractable to analyze, and therefore useful. We summarize these properties here:

- Given $f \in L^1(\mathbf{T}^d)$, define $\text{Con}f, \text{Ref}f \in L^1(\mathbf{T}^d)$ by setting $\text{Con}f(x) = \overline{f(x)}$ and $\text{Ref}f(x) = f(-x)$. Then

$$\widehat{\text{Con}f} = (\text{Con} \circ \text{Ref})\widehat{f}.$$

and

$$\widehat{\text{Ref}f} = \text{Ref}\widehat{f}.$$

As a corollary, if f is real-valued, then

$$\widehat{f} = \widehat{\text{Con}f} = (\text{Con} \circ \text{Ref})\widehat{f}$$

In other words, for each $n \in \mathbf{Z}^d$,

$$\widehat{f}(n) = \overline{\widehat{f}(-n)}.$$

It also follows from the reflection symmetry that if $f \in L^1(\mathbf{T}^d)$ is odd, then \widehat{f} is odd, and if $f \in L^1(\mathbf{T}^d)$ is even, \widehat{f} is even.

- For each $s \in \mathbf{R}^d$, and $m \in \mathbf{Z}^d$, and any $f \in L^1(\mathbf{T}^d)$, define the translation and frequency modulation operators Trans_s and Mod_m by setting

$$(\text{Trans}_s f)(t) = f(t - s) \quad \text{and} \quad (\text{Mod}_m f)(t) = e_m(t) f(t).$$

Similarly, for each function $C : \mathbf{Z}^d \rightarrow \mathbf{C}$, for each $m \in \mathbf{Z}^d$ and $\xi \in \mathbf{R}$, define

$$(\text{Trans}_m C)(n) = C(n - m) \quad \text{and} \quad (\text{Mod}_\xi C)(n) = e_\xi(n) C(n).$$

Then for any $f \in L^1(\mathbf{T}^d)$, $\widehat{\text{Trans}_s f} = \text{Mod}_{-s} \hat{f}$, and $\widehat{\text{Mod}_m f} = \text{Trans}_{-m} \hat{f}$.

- An easy integration by parts shows that if $f \in C^\infty(\mathbf{T}^d)$, then for any $k \in \{1, \dots, d\}$,

$$\widehat{D^k f}(n) = 2\pi i n_k \hat{f}(n)$$

for each $n \in \mathbf{Z}^d$. The proof follows from an easy integration by parts, so the claim is actually true for any $f \in L^1(\mathbf{T}^d)$ with a weak derivative $D^k f$ in $L^1(\mathbf{T}^d)$. Iterating this argument shows that, assuming the required weak derivatives exist,

$$\widehat{D^\alpha f}(n) = (2\pi i n)^\alpha \hat{f}(n).$$

Remark. We note that if $f \in L^1(\mathbf{T})$ is even, then \hat{f} is even, so formally

$$f(t) \sim \hat{f}(0) + \sum_{m=1}^{\infty} \hat{f}(m) [e_m(t) + e_{-m}(t)] \sim \hat{f}(0) + 2 \sum_{m=1}^{\infty} \hat{f}(m) \cos(mt).$$

Moreover,

$$\hat{f}(m) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \cos(mt) dt$$

If f is an odd function, then the fact that \hat{f} is odd implies formally that

$$f(t) \sim \sum_{m=1}^{\infty} \hat{f}(m) [e_m(t) - e_{-m}(t)] = 2i \sum_{m=1}^{\infty} \hat{f}(m) \sin(mt).$$

Thus we get a sine expansion, and moreover,

$$\hat{f}(m) = \frac{1}{2\pi i} \int_{-\pi}^{\pi} f(t) \sin(mt) dt.$$

This is one way to reduce the study of complex exponentials back to the study of sines and cosines, since every function can be written as a sum of an even and an odd function.

2.2 Unique Representation of a Function?

To study the convergence properties of Fourier series, we begin by studying whether a function is uniquely determined by its Fourier coefficients, which would be certainly true if the Fourier series held. However, such a statement is clearly cannot be true for all $f \in L^1(\mathbf{T}^d)$, since the Fourier coefficients of a function depend only on the *distributional* properties of f , i.e. those that can be obtained through integration. In particular, if two integrable functions f and g agree on a set of measure zero, then they have the same Fourier coefficients depends only on the equivalence class of f in $L^1(\mathbf{T}^d)$, with functions identified if they are equal almost everywhere. Nonetheless, if $f \in C(\mathbf{T}^d)$ then there is no way to edit f on a set of measure zero while preserving continuity. Thus we can hope for unique Fourier coefficients in the setting of continuous functions.

Theorem 2.1. *Suppose $f \in L^1(\mathbf{T}^d)$. If $\hat{f}(n) = 0$ for all $n \in \mathbf{Z}^d$, then f vanishes at all its continuity points.*

Proof. It suffices to prove that if $f \in L^1(\mathbf{T}^d)$ is continuous at the origin, then $f(0) = 0$. We treat the real-valued case first. For every trigonometric polynomial $g(x) = \sum a_n e_n(-x)$, we have

$$\int_{\mathbf{T}} f(x)g(x)dx = \sum a_n \hat{f}(n) = 0.$$

Suppose that f is continuous at zero, and assume without loss of generality that $f(0) > 0$. Pick $\delta > 0$ such that if $|x| \leq \delta$, $f(x) > f(0)/2$. Consider the trigonometric polynomial

$$g(x) = \prod_{k=1}^d [\varepsilon + \cos(2\pi x_k)] = \prod_{k=1}^d \left[\varepsilon + \frac{e^{2\pi i x_k} + e^{-2\pi i x_k}}{2} \right],$$

and where $\varepsilon > 0$ is small enough that if $|x| \geq \delta$, then $g(x) \leq B < 1$. We can then choose $0 < \eta < \delta$ such that if $|x| < \eta$, $g(x) \geq A > 1$. Finally, if

δ is sufficiently small, we also have $g(x) > 0$ if $0 \leq |x| \leq \delta$. The series of trigonometric polynomials $g_n(x) = g(x)^n$ satisfy

$$\left| \int_{\mathbf{T}^d} g_n(x) f(x) dx \right| \geq \int_{|x| \leq \delta} g_n(x) f(x) dx - \left| \int_{|x| \geq \delta} g_n(x) f(x) dx \right|.$$

Hölder's inequality guarantees that as $n \rightarrow \infty$,

$$\left| \int_{|x| \geq \delta} g_n(x) f(x) dx \right| \lesssim B^n.$$

On the other hand,

$$\left| \int_{|x| \leq \delta} g_n(x) f(x) dx \right| \geq \int_{|x| < \delta/2} g_n(x) f(x) \gtrsim A^n.$$

Thus we conclude

$$0 = \left| \int_0^1 g_n(x) f(x) dx \right| \gtrsim A^n - B^n.$$

For suitably large values of n , the right hand side is positive, whereas the left hand side is zero, which is impossible. By contradiction, we conclude $f(0) = 0$. In general, if f is complex valued, then we may write $f = u + iv$, where

$$u(x) = \frac{f(x) + \overline{f(x)}}{2} \quad v(x) = \frac{f(x) - \overline{f(x)}}{2i}.$$

The Fourier coefficients of \overline{f} all vanish, because the coefficients of f vanish, and so we conclude the coefficients of u and v vanish. f is continuous at x if and only if u and v are continuous at x , so we can apply the real-valued case to complete the proof in the case of complex values. \square

Corollary 2.2. *If $f, g \in C(\mathbf{T}^d)$ and $\hat{f} = \hat{g}$, then $f = g$.*

Proof. Then $f - g$ is continuous with vanishing Fourier coefficients. \square

Corollary 2.3. *If $f \in C(\mathbf{T}^d)$ and $\hat{f} \in L^1(\mathbf{Z}^d)$, then for each $x \in \mathbf{Z}^d$,*

$$f(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}$$

Proof. Since $\hat{f} \in L^1(\mathbf{Z}^d)$, the sum

$$g(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}$$

converges *uniformly*. In particular, this implies that g is a continuous function. Moreover, it allows us to conclude that $\hat{g}(n) = \hat{f}(n)$ for each $n \in \mathbf{Z}$. But this means $f = g$. \square

In the next section we will show that if $f \in C^m(\mathbf{T}^d)$, then

$$\hat{f}(n) = O(\langle n \rangle^{-m}).$$

In particular, if $m \geq d + 1$, then the Fourier series of f is integrable. Moreover, if $f \in C^\infty(\mathbf{T})$, then using the Fourier series equation for the derivative of a function, for each multi-index α , we conclude that for all $x \in \mathbf{R}^d$

$$(D^\alpha f)(x) = \sum_{n \in \mathbf{Z}^d} (2\pi i n)^\alpha \hat{f}(n) e^{2\pi i n \cdot x}.$$

On the other hand, suppose $\{a_n : n \in \mathbf{Z}^d\}$ such that $|a_n| \lesssim_m \langle n \rangle^{-m}$ for all $m > 0$, then the infinite sum

$$\sum_{n \in \mathbf{Z}^d} a_n e^{2\pi i n \cdot x}$$

and all its derivatives converge uniformly to an infinitely differentiable function with the Fourier coefficients $\{a_n\}$. Thus there is a perfect duality between infinitely differentiable functions and arbitrarily fast decaying sequences of integers. In more advanced contexts, like distribution theory, this duality is very useful for studying the Fourier transform in a much more general setting.

2.3 Quantitative Bounds on Fourier Coefficients

There are various reasons why one would not be completely satisfied by the convergence result above. Unlike with the case of a Taylor series, the Fourier series can be applied to a much more general family of situations. There is no hope of the Fourier series being integrable *and* obtaining a

pointwise convergence result unless we are dealing with continuous functions, because any absolutely summable trigonometric series sums up to a continuous function. Thus we must analyze non-integrable families of coefficients if we are to obtain deeper convergence properties of the Fourier series for non-continuous functions.

On the other hand, in practical contexts, one might argue that the functions dealt with can be assumed arbitrarily smooth, so the picture established in the last section seems rather complete. However, even if this is true it is still important to study more *qualitative questions* about the Fourier series. Instead of taking the infinite Fourier series, we take a finite sum. For a function $f \in L^1(\mathbf{T})$, it is most natural to consider the partial sums

$$S_N f(x) = \sum_{n=-N}^N \hat{f}(n) e^{2\pi i n x}.$$

In higher dimensions, no canonical ‘cutoff’ exists. Two possible options are *spherical summation*

$$\sum_{|n| \leq N} \hat{f}(n) e_n$$

and *square summation*

$$\sum_{n_1, \dots, n_d = -N}^N \hat{f}(n) e_n.$$

There are subtle differences in these operators which cause problems in the higher dimensional theory. Right now, all we assume is that we are consider an increasing family of sets $\{E_N\}$ in \mathbf{Z}^d with $\lim_{N \rightarrow \infty} E_N = \mathbf{Z}^d$, and we then define

$$S_N f = \sum_{n \in E_N} \hat{f}(n) e_n$$

for $f \in L^1(\mathbf{T}^d)$. A natural question now is whether $S_N f$ is *qualitatively similar* to the function f globally rather than just pointwise. The most natural way to measure how similar two functions are from the perspective of analysis is via measuring the differences with respect to a suitable *norm*. For instance, under the assumptions of the last section, we not only get pointwise convergence at each point, but *uniform convergence*.

Theorem 2.4. Suppose $f \in C(\mathbf{T})$ and $\hat{f} \in L^1(\mathbf{Z})$. Then

$$\lim_{N \rightarrow \infty} \|S_N f - f\|_{L^\infty(\mathbf{R})}.$$

In other words, $S_N f$ converges uniformly to f instead of pointwise.

Proof. We know that for each $x \in \mathbf{T}^d$,

$$f(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}.$$

A simple application of the triangle inequality shows that

$$|f(x) - S_N f(x)| \leq \sum_{n \notin E_N} |\hat{f}(n)|.$$

Since the Fourier coefficients are absolutely summable, for each $\varepsilon > 0$, there is N_0 such that for $N \geq N_0$,

$$\sum_{n \notin E_N} |\hat{f}(n)| \leq \varepsilon,$$

and thus $\|f - S_N f\|_{L^\infty(\mathbf{T}^d)} \leq \varepsilon$. □

Another question one might ask is the *rate of convergence* of the function f . In this situation, things are quite bad even in the setting of the previous setting. For general elements of $C(\mathbf{T})$ with integrable Fourier coefficients, the convergence of $\|f - S_N f\|_{L^\infty(\mathbf{T}^d)}$ as $N \rightarrow \infty$ can be as slow as any convergent sequence.

Theorem 2.5. Let $\{a_n : n \in \mathbf{Z}^d\}$ be any sequence of coefficients with $\lim_{|n| \rightarrow \infty} a_n = 0$. Then there exists $f \in C(\mathbf{T}^d)$ such that $\hat{f} \in L^1(\mathbf{Z})$, but $\hat{f}(n) = a_n$ for infinitely many $n \in \mathbf{Z}^d$.

Proof. For each $1 \leq k < \infty$, pick n_k such that $|a_{n_k}| \leq 1/2^k$ and such that the family $\{n_k\}$ is distinct. Then define

$$f(x) = \sum_{k=1}^{\infty} a_{n_k} e^{2\pi i n_k \cdot x}.$$

The absolute convergence of the right hand side shows $f \in C(\mathbf{T}^d)$, and that $\hat{f}(n_k) = a_{n_k}$ for each $1 \leq k \leq \infty$. □

A natural question is whether we *can* get quantitative convergence results for functions under additional assumptions. For instance, do we get faster convergence rates if $\|f\|_{L^\infty(\mathbf{T}^d)}$ is small (i.e. we have uniform control on the magnitude of f) rather than just if $\|f\|_{L^1(\mathbf{T}^d)}$ is small.

Example. *If we consider a square wave χ_I for some interval I , then the techniques of the following section allow us to prove that*

$$\|\chi_I - S_N \chi_I\|_{L^2(\mathbf{T})} \sim 1/\sqrt{N},$$

independently of I . This means that if we want to simulate square waves with a musical instrument up to some square mean error ε , then we will need about $1/\varepsilon^2$ different notes to represent the sound accurately. Thus a piano with 88 keys can only approximate square waves slightly better than a keyboard with 20 keys. If $f \in C^{m+1}(\mathbf{T}^d)$, then we will see

$$\|f - S_N f\|_{L^2(\mathbf{T})} \lesssim 1/N^{m/2},$$

so we require significantly less notes to simulate this sound, i.e. $\varepsilon^{-2/m}$. In this case a piano can simulate these sounds much more accurately.

Another question is whether $S_N f$ is stable under perturbations. For instance, if we replace f with a function g close to f the original function, is $S_N f$ close to $S_N g$? This is of interest in many practical applications, where error terms are inherently present. If an operator is unstable under perturbations that it is impractical to use it in an application to a real life situation. Again, the best way to measure the error terms are using an appropriate norm space.

These examples show that working with certain norms is an important way to understand the deeper properties of the Fourier series. It is an important property of norm spaces that most questions are equivalent to questions in the *completion* of that norm space. For instance, if one wants to use the norm $\|\cdot\|_{L^1(\mathbf{T}^d)}$ to analyze the space $C(\mathbf{T}^d)$, most questions are equivalent to questions about the completion of $C(\mathbf{T}^d)$, i.e. the space $L^1(\mathbf{T}^d)$ of all integrable functions. Moreover, working in the completion of a space enables us to employ many functional analysis arguments which make working with the more general space essential to many modern arguments. Despite the fact that we will be analyzing functions that one never deals with in ‘practical situations’, using these functions is a useful tool to determine the quantitative behaviour of more regular functions with respect to a norm.

2.4 Boundedness of Partial Sums

One initial equation which might summarize how well behaved the Fourier series is with respect to suitable norms would be to obtain an estimate of the form $\|\hat{f}\|_{L^q(\mathbf{Z}^d)} \lesssim \|f\|_{L^p(\mathbf{T}^d)}$ for particular values of p and q . This does not explicitly answer a question about convergence, but still shows that the Fourier series is stable under small perturbations in the norm on $L^p(\mathbf{T}^d)$. The first inequality we give is trivial, but is certainly tight, e.g. for $f(t) = e_n(t)$.

Theorem 2.6. *For any $f \in L^1(\mathbf{T}^d)$, $\|\hat{f}\|_{L^\infty(\mathbf{T}^d)} \leq \|f\|_{L^1(\mathbf{T}^d)}$.*

Proof. We just take absolute values into the oscillatory integral defining the Fourier coefficients, calculating that for any $n \in \mathbf{Z}^d$,

$$|\hat{f}(n)| = \left| \int_{\mathbf{T}^d} f(t) \overline{e_n(t)} \right| \leq \int_{\mathbf{T}^d} |f(t)| = \|f\|_{L^1(\mathbf{T}^d)},$$

which was the required bound. \square

This proof doesn't really take any deep features of the Fourier coefficients. The same bound holds for any integral

$$\int_{\mathbf{T}} f(t) K(t) dt,$$

where $|K(t)| \leq 1$ for all t . But the bound is still tight, which might be explained by the fact that the Fourier series gives oscillatory information which is not immediately present in the L^1 norms of the phase spaces, other than by taking a naive absolute bound into the L^1 norm. The only L^p norm where we can get a completely satisfactory bound is for $p = 2$, where we can use Hilbert space techniques; this should be expected to be very useful since orthogonality was implicitly used to define the Fourier series.

Theorem 2.7. *For any function $f \in L^2(\mathbf{T}^d)$, $\|\hat{f}\|_{L^2(\mathbf{Z}^d)} = \|f\|_{L^2(\mathbf{T}^d)}$.*

Proof. With respect to the normalized inner product on the space $L^2(\mathbf{T}^d)$, the calculations of the last chapter tell us that the exponentials $\{e_n : n \in \mathbf{Z}^d\}$ are an orthonormal family of functions, in the sense that for distinct pair

$n, m \in \mathbf{Z}^d$, $(e_n, e_m) = 0$ and $(e_n, e_n) = 1$. Since $\hat{f}(n) = (f, e_n)$, we apply Bessel's inequality to conclude

$$\|\hat{f}\|_{L^2(\mathbf{Z}^d)} \leq \|f\|_{L^2(\mathbf{T}^d)}.$$

The exponentials $\{e_n\}$ are actually an orthonormal basis for $L^2(\mathbf{T}^d)$; there are many ways to see this (the Stone-Weierstrass theorem, for instance). The most convenient way for us will be to note that if $f \in C^\infty(\mathbf{T}^d)$, then we have shown that if $(f, e_n) = 0$ for all $n \in \mathbf{Z}^d$, then $f = 0$. But $S_N f$ converges to f in $L^2(\mathbf{T}^d)$ (it actually converges uniformly), and so

$$\begin{aligned} \|f\|_{L^2(\mathbf{T}^d)} &= \lim_{N \rightarrow \infty} \|S_N f\|_{L^2(\mathbf{T}^d)} = \lim_{N \rightarrow \infty} \left(\sum_{n \in E_N} |\hat{f}(n)|^2 \right)^{1/2} \\ &= \left(\sum_{n \in \mathbf{Z}^d} |\hat{f}(n)|^2 \right)^{1/2} = \|\hat{f}\|_{L^2(\mathbf{Z}^d)}. \end{aligned}$$

This is Parseval's inequality for $C^\infty(\mathbf{T}^d)$. Now a density argument will give the general result. If $f \in L^2(\mathbf{T}^d)$ is a general element, then for each $\varepsilon > 0$ we can find $f_\varepsilon \in C^\infty(\mathbf{T}^d)$ such that $\|f_\varepsilon - f\|_{L^2(\mathbf{T}^d)} \leq \varepsilon$. Then Bessel's inequality

$$|\|f\|_{L^2(\mathbf{T}^d)} - \|\hat{f}\|_{L^2(\mathbf{Z}^d)}| \leq |\|f\|_{L^2(\mathbf{T}^d)} - \|f_\varepsilon\|_{L^2(\mathbf{T}^d)}| + |\|\hat{f}_\varepsilon\|_{L^2(\mathbf{Z}^d)} - \|\hat{f}\|_{L^2(\mathbf{Z}^d)}| \leq 2\varepsilon.$$

Taking $\varepsilon \rightarrow 0$ completes the proof. \square

This equality makes the Hilbert space $L^2(\mathbf{T}^d)$ often the best place to understand Fourier expansion techniques, and general results are often achieved by reduction to this well understood case. For instance, the inequality above, combined with the trivial inequality, is easily interpolated using the Riesz-Thorin technique to give the Hausdorff Young inequality.

Theorem 2.8. *If $1 \leq p \leq 2$, and $f \in L^p(\mathbf{T}^d)$, then $\|\hat{f}\|_{L^{p^*}(\mathbf{Z}^d)} \leq \|f\|_{L^p(\mathbf{T}^d)}$.*

It might be surprising to note that the Hausdorff Young inequality essentially completes the bounds on the Fourier series with respect to the L^p norms. There is no interesting result one can obtain for $p > 2$ other than the obvious inequality

$$\|\hat{f}\|_{L^2(\mathbf{Z}^d)} \leq \|f\|_{L^2(\mathbf{T}^d)} \leq \|f\|_{L^p(\mathbf{T}^d)}.$$

Thus we can control the magnitude of the Fourier coefficients in terms of the width of the original function, but we are limited in our ability to control the width of the Fourier coefficients in terms of the magnitudes of the original function. This makes sense, because the L^p norm of f measures fairly different aspects of the function than the L^q norm of the Fourier transform of f . It is only in the case of the L^2 norm where results are precise, and where p is small that we can take a trivial bound, that we get an inequality like the Hausdorff Young result.

2.5 Asymptotic Decay of Fourier Series

The next result, known as Riemann-Lebesgue lemma, shows that the Fourier series of any integrable function decays, albeit arbitrarily slowly. The proof we give is an instance of an important principle in Functional analysis that we will use over and over again. Suppose for each n , we have a bounded operator $T_n : X \rightarrow Y$ between norm spaces, and we want to show that for each $x \in X$, $\lim_{n \rightarrow \infty} T_n(x) = T(x)$, where T is another bounded operator. Suppose there is a dense set $X_0 \subset X$ such that for each $x_0 \in X_0$, $\lim_{n \rightarrow \infty} T_n(x_0) = T(x_0)$, and the family of operators $\{T_n\}$ are *uniformly* bounded in operator norm. Then for any $x \in X$,

$$\|T_n(x) - T(x)\| \leq \|T_n(x) - T_n(x_0)\| + \|T_n(x_0) - T(x_0)\| + \|T(x_0) - T(x)\|.$$

If we choose x_0 such that $\|x - x_0\| \leq \varepsilon$, then for n large enough we find that $\|T_n(x) - T(x)\| \leq \varepsilon$. Since ε was arbitrary, this means that $T_n(x) \rightarrow T(x)$ as $n \rightarrow \infty$. If we are working in a Banach space, the uniform boundedness says obtaining a uniform operator norm bound on $\{T_n\}$ is the *only* way to obtain this convergence.

The advantage of the principle is that it is suitably abstract, and can thus be used very flexibly. But the disadvantage is that it is a very soft analytical argument, and cannot be used to obtain results on the rate of convergence of $T_n(x)$ to $T(x)$. Here is a simple application.

Lemma 2.9 (Riemann-Lebesgue). *If $f \in L^1(\mathbf{T}^d)$, then $\widehat{f}(n) \rightarrow 0$ as $|n| \rightarrow \infty$.*

Proof. We claim the lemma is true for the characteristic function χ_I of a cube I . If $I = [a_1, b_1] \times \cdots \times [a_d, b_d]$, then it is simple to calculate that

$$\widehat{\chi_I}(n) = \prod_{k=1}^d \frac{e_n(-b_k) - e_n(-a_k)}{-in} = O(1/n)$$

By linearity of the integral, the Fourier transform of any step function vanishes at ∞ . But if

$$\Lambda_n(f) = \widehat{f}(n),$$

then

$$|\Lambda_n f| \leq \|\widehat{f}\|_{L^\infty(\mathbf{T})} \leq \|f\|_{L^1(\mathbf{T})},$$

which shows that the sequence of functionals $\{\Lambda_n\}$ are uniformly bounded as linear functionals on $L^1(\mathbf{T}^d)$. Since $\lim_{|n| \rightarrow \infty} \Lambda_n(f) = 0$ for any step function f , and the step functions are dense in $L^1(\mathbf{T}^d)$, we conclude that

$$\lim_{|n| \rightarrow \infty} \Lambda_n(f) = 0$$

for all $f \in L^1(\mathbf{T}^d)$. □

Even though the Fourier series of any step function decays at a rate $O(1/n)$, it is *not* true that a general Fourier series decays at a rate of $O(1/n)$. For instance, we have shown that there are continuous functions whose Fourier decay is arbitrarily slow. This is precisely the penalty for using a soft type analytical argument. Nonetheless, for smoother functions, we can obtain a uniform decay rate, which is our goal in the next section.

2.6 Smoothness and Decay

The next theorem obtains sharper bounds for smoother functions, and is an instance of a general phenomenon relating the duality between decay and smoothness in phase and frequency space.

Theorem 2.10. *If $f \in C^m(\mathbf{T}^d)$, then for each $n \in \mathbf{Z}^d$,*

$$|\widehat{f}(n)| \lesssim_{d,m} |n|^{-m} \max_{1 \leq i \leq d} \|\partial_i^m f\|_{L^1(\mathbf{T}^d)}.$$

Proof. We have

$$\widehat{\partial_i^m f}(\xi) = (2\pi i \xi_i)^m \widehat{f}(\xi).$$

Thus

$$|\widehat{f}(\xi)| \leq \frac{|\partial_i^m f|(\xi)|}{(2\pi |\xi_i|)^m} \leq \frac{\|\partial_i^m f\|_{L^1(\mathbf{T}^d)}}{(2\pi |\xi_i|)^m}.$$

But taking infima over all $1 \leq i \leq d$, we find

$$|\hat{f}(\xi)| \leq \frac{\max_{1 \leq k \leq d} \|\partial_i^m f\|_{L^1(\mathbf{T}^d)}}{[2\pi \max |\xi_i|]^m} \leq \frac{d^{1/2}}{(2\pi)^m} \frac{\max_{1 \leq i \leq d} \|\partial_i^m f\|_{L^1(\mathbf{T}^d)}}{|\xi|^m}. \quad \square$$

On the other hand, if $|\hat{f}(n)| \lesssim 1/|n|^{d+m}$, it is easy to see from the point-wise convergence of the Fourier series that $f \in C^m(\mathbf{R}^d)$. Note, however, that the introduction of the factor of d here gives a large gap between obtaining decay from smoothness and smoothness and decay when d is large, which is often a tricky problem to control when studying problems using harmonic analysis.

If $0 < \alpha < 1$, we say a function f is *Hölder continuous* of order α if there exists a constant A such that $|f(x+h) - f(x)| \leq A|h|^\alpha$ for all $x, h \in \mathbf{T}^d$. We define

$$\|f\|_{C^{0,\alpha}(\mathbf{T}^d)} = \sup_{x,h \in \mathbf{T}^d} \frac{|f(x+h) - f(x)|}{|h|^\alpha}.$$

Then the space $C^{0,\alpha}(\mathbf{T}^d)$ of all functions satisfying a Hölder condition of order α forms a Banach space.

Theorem 2.11. *If $f \in C^{0,\alpha}(\mathbf{T}^d)$, then $|\hat{f}(n)| \lesssim_d \|f\|_{C^{0,\alpha}(\mathbf{T}^d)} |n|^{-\alpha}$ for all $n \in \mathbf{Z}^d$.*

Proof. Fix $n \in \mathbf{Z}^d$. Then there is some $k \in \{1, \dots, d\}$ such that $|n_k| \gtrsim_d |n|$. We calculate that by periodicity,

$$\hat{f}(n) = - \int_{\mathbf{T}^d} f(x + e_k/n_k) \overline{e_n(x)} dx,$$

so

$$\hat{f}(n) = \frac{1}{2} \int_{\mathbf{T}^d} [f(x) - f(x + e_k/n_k)] \overline{e_n(x)} dx.$$

Thus taking in absolute values and applying Hölder continuity gives

$$|\hat{f}(n)| \leq \frac{\|f\|_{C^{0,\alpha}(\mathbf{T}^d)}}{2|n_k|^\alpha} \lesssim_d \frac{\|f\|_{C^{0,\alpha}(\mathbf{T}^d)}}{|n|^\alpha}. \quad \square$$

We also have a weaker converse statement, which shows f is Hölder continuous if it's Fourier series decays fast enough.

Theorem 2.12. Fix $f \in L^1(\mathbf{T}^d)$. Then

$$\|f\|_{C^{0,\alpha}(\mathbf{T}^d)} \lesssim_d \sup_{n \in \mathbf{Z}^d} |n|^{d+\alpha} |\hat{f}(n)|.$$

Proof. Let $A = \sup_{n \in \mathbf{Z}^d} |n|^{d+\alpha} |\hat{f}(n)|$. Then $\hat{f} \in L^1(\mathbf{Z}^d)$, so the Fourier inversion formula implies that for almost every $x \in \mathbf{T}^d$,

$$f(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}.$$

Then for $|h| < 1$,

$$f(x+h) - f(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x} (e^{2\pi i n \cdot h} - 1).$$

Now $|e^{2\pi i n \cdot h} - 1| \lesssim \min(1, |n||h|)$, so

$$\left| \sum_{|n| \leq 1/|h|} \hat{f}(n) e^{2\pi i n \cdot x} (e^{2\pi i n \cdot h} - 1) \right| \leq A|h| \sum_{|n| \leq 1/|h|} \frac{1}{|n|^{d-1+\alpha}} \lesssim_d A|h||h|^{\alpha-1} = A|h|^\alpha$$

and

$$\begin{aligned} \left| \sum_{|n| \geq 1/|h|} \hat{f}(n) e^{2\pi i n \cdot x} (e^{2\pi i n \cdot h} - 1) \right| &\leq 2A \sum_{|n| \geq 1/|h|} 1/|n|^{d+\alpha} \\ &\lesssim_d 2A|h|^\alpha. \end{aligned}$$

Combining these two calculations shows that

$$|f(x+h) - f(x)| \lesssim_d A|h|^\alpha,$$

so $\|f\|_{C^{0,\alpha}(\mathbf{T}^d)} \lesssim_d A$. □

Remark. Suppose that μ is a finite Borel measure on \mathbf{T}^d , for which we write $\mu \in M(\mathbf{T}^d)$. Then one can define the Fourier series of μ by setting

$$\hat{\mu}(n) = \int_{\mathbf{T}^d} e^{-2\pi i n \cdot x} d\mu(x).$$

If μ is absolutely continuous with respect to the normalized Lebesgue measure on \mathbf{T} , and $d\mu = f dx$, then $\hat{\mu} = \hat{f}$, so this is an extension of the Fourier series from integrable functions to finite measures. One can verify that

$$\|\hat{\mu}\|_{L^\infty(\mathbf{Z}^d)} \leq \|\mu\|_{M(\mathbf{T}^d)}.$$

If δ is the Dirac delta measure at the origin, i.e. $\mu(E) = 1$ if $0 \in E$, and $\mu(E) = 0$ otherwise, then for all n ,

$$\hat{\delta}(n) = 1.$$

Thus the Fourier series of δ has no decay at all. One can view this as saying functions are ‘smoother’ than measures, and therefore have a Fourier decay. Indeed, it is not too difficult to prove that a finite Borel measure μ on \mathbf{T}^d is absolutely continuous with respect to the Lebesgue measure if and only if

$$\lim_{|y| \rightarrow 0} \int_{\mathbf{R}^d} d|\mu(x+y) - \mu(x)| \rightarrow 0,$$

(we show that integrable functions satisfy this property in the next section) which shows that integrable functions are precisely the measures such that, in a certain sense, $\mu(x+y) \approx \mu(x)$ for small y .

2.7 Convolution and Kernel Methods

The notion of the convolution of two functions f and g is a key tool in Fourier analysis, both as a way to regularize functions, and as an operator that transforms nicely when we take Fourier series. Given $f, g \in L^1(\mathbf{T}^d)$, we define

$$(f * g)(x) = \int_{\mathbf{T}^d} f(y)g(x-y) dy.$$

Thus we smear the values of g with respect to a density function f .

Lemma 2.13. *For any $1 \leq p < \infty$, and $f \in L^p(\mathbf{T}^d)$,*

$$\lim_{h \rightarrow 0} \text{Trans}_h f = f$$

in $L^p(\mathbf{T}^d)$.

Proof. If f is $C^1(\mathbf{T}^d)$, then $|f(x+h) - f(x)| \lesssim_f h$ uniformly in x , implying that $\|\text{Trans}_h f - f\|_{L^p(\mathbf{T}^d)} \leq \|\text{Trans}_h f - f\|_{L^\infty(\mathbf{T}^d)} \lesssim_f h$, and so $\text{Trans}_h f \rightarrow f$ in all the spaces $L^p(\mathbf{T}^d)$. We have $\|\text{Trans}_h f\|_{L^p(\mathbf{T}^d)} = \|f\|_{L^p(\mathbf{T}^d)}$, so the operators $\{\text{Trans}_h\}$ are uniformly bounded. Since $C^1(\mathbf{T}^d)$ is dense in $L^p(\mathbf{T}^d)$ for $1 \leq p < \infty$, we conclude that $\lim_{h \rightarrow 0} \text{Trans}_h f = f$ for all $f \in L^p(\mathbf{T}^d)$. \square

Theorem 2.14. *Convolution has the following properties:*

- If $f \in L^p(\mathbf{T}^d)$ and $g \in L^q(\mathbf{T}^d)$, for $1/p + 1/q = 1$, then $f * g$ is uniformly continuous.
- If $f \in L^p(\mathbf{T}^d)$ and $g \in L^q(\mathbf{T}^d)$, and if we define r so that $1/r = 1/p + 1/q - 1$, with $1 \leq r \leq \infty$, then $f * g$ is well-defined by the convolution integral formula almost everywhere, and

$$\|f * g\|_{L^r(\mathbf{T}^d)} \leq \|f\|_{L^p(\mathbf{T}^d)} \|g\|_{L^q(\mathbf{T}^d)}.$$

This is known as Young's inequality for convolutions.

- Convolution is a commutative, associative, bilinear operation.
- If $f, g \in L^1(\mathbf{T})$, then $\widehat{f * g} = \widehat{f} \widehat{g}$.
- If f has a weak derivative $D^k f$ in $L^1(\mathbf{T}^d)$, then $f * g$ has a weak derivative in $L^1(\mathbf{T}^d)$, and $D^k(f * g) = D^k f * g$. Thus convolution is 'additively smoothing'. In particular, if $f \in C^k(\mathbf{T}^d)$ and $g \in C^l(\mathbf{T}^d)$, then $f * g \in C^{k+l}(\mathbf{T}^d)$.
- If f is supported on $E \subset \mathbf{T}^d$, and g on $F \subset \mathbf{T}^d$, then $f * g$ is supported on $E + F$.

Proof. Suppose $f \in L^p(\mathbf{T}^d)$, and $g \in L^q(\mathbf{T}^d)$, then

$$\begin{aligned} |(f * g)(t-h) - (f * g)(t)| &\leq \int_{\mathbf{T}^d} |f(t-h-s) - f(t-s)| |g(s)| ds \\ &\leq \|f_h - f\|_{L^p(\mathbf{T}^d)} \|g\|_{L^q(\mathbf{T}^d)}. \end{aligned}$$

The right hand side is a bound independent of t and converges to zero as $h \rightarrow 0$, so $f * g$ is uniformly continuous. Applying Hölder's inequality again

gives that $\|f * g\|_{L^\infty(\mathbf{T}^d)} \leq \|f\|_{L^p(\mathbf{T}^d)} \|g\|_{L^q(\mathbf{T}^d)}$. If $f \in L^p(\mathbf{T}^d)$, and $g \in L^1(\mathbf{T}^d)$, we use Minkowski's inequality to conclude that

$$\begin{aligned} \|f * g\|_{L^p(\mathbf{T}^d)} &= \left(\int_{\mathbf{T}^d} \left| \int_{\mathbf{T}^d} f(t-s)g(s) ds \right|^p dt \right)^{1/p} \\ &\leq \int_{\mathbf{T}^d} \left(\int_{\mathbf{T}^d} |f(t-s)g(s)|^p dt \right)^{1/p} ds \\ &= \int_{\mathbf{T}^d} g(s) \|f\|_{L^p(\mathbf{T}^d)} ds = \|f\|_{L^p(\mathbf{T}^d)} \|g\|_{L^1(\mathbf{T}^d)}. \end{aligned}$$

Thus $f * g$ is finite almost everywhere. The inequality also implies that

$$\|f * g\|_{L^p(\mathbf{T}^d)} \leq \|f\|_{L^1(\mathbf{T}^d)} \|g\|_{L^p(\mathbf{T}^d)}$$

if $f \in L^1(\mathbf{T}^d)$, and $g \in L^p(\mathbf{T}^d)$. But now implying Riesz-Thorin interpolation gives the general Young's inequality. Elementary applications of change of coordinates and Fubini's theorem establish the commutativity and associativity of convolution for functions $f, g \in L^1(\mathbf{T}^d)$. Similarly, one can apply Fubini's theorem to obtain associativity for $f, g, h \in L^1(\mathbf{T}^d)$. To obtain the product identity for the Fourier series, we can apply Fubini's theorem to write

$$\begin{aligned} \widehat{f * g}(n) &= \int_{\mathbf{T}^d} (f * g)(t) e_n(-t) dt \\ &= \int_{\mathbf{T}^d} \int_{\mathbf{T}^d} f(s) g(t-s) e_n(-t) ds dt \\ &= \int_{\mathbf{T}^d} f(s) \int_{\mathbf{T}^d} (L_{-s}g)(t) e_n(-t) dt ds \\ &= \int_{\mathbf{T}^d} f(s) e_n(-s) \widehat{g}(n) ds \\ &= \widehat{f}(n) \widehat{g}(n), \end{aligned}$$

and this is exactly the identity required. To calculate the weak derivative of $f * g$, we fix $\phi \in C^\infty(\mathbf{T}^d)$, and calculate using two applications of Fubini's

theorem that

$$\begin{aligned}
\int_{\mathbf{T}^d} (f' * g)(t) \phi(t) dt &= \int_{\mathbf{T}^d} \int_{\mathbf{T}^d} f'(t-s) g(s) \phi(t) ds dt \\
&= \int_{\mathbf{T}^d} g(s) \int_{\mathbf{T}^d} f'(t-s) \phi(t) dt ds \\
&= - \int_{\mathbf{T}^d} g(s) \int_{\mathbf{T}^d} f(t-s) \phi'(t) dt ds \\
&= - \int_{\mathbf{T}^d} \left(\int_{\mathbf{T}^d} g(s) f(t-s) ds \right) \phi'(t) dt \\
&= - \int_{\mathbf{T}^d} (f * g)(t) \phi'(t) dt.
\end{aligned}$$

If $f = 0$ a.e outside E , and $g = 0$ a.e. outside F , then $(f * g)(t)$ can be nonzero only when there is a set G of positive measure such that for any $s \in G$, $f(s) \neq 0$ and $g(t-s) \neq 0$. But this means that $E \cap G \cap (t-F)$ has positive measure, so that there is $s \in E$ such that $t-s \in F$, meaning that $t \in E + F$. \square

Example. Given a function $f \in L^1(\mathbf{T}^d)$ we define the autocorrelation function

$$R(\tau) = \int_{\mathbf{T}^d} f(t+\tau) \overline{f(t)} dt.$$

Then R is the convolution of $f(t)$ with $g(t) = \overline{f(-t)}$. Thus for $f \in L^1(\mathbf{T}^d)$, $R \in L^1(\mathbf{T}^d)$, and

$$\widehat{R}(n) = \widehat{f}(n) \overline{\widehat{f}(n)} = |\widehat{f}(n)|^2.$$

The function \widehat{R} is known as the power spectrum of f .

We know that suitably smooth functions have convergent Fourier series. The advantage of convolution is if we want to study the properties of any integrable function f , then for any function $u \in C^\infty(\mathbf{T})$, $f * u$ will be smooth, and if we choose u appropriately, we can let $f * u$ approximate f . In terms of Fourier analysis, provided that \widehat{u} is close to one, $(f * u)^\wedge$ will be close to \widehat{f} , and so we might expect $f * u$ to be close to f . If we can establish the convergence properties on the convolution $f * g$, then we can probably obtain results about f . From the frequency side, $\sum \widehat{f}(n) e_n$ might not converge, but $\sum a_n \widehat{f}(n) e_n$ might converge for a suitably fast decaying sequence

a_n . But if a_n is close to one, this sequence might still reflect properties of the original sequence.

To make rigorous the idea of approximating the Fourier series of a function, we introduce families of *good kernels*. A good kernel is a sequence of integrable functions $\{K_n\}$ on \mathbf{T} bounded in L^1 norm, for which

$$\int_{\mathbf{T}} K_n(t) = 1.$$

so that integration against K_n operates essentially like an average, and for any $\delta > 0$,

$$\lim_{n \rightarrow \infty} \int_{|t| > \delta} |K_n(t)| \rightarrow 0. \quad (2.1)$$

Thus the functions $\{K_n\}$ become concentrated at the origin as $n \rightarrow \infty$. If in addition, we have an estimate $\|K_n\|_{L^\infty(\mathbf{T}^d)} \lesssim n^d$, we say it is an *approximation to the identity*.

Example. The simplest way to obtain a good kernel is to fix $K \in L^1(\mathbf{T}^d)$ with

$$\int_{\mathbf{T}^d} K(x) dx = 1,$$

and to define

$$K_n(x) = \begin{cases} n^d \cdot K(nx) & : |x_1|, \dots, |x_d| \leq 1/n, \\ 0 & : \text{otherwise.} \end{cases}$$

Then $\|K_n\|_{L^1(\mathbf{T})} = 1$ for all $n > 0$, and K_n is eventually supported on every small ball around the origin, which implies (2.1). If $K \in L^\infty(\mathbf{T}^d)$, then the resulting sequence $\{K_n\}$ is also an approximation to the identity.

Theorem 2.15. Let $\{K_n\}$ be a good kernel. Then

- $(K_n * f)(t) \rightarrow f(t)$ for any continuity point t of f .
- $(K_n * f) \rightarrow f$ uniformly if $f \in C(\mathbf{T}^d)$, and $K_n * f$ converges to f in $L^p(\mathbf{T}^d)$ if $f \in L^p(\mathbf{T}^d)$, for $1 \leq p < \infty$.
- If K_n is an approximation to the identity, $(K_n * f)(t) \rightarrow f(t)$ for all t in the Lebesgue set of f .

Proof. The operators $T_n f = K_n * f$ are uniformly bounded as operators on $L^p(\mathbf{T})$. Basic analysis shows that $(K_n * f)(t) \rightarrow f(t)$ at each point t where f is continuous, and converges uniformly to f if f is in $C(\mathbf{T}^d)$. But a density argument allows us to conclude that $K_n * f \rightarrow f$ in $L^p(\mathbf{T})$ for each $f \in L^p(\mathbf{T}^d)$, for $1 \leq p < \infty$. To obtain pointwise convergence for t in the Lebesgue set of f , we calculate

$$|(K_n * f)(t) - f(t)| \leq \int_{\mathbf{T}^d} |f(t-s) - f(t)| |K_n(s)| ds.$$

Let $A(\delta) = \delta^{-d} \int_{|s| < \delta} |f(t-s) - f(t)|$. Then as $\delta \rightarrow 0$, $A(\delta) \rightarrow 0$ because t is in the Lebesgue set of f . And we find that for each k , since $|K_n(s)| \lesssim n^d$,

$$\int_{2^k/n < |t| < 2^{k+1}/n} |f(t-s) - f(t)| |K_n(s)| \lesssim \frac{A(2^{k+1}/n)}{2^{d(k+1)}}.$$

Thus we have a bound

$$|(K_n * f)(t) - f(t)| \lesssim_d \sum_{k=0}^{\infty} \frac{A(2^k/n)}{2^{dk}}.$$

Because f is integrable, A is continuous, and hence bounded. This means that for each m ,

$$|(K_n * f)(t) - f(t)| \lesssim_d \sum_{k=0}^m \frac{A(2^k/n)}{2^{dk}} + \|A\|_{\infty} \sum_{k=m}^{\infty} \frac{1}{2^{dk}} = \sum_{k=0}^m \frac{A(2^k/n)}{2^{dk}} + O_d(1/2^{dm}).$$

For any fixed m , the finite sum tends to zero as $n \rightarrow \infty$, so we obtain that $|(K_n * f)(t) - f(t)| = o(1) + O_d(1/2^m)$. Taking $m \rightarrow \infty$ proves the result. \square

As mentioned above, if $u \in C^{\infty}(\mathbf{T})$, then for any $f \in L^1(\mathbf{T})$, $f * u \in C^{\infty}(\mathbf{T})$, and we have $D^{\alpha}(f * u) = f * D^{\alpha}u$. Thus we can use Young's inequality to bound $D^{\alpha}(f * u)$ in terms of f and $D^{\alpha}u$. Thus in some contexts, it is useful to find out how smooth we can make u , while still being able to approximate a non-smooth function f with $f * u$. If we fix $u \in C^{\infty}(\mathbf{T})$ supported on an interval of length $1/4$ centered at the origin, then for $0 < \varepsilon < 1$, we can define $u_{\varepsilon}(t) = u(t/\varepsilon)$ for $|t| \leq \varepsilon$, and $u_{\varepsilon}(t) = 0$ for $|t| > \varepsilon$. Such a function will lie in $C^{\infty}(\mathbf{T})$, be supported on an interval of length 2ε interval centered at the origin. Such a function satisfies estimates of the

form $\|D^k u_\varepsilon\|_{L^\infty(\mathbf{T})} \lesssim_k \varepsilon^{-k}$. Roughly speaking, this is pretty much the best we can do for a function with such small support. $u \in C^\infty(\mathbf{T})$ is supported on $|t| \leq \varepsilon$, and $\int_{\mathbf{T}} u(t) dt = 1$, then the mean value theorem can be used to show that $\|D^k u\|_{L^\infty(\mathbf{T})} \gtrsim_k \varepsilon^{-k}$ for all $k > 0$.

2.8 The Dirichlet Kernel

For simplicity, let us now focus exclusively on the case $d = 1$ with the canonical summation operators S_N . For $f \in L^1(\mathbf{T})$, we calculate that

$$(S_N f)(t) = \sum_{n=-N}^N \hat{f}(n) e_n(t) = \int_{\mathbf{T}^d} f(x) \left(\sum_{n=-N}^N e_n(t-x) \right) dx.$$

The bracketed part of the final term in the equation is independant of the function f , and is therefore key to understanding the behaviour of the sums S_N . We call it the *Dirichlet kernel*, denoted D_N . Thus

$$D_N(t) = \sum_{n=-N}^N e_n(t)$$

and so $S_N f = f * D_N$. Thus analyzing this convolution is *key* to understanding the partial summation operators.

Remark. In the higher dimensional case, we can consider the operators

$$K_N(t) = \sum_{n \in E_N} e_n(t).$$

The behaviour of these functions is highly dependant on the choice of the sets E_N , and we thus leave the higher dimensional analysis to a different time.

Theorem 2.16. *For any integer N and $t \in \mathbf{T}$,*

$$D_N(t) = \frac{\sin(2\pi(N + 1/2)t)}{\sin(\pi t)}.$$

Proof. By the geometric series summation formula, we may write

$$\begin{aligned}
D_N(t) &= 1 + \sum_{n=1}^N e_n(t) + e_n(-t) = 1 + e(t) \frac{e_N(t) - 1}{e(t) - 1} + e(-t) \frac{e_N(-t) - 1}{e(-t) - 1} \\
&= 1 + e(t) \frac{e_N(t) - 1}{e(t) - 1} + \frac{e_N(-t) - 1}{1 - e(t)} = \frac{e_{N+1}(t) - e_N(-t)}{e(t) - 1} \\
&= \frac{e_{N+1/2}(t) - e_{N+1/2}(-t)}{e_{1/2}(t) - e_{1/2}(-t)} = \frac{\sin(2\pi(N + 1/2)t)}{\sin(\pi t)}. \quad \square
\end{aligned}$$

If D_N was a good kernel, then we would obtain that the partial sums of S_N converge uniformly. This initially seems a good strategy, because $\int D_N(t) = 1$. However, we find

$$\begin{aligned}
\int_{\mathbf{T}^d} |D_N(t)| &= \int_0^1 \left| \frac{\sin(2\pi(N + 1/2)t)}{\sin(\pi t)} \right| dt \\
&\gtrsim \int_0^1 \frac{|\sin(2\pi(N + 1/2)t)|}{\sin(\pi t)} dt \\
&\gtrsim \int_0^1 \frac{|\sin(2\pi(N + 1/2)t)|}{t} dt \\
&= \int_0^{2\pi N + \pi} \frac{|\sin(t)|}{t} dt \\
&\gtrsim \sum_{n=0}^N \frac{1}{t} dt \gtrsim \log(N).
\end{aligned}$$

Thus the L^1 norm of D_N grows, albeit slowly, to ∞ . This reflects the fact that D_N oscillates very frequently, and also that the pointwise convergence of the Fourier series is much more subtle than that provided by good kernels. In fact, a simple functional analysis argument shows that pointwise convergence of Fourier series fails for continuous functions.

Theorem 2.17. *There exists $f \in C(\mathbf{T})$ such that $(S_N f)(0)$ diverges as $N \rightarrow \infty$.*

Proof. If we consider the linear functionals $\Lambda_N f = (S_N f)(0) = (f * D_N)(0)$ on $C(\mathbf{T})$. If we let f_N be a continuous function approximating $\text{sgn}(D_N)$ for each N , then $|\Lambda_N f_N| \gtrsim \log N \cdot \|f_N\|_{L^\infty(\mathbf{T})}$. This implies that $\|\Lambda_N\| \rightarrow \infty$ as $N \rightarrow \infty$. The uniform boundedness principle thus implies that there exists a *single* function $f \in C(\mathbf{T})$ such that $\sup |\Lambda_N f| = \infty$, so $(S_N f)(0)$ diverges as $N \rightarrow \infty$. \square

The situation is even worse than this for general integrable functions. In 1927, Andrey Kolmogorov constructed an integrable function whose Fourier series diverges everywhere. But there is some hope. In 1928, Marcel Riesz showed, using methods we will develop in these notes, that if $1 < p < \infty$, and $f \in L^p(\mathbf{T})$, that $S_N f$ converges in the L^p norm to f , by showing the Hilbert transform was bounded from $L^p(\mathbf{T})$ to $L^p(\mathbf{T})$. And after a half century of the development of techniques in harmonic analysis, in 1966, Carleson proved that for each $f \in L^p(\mathbf{T})$, for $1 < p \leq \infty$, the Fourier series of f converges almost everywhere to f . The multivariate picture is more complicated and many questions remain open today; tensoring shows that $S_N f$ converges to f in $L^p(\mathbf{T}^d)$ if $f \in L^p(\mathbf{T}^d)$, *provided that we interpret $S_N f$ as a square summation*, and in 1970 Charles Fefferman showed that for square summation $S_N f$ converges to f almost everywhere. On the other hand, in 1971 Charles Fefferman showed that for spherical summation the *only* place we have norm convergence is in $L^2(\mathbf{T}^d)$. It remains an open question whether the partial spherical summation $S_N f$ converges to f almost everywhere.

2.9 Countercultural Methods of Summation

We now interpret our convergence of series according to a different kernel, so we do get a family of good kernels, and therefore we obtain pointwise convergence for suitable reinterpretations of partial sums. One reason why the Dirichlet kernel fails to be a good kernel is that the Fourier coefficients of the kernel have a sharp drop – the coefficients are either equal to one or to zero. If we mollify, then we will obtain a family of good kernels. And the best way to do this is to alter our summation methods slightly.

The standard method of summation suffices for much of analysis. Given a sequence a_0, a_1, \dots , we define the infinite sum as the limit of partial sums. Some sums, like $\sum_{k=1}^{\infty} k$, obviously diverge, whereas other sums, like $\sum 1/n$, ‘just’ fail to converge because they grow suitably slowly towards infinity over time. Since the time of Euler, a new method of summation developed by Cesaro was introduced which ‘regularized’ certain terms by considering averaging the sums over time. Rather than considering limits of partial sums, we consider limits of averages of sums, known

as Cesaro means. Letting $s_n = \sum_{k=0}^n a_k$, we define the Cesaro means

$$\frac{s_0 + \cdots + s_n}{n+1},$$

A sequence is Cesaro summable to some value if these averages converge. If the normal summation exists, then the Cesaro limit exists, and is equal to the original sum. However, the Cesaro summation is stronger than normal convergence.

Example. *In the sense of Cesaro, we have $1 - 1 + 1 - 1 + \cdots = 1/2$, which reflects the fact that the partials sums do ‘converge’, but to two different numbers 0 and 1, which the series oscillates between, and the Cesaro means average these two points of convergence out to give a single method of convergence.*

Another notion of regularization sums emerged from Complex analysis, called Abel summation. Given a sequence $\{a_i\}$, we can consider the power series $\sum a_k r^k$. If this is well defined for $|r| < 1$, we can consider the Abel means $A_r = \sum a_k r^k$, and ask if $\lim_{r \rightarrow 1} A_r$ exists, which should be ‘almost like’ $\sum a_k$. If this limit exists, we call it the Abel sum of the sequence.

Example. *In the Abel sense, we have $1 - 2 + 3 - 4 + 5 - \cdots = 1/4$, because*

$$\sum_{k=0}^{\infty} (-1)^k (k+1) z^k = \frac{1}{(1+z)^2}.$$

The coefficients here are $\Omega(N)$, so they can’t be Cesaro summable.

Abel summation is even more general than Cesaro summation, as the following theorem shows.

Theorem 2.18. *A Cesaro summable sequence is Abel summable.*

Proof. Let $\{a_i\}$ be a Cesaro summable sequence, which we may without loss of generality assume converges to 0. Now $(n+1)\sigma_n - n\sigma_{n-1} = s_n$, so

$$(1-r)^2 \sum_{k=0}^n (k+1)\sigma_k r^k = (1-r) \sum_{k=0}^n s_k r^k = \sum_{k=0}^n a_k r^k$$

As $n \rightarrow \infty$, the left side tends to a well defined value for $r < 1$, hence the same is true for $\sum_{k=0}^n a_k r^k$. Given $\varepsilon > 0$, let N be large enough that $|\sigma_n| < \varepsilon$ for $n > N$, and let M be a bound for all $|\sigma_n|$. Then

$$\begin{aligned}
\left| (1-r)^2 \sum_{k=0}^{\infty} (k+1) \sigma_k r^k \right| &\leq (1-r)^2 \left(\sum_{k=0}^N (k+1) |\sigma_k| r^k + \varepsilon \sum_{k=N+1}^{\infty} (k+1) r^k \right) \\
&= (1-r)^2 \left(\sum_{k=0}^N (k+1) (|\sigma_k| - \varepsilon) r^k + \varepsilon \left[\frac{r^{N+1}}{1-r} + \frac{1}{(1-r)^2} \right] \right) \\
&\leq (1-r)^2 M \sum_{k=0}^N (k+1) r^k + \varepsilon r^{N+1} (1-r) + \varepsilon \\
&\leq (1-r)^2 M \frac{(N+1)(N+2)}{2} + \varepsilon r^{N+1} (1-r) + \varepsilon
\end{aligned}$$

Fixing N , and letting $r \rightarrow 1$, we may make the complicated sum on the end as small as possible, so the absolute value of the infinite sum is less than ε . Thus the Abel limit converges to zero. \square

2.10 Fejer Summation

Note that the Cesaro means of the Fourier series of f are given by

$$\sigma_N(f) = \frac{S_0(f) + \cdots + S_{N-1}(f)}{N} = f * \left(\frac{D_0 + \cdots + D_{N-1}}{N} \right) = f * F_N,$$

where we have introduced a new kernel F_N , called the *Fejer kernel*. Here, we have a simple formula for the Cesaro means, i.e.

$$F_N(x) = \sum_{n=-N}^N \left(1 - \frac{|n|}{N} \right) e_n(t) = \frac{1}{N} \frac{\sin^2(Nx/2)}{\sin^2(x/2)}.$$

Thus the oscillations of the Dirichlet kernel are slightly dampened, and as a result, we can easily see that F_N is an approximation to the identity.

Theorem 2.19 (Fejér's Theorem). *For any $f \in L^1(\mathbf{T})$,*

- $(\sigma_N f)(x) \rightarrow f(x)$ for all x in the Lebesgue set of f .

- $\sigma_N f \rightarrow f$ uniformly if $f \in C(\mathbf{T})$.
- $\sigma_N f \rightarrow f$ in the L^p norm for $1 \leq p < \infty$, if $f \in L^p(\mathbf{T})$.

If we look at the Fourier expansion of the trigonometric polynomial $\sigma_N(f)$, viewing σ_N as a *Fourier multiplier operator*, we see that

$$\sigma_N f = \sum_{n=-N}^N \left(1 - \frac{|n|}{N}\right) \hat{f}(n) e_n.$$

Thus the Fourier coefficients are slowly added to the expansion, rather than a sharp cutoff as with ordinary Dirichlet summation. This is one reason for the nice convergence properties the kernel has as compared to the Dirichlet kernel.

Corollary 2.20. *If $f \in L^1(\mathbf{T})$ and $\hat{f} = 0$, then $f = 0$ almost everywhere.*

Proof. If $\hat{f} = 0$, then $\sigma_N f = 0$ for all N . But $\sigma_N f \rightarrow f$ in $L^1(\mathbf{T})$, which means that $f = 0$ in $L^1(\mathbf{T})$, so $f = 0$ almost everywhere. \square

This corollary is often more useful than the more technical convergence statements due to its relative simplicity. We will later see this result is also true for $d > 1$, via use of the Poisson summation formula for the Fourier transform.

Example. We say $f \in L^1(\mathbf{T}^d)$ is *band limited* if its Fourier series is supported on finitely many points. If $\{S_N\}$ is defined as before, and N is suitably large that E_N contains the support of \hat{f} , then

$$\hat{f} = \hat{f} \cdot \mathbf{I}_{E_N} = \widehat{f K_N} = \widehat{f * K_N}.$$

It thus follows from the previous result that $f = f * K_N$ almost everywhere. But this means we can adjust f on a set of measure zero such that $f \in C^\infty(\mathbf{T}^d)$.

2.11 Abel Summation

Let us now consider the Abel sum of the Fourier integrals. We begin by focusing on the one-dimensional case, as in the last section. Thus for $f \in L^1(\mathbf{T})$ we have

$$A_r(f) = \sum_{n=-\infty}^{\infty} \hat{f}(n) r^n e_n(t).$$

Thus, if we define the *Poisson kernel*

$$P_r(t) = \sum_{n=-\infty}^{\infty} r^{|n|} e_n(t)$$

For each $r < 1$, this series converges uniformly for $t \in \mathbf{T}$, so P_r is a well-defined continuous function, and the uniform convergence shows that $A_r(f) = P_r * f$. As with the Fejer kernel, the family $\{P_r\}$ is *also* a good kernel as $r \rightarrow 1$. To see this, we can apply an infinite geometric series summation to obtain that

$$\begin{aligned} \sum r^{|n|} e_n(t) &= 1 + \frac{re(t)}{1 - re(t)} + \frac{re(-t)}{1 - re(-t)} = 1 + \frac{2r \cos 2\pi t - 2r^2}{(1 - re(t))(1 - re(-t))} \\ &= 1 + \frac{2r \cos 2\pi t - 2r^2}{1 - 2r \cos 2\pi t + r^2} = \frac{1 - r^2}{1 - 2r \cos 2\pi t + r^2}. \end{aligned}$$

As $r \rightarrow 1$, the function concentrates at the origin, because as $r \rightarrow 1$, if $\delta \leq |t| \leq \pi$, then $1 - \cos 2\pi t$ is bounded away from the origin, so

$$\begin{aligned} \left| \frac{1 - r^2}{1 - 2r \cos 2\pi t + r^2} \right| &= \left| \frac{1 + r}{(1 + (1 - 2 \cos 2\pi t)r) + 2(1 - \cos 2\pi t)r^2/(1 - r)} \right| \\ &= O\left(\frac{1 - r}{1 - \cos 2\pi t}\right) = O_\delta(1 - r). \end{aligned}$$

Moreover,

$$\|P_r\|_{L^\infty(\mathbf{T})} \leq \frac{1 - r^2}{1 - 2r + r^2} \leq \frac{2}{1 - r}.$$

Thus the Poisson kernel is an approximation to the identity; the oscillation in the kernel cancels out as $r \rightarrow 1$.

Theorem 2.21. *For any $f \in L^1(\mathbf{T})$,*

- $(A_r f)(t) \rightarrow f(t)$ for all x in the Lebesgue set of f .
- $A_r f \rightarrow f$ uniformly if $f \in C(\mathbf{T})$.
- $A_r f \rightarrow f$ in the L^p norm for $1 \leq p < \infty$, if $f \in L^p(\mathbf{T})$.

The Poisson kernel is not a trigonometric polynomial, and therefore not quite as easy to work with as the Féjer kernel. However, it is the real part of the Cauchy kernel

$$\frac{1 + re^{2\pi it}}{1 - re^{2\pi it}},$$

and therefore links the study of trigonometric series and the theory of analytic functions. We will see the kernel return when we study application of harmonic analysis to partial differential equations.

2.12 The De la Vallée Poisson Kernel

By taking a kernel halfway between the Dirichlet kernel and the Fejer kernel, we can actually obtain important results about ordinary summation. For two integers $M > N$, we define

$$\sigma_{N,M}(f) = \frac{M\sigma_M(f) - N\sigma_N(f)}{M - N}.$$

If we take a look at the Fourier expansion of $\sigma_{N,M}f$, we find

$$\sigma_{N,M}f = \sum_{n=-M}^M \frac{M - |n|}{M - N} e_n - \sum_{n=-N}^N \frac{N - |n|}{M - N} e_n = S_N f + \sum_{|n|=N+1}^M \frac{M - |n|}{M - N} e_n.$$

So we still have a slow decay in the Fourier coefficients. And as a result, if we look at the associated De la Vallée Poisson kernel, we find that a suitable subsequence is an approximation to the identity. In particular, for any fixed integer k , the sequence $\sigma_{kN, (k+1)N}$ leads to a good kernel. More interestingly, if the Fourier coefficients of f have some decay, then the De la Vallée does not differ that much from the ordinary sum, which gives useful results.

Theorem 2.22. *If $\hat{f}(n) = O(|n|^{-1})$, then for any integers N and k , if*

$$kN \leq M < (k+1)N,$$

then

$$\|\sigma_{kN, (k+1)N}f - S_M f\|_{L^\infty(\mathbf{T})} \lesssim 1/k.$$

Where the implicit constant is independant of N and k .

Proof. We just calculate that, since the Poisson sum has essentially the same weight for low term coefficients as the sum $S_M f$,

$$\|\sigma_{kN, (k+1)N} f - S_M f\|_{L^\infty(\mathbf{T})} \lesssim \sum_{kN \leq |n| < (k+1)N} |\hat{f}(n)| \lesssim \sum_{n=kN}^{(k+1)N} \frac{1}{n} \leq \frac{N}{kN} = \frac{1}{k}. \quad \square$$

Corollary 2.23. *If $f \in L^1(\mathbf{T})$ with $\hat{f}(n) = O(|n|^{-1})$,*

- $S_N f$ converges to f in the L^p norm for $1 \leq p < \infty$.
- $S_N f$ converges uniformly to f if $f \in C(\mathbf{T})$.
- $(S_N f)(x) \rightarrow f(x)$ for each Lebesgue point x of f .

Proof. The idea is quite simple. Fix N . Given any ε , we can use the last theorem to find k large enough such that if $kN \leq M < k(N+1)$,

$$\|\sigma_{kN, (k+1)N} f - S_M f\|_{L^\infty(\mathbf{T})} \leq \varepsilon.$$

But this gives the first and second result, up to perhaps a ε of error. The latter result is given by similar techniques. \square

2.13 Pointwise Convergence

One way around around the blowup in the L^1 norm of D_N is to consider only functions f which provide a suitable dampening condition on the oscillation of D_N near the origin. This is provided by smoothness of f , manifested in various ways. The first thing we note is that the convergence of $(S_N f)(t)$ for a fixed x_0 depends only *locally* on the function f .

Lemma 2.24 (Riemann Localization Principle). *If f_0 and f_1 agree in an interval around t_0 , then*

$$(S_N f_0)(t_0) = (S_N f_1)(t_0) + o(1).$$

Proof. Let

$$X = \{f \in L^1(\mathbf{T}) : f(x) = 0 \text{ for almost every } x \in (t_0 - \varepsilon, t_0 + \varepsilon)\}.$$

Then X is a closed subset of $L^1(\mathbf{T})$. Note that for all $x \in [-\pi, \pi]$,

$$\sin(t/2) \gtrsim t \quad \text{and} \quad \sin((N + 1/2)t) \leq 1.$$

Thus if $|t| \geq \varepsilon$,

$$|D_N(t)| = \frac{|\sin(2\pi(N + 1/2)t)|}{|\sin(\pi t)|} \lesssim 1/\varepsilon.$$

In particular, by Hölder's inequality, the functionals $T_N f = (S_N f)(t_0)$ are uniformly bounded on X , i.e. $\|T_N\| \lesssim 1/\varepsilon$. If f is smooth, and vanishes on $(t_0 - \varepsilon, t_0 + \varepsilon)$, then $T_N f \rightarrow 0$ as $N \rightarrow \infty$. But the space of such functions is dense in X , which implies that $T_N f \rightarrow 0$ for *any* $f \in X$. Thus if f_0, f_1 are two functions that agree in $(t_0 - \varepsilon, t_0 + \varepsilon)$, then $f_0 - f_1 \in X$, so $(S_N f_0)(t_0) = (S_N f_1)(t_0) + o(1)$. In particular, the pointwise convergence properties of f_0 and f_1 are equivalent at t_0 . \square

Thus any result about the pointwise convergence of Fourier series must depend on the local properties of a function f . Here, we give two of the main criteria, which corresponds to the smoothness of a function about a point x : either f is in a sense, 'locally Lipschitz', or 'locally of bounded variation'.

Theorem 2.25 (Dini's Criterion). *If there exists δ such that*

$$\int_{|t| < \delta} \left| \frac{f(x+t) - f(x)}{t} \right| dt < \infty,$$

then $(S_N f)(x) \rightarrow f(x)$.

Proof. Assume without loss of generality that $x = 0$ and $f(x) = 0$. Fix $\varepsilon > 0$, and pick δ_0 such that

$$\int_{|t| < \delta_0} \left| \frac{f(t)}{t} \right| dt < \varepsilon.$$

We have

$$|(S_N f)(0)| = \left| \left(\int_{|t| < \delta_0} + \int_{|t| \geq \delta_0} \right) f(t) D_N(t) dt \right|.$$

Now

$$\int_{|t| \geq \delta_0} f(t) D_N(t) dt = (D_N * (\mathbf{I}_{|t| \geq \delta_0} f))(0) = S_N(\mathbf{I}_{|t| \geq \delta_0} f)(0) = o(1)$$

since $f \mathbf{I}_{|t| \geq \delta_0}$ vanishes in a neighbourhood of the origin. On the other hand, we note that $t/\sin(\pi t)$ is a bounded function on \mathbf{T} , so

$$\begin{aligned} \int_{|t| < \delta_0} f(t) D_N(t) dt &= \int_{|t| < \delta_0} \left(\sin(2\pi(N + 1/2)t) \frac{f(t)}{t} \right) \left(\frac{t}{\sin(\pi t)} \right) dt \\ &\lesssim \|f(t)/t\|_{L^1[-\delta_0, \delta_0]} \leq \varepsilon. \end{aligned}$$

Thus, for suitably large N , $|(S_N f)(0)| \lesssim \varepsilon$. Since ε was arbitrary, the proof is complete. \square

This proof applies, in particular, if f is locally Lipschitz at x . Note the application of the Riemann Lebesgue lemma to show that to analyze the pointwise convergence of $(S_N f)(x)$, it suffices to analyze

$$\lim_{N \rightarrow \infty} \int_{|t| < \delta} f(x+t) D_N(t) dt$$

for any fixed $\delta > 0$.

Lemma 2.26 (Jordan's Criterion). *If $f \in L^1(\mathbf{T})$ locally has bounded variation about x , then*

$$(S_N f)(x) \rightarrow \frac{f(x^+) + f(x^-)}{2}.$$

Proof. By Riemann's localization principle, we may assume f has bounded variation everywhere. Then without loss of generality, we may assume f is an increasing function, since a bounded variation function is the difference of two monotonic functions. Since

$$\int_{-1/2}^{1/2} D_N(t) dt = \int_0^{1/2} [f(x+t) + f(x-t)] D_N(t) dt,$$

it suffices without loss of generality to show that

$$\lim_{N \rightarrow \infty} \int_0^{1/2} f(x+t) D_N(t) dt = \frac{f(x+)}{2}.$$

Since $\int_0^{1/2} D_N(t) dt = 1/2$, this is equivalent to showing

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi} \int_0^\pi [f(x+t) - f(x+)] D_N(t) dt = 0.$$

Because of this, we may assume without loss of generality that $x = 0$ and $f(x+) = 0$. Then by the mean value theorem for integrals (which only applies for monotonic functions), for each N , there exists $0 \leq \nu_N \leq 1/2$ such that

$$\int_0^{1/2} f(t) D_N(t) dt = \|f\|_\infty \int_{\nu_N}^{1/2} D_N(t) dt.$$

Now an integration by parts gives

$$\int_{\nu_N}^{1/2} D_N(t) dt \lesssim \int_{\nu_N}^{1/2} \frac{\sin((N+1/2)t)}{t} dt = \int_{\nu_N/(N+1/2)}^{1/2(N+1/2)} \frac{\sin(t)}{t} dt \lesssim \frac{1}{N+1/2}.$$

Thus

$$\int_0^{1/2} f(t) D_N(t) dt \lesssim \frac{1}{N+1/2} \rightarrow 0. \quad \square$$

Remark. The calculations in this proof also show that if $f \in L^1(\mathbf{T})$ has bounded variation, then

$$\hat{f}(n) = O(1/|n|).$$

We have seen that this implies $S_N f$ converges to f at every point on the Lebesgue set of f , $S_N f$ converges uniformly to f if $f \in C(\mathbf{T})$, and for any $1 \leq p < \infty$, if $f \in L^p(\mathbf{T})$, $S_N f$ converges to f in $L^p(\mathbf{T})$. Dirichlet's theorem says that the Fourier series of a continuous function f with only finitely many maxima and minima converges uniformly to f everywhere. Such a function has bounded variation, and so Dirichlet's theorem is an easy consequence of our discussion.

Of course, applying various better decay rates leads to a more uniform version of this theorem. The decay of the Fourier series depends on the decay of the Fourier coefficients of $yg(y)$ and $g(y) \cos(y/2)(y/\sin(y/2))$. In particular, if these coefficients is $O(|n|^{-m})$, then the convergence rate is also $O(|n|^{-m})$. If this decay rate is independent of x for suitable values of x , the convergence will be uniform over these values of x .

Example. Consider the sawtooth function defined on $[-1/2, 1/2)$ by $s(t) = t$, and then made periodic on the entire real line. We can easily calculate the Fourier series here, obtaining that

$$s(t) = i \sum_{n \neq 0} \frac{(-1)^n e_n(t)}{2\pi n} = -2 \sum_{n=1}^{\infty} \frac{(-1)^n \sin(2\pi n t)}{n}.$$

Thus for any $t \in (-1/2, 1/2)$,

$$\sum_{n=1}^{\infty} \frac{(-1)^n \sin(2\pi n t)}{n} = -t/2.$$

Theorem 2.27. If $\hat{f}(n) = O(|n|^{-1})$, and $f(t_0-)$ and $f(t_0+)$ exist, then

$$(S_N f)(t_0) \rightarrow \frac{f(t_0-) + f(t_0+)}{2}.$$

Proof. The idea of our proof is to break f into a nice continuous function, and the sawtooth function, where we already understand the convergence of Fourier series. Without loss of generality, let $t_0 = 1/2$. Define $g(t) = f(t) + (f(1/2+) - f(1/2-))s(t)/2$ on $(-1/2, 1/2)$, where s is the sawtooth function. Then

$$\lim_{t \uparrow 1/2} g(t) = \lim_{t \downarrow -1/2} g(t) = \frac{f(1/2+) + f(1/2-)}{2}.$$

Thus g can be defined on \mathbf{T} so it is continuous at t_0 . Now we find $|\hat{g}| \lesssim |\hat{f}| + |\hat{s}| = O(|n|^{-1})$, and so

$$(S_N g)(1/2) \rightarrow \frac{f(1/2+) + f(1/2-)}{2}.$$

We also have $(S_N s)(1/2) \rightarrow 0$. Thus

$$(S_N f)(1/2) = (S_N g)(1/2) - (S_N s)(1/2) \rightarrow \frac{f(1/2+) + f(1/2-)}{2}. \quad \square$$

2.14 Gibbs Phenomenon

This isn't the end of our discussion about points of discontinuity. There is an interesting phenomenon which occurs locally around the point of

discontinuity. If f is continuous locally around a discontinuity point t_0 , $S_N f \rightarrow f$ pointwise locally around t_0 . Thus, being continuous, $S_N f$ must ‘jump’ from $(S_N f)(t_0-)$ to $(S_N f)(t_0+)$ locally around t_0 . Interestingly enough, we find that the jump is not precise, the jump is overshoot and then must be corrected to the left and right of t_0 . This is known as the *Gibb’s phenomenon*, after the man who clarified the reason for why this phenomenon occurred in physical measurements where first thought to be a defect in the equipment used to take the measurements. Gibb’s phenomenon is one instance where a series of functions $\{f_k\}$ converges pointwise to some function f , whereas qualitatively with respect to the L^∞ norm, the sequence $\{f_k\}$ does not converge to f .

Theorem 2.28. *Given f with finitely many discontinuity points and with $\hat{f}(n) = O(|n|^{-1})$, in particular one at t_0 , we find*

$$\lim_{N \rightarrow \infty} (S_N f)(t_0 \pm 1/N) = f(t_0 \pm) \pm C \cdot \frac{f(t_0+) - f(t_0-)}{2},$$

where

$$C = 2\pi \int_0^\pi \frac{\sin x}{x} \approx 16.610.$$

Proof. First consider the jump function s , with $t_0 = 1/2$. Then

$$(S_N s)(1/2 + 1/N) = -2 \sum_{n=1}^N \frac{\sin(2\pi n/N)}{n} = -2 \sum_{n=1}^N \frac{2\pi}{N} \left(\frac{\sin(2\pi n/N)}{2\pi n/N} \right).$$

Here we’re just taking averages of values of $\sin(x)/x$ at $x = 2\pi/N$, $x = 4\pi/N$, and so on and so forth up to $x = 2\pi$. Thus is a Riemann sum, so as $N \rightarrow \infty$, we get that

$$(S_N s)(\pi + 1/N) \rightarrow -2 \int_0^{2\pi} \frac{\sin x}{x}.$$

The same calculations give

$$(S_N s)(\pi - \pi/N) \rightarrow 2\pi \int_0^\pi \frac{\sin x}{x}.$$

In general, given f , we can write $f = g + \sum \lambda_j h_j$, where g is continuous, and h_j is a translate of the sawtooth function. Then $S_N g$ converges to g

uniformly, and $S_N h_j \rightarrow 0$ for all h_j uniformly in an interval outside of their discontinuity point. To see this, we note that an integration by parts gives

$$\left| \int_{-\pi}^{\pi} D_N(y)[s(x-y) - s(x)] dy \right| \leq |G_N(x - \pi)|,$$

where $G_N(y) = -i \sum_{|n| \leq N} e_n(t)/n$, so $G'_N = D_N$. It now suffices to show $G_N(x - \pi) \rightarrow 0$ outside a neighbourhood of π . But if $A(u, t) = \sum_{|n| \leq u} e_n(t)$, summation by parts gives

$$\sum_{|n| \leq N} \frac{e_n(t)}{n} = \frac{A(N, t)}{N} + \int_1^N \frac{A(u, t)}{u^2}.$$

Now a simple geometric sum shows $A(u, t) \lesssim 1/|e(t) - 1|$, so provided $d(t, 2\pi\mathbf{Z})$ is bounded below, the quantity above tends to zero uniformly. This gives the required result. \square

Chapter 3

Applications of Fourier Series

3.1 Tchebychev Polynomials

If f is everywhere continuous, then for every ε , Fejér's theorem says that we can find N such that $\|\sigma_N(f) - f\| \leq \varepsilon$. But $\sigma_N f$ is just a trigonometric polynomial, and so we have shown that with respect to the L^∞ norm, the space of trigonometric polynomials is dense in the space of all continuous functions. Now if f is a continuous function on $[0, \pi]$, then we can extend it to be even and 2π periodic, and then the trigonometric series $S_N(f)$ of f will be a cosine series, hence $\sigma_N(f)$ will also be a cosine series, and so for each ε , we can find N and coefficients a_1, \dots, a_N such that

$$\left| f(x) - \sum_{n=1}^N a_n \cos(nx) \right| < \varepsilon.$$

Now we use a surprising fact. For each n , there exists a degree n polynomial T_n such that $\cos(nx) = T_n(\cos x)$. This is clear for $n = 0$ and $n = 1$. More generally, we can write

$$\begin{aligned} \cos((m+1)x) &= \cos((m+1)x) + \cos((m-1)x) - \cos((m-1)x) \\ &= \cos(mx+x) + \cos(mx-x) - \cos((m-1)x) \\ &= 2\cos x \cos(mx) - \cos((m-1)x). \end{aligned}$$

Thus we have the relation $T_{m+1}(x) = 2xT_m(x) - T_{m-1}(x)$. These polynomials are known as Tchebyshev polynomials, enabling us to move between 'periodic coordinates' and standard Euclidean coordinates.

Corollary 3.1 (Weirstrass). *The polynomials are uniformly dense in $C[0, 1]$.*

Proof. If f is a continuous function on $[0, 1]$, we can define $g(t) = f(|\cos(t)|)$. Then g is even, and so for every $\varepsilon > 0$, we can find a_1, \dots, a_N such that

$$\left| g(t) - \sum_{n=1}^N a_n \cos(nt) \right| = \left| g(t) - \sum_{n=1}^N a_n T_n(\cos t) \right| < \varepsilon.$$

But if $x = \cos t$, for $\cos t \geq 0$, this equation says

$$\left| f(x) - \sum_{n=1}^N a_n T_n(x) \right| < \varepsilon,$$

and so we have uniformly approximated f by a polynomial. \square

Another proof uses the family of *Landau kernels*

$$L_n(x) = c_n \cdot \begin{cases} (1 - x^2)^n & : -1 \leq x \leq 1 \\ 0 & : |x| \geq 1 \end{cases}$$

where c_n is chosen so that $\int_{-1}^1 L_n(x) dx = 1$. It is simple to show that the family $\{L_n\}$ is a

$$\begin{aligned} \int_{-1}^1 (1 - x^2)^n dx &\geq \sum_{k=0}^{\infty} \int_{1/2^{k+1} \leq |x| \leq 1/2^k} (1 - x^2)^n \\ &\gtrsim \sum_{k=0}^{\infty} \frac{(1 - 1/4^{k+1})^n}{2^k} \\ &= \sum_{k=0}^{\infty} \exp \left(n \log(1 - 1/4^{k+1}) - k \log(2) \right) \\ &\geq \sum_{k=0}^{\infty} \exp \left(-n/4^k \right) / 2^k \\ &\gtrsim \sum_{k=\log_4 n}^{\infty} 1/2^k \gtrsim 1/2^{\log_4 n} = 1/n^{1/2} \end{aligned}$$

Thus $\|L_n\|_{L^\infty(\mathbb{R})} \leq n^{1/2}$ which can be used to show the family $\{L_n\}$ is an approximation to the identity. An important fact here is that if f is supported on $[-1/2, 1/2]$, then $L_N * f$ agrees with a polynomial on $[-1/2, 1/2]$, which can be used to approximate f by a polynomial on this region.

3.2 Exponential Sums and Equidistribution

The next result uses Fourier analysis to characterize the asymptotic distribution of a certain sequence a_1, a_2, \dots . In particular, it is most useful in determining when this distribution is distributed when we consider $2\pi a_1, 2\pi a_2, \dots$ as elements of \mathbf{T} , i.e. so we only care about the fractional part of the numbers, or in other terms their behaviour modulo one. We say the sequence is *uniformly distributed* if for any interval $I \subset \mathbf{T}$, $\#\{2\pi a_n \in I : n \leq N\} \sim N|I|$ as $N \rightarrow \infty$. By approximating continuous functions by step functions, this implies that if $f : \mathbf{T} \rightarrow \mathbf{C}$ is continuous, then

$$\frac{f(2\pi a_1) + \dots + f(2\pi a_N)}{N} \rightarrow \int_{\mathbf{T}} f(t) dt.$$

It is the right hand side to which we can apply Fourier summation to obtain a very useful condition. We let $S_N f$ denote the left hand side of the equation, and Tf the right hand side.

Theorem 3.2 (Weyl Condition). *A sequence $a_1, a_2, \dots \in \mathbf{T}$ is uniformly distributed if and only if for every n , as $N \rightarrow \infty$, $e_n(2\pi a_1) + \dots + e_n(2\pi a_N) = o(N)$.*

Proof. The condition in the theorem implies that for any trigonometric polynomial f , $S_N f \rightarrow Tf$. The S_N are uniformly bounded as functions on $L^\infty(\mathbf{T})$, and T is a bounded functional on this space as well. But this means that $\lim S_N f = Tf$ for all f in $C(\mathbf{T})$, since this equation holds on the dense subset of trigonometric polynomials. \square

This technique enables us to completely characterize the equidistribution behaviour of arithmetic sequences. Given a particular γ , we consider the equidistribution of the sequence $\gamma, 2\gamma, \dots$, which depends on the irrationality of γ .

Example. *Let γ be an arbitrary real number. Then for any n , if $e_n(2\pi\gamma) \neq 1$,*

$$\sum_{m=1}^N e_n(2\pi m\gamma) = \frac{e_n(2\pi(N+1)\gamma) - 1}{e_n(2\pi\gamma) - 1} \lesssim 1 = o(N).$$

If γ is an irrational number, then $e_n(2\pi\gamma) \neq 1$ for all n , which implies that $\gamma, 2\gamma, \dots$ is equidistributed. Conversely, if $e_n(2\pi\gamma) = 1$ for some n , we have

$$\sum_{m=1}^N e_n(a_m) = N.$$

which is not $o(N)$, so the sequence $\gamma, 2\gamma, \dots$ is not equidistributed. If γ is rational, there certainly is n such that $n\gamma \in \mathbf{Z}$, and so $e_n(2\pi\gamma) = 1$.

On the other hand, it is still an open research to characterize, for which γ the sequence $\gamma, \gamma^2, \gamma^3, \dots$ is equidistributed. Here is an example showing that there are γ for which the sequence is not equidistributed.

Example. Let γ be the golden ratio $(1 + \sqrt{5})/2$. Consider the sequence

$$a_n = \left(\frac{1 + \sqrt{5}}{2}\right)^n + \left(\frac{1 - \sqrt{5}}{2}\right)^n = b_n + c_n.$$

Then one checks that a_n is a kind of Fibonacci sequence, with $a_{n+1} = a_n + a_{n-1}$, and initial conditions $a_0 = 2, a_1 = 1$. One checks that c_n is always negative for odd n , and positive for even n , and tends to zero as $n \rightarrow \infty$. Since a_n is an integer, this means that $d(b_n, \mathbf{Z}) = d(\gamma^n, \mathbf{Z}) \rightarrow 0$. But this means that the average distribution of the γ^n modulo one is concentrated at the origin.

3.3 The Isoperimetric Inequality

TODO

3.4 The Poisson Equation

Consider Poisson's equation

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

on the unit disk. Solutions are called *harmonic*. We can reduce this equation to a problem about Fourier series by writing

$$u(re^{2\pi it}) = \sum_{n=0}^{\infty} a_n(r) e^{2\pi nit}.$$

We consider a boundary condition, that $u(e^{2\pi it}) = f(t)$ for some function $f(t)$ on \mathbf{T} . Working formally, noting that in radial coordinates,

$$\Delta u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial t^2}$$

and then taking Fourier series on each side, we find that for each $n \in \mathbf{Z}$,

$$a_n''(r) + a_n'(r)/r - 4\pi^2 n^2 a_n(r)/r^2 = 0.$$

The only *bounded* solution to this differential equation subject to the initial condition $a_n(1) = \hat{f}(n)$ is $a_n(r) = \hat{f}(n)r^{|n|}$. Thus we might guess that

$$u(re^{2\pi it}) = \sum_{n \in \mathbf{Z}} \hat{f}(n)r^{|n|}e^{2\pi nit} = (P_r * f)(x),$$

where P_r is the Poisson kernel. Working backwards through this calculation shows that if $f \in L^1(\mathbf{T})$, then the function $u(re^{2\pi it}) = (P_r * f)(t)$ lies in $C^\infty(\mathbf{D})$ and

$$\lim_{r \rightarrow 1} \int_{\mathbf{T}} |u(re^{2\pi it}) - f(t)| dt = 0.$$

The next theorem shows this is the *only* harmonic function with this property.

Theorem 3.3. *Suppose $f \in L^1(\mathbf{T})$. Then the function $u : \mathbf{D}^\circ \rightarrow \mathbf{C}$ defined for $r > 0$ and $t \in \mathbf{T}$ by setting*

$$u(re^{2\pi it}) = (A_r f)(t)$$

is the unique harmonic function in $C^2(\mathbf{D}^\circ)$ such that

$$\lim_{r \rightarrow 1} \int_{\mathbf{T}} |u(re^{2\pi it}) - f(t)| dt = 0.$$

Proof. Suppose $u \in C^2(\mathbf{D})$ is harmonic. Then we can find functions $a_n(r)$ for each $n \in \mathbf{Z}$ such that

$$u(re^{it}) = \sum_{n=-\infty}^{\infty} a_n(r)e_n(t),$$

where

$$a_n(r) = \int_{\mathbf{T}} u(re^{it}) \overline{e_n(t)} dt.$$

Because $u \in C^2(\mathbf{D})$, we see that $a_n \in C^2((0,1))$ and $a_n(r)$ is bounded as $r \rightarrow 0$. Interchanging integrals shows that

$$a_n''(r) + (1/r)a_n'(r) - (n^2/r^2)a_n(r) = 0.$$

This is an ordinary differential equation, whose only bounded solutions are given by $a_n(r) = A_n r^{|n|}$. If $u(re^{it}) \rightarrow f$ in the L^1 norm as $r \rightarrow 1$, then we conclude

$$A_n = \lim_{r \rightarrow 1} \int_{\mathbf{T}} u(re^{it}) \overline{e_n(t)} dt = \int_{\mathbf{T}} f(t) \overline{e_n(t)} dt = \hat{f}(n),$$

so

$$u(re^{it}) = \sum \hat{f}(n) r^{|n|} e_n(t). \quad \square$$

In particular, the theorem above gives us a map from $L^1(\mathbf{T})$ to the space of harmonic functions on the interior of the unit disk. This is a very handy idea in classical harmonic analysis, and is exploited to it's fullest extent in the theory of Hardy spaces.

3.5 The Heat Equation on a Torus

Recall the heat equation. We are given an initial temperature distribution on \mathbf{T}^d . We wish to study the propagation of this temperature over time. If we let $u(x, t)$ denote the temperature density at $x \in \mathbf{T}^d$ and at time t , then this temperature evolves under the heat equation

$$\frac{\partial u}{\partial t} = \Delta u.$$

We let $f(x) = u(x, 0)$ denote the initial heat distribution. To solve this heat equation, we expand u in a Fourier series, i.e. writing

$$u(x, t) = \sum_{n \in \mathbf{Z}^d} a_n(t) e^{2\pi i n \cdot x}.$$

We then formally find that for each $n \in \mathbf{Z}^d$,

$$a'_n(t) = -4\pi^2 |n|^2 a_n(t),$$

which we can solve to give

$$a_n(t) = \hat{f}(n) e^{-4\pi^2 |n|^2 t}.$$

In particular, we would expect the solution to the heat equation would be given by letting

$$u(x, t) = \sum_{n \in \mathbf{Z}^d} \widehat{f}(n) e^{-4\pi^2 |n|^2 t} e^{2\pi n i t}.$$

As with Poisson's equation on the disk, we can write this as

$$u(x, t) = (H_t * f)(x)$$

where H_t is the *heat kernel*

$$H_t(x) = \sum_{n \in \mathbf{Z}^d} e^{-4\pi^2 |n|^2 t} e^{2\pi n i t}.$$

The rapid convergence of this sum implies that $H_t \in C^\infty(\mathbf{T}^d)$ and that $\widehat{H_t}(n) = e^{-4\pi^2 |n|^2 t}$ for each $n \in \mathbf{Z}^d$. To study this partial differential equation, it suffices to study the heat kernel H_t . Unlike in the case of the Poisson kernel however, we have no explicit formula for the heat kernel, which makes the kernel a little harder to work with.

Lemma 3.4. *The family $\{H_t : t > 0\}$ is an approximation to the identity.*

Proof. The Poisson summation formula implies that

$$H_t(x) = \frac{1}{(4\pi t)^{d/2}} \sum_{n \in \mathbf{Z}^d} e^{-|x+n|^2/4t}.$$

This shows that $H_t(x) \geq 0$, and that

$$\int_{\mathbf{T}^d} H_t(x) = \frac{1}{(4\pi t)^{d/2}} \int_{\mathbf{R}^d} e^{-|x|^2/4t} dx = \int_{\mathbf{R}^d} e^{-\pi |x|^2} dx = 1.$$

We claim that for $|x| \leq 1/2$,

$$\left| H_t(x) - \frac{e^{-x^2/4t}}{(4\pi t)^{d/2}} \right| \lesssim_d e^{-c/t},$$

where $c > 0$ is a universal constant. To prove this, we note this difference is equal to

$$\begin{aligned}
(4\pi t)^{-d/2} \left| \sum_{n \neq 0} e^{-|x+n|^2/4t} \right| &\lesssim t^{-d/2} \sum_{n \neq 0} e^{-c'|n|^2/4t} \\
&\lesssim t^{-d/2} e^{-c'/2t} \sum_{n \neq 0} e^{-c'|n|^2/2} \\
&\lesssim_d t^{-d/2} e^{-c'/2t} \lesssim_d e^{-c/t}.
\end{aligned}$$

This implies that for any fixed $\delta > 0$,

$$\begin{aligned}
\int_{|x|>\delta} H_t(x) &\lesssim t^{-d/2} \int_{|x|>\delta} e^{-|x|^2/4t} dx + e^{-c/t} \\
&\lesssim_d t^{-d/2} e^{-\delta^2/4t} + e^{-c/t}
\end{aligned}$$

which tends to zero as $t \rightarrow \infty$. Thus we have proved that H_t is an approximation to the identity. \square

Theorem 3.5. *For any $f \in L^1(\mathbf{T}^d)$, for $1 \leq p < \infty$. Then the function*

$$u(x, t) = (H_t * f)(x)$$

lies in $C^\infty(\mathbf{T}^d \times (0, \infty))$, and for $t > 0$ solves the heat equation. Moreover, u is the unique solution to the heat equation in $C^2(\mathbf{T}^d \times (0, \infty))$ such that

$$\lim_{t \rightarrow 0^+} \int_{\mathbf{T}^d} |u(t, x) - f(x)| dx = 0.$$

Proof. We have already shown the former statement by the fact that $\{H_t : t > 0\}$ is an approximation to the identity. To prove the latter statement, given $u \in C^2(\mathbf{T}^d \times (0, \infty))$, we can take a Fourier series, letting

$$a_n(t) = \int_{\mathbf{T}^d} u(x, t) e^{-2\pi i n \cdot x} dx.$$

Then $a_n \in C^2((0, \infty))$ and differentiation under the integral sign shows that $a'_n(t) = -4\pi^2 a_n(t)$, so that $a_n(t) = c_n e^{-4\pi^2 t}$ for some c_n . But $a_n(t) \rightarrow \hat{f}(n)$ as $t \rightarrow 0$ uniformly in n by the convergence assumption, so $c_n = \hat{f}(n)$. But this implies that $u(x, t) = (H_t * f)(x)$ for each $x \in \mathbf{T}^d$, since both sides have the same Fourier series for all $t > 0$. \square

Chapter 4

The Fourier Transform

In the last few chapters, we discussed the role of analyzing the frequency decomposition of a periodic function on the real line. In this chapter, we explore the ways in which we may extend this construction to perform frequency analysis for not necessarily periodic functions on the real line, and more generally, in higher dimensional Euclidean space. The only periodic trigonometric functions on $[0, 1]$ on the real line had integer frequencies of the form $2\pi n$, whereas on the real line periodic functions can have frequencies corresponding to any real number. The analogue of the discrete Fourier series formula

$$f(x) = \sum_{k=-\infty}^{\infty} \hat{f}(k) e^{2\pi i k x}$$

is the Fourier inversion formula

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x} d\xi,$$

where for each real number ξ , we define

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i \xi x} dx.$$

The function \hat{f} is known as the *Fourier transform* of the function f . It is also denoted by $\mathcal{F}(f)$. The role to which we can justify this formula is the main focus of this chapter. The fact that \mathbf{R} is non-compact and has infinite

measure adds some difficulty to the study of the Fourier transform over the Fourier series. For instance, since $L^p(\mathbf{R}^d)$ is not included in $L^q(\mathbf{R}^d)$ for $p \neq q$, which makes it more difficult to perform a qualitative analysis of convergence in this setting. Nonetheless, the Fourier transform has many properties as the Fourier series. We add an additional difficulty by also analyzing the Fourier transform on \mathbf{R}^d , which, given $f : \mathbf{R}^d \rightarrow \mathbf{C}$, considers the quantities

$$f(x) \sim \int_{\mathbf{R}^d} \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi, \quad \text{where} \quad \hat{f}(\xi) = \int_{\mathbf{R}^d} f(x) e^{-2\pi i \xi \cdot x} dx$$

for $\xi \in \mathbf{R}^d$. The basic theory of the Fourier transform in one dimension is essentially the same as the theory of the Fourier transform in d dimensions, though as d increases certain more technical considerations such as pointwise convergence become more difficult to understand.

4.1 Basic Calculations

In order to interpret the Fourier transform as an absolutely convergent integral, we require that we are dealing with integrable assumptions. Thus we analyze functions in $L^1(\mathbf{R}^d)$. During arguments, we can often assume additional regularity properties of f , and then apply density arguments to get the result in general. Most of the properties of the Fourier transform are exactly the same as for Fourier series. However, one novel phenomenon in the basic theory is that the Fourier transform of an integrable function is continuous and vanishes at ∞ .

Theorem 4.1. *For any $f \in L^1(\mathbf{R}^d)$, $\|\hat{f}\|_{L^\infty(\mathbf{R}^d)} \leq \|f\|_{L^1(\mathbf{R}^d)}$, and $\hat{f} \in C_0(\mathbf{R}^d)$.*

Proof. For any $\xi \in \mathbf{R}^d$,

$$|\hat{f}(\xi)| = \left| \int f(x) e(-\xi \cdot x) dx \right| \leq \int |f(x)| |e(-\xi \cdot x)| dx = \|f\|_{L^1(\mathbf{R}^d)}.$$

If χ_I is the characteristic function of an n dimensional box, i.e.

$$I = [a_1, b_1] \times \cdots \times [a_n, b_n] = I_1 \times \cdots \times I_n,$$

then

$$\widehat{\chi_I}(\xi) = \int_I e(-\xi \cdot x) = \prod_{k=1}^n \int_{a_k}^{b_k} e(-\xi_k x_k) = \prod_{k=1}^n \widehat{\chi_{I_k}}(\xi_k).$$

where

$$\widehat{\chi}_{I_k}(\xi_k) = \begin{cases} \frac{e(-\xi_k a_k) - e(-\xi_k b_k)}{2\pi i \xi_k} & \xi_k \neq 0, \\ b_k - a_k & \xi_k = 0. \end{cases}$$

L'Hopital's rule shows $\widehat{\chi}_{I_k}$ is a continuous function. We also have the upper bound

$$\widehat{\chi}_{I_k}(\xi_k) \lesssim_{I_k} (1 + |\xi_k|)^{-1}$$

for all $\xi_k \in \mathbf{R}$, which implies that

$$\widehat{\chi}_I(\xi) = \prod \widehat{\chi}_{I_k}(\xi_k) \lesssim_I \prod \frac{1}{1 + |\xi_k|} \lesssim_n \frac{1}{1 + |\xi|}.$$

Thus $\widehat{\chi}_I(\xi) \rightarrow 0$ as $|\xi| \rightarrow \infty$. But this implies the Fourier transform of any step function is continuous and vanishes at ∞ . Since step functions are dense in $L^1(\mathbf{R}^d)$, a density argument then gives the result for all integrable functions. \square

Elementary properties of integration give the following relations among the Fourier transforms of functions on \mathbf{R}^d . They are strongly related to the translation invariance of the Lebesgue integral on \mathbf{R}^d :

- If $f^*(x) = \overline{f(x)}$ is the conjugate of a function f , then

$$\widehat{f^*}(\xi) = \int \overline{f(x)} e^{-2\pi i x \cdot \xi} dx = \overline{\int f(x) e^{2\pi i \xi \cdot x} dx} = \widehat{f}(-\xi).$$

If f is real, the formula above says $\widehat{f}(\xi) = \overline{\widehat{f}(-\xi)}$, and so if we define $a(\xi) = \operatorname{Re}(\widehat{f}(\xi))$, $b(\xi) = \operatorname{Im}(\widehat{f}(\xi))$, then formally we have

$$\int_{-\infty}^{\infty} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi = 2 \int_0^{\infty} a(\xi) \cos(2\pi \xi \cdot x) - b(\xi) \sin(2\pi \xi \cdot x) d\xi.$$

Thus the Fourier representation formula expresses the function f as an integral in sines and cosines.

- There is a duality between translation and frequency modulation. For $y \in \mathbf{R}^d$, we define $(\operatorname{Trans}_y f)(x) = f(x - y)$. If $\xi \in \mathbf{R}^d$, then we define $(\operatorname{Mod}_\xi f)(x) = e^{2\pi i \xi \cdot x} f(x)$. We then find that

$$\begin{aligned} \widehat{\operatorname{Trans}_y f}(\xi) &= \int f(x - y) e^{-2\pi i \xi \cdot x} dx \\ &= e^{-2\pi i \xi \cdot y} \int f(x) e^{-2\pi i \xi \cdot x} dx = (\operatorname{Mod}_{-y} \widehat{f})(\xi). \end{aligned}$$

and

$$\widehat{\text{Mod}_\xi f}(\eta) = \int e^{2\pi i \xi \cdot x} f(x) e(-\eta \cdot x) dx = \widehat{f}(\eta - \xi) = (\text{Trans}_\xi \widehat{f})(\eta).$$

Thus we conclude $\mathcal{F} \circ \text{Trans}_y = \text{Mod}_{-y} \circ \mathcal{F}$, and $\mathcal{F} \circ \text{Mod}_\xi = \text{Trans}_\xi \circ \mathcal{F}$.

- A very related property to the translational symmetry of the Fourier transform is related to the convolution

$$(f * g)(x) = \int f(y) g(x - y) dy$$

of two functions $f, g \in L^1(\mathbf{R}^d)$. This convolution possesses precisely the same properties as convolution on \mathbf{T} . Most importantly for us,

$$\mathcal{F}(f * g) = \mathcal{F}(f) \cdot \mathcal{F}(g),$$

so convolution in phase space is just a product in frequency space.

Another related property to the translation symmetry is that the Fourier transform behaves well with respect to differentiation. If $f \in L^1(\mathbf{R}^d)$ has a weak derivative $D^\alpha f \in L^1(\mathbf{R}^d)$, then

$$\widehat{D^\alpha f}(\xi) = (2\pi i \xi)^\alpha \widehat{f}(\xi).$$

In particular, this is true if f is a *Schwartz function*, i.e. an element of

$$\mathcal{S}(\mathbf{R}^d) = \{f \in C^\infty(\mathbf{R}^d) : |(D_\alpha f)(x)| \lesssim_{\alpha, N} |x|^{-N} \text{ for all } N, \alpha, x\}$$

which is often a natural place to consider the Fourier transform. Conversely, if $f \in L^1(\mathbf{R}^d)$, and $x^\alpha f \in L^1(\mathbf{R}^d)$ for some multi-index α , then \widehat{f} has a weak derivative $D^\alpha \widehat{f}$ in $L^1(\mathbf{R}^d)$, and

$$D^\alpha \widehat{f}(\xi) = (-2\pi i x)^\alpha f(\xi).$$

In particular, this means that the Fourier transform of a compactly supported element of $L^1(\mathbf{R}^d)$ lies in $C^\infty(\mathbf{R}^d)$, and all derivatives of the Fourier transform are integrable.

- Let $T : \mathbf{R}^d \rightarrow \mathbf{R}^d$ be an invertible linear transformation. Then a change of variables $y = Tx$ gives

$$\begin{aligned}
\widehat{f \circ T}(\xi) &= \int f(Tx) e^{-2\pi i \xi \cdot x} dx \\
&= \frac{1}{|\det(T)|} \int f(y) e^{-2\pi i \xi \cdot T^{-1}y} dy \\
&= \frac{1}{|\det(T)|} \int f(y) e^{-2\pi i T^{-T} \xi \cdot y} dy \\
&= \frac{1}{|\det(T)|} (\widehat{f} \circ T^{-T})(\xi).
\end{aligned}$$

Thus we conclude that if $T^* : L^1(\mathbf{R}^d) \rightarrow L^1(\mathbf{R}^d)$ is the operator defined by setting $T^*(f) = f \circ T$, then

$$\mathcal{F} \circ T^* = \frac{1}{|\det(T)|} \cdot (T^{-T})^* \circ \mathcal{F}.$$

This property indicates the ‘cotangent’ and symplectic properties of the Fourier transform. One way to think of this property is that one can think of the frequency variable as ‘cotangent’ to the spatial variable, since if we have a coordinate change $y = Tx$, and we define the ‘coordinatized’ Fourier transforms \mathcal{F}_x and \mathcal{F}_y by setting

$$\mathcal{F}_x f(\xi) = \int f(x) e^{-2\pi i x \cdot \xi} dx$$

and

$$\mathcal{F}_y f(\eta) = \int f(T^{-1}y) e^{-2\pi i y \cdot \eta} dy,$$

then the transformation formula tells us that

$$\mathcal{F}_y = \mathcal{F}_x \circ (T^{-1})^* = |\det(T)| \cdot (T^T)^* \circ \mathcal{F}_x,$$

i.e. $\mathcal{F}_y f(\eta) = |\det(T)| \cdot \mathcal{F}_x f(\xi)$, where $\eta = T^T \xi$. By interpreting ξ and η as *cotangent* vectors to the x and y coordinates respectively, one can therefore use this symmetry property to define a version of the Fourier transform that is invariant under area preserving changes of coordinates.

- As a special case of the theorem above, if $a \in \mathbf{R}$ and $\text{Dil}_a : L^1(\mathbf{R}^d) \rightarrow L^1(\mathbf{R}^d)$ is the operator defined by setting

$$(\text{Dil}_a f)(x) = f(x/a),$$

then

$$\widehat{\text{Dil}_a f} = a^d \cdot \text{Dil}_{1/a} \hat{f}$$

As we increase a , the values of f are traced over more quickly, and so it is natural for the support of f to lie over larger frequencies. Note that this dilation preserves the L^∞ norm of f , but the Fourier transform preserves the L^1 norm. We could have alternatively consider the L^1 preserving dilation $f \mapsto a^{-d} f(x/a)$, which on the frequency side of things preserves the L^∞ norm. We can consider various magnitude adjustments; for instance, $f \mapsto a^{-d/2} f(x/a)$ preserves the L^2 norm in both space and frequency. But regardless, as we concentrate f in a smaller neighborhood, \hat{f} is dilated so its support lies in a larger and larger neighborhood.

- Another special case is that if $R \in O_n(\mathbf{R})$, then $\widehat{f \circ R}(\xi) = \hat{f}(R\xi)$, i.e. $\mathcal{F} \circ R^* = R^* \circ \mathcal{F}$. In particular, if f is a radial function, so $f \circ R = f$ for any R , then $\hat{f}(R\xi) = \hat{f}(\xi)$ for any $R \in O_n(\mathbf{R})$, so \hat{f} is also a radial function. If f is even, so $f(x) = f(-x)$ for all x , then $\hat{f}(\xi) = \hat{f}(-\xi)$ for all ξ , so \hat{f} is even. Similarly, if f is odd, then \hat{f} is odd. In particular, the space of radial functions is an *invariant subspace* with respect to the Fourier transform, which becomes important in more advanced theories, such as the theory of spherical harmonics.

4.2 The Fourier Algebra

The space

$$\mathbf{A}(\mathbf{R}^d) = \left\{ \hat{f} : f \in L^1(\mathbf{R}^d) \right\}$$

is called the *Fourier algebra*. The last theorem shows $\mathbf{A}(\mathbf{R}^d) \subset C_0(\mathbf{R}^d)$, but it is *not* the case that $\mathbf{A}(\mathbf{R}^d) = C_0(\mathbf{R}^d)$. Current research cannot give a satisfactory description of the elements of $\mathbf{A}(\mathbf{R}^d)$, and a simple characterization is unlikely. The next lemma will be used to show $\mathbf{A}(\mathbf{R}^d) \neq C_0(\mathbf{R}^d)$.

Lemma 4.2. For any $0 \leq a < b < \infty$, independantly of a and b ,

$$\left| \int_a^b \frac{\sin x}{x} \right| = O(1).$$

Proof. Since $\|\sin(x)/x\|_{L^\infty(\mathbf{R})} \leq 1$, we may assume $b > 1$, for otherwise we obtain a trivial bound. This also implies

$$\left| \int_a^b \frac{\sin x}{x} dx \right| \leq 1 + \left| \int_1^b \frac{\sin x}{x} dx \right|.$$

An integration by parts then shows that

$$\left| \int_1^b \frac{\sin x}{x} dx \right| \leq \left| \left(\cos 1 - \frac{\cos b}{b} \right) \right| + \left| \int_1^b \frac{\cos x}{x^2} dx \right| \lesssim 1. \quad \square$$

Theorem 4.3. $\mathbf{A}(\mathbf{R}) \neq C_0(\mathbf{R})$. In particular, $\mathbf{A}(\mathbf{R})$ does not contain any odd functions g in $C_0(\mathbf{R})$ such that

$$\limsup_{b \rightarrow \infty} \left| \int_1^b \frac{g(\xi)}{\xi} d\xi \right| = \infty.$$

Proof. Suppose $f \in L^1(\mathbf{R})$, and $\hat{f} \in C_0(\mathbf{R})$ is an odd function. Then we know

$$\hat{f}(\xi) = -i \int_{-\infty}^{\infty} f(x) \sin(2\pi \xi x) dx.$$

If $b \geq 1$, an application of Fubini's theorem shows that

$$\left| \int_1^b \frac{\hat{f}(\xi)}{\xi} d\xi \right| = \left| \int_{-\infty}^{\infty} f(x) \left(\int_1^b \frac{\sin(2\pi \xi x)}{\xi} d\xi \right) dx \right|.$$

But

$$\left| \int_1^b \frac{\sin(2\pi \xi x)}{\xi} d\xi \right| = \left| \int_{2\pi x}^{2\pi b x} \frac{\sin \xi}{\xi} d\xi \right| \lesssim 1.$$

Thus we obtain that

$$\left| \int_1^b \frac{\hat{f}(\xi)}{\xi} d\xi \right| \lesssim \|f\|_{L^1(\mathbf{R})}.$$

For instance, this implies that there is no $f \in L^1(\mathbf{R})$ such that

$$\hat{f}(\xi) = \operatorname{sgn}(\xi) \frac{|\sin(2\pi\xi)|}{\log|\xi|}$$

for all $\xi \in \mathbf{R}$, since

$$\lim_{b \rightarrow \infty} \int_1^b \frac{|\sin(2\pi\xi)|}{\xi \log|\xi|} = \infty. \quad \square$$

On the other hand, for a finite measure μ on \mathbf{R}^d , we can define the Fourier transform to be the continuous function

$$\hat{\mu}(\xi) = \int_{\mathbf{R}^d} e^{-2\pi i \xi \cdot x} d\mu(x).$$

In this case, the family of continuous functions which are the Fourier transforms of finite measures is precisely the family of $f \in C(\mathbf{R}^d)$ which are *positive definite*, in the sense that for each $x_1, \dots, x_N \in \mathbf{R}^d$ and $\xi_1, \dots, \xi_N \in \mathbf{C}$,

$$\sum_{i=1}^N \sum_{j=1}^N f(x_i - x_j) \xi_i \bar{\xi}_j \geq 0.$$

The theorem, proved by Bochner, is best addressed in the more general case of harmonic analysis on locally compact abelian groups, and so we leave the proof of this for another time.

4.3 Basic Convergence Properties

As we might expect from the Fourier series theory, if $f \in C(\mathbf{R}) \cap L^1(\mathbf{R}^d)$ and $\hat{f} \in L^1(\mathbf{R}^d)$, then the formula

$$f(x) = \int_{\mathbf{R}^d} \hat{f}(\xi) e(\xi \cdot x) dx$$

holds for all $x \in \mathbf{R}^d$. Unlike in the case of the Fourier series, we cannot test our function against trigonometric polynomials. On the other hand, we have a multiplication formula which often comes in useful.

Theorem 4.4 (The Multiplication Formula). *If $f, g \in L^1(\mathbf{R}^d)$,*

$$\int f(x) \hat{g}(x) dx = \int \hat{f}(\xi) g(\xi) d\xi.$$

Proof. If $f, g \in L^1(\mathbf{R}^d)$, then \widehat{f} and \widehat{g} are bounded, continuous functions on \mathbf{R}^d . In particular, $\widehat{f}g$ and $f\widehat{g}$ are integrable. A simple use of Fubini's theorem gives

$$\int f(x)\widehat{g}(x) dx = \int \int f(x)g(\xi)e(-\xi \cdot x) dx d\xi = \int g(\xi)\widehat{f}(\xi) d\xi. \quad \square$$

In particular, if $f \in L^1(\mathbf{R}^d)$ and $\widehat{f} = 0$, then for any $g \in L^1(\mathbf{R}^d)$,

$$\int_{\mathbf{R}^d} f(x)\widehat{g}(x) dx = 0.$$

If x_0 is a continuity point of f , then it suffices to choose a function g such that the majority of the mass of \widehat{g} is concentrated at the point x_0 . A natural choice here is to use a *Gaussian function*.

Let $g(x) = e^{-\pi x^2}$. Then $g \in L^1(\mathbf{R})$. Then $g'(x) = -2\pi xg(x)$, and since $xg \in L^1(\mathbf{R})$, we conclude that

$$\frac{d\widehat{g}(\xi)}{d\xi} = -2\pi\xi\widehat{g}(\xi).$$

Since $\widehat{g}(0) = \int_{-\infty}^{\infty} e^{-\pi x^2} dx = 1$, we conclude from solving the ordinary differential equation that

$$\widehat{g}(\xi) = e^{-\pi\xi^2} = g(\xi).$$

Tensorizing, it follows that if $g(x) = e^{-\pi|x|^2}$ is the element of $L^1(\mathbf{R}^d)$, then

$$\widehat{g}(\xi) = e^{-\pi|\xi|^2} = g(x).$$

In particular, if for $x_0 \in \mathbf{R}^d$ and $\delta > 0$, we define

$$g_{x_0, \delta}(\xi) = e^{-2\pi i x_0 \cdot \xi} g(\delta\xi)$$

then the symmetries of the Fourier transform imply that

$$\widehat{g_{x_0, \delta}}(\xi) = \delta^{-d} e^{-(\pi/\delta^2)|\xi - x_0|^2}.$$

Thus we conclude that if $\widehat{f} = 0$, then for any x_0 and δ ,

$$\delta^{-d} \int_{\mathbf{R}^d} f(x) e^{-(\pi/\delta^2)|\xi - x_0|^2} dx = 0.$$

A simple approximation as $\delta \rightarrow 0$ gives the following result.

Theorem 4.5. Suppose $f \in L^1(\mathbf{R}^d)$ and $\hat{f} = 0$. Then f vanishes at any of its continuity points. In particular, if $f \in L^1(\mathbf{R}^d) \cap C(\mathbf{R}^d)$ and $\hat{f} = 0$, then $f = 0$.

As in the case of the Fourier series, if $f \in L^1(\mathbf{R}^d)$ and $\hat{f} \in L^1(\mathbf{R}^d)$, then the multiplication formula implies that for any $g \in L^1(\mathbf{R}^d)$ with $\hat{g} \in L^1(\mathbf{R}^d)$,

$$\int_{\mathbf{R}^d} \hat{f}(x)g(x) dx = \int_{\mathbf{R}^d} \hat{f}(\xi)\hat{g}(\xi) d\xi = \int_{\mathbf{R}^d} f(x)\hat{g}(x) dx.$$

If $g = g_{x_0, \delta}$ for some x_0 and δ , then it is simple to calculate that $\hat{g}(x) = g(-x)$. Thus we conclude that for any such function,

$$\int_{\mathbf{R}^d} \hat{f}(x)g(x) dx = \int_{\mathbf{R}^d} f(-x)g(x) dx.$$

In particular, a similar approximation technique to the last theorem gives the Fourier inversion theorem.

Theorem 4.6. Suppose $f \in L^1(\mathbf{R}^d)$ and $\hat{f} \in L^1(\mathbf{R}^d)$. Then for any continuity point x of f ,

$$f(x) = \int_{\mathbf{R}^d} \hat{f}(\xi)e^{2\pi i \xi \cdot x} d\xi.$$

In particular, if we also assume $f \in C(\mathbf{R}^d)$, then the inversion formula holds everywhere.

4.4 Alternative Summation Methods

As with the Fourier series, we can obtain results for more general functions by ‘dampening’ the integration factor. To do this, we consider ‘alternate integral’ methods which can define the integral of a measurable function that is not necessarily absolutely integrable.

Example. Even if f is a non integrable function, the functions $f(x)e^{-\delta|x|}$ may be integrable for $\delta > 0$. If this is the case, we say f is Abel summable to a value A if

$$\lim_{\delta \rightarrow 0} \int_{\mathbf{R}^d} f(x)e^{-\delta|x|} dx = A$$

For each $\delta > 0$ and $f \in L^1(\mathbf{R}^d)$, we let

$$(A_\delta f)(x) = \int_{\mathbf{R}^d} \hat{f}(\xi) e(\xi \cdot x) e^{-\delta|\xi|} d\xi.$$

Thus $A_\delta f$ represents the Abel sums of the Fourier inversion formula.

If $f \in L^1(\mathbf{R}^d)$, then the dominated convergence theorem implies that

$$\int_{\mathbf{R}^d} f(x) e^{-\delta|x|} dx \rightarrow \int_{\mathbf{R}^d} f(x) dx.$$

so f is Abel summable. However, f may be Abel summable even if f is not integrable. For instance, if $f(x) = \sin(x)/x$, then f is not integrable, yet f is Abel summable to π over the real line.

Example. Similarly, we can consider the Gauss sums

$$\int f(x) e^{-\delta|x|^2} dx$$

We say f is Gauss summable to if these values converge as $\delta \rightarrow 0$. For $f \in L^1(\mathbf{R}^d)$, we let

$$(G_\delta f)(x) = \int \hat{f}(\xi) e(\xi \cdot x) e^{-\delta|\xi|^2} d\xi.$$

Then as $\delta \rightarrow 0$, $G_\delta f$ represents the Gauss sums of the Fourier inversion formula.

Example. For $d = 1$, we can also consider the Fejér sums

$$(\sigma_\delta f)(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e(\xi \cdot x) \left(\frac{\sin(\delta\pi\xi)}{\delta\pi\xi} \right)^2 d\xi,$$

which are analogous to the Fejér sums in the periodic setting.

Example. In basic calculus, the integral of a function f over the entire real line is defined as

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx.$$

These integrals can be written as the integral of $f \chi_{[-R,R]}$, and so in a generalized sense, we can integrate a function f if $f \chi_{[-R,R]}$ is integrable for each N , and the integrals of these functions converge as $t \rightarrow \infty$. Thus we study

$$(S_R f)(x) = \int_{-R}^R \hat{f}(\xi) e(\xi \cdot x) d\xi.$$

Abel summability is more general than the piecewise limit integral considered in the last example, as the next lemma proves.

Lemma 4.7. Suppose $f \in L^1_{loc}(\mathbf{R})$, that

$$\lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$$

exists, and that $f(x)e^{-\delta x^2}$ is absolutely integrable for each $\delta > 0$. Then f is Abel summable, and

$$\lim_{\delta \rightarrow 0} \int_{-\infty}^{\infty} f(x)e^{-\delta|x|^2} dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx.$$

Proof. Let

$$\lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx = A.$$

For each $x \geq 0$, write

$$F(x) = \int_{-x}^x f(x) dx.$$

Then F is continuous and differentiable almost everywhere, and $F(x) \rightarrow A$ as $x \rightarrow \infty$. We know that $F'(x) = f(x) + f(-x)$, and an integration by parts gives for each $s > 0$,

$$\begin{aligned} \int_{-s}^s f(x)e^{-\delta x^2} dx &= \int_0^s [f(x) + f(-x)]e^{-\delta x^2} dx \\ &= F(s)e^{-\delta s^2} + 2\delta \int_0^s xF(x)e^{-\delta x^2} dx. \end{aligned}$$

Taking $s \rightarrow \infty$, using the fact that F is bounded so that $F(s)e^{-\delta s^2} \rightarrow 0$, we conclude

$$\int_{-\infty}^{\infty} f(x)e^{-\delta x^2} dx = 2\delta \int_0^{\infty} xF(x)e^{-\delta x^2} dx.$$

Given $\varepsilon > 0$, fix t such that $|F(s) - A| \leq \varepsilon$ for $s \geq t$. Then

$$\begin{aligned} \left| \int f(x)e^{-\delta x^2} dx - A \right| &\leq 2\delta \left| \int_0^t xF(x)e^{-\delta x^2} dx \right| \\ &\quad + 2\delta\varepsilon \left| \int_t^{\infty} xe^{-\delta x^2} dx \right| \\ &\quad + \left| 2\delta A \int_t^{\infty} xe^{-\delta x^2} dx - A \right|. \end{aligned}$$

The first and second components of this upper bound can each be made smaller than ε for small enough δ . And

$$2\delta \int_t^\infty x e^{-\delta x^2} dx = e^{-\delta t^2}$$

So the third term is equal to $|A||1 - e^{-\delta t^2}|$ and so for small enough δ , we can also bound this by ε . Thus we have shown for small enough δ that

$$\left| \int f(x) e^{-\delta x^2} dx - A \right| \leq 3\varepsilon.$$

It now suffices to take $\varepsilon \rightarrow 0$. □

Abel summation is even more general than Gauss summation.

Lemma 4.8. *If f is Gauss summable, and $f(x)e^{-\delta|x|}$ is absolutely integrable for each $\delta > 0$, then f is Abel summable, and*

$$\lim_{\delta \rightarrow 0} \int f(x) e^{-\delta|x|^2} dx = \lim_{\delta \rightarrow 0} \int f(x) e^{-\delta|x|} dx.$$

Proof. Let

$$\lim_{\delta \rightarrow 0} \int f(x) e^{-\delta|x|^2} dx = A.$$

If there existed constants c_n and λ_n such that $e^{-\delta|x|} = \sum c_n e^{-(\lambda_n \delta|x|)^2}$, this theorem would be easy. This is not exactly true, but we do have the *subordination principle*, which says

$$e^{-\delta|x|} = \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} e^{-\delta^2|x|^2/4u} du.$$

This formula, which is proved using basic complex analysis, is shown later on in this chapter. Applying Fubini's theorem, this means that

$$\int f(x) e^{-\delta|x|} dx = \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} \int f(x) e^{-\delta^2|x|^2/4u} dx du.$$

For any fixed $t > 0$, we certainly have

$$\lim_{\delta \rightarrow 0} \int_t^\infty \frac{e^{-u}}{\sqrt{\pi u}} \int f(x) e^{-\delta^2|x|^2/4u} dx du = A \int_t^\infty \frac{e^{-u}}{\sqrt{\pi u}}$$

And this is equal to $A(1 + o(1))$ as $t \rightarrow 0$. And now we calculate

$$\int_0^t \frac{e^{-u}}{\sqrt{\pi u}} \int f(x) e^{-\delta^2 |x|^2 / 4u} du \leq \left\| \frac{e^{-u}}{\sqrt{\pi u}} \right\|_{L^1[0,t]} \left\| \int f(x) e^{-\delta^2 |x|^2 / 4u} \right\|_{L^\infty[0,t]}$$

The left norm tends to zero as $t \rightarrow 0$. And as $u \downarrow 0$, the dominated convergence theorem implies that

$$\int f(x) e^{-\delta |x|^2 / 4u} \rightarrow 0.$$

This completes the proof. □

For any family of functions Φ_δ , we can consider the ‘ Φ sums’

$$\int f(x) \Phi_\delta(x) d\xi$$

and the corresponding Fourier transform operators

$$S_\delta(f, \Phi)(x) = \int \hat{f}(\xi) e(\xi \cdot x) \Phi_\delta(\xi) d\xi.$$

We say f is Φ summable to a value if

$$\int f(x) \Phi_\delta(x) d\xi$$

converges. In all the examples we will consider, we construct Φ sums by fixing a function $\Phi \in C_0(\mathbf{R}^d)$ with $\Phi(0) = 1$, and defining $\Phi_\delta(x) = \Phi(\delta x)$. When this is the case $f(x) \Phi_\delta(x)$ converges to $f(x)$ pointwise for each x as $\delta \rightarrow 0$. Thus if $f \in L^1(\mathbf{R}^d)$, the dominated convergence theorem implies that f is Φ summable to its usual integral. We now use these summability kernels to understand the Fourier summation formula.

Theorem 4.9 (The Multiplication Formula). *If $f, g \in L^1(\mathbf{R}^d)$,*

$$\int f(x) \hat{g}(x) dx = \int \hat{f}(\xi) g(\xi) d\xi.$$

Proof. If $f, g \in L^1(\mathbf{R}^d)$, then \hat{f} and \hat{g} are bounded, continuous functions on \mathbf{R}^d . In particular, $\hat{f}g$ and $f\hat{g}$ are integrable. A simple use of Fubini’s theorem gives

$$\int f(x) \hat{g}(x) dx = \int \int f(x) g(\xi) e(-\xi \cdot x) dx d\xi = \int g(\xi) \hat{f}(\xi) d\xi. \quad \square$$

If Φ is integrable, then the multiplication formula shows

$$\begin{aligned} S_\delta(f, \Phi) &= \int \hat{f}(\xi) e(\xi \cdot x) \Phi(\delta \xi) d\xi \\ &= \int f(x) (\text{Mod}_x(\delta_\delta \Phi))^\wedge(x) dx = \delta^{-n} \int f(x) \cdot \hat{\Phi}\left(\frac{x-y}{\delta}\right) dx. \end{aligned}$$

Thus if we define $K_\delta^\Phi(x) = \delta^{-d} \hat{\Phi}(-x/\delta)$, then $S_\delta(f, \Phi) = K_\delta^\Phi * f$. Thus we have expressed the summation operators as convolution operations.

We now recall some notions of convolution kernels that help us approximate functions. Recall that if a family of kernels $\{K_\delta\}$ satisfies

- For any $\delta > 0$,

$$\int K_\delta(\xi) d\xi = 1.$$

- The values $\{\|K_\delta\|_{L^1(\mathbf{R}^d)}\}$ are uniformly bounded in δ .
- For any $\varepsilon > 0$,

$$\lim_{\delta \rightarrow 0} \int_{|\xi| \geq \varepsilon} |K_\delta(\xi)| d\xi \rightarrow 0.$$

then the family forms a *good kernel*. If this is the case, then $f * K_\delta$ converges to f in the L^p norms if $f \in L^p(\mathbf{R}^d)$, and converges to f uniformly if f is continuous and bounded. If we have the stronger conditions that

- For any $\delta > 0$,

$$\int K_\delta(\xi) d\xi = 1.$$

- $\|K_\delta\|_{L^\infty(\mathbf{R}^d)} \lesssim 1/\delta^d$.
- For any $\delta > 0$ and $\xi \in \mathbf{R}^d$,

$$|K_\delta(\xi)| \lesssim \frac{\delta}{|\xi|^{d+1}}.$$

then the family $\{K_\delta\}$ is an approximation to the identity, and so $(K_\delta * f)(x)$ converges to $f(x)$ for any x in the Lebesgue set of f . For a particular function Φ , the family $\{K_\delta^\Phi\}$ forms a good kernel as $\delta \rightarrow 0$ if $\hat{\Phi} \in L^1(\mathbf{R}^d)$ and

$\int \hat{\Phi}(\xi) d\xi = 1$, and forms an approximation to the identity if we assume in addition that $\Phi \in C^{d+1}(\mathbf{R}^d)$. Thus we conclude that as $\delta \rightarrow 0$, if Φ satisfies the appropriate conditions then as $\delta \rightarrow 0$, the summations $S_\delta(f, \Phi)$ converge to f in the appropriate sense as considered above.

Example. We obtain the Fejér kernel F_δ from the initial function

$$F(x) = \left(\frac{\sin \pi x}{\pi x} \right)^2$$

Using contour integration, we now show

$$\hat{F}(\xi) = \begin{cases} 1 - |\xi| & : |\xi| \leq 1 \\ 0 & : |\xi| > 1 \end{cases}$$

Since this functions is compactly supported, with total mass one, it is easy to see the corresponding Kernel K_δ^F are an approximation to the identity. Thus $\sigma_\delta f$ converges to f in all the manners described above.

Since F is an even function, \hat{F} is even, and so we may assume $\xi \geq 0$. We initially calculate

$$\hat{F}(\xi) = \int_{-\infty}^{\infty} \left(\frac{\sin(\pi x)}{\pi x} \right)^2 e(-\xi x) dx = \frac{1}{\pi} \int_{-\infty}^{\infty} \left(\frac{\sin x}{x} \right)^2 e(-2\xi x) dx.$$

Now we have

$$(\sin z)^2 = \left(\frac{e(z) - e(-z)}{2i} \right)^2 = \frac{(2 - e^{2iz}) - e^{-2iz}}{4}.$$

This means

$$\frac{(\sin z)^2}{z^2} e^{-2i\xi z} = \frac{2e^{-2i\xi z} - e^{-2(\xi+1)iz} - e^{-2(\xi-1)iz}}{4z^2} = \frac{f_\xi(z) + g_\xi(z)}{4}.$$

For $\xi \geq 0$, $f_\xi(z)$ is $O_\xi(1/|z|^2)$ in the lower half plane, because if $\text{Im}(z) \leq 0$,

$$|2e^{-2i\xi z} - e^{-2(\xi+1)iz}| \leq 2e^{2\xi} + e^{2(\xi+1)} = O_\xi(1).$$

For $\xi \geq 1$, $g_\xi(z)$ is also $O_\xi(1/|z|^2)$ in the lower half plane, because

$$|e^{-2(\xi-1)iz}| \leq e^{2(\xi-1)}.$$

Now since $(\sin x/x)^2 e^{-2i\xi x}$ can be extended to an entire function on the entire complex plane, which is bounded on any horizontal strip, we can apply Cauchy's theorem and take limits to conclude that

$$\begin{aligned}\widehat{F}(\xi) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{(\sin x)^2}{x^2} e^{-2i\xi x} dx = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{(\sin(x-iy))^2}{(x-iy)^2} e^{-2i\xi x - 2\xi y} dx \\ &= \frac{1}{4\pi} \int_{-\infty}^{\infty} f_{\xi}(x-iy) + g_{\xi}(x-iy) dx.\end{aligned}$$

If $\xi \geq 1$, the functions f_{ξ} and g_{ξ} are both negligible in the lower half plane, and have no poles in the lower half plane, so if we let γ denote the curve of length $2\pi n$ travelling anticlockwise along the lower semicircle with vertices $-n-iy$ and $n-iy$, then because $|z| \geq n$ on γ ,

$$\begin{aligned}\int_{-n}^n f_{\xi}(x-iy) + g_{\xi}(x-iy) dx &= \int_{\gamma} f_{\xi}(z) + g_{\xi}(z) dz \\ &= \text{length}(\gamma) \|f_{\xi} + g_{\xi}\|_{L^{\infty}(\gamma)} \\ &= (2\pi n) O_{\xi}(1/n^2) = O_{\xi}(1/n),\end{aligned}$$

and so we conclude that

$$\int_{-\infty}^{\infty} f_{\xi}(x-iy) + g_{\xi}(x-iy) dx = 0.$$

This means $\widehat{F}(\xi) = 0$. If $0 \leq \xi \leq 1$, then f_{ξ} is still small in the lower half plane, so we can conclude that

$$\int_{-\infty}^{\infty} f_{\xi}(x-iy) dx = 0.$$

But g_{ξ} is now small in the upper half plane. For $\text{Im}(z) \geq -y$,

$$|e^{-2(\xi-1)iz}| = |e^{2(1-\xi)iz}| \leq e^{2(1-\xi)y},$$

so $g_{\xi}(z) = O_{\xi}(1/|z|^2)$ in the half plane above the line $\mathbf{R}-iy$. The only problem now is that g_{ξ} has a pole in this upper half plane, at the origin. Taking Laurent series here, we find that the residue at this point is $2i(\xi-1)$. Thus, if we let γ be the curve obtained from travelling anticlockwise about the upper semicircle

with vertices $-n - iy$ and $n - iy$, then $|z| \geq n - y$ on this curve, and the residue theorem tells us that

$$\int_{-n}^n g_{\xi}(x - iy) dx + \int_{\gamma} g_{\xi}(z) dz = 2\pi i(2i(\xi - 1)) = 4\pi(1 - \xi),$$

and we now find that, as with the evaluation of the previous case,

$$\int_{\gamma} g_{\xi}(z) dz \leq (2\pi n)O_{\xi,y}(1/n^2) = O_{\xi,y}(1/n).$$

Taking $n \rightarrow \infty$, we conclude

$$\int_{-\infty}^{\infty} g_{\xi}(x - iy) dx = 4\pi(1 - \xi),$$

and putting this all together, we conclude that $\hat{F}(\xi) = 1 - \xi$.

Example. In the next paragraph, we calculate that if $\Phi(x) = e^{-\pi|x|^2}$, then $\hat{\Phi} = \Phi$. Thus if we define the Weierstrass kernel by

$$W_{\delta}(\xi) = \delta^{-d} e^{-\pi|x|^2/\delta^2},$$

then $G_{\delta}(f) = W_{\delta} * f$. Since the family $\{W_{\delta}\}$ is an approximation to the identity, this shows $G_{\delta}(f)$ converges to f in all the appropriate senses.

Since Φ breaks onto products of exponentials over each coordinate, it suffices to calculate the Fourier transform in one dimension, from which we can obtain the general transform by taking products. In the one dimensional case, since $\Phi'(x) = -2\pi x e^{-\pi x^2}$ is integrable, we conclude that $\hat{\Phi}$ is differentiable, and

$$(\hat{\Phi})'(\xi) = (-2\pi i \xi \Phi)^{\wedge}(\xi) = i(\Phi')^{\wedge}(\xi) = i(2\pi i \xi) \hat{\Phi}(\xi) = -2\pi \xi \hat{\Phi}(\xi)$$

The uniqueness theorem for ordinary differential equations says that since

$$\hat{\Phi}(0) = \int_{-\infty}^{\infty} e^{-\pi x^2} = 1 = \Phi(0)$$

Thus we must have $\hat{\Phi} = \Phi$.

Example. The Fourier transform of the function $e^{-|x|}$ is the Poisson kernel

$$P(\xi) = \frac{\Gamma((d+1)/2)}{\pi^{(d+1)/2}(1+|\xi|^2)^{(d+1)/2}}$$

Later on we show the corresponding scaled kernel $\{P_\delta\}$ is an approximation to the identity, and thus $A_\delta f = P_\delta * f$ converges to f in all appropriate senses.

The Abel kernel A_δ on \mathbf{R}^d is obtained from the initial function $A(x) = e^{-2\pi|x|}$. The calculation of the Fourier transform of this function indicates a useful principle in analysis: one can reduce expressions involving e^{-x} into expressions involving e^{-x^2} using the subordination principle. In particular, for $\beta > 0$ we have the formula

$$e^{-\beta} = \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} e^{-\beta^2/4u} du$$

We establish this by letting $v = \sqrt{u}$, so

$$\int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} e^{-\beta^2/4u} du = \frac{2}{\sqrt{\pi}} \int_0^\infty e^{-v^2 - \beta^2/4v^2} dv = \frac{2e^{-\beta}}{\sqrt{\pi}} \int_0^\infty e^{-(v - \beta/2v)^2} dv$$

But the map $v \mapsto v - \beta/2v$ is measure preserving by Glasser's master theorem, so this integral is

$$\frac{2e^{-\beta}}{\sqrt{\pi}} \int_0^\infty e^{-v^2} dv = e^{-\beta}$$

In tandem with Fubini's theorem, this formula implies

$$\begin{aligned} \hat{A}(\xi) &= \int_{\mathbf{R}^d} e^{-2\pi|x|} e^{-2\pi i \xi \cdot x} dx = \int_{\mathbf{R}^d} \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} e^{-|\pi x|^2/u} e^{-2\pi i \xi \cdot x} du dx \\ &= \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} \int_{\mathbf{R}^d} e^{-|\pi x|^2/u} e^{-2\pi i \xi \cdot x} dx du = \int_0^\infty \frac{e^{-u}}{\sqrt{\pi u}} (\text{Dil}_{(\pi^{1/2}/u^{1/2})} \Phi)^\wedge(\xi) du \\ &= \frac{1}{\pi^{(d+1)/2}} \int_0^\infty e^{-u} u^{(d-1)/2} e^{-u|\xi|^2} du \end{aligned}$$

Setting $v = (1 + |\xi|^2)u$, we conclude that since by definition,

$$\int_0^\infty e^{-v} v^{(d-1)/2} dv = \Gamma\left(\frac{d+1}{2}\right)$$

$$\hat{A}(\xi) = \frac{\Gamma((d+1)/2)}{[\pi(1+|\xi|^2)]^{(d+1)/2}}$$

Thus the Abel mean is the Fourier inverse of the Poisson kernel on the upper half plane \mathbf{H}^{d+1} . We note that the Poisson summation formula shows that for $d = 1$, the Poisson kernel on \mathbf{T} is the periodization of the Poisson kernel on \mathbf{R} .

In order to conclude $\{P_\delta\}$ is a good kernel, it now suffices to verify that

$$\int_{\mathbf{R}^d} \frac{d\xi}{(1+|\xi|^2)^{(d+1)/2}} = \frac{\pi^{(d+1)/2}}{\Gamma((d+1)/2)}$$

The right hand side is half the surface area of the unit sphere in \mathbf{R}^{d+1} . Denoting the surface area of the unit sphere in \mathbf{R}^{d+1} by S_d , and switching to polar coordinates, we find that

$$\int_{\mathbf{R}^d} \frac{d\xi}{(1+|\xi|^2)^{(d+1)/2}} = S_{d-1} \int_0^\infty \frac{r^{d-1}}{(1+r^2)^{(d+1)/2}} dr$$

Setting $r = \tan u$, we find

$$\int_0^\infty \frac{r^{d-1}}{(1+r^2)^{(d+1)/2}} dr = \int_0^{\pi/2} (\sin u)^{d-1} du$$

But we can now show by induction that

$$\frac{S_d}{2} = S_{d-1} \int_0^{\pi/2} (\sin u)^{d-1} du.$$

Using the values $S_0 = 2$, $S_1 = 2\pi$, and $S_2 = 4\pi$, the theorem certainly holds for $d = 1$ and $d = 2$. For $d > 2$, integration by parts and induction shows that

$$\begin{aligned} S_{d-1} \int_0^{\pi/2} (\sin u)^{d-1} du &= S_{d-1} \frac{d-2}{d-1} \int_0^{\pi/2} (\sin u)^{d-3} du \\ &= \frac{d-2}{d-1} \frac{S_{d-1} S_{d-2}}{2 S_{d-3}} \\ &= \frac{d-2}{d-1} \frac{\pi^{d/2} \pi^{d/2-1/2}}{\pi^{d/2-1}} \frac{\Gamma(d/2-1)}{\Gamma(d/2) \Gamma(d/2-1/2)} \\ &= \frac{\pi^{d/2+1/2}}{\Gamma(d/2+1/2)} = \frac{S_d}{2}. \end{aligned}$$

Thus our theorem is complete.

Example. We note that

$$\int_{-R}^R e^{-2\pi i \xi x} dx = \frac{e^{-2\pi i \xi R} - e^{2\pi i \xi R}}{-2\pi i \xi} = \frac{\sin(2\pi \xi R)}{\pi \xi}.$$

so the Fourier transform of $\chi_{[-R,R]}$ is the Dirichlet kernel

$$D_R(\xi) = \frac{\sin(2\pi \xi R)}{\pi \xi}$$

We note that $D_R \notin L^1(\mathbf{R})$. Thus D_R is not a good kernel, which makes the convergence rates of $S_R f$ more subtle. Nonetheless, D_R does lie in $L^p(\mathbf{R})$ for all $p \in (1, \infty]$, and is uniformly bounded in $L^p(\mathbf{R})$ for all $p \in (1, \infty)$, a fact we will prove later. This is enough to conclude that for all $p \in (1, \infty)$, $S_R f \rightarrow f$ in $L^p(\mathbf{R})$.

Thus we now know there are a large examples of functions $\Phi \in C_0(\mathbf{R}^d)$ with $\Phi(0) = 1$, and such that for any x in the Lebesgue set of f ,

$$f(x) = \lim_{\delta \rightarrow 0} \int \hat{f}(\xi) e^{2\pi i \xi \cdot x} \Phi(\delta x) d\xi.$$

If \hat{f} is integrable, then the bound $|\hat{f}(\xi) e^{2\pi i \xi \cdot x} \Phi(\delta \xi)| \leq \|\Phi\|_\infty |\hat{f}(\xi)|$ implies that we can use the dominated convergence theorem to conclude that for any point x in the Lebesgue set of f ,

$$f(x) = \lim_{\delta \rightarrow 0} \int \hat{f}(\xi) e(\xi \cdot x) \Phi(\delta \xi) d\xi = \int \hat{f}(\xi) e(\xi \cdot x) d\xi$$

Thus the inversion theorem holds pointwise almost everywhere.

Theorem 4.10. If f and \hat{f} are elements of $L^1(\mathbf{R}^d)$, then for any x in the Lebesgue set of f ,

$$f(x) = \int_{\mathbf{R}^d} \hat{f}(\xi) e(\xi \cdot x) d\xi.$$

Remark. We note that if $f \in L^1(\mathbf{R}^d)$, $\hat{f} \geq 0$, and f is continuous at the origin, then the Fourier inversion formula and the monotone convergence theorem implies that

$$f(0) = \lim_{\delta \rightarrow 0} \int_{\mathbf{R}^d} \hat{f}(\xi) e^{-\delta \xi} d\xi = \int_{\mathbf{R}^d} \hat{f}(\xi) d\xi.$$

Thus \hat{f} is integrable, and so the Fourier inversion theorem holds.

As a particular example of this remark, if $f \in L^1(\mathbf{R}^d)$ then we can define the autocorrelation function

$$R(x) = \int_{\mathbf{R}^d} f(y+x)f(y) dy,$$

then $R \in L^1(\mathbf{R}^d)$ and $\hat{R}(\xi) = |\hat{f}(\xi)|^2$. Thus R is continuous at the origin if and only if \hat{R} is integrable, which, using the L^2 theory we develop in the next section, holds if and only if $f \in L^2(\mathbf{R}^d)$.

It is often useful to note that if the Fourier transform of an integrable function is non-negative, then it's Fourier transform is automatically integrable.

Theorem 4.11. *If $f \in L^1(\mathbf{R})$ is continuous at the origin, and $\hat{f} \geq 0$, then \hat{f} is integrable.*

Proof. This follows because

$$f(0) = \lim_{\delta \rightarrow 0} \int \hat{f}(\xi) e^{-\delta|\xi|} d\xi$$

By Fatou's lemma,

$$f(0) = \lim_{\delta \rightarrow 0} \int \hat{f}(\xi) e^{-\delta|\xi|} d\xi \geq \int \liminf_{\delta \rightarrow 0} \hat{f}(\xi) e^{-\delta|\xi|} d\xi = \int \hat{f}(\xi) d\xi$$

so \hat{f} is finitely integrable. □

Note that this implies that we obtain the general inversion theorem, so in particular, it is only continuous functions, and functions almost everywhere equal to continuous functions, which can have non-negative Fourier transforms. Here is a related result. Here, without integrability constraints, the Fourier transform of $m \in L^\infty(\mathbf{R}^d)$ will be taken in the sense of tempered distributions, introduced later on in the notes.

Lemma 4.12. *If $m \in L^\infty(\mathbf{R}^d)$, and $\hat{m} \geq 0$, then $\|m\|_{L^\infty(\mathbf{R}^d)} = \|\hat{m}\|_{L^1(\mathbf{R}^d)}$.*

Proof. Set $k = \hat{m}$. The result is equivalent to showing that if $m \in L^\infty(\mathbf{R}^d)$ and $\hat{m} \geq 0$, then $\|m\|_{L^\infty(\mathbf{R}^d)} = \|k\|_{L^1(\mathbf{R}^d)}$. From this it then follows that for a general p ,

$$\|k\|_{L^1(\mathbf{R}^d)} = \|m\|_{M^2} \leq \|m\|_{M^p} \leq \|m\|_{M^1} = \|k\|_{L^1(\mathbf{R}^d)}.$$

Suppose first that m is continuous. If Φ is the Gaussian function, and $\Phi_\delta(\xi) = \delta^{-d}\Phi(\xi/\delta)$ then the multiplication formula implies that

$$\int m(\xi)\Phi_\delta(\xi - \xi_0) = \int k(x)\Phi(\delta\xi)e^{2\pi i\xi_0 \cdot x} dx.$$

If m is continuous, taking $\delta \rightarrow 0$ gives

$$|m(\xi_0)| \leq \limsup_{\delta \rightarrow 0} \left| \int k(x)\Phi(\delta\xi)e^{2\pi i\xi_0 \cdot x} dx \right| \leq \int k(x); dx$$

and by monotone convergence,

$$m(0) = \lim_{\delta \rightarrow 0} \int k(x)\Phi(\delta\xi) = \int k(x) dx.$$

Thus $\|m\|_{L^\infty(\mathbf{R}^d)} = \|k\|_{L^1(\mathbf{R}^d)}$.

But now if we let $k_\varepsilon(x) = k(x)\Phi(\varepsilon x)$, and $m_\varepsilon = \widehat{k_\varepsilon} = m * \Phi_\varepsilon$, then m_ε converges distributionally to m as $\varepsilon \rightarrow 0$. Now k_ε is non-negative, and Young's inequality implies $\|m_\varepsilon\|_{L^\infty(\mathbf{R}^d)} \leq \|m\|_{L^\infty(\mathbf{R}^d)}$. Thus

$$\|m_\varepsilon\|_{L^\infty(\mathbf{R}^d)} = \|k_\varepsilon\|_{L^1(\mathbf{R}^d)}.$$

Taking $\varepsilon \rightarrow 0$ gives

$$\lim_{\varepsilon \rightarrow 0} \|m_\varepsilon\|_{L^\infty(\mathbf{R}^d)} = \|k\|_{L^1(\mathbf{R}^d)}.$$

But now

$$\begin{aligned} \|m\|_{L^\infty(\mathbf{R}^d)} &\geq \lim_{\delta \rightarrow 0} \delta^{d/2} \|k * \Phi_\delta\|_{L^2(\mathbf{R}^d)} \\ &= \lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \delta^{d/2} \|k_\varepsilon * \Phi_\delta\|_{L^2(\mathbf{R}^d)}. \end{aligned}$$

These limits are both monotonic, so we can interchange them, thus getting that

$$\begin{aligned} \lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \delta^{d/2} \|k_\varepsilon * \Phi_\delta\|_{L^2(\mathbf{R}^d)} &= \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \delta^{d/2} \|k_\varepsilon * \Phi_\delta\|_{L^2(\mathbf{R}^d)} \\ &= \lim_{\varepsilon \rightarrow 0} \|k_\varepsilon\|_{L^1(\mathbf{R}^d)} \\ &= \|k\|_{L^1(\mathbf{R}^d)}. \end{aligned}$$

This completes the proof. □

We define, for any integrable $f : \mathbf{R}^n \rightarrow \mathbf{R}$, the *inverse* Fourier transform

$$\check{f}(x) = \int f(\xi) e(\xi \cdot x) d\xi$$

The inverse transform is also denoted by $\mathcal{F}^{-1}(f)$. The last theorem says that \mathcal{F}^{-1} really is the inverse operator to the operator \mathcal{F} , at least on the set of functions f where \hat{f} is integrable. In particular, this is true if f has weak derivatives in the L^1 norm for any multi-index $|\alpha| \leq n+1$, and so the Fourier inversion formula holds for sufficiently smooth functions.

Corollary 4.13. *If $f \in C(\mathbf{R})$ is integrable and $\hat{f} \in L^1(\mathbf{R})$, $S_R f \rightarrow f$ uniformly.*

Proof. The dominated convergence theorem implies that for each $x \in \mathbf{R}$,

$$f(x) = \int_{\mathbf{R}} \hat{f}(\xi) e(\xi \cdot x) = \lim_{R \rightarrow \infty} \int_{-R}^R \hat{f}(\xi) e(\xi \cdot x) = \lim_{R \rightarrow \infty} (S_R f)(x).$$

And

$$\int_{|x| \geq R} \hat{f}(\xi) e(\xi \cdot x) \leq \int_{|x| \geq R} |\hat{f}(\xi)| d\xi = o(1).$$

so the pointwise convergence is uniform in x . □

Remark. This theorem also generalizes to \mathbf{R}^d . Here, the operators S_R are no longer canonically defined, but if we consider any increasing nested family of sets B_R with $\lim B_R = \mathbf{R}^n$, then the corresponding operators

$$S_R f = \int_{B_R} \hat{f}(\xi) e(\xi \cdot x)$$

also converge uniformly to f .

Corollary 4.14. *The map $\mathcal{F} : L^1(\mathbf{R}^d) \rightarrow C_0(\mathbf{R}^d)$ is injective.*

Proof. If $\hat{f} = 0$, then \hat{f} is certainly integrable. But this means that the Fourier inversion theorem can apply, giving that for almost every point x ,

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(x) e(\xi \cdot x) = 0.$$

Thus $f = 0$ almost everywhere. □

The corollary above is often underestimated in utility. Even if the Fourier inversion theorem doesn't hold, we can still view the Fourier transform as another way to represent a function, since the Fourier transform does not lose any information. For instance, it can be used very easily to verify identities involving convolutions.

Corollary 4.15. *For any δ_1, δ_2 ,*

$$W_{\delta_1+\delta_2} = W_{\delta_1} * W_{\delta_2} \quad \text{and} \quad P_{\delta_1+\delta_2} = P_{\delta_1} * P_{\delta_2}.$$

Proof. We recall that

$$W_{\delta_1+\delta_2} = \mathcal{F}(e^{-(\delta_1+\delta_2)|x|^2}).$$

But $e^{-(\delta_1+\delta_2)|x|^2} = e^{-\delta_1|x|^2} e^{-\delta_2|x|^2}$ breaks into a product, which allows us to calculate

$$\mathcal{F}(e^{-\pi\delta_1|x|^2} e^{-\pi\delta_2|x|^2}) = \mathcal{F}(e^{-\pi\delta_1|x|^2}) * \mathcal{F}(e^{-\pi\delta_2|x|^2}) = W_{\delta_1} * W_{\delta_2}.$$

Thus $W_{\delta_1} * W_{\delta_2} = W_{\delta_1+\delta_2}$. Similarly, $P_{\delta_1+\delta_2}$ is the Fourier transform of $e^{-(\delta_1+\delta_2)|x|}$, which breaks into a product, whose individual Fourier transforms are P_{δ_1} and P_{δ_2} . \square

Many of the other convergence statements for Fourier series hold in the case of the Fourier transform. For instance, a non-periodic variant of the De la Vallee Poisson kernel shows that if $f \in L^1(\mathbf{R})$ and $\hat{f}(\xi) = O(1/|\xi|)$, then $S_R f$ converges uniformly to f . But for the purpose of novelty, we move on to other concepts.

4.5 The L^2 Theory

There are various differences in the L^2 for the Fourier transform vs the case of Fourier series. In the compact, periodic case, $L^2(\mathbf{T}^d)$ is contained in $L^1(\mathbf{T}^d)$ and can thus be viewed as a *more regular* family of functions than the square integrable functions. In the noncompact case, $L^2(\mathbf{R}^d)$ is not contained in $L^1(\mathbf{R}^d)$, and thus is *more regular* in some respects (we have more control over singularities), but we have less control on how spread out the function is. In particular, we often have to rely on density arguments, working in the space $L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$, which is a dense subspace of $L^2(\mathbf{R}^d)$.

One integral component of Fourier series on $L^2(\mathbf{T}^d)$ is Plancherel's equality

$$\sum_{n \in \mathbf{Z}^d} |\hat{f}(n)|^2 = \int_{\mathbf{T}^d} |f(x)|^2 dx$$

Let us try and extend this to \mathbf{R}^d . A natural formula is to expect that

$$\int_{\mathbf{R}^d} |\hat{f}(\xi)|^2 d\xi = \int_{\mathbf{R}^d} |f(x)|^2 dx.$$

In order to interpret the right hand side as a finite quantity, we must assume $f \in L^2(\mathbf{R}^d)$, and to interpret the left hand side, we must assume $f \in L^1(\mathbf{R}^d)$. A result of our calculation will show that under these assumptions, $\hat{f} \in L^2(\mathbf{R}^d)$, and that the formula holds.

Theorem 4.16. *If $f \in L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$, then $\|\hat{f}\|_{L^2(\mathbf{R}^d)} = \|f\|_{L^2(\mathbf{R}^d)}$.*

Proof. The theorem is an easy consequence of the multiplication formula, since

$$|\hat{f}(\xi)| = \hat{f}(\xi) \overline{\hat{f}(\xi)},$$

and

$$\left(\overline{\hat{f}} \right)^\wedge (\xi) = \overline{(f^\wedge)^\wedge(-\xi)} = \overline{f(\xi)}.$$

This implies

$$\int_{\mathbf{R}^d} |\hat{f}(\xi)|^2 d\xi = \int_{\mathbf{R}^d} \hat{f}(\xi) \overline{\hat{f}(\xi)} d\xi = \int_{\mathbf{R}^d} f(x) \overline{f(x)} dx = \int_{\mathbf{R}^d} |f(x)|^2 dx. \quad \square$$

A simple interpolation argument leads to the following corollary, which is a variant of the Hausdorff-Young inequality for functions on \mathbf{R}^d .

Corollary 4.17. *If $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$ for $1 \leq p \leq 2$, then*

$$\|\hat{f}\|_{L^q(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)},$$

where $2 \leq q \leq \infty$ is the conjugate of p .

Though the integral formula of an element of $L^2(\mathbf{R}^d)$ does not make sense, the bounds above provide a canonical way to define the Fourier transform of an element of $L^p(\mathbf{R}^d)$, for $1 \leq p \leq 2$. The space $L^1(\mathbf{R}^d) \cap$

$L^p(\mathbf{R}^d)$ is a dense subset of $L^p(\mathbf{R}^d)$, so we can use the Hahn-Banach theorem to define the Fourier transform $\mathcal{F} : L^p(\mathbf{R}^d) \rightarrow L^q(\mathbf{R}^d)$ as the *unique* bounded operator agreeing with the integral formula on the common domain. A more explicit way to define the Fourier transform is as the L^2 limit of the bounded Fourier transform operators; for each $f \in L^2(\mathbf{R}^d)$, and $R > 0$, $f \mathbf{I}_{B_R} \in L^1(\mathbf{R}^d)$, where B_R is the ball of radius R about the origin. It follows that if we define

$$\mathcal{F}_R f(\xi) = \int_{|x| \leq R} f(x) e^{-2\pi i \xi \cdot x} dx.$$

then since $\lim_{R \rightarrow \infty} \|\mathbf{I}_{B_R} f - f\|_{L^2(\mathbf{R}^d)} = 0$, the L^2 continuity of the Fourier transform implies that $\mathcal{F}_R f$ converges to \hat{f} in $L^2(\mathbf{R}^d)$.

The main way to obtain results about the Fourier transform of square integrable functions is by a density argument. For instance, suppose we wish to prove that for $f, g \in L^2(\mathbf{R}^d)$,

$$\int_{\mathbf{R}^d} \hat{f}(\xi) \hat{g}(\xi) d\xi = \int_{\mathbf{R}^d} f(x) g(x) dx.$$

This equality certainly holds by the multiplication formula if $f, \hat{g} \in L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$. We also find that both sides are continuous as bilinear functionals, by applying the Cauchy-Schwartz inequality and the isometry of the Fourier transform. Since any element f of $L^2(\mathbf{R}^d)$ can be approximated in the L^2 norm by an element of $L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$, and since any element g of $L^2(\mathbf{R}^d)$ can be approximated by functions with $\hat{g} \in L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$, the theorem holds in general. In particular, this shows that the extension of the Fourier transform to $L^2(\mathbf{R}^d)$ remains unitary.

Another approximation argument can be used to obtain convergence results in $L^2(\mathbf{R}^d)$ for the Fourier transform. If we let

$$S_\delta(f, \Phi) = \int \hat{f}(\xi) e^{2\pi i \xi \cdot x} \Phi(\delta \xi) d\xi$$

which is well defined for a particular function $\Phi \in L^2(\mathbf{R}^d)$, then a density argument again shows that $S_\delta(f, \Phi) = K_\delta^\Phi * f$, where K_δ^Φ is defined as in the last section. Provided that we also have $\Phi \in L^1(\mathbf{R}^d)$ and $\int \hat{\Phi}(\xi) d\xi = 1$, then we conclude that $S_\delta(f, \Phi) \rightarrow f$ in $L^2(\mathbf{R}^d)$, and that if $\Phi \in C^{d+1}(\mathbf{R}^d)$, then

$S_\delta(f, \Phi) \rightarrow f$ almost everywhere. In particular, one can use the Gauss, Abel, and Fejer sums here to get L^2 convergence.

Unlike in the case of Fourier series, where the L^2 theory gives an isometry between $L^2(\mathbf{T}^d)$ and $L^2(\mathbf{Z}^d)$, in the case of the Fourier transform the Fourier transform gives a unitary operator from $L^2(\mathbf{R}^d)$ to itself, and thus we can consider the spectral theory of such an operator. The Fourier inversion formula implies that the Fourier transform has order four. Thus the only eigenfunctions of the Fourier transform correspond to eigenvalues in $\{1, -1, i, -i\}$. We have seen $e^{-\pi x^2}$ is an eigenfunction with eigenvalue one. If we consider the family of all *Hermite polynomials*

$$H_n(x) = \frac{(-1)^n}{n!} e^{\pi x^2} \frac{d^n}{dx^n} (e^{-\pi x^2}).$$

One can also see that

$$\sum_{n=0}^{\infty} (-t)^n / n! \\ \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} = e^{-\pi x^2 - (2\pi)^{1/2} tx + t^2}$$

TODO PROVE ORTHOGONALITY AND COMPLETENESS. which satisfy $\widehat{H}_n = (-i)^n H_n$, then we obtain an orthonormal basis of eigenfunctions. In higher dimensions, a basis of eigenfunctions for $L^2(\mathbf{R}^d)$ is given by taking tensor products of Hermite polynomials.

4.6 The Hausdorff-Young Inequality

For functions on \mathbf{T} , it is unclear how to provide examples which show why the Hausdorff-Young inequality cannot be extended to give results for $p > 2$. Over \mathbf{R} , we can provide examples which explicitly indicate the tightness of the appropriate constants by applying symmetry arguments.

Example. Given $f \in L^1(\mathbf{R})$, let $f_r(x) = f(rx)$. Then we find $\widehat{f}_r(\xi) = r^{-d} \widehat{f}(\xi/r)$, and so

$$\|f_r\|_{L^p(\mathbf{R}^d)} = r^{-d/p} \|f\|_{L^p(\mathbf{R}^d)} \quad \text{and} \quad \|\widehat{f}_r\|_{L^q(\mathbf{R}^d)} = r^{d/q-d} \|\widehat{f}\|_{L^q(\mathbf{R}^d)}.$$

In order for a bound to hold in terms of p and q uniformly for all values of r , we need $r^{-d/p} = r^{d/q-d}$, which means $1/q + 1/p = 1$, so p and q must be

conjugates of one another. In the case of \mathbf{T} , a function analogous to f_r can only be defined for small value of r , and a uniform estimate can then only hold if $1/p + 1/q \geq 1$.

If $p > 2$, then for the value q with $1/p + 1/q = 1$, we have $q < p$. It is a principle of Littlewood that translation invariant operators cannot satisfy a L^p to L^q bound.

Theorem 4.18. Suppose $T : L^p(\mathbf{R}^d) \rightarrow L^q(\mathbf{R}^d)$ is a translation invariant continuous operator, where $q < p$. Then $T = 0$.

Proof. If T was nonzero, we could pick some $f_0 \in L^p(\mathbf{R}^d)$ such that $Tf_0 \neq 0$. Rescaling T and f_0 , we may assume without loss of generality that $\|f_0\|_{L^p(\mathbf{R}^d)} = \|Tf_0\|_{L^q(\mathbf{R}^d)} = 1$. Furthermore, by truncation we may assume that f_0 has compact support on some ball B_R . But then the supports of the functions $\text{Trans}_{2Rn}f_0$ and f_0 are disjoint for $n \in \mathbf{Z}^d$, so for any choice of coefficients $\{a_n\}$,

$$\left\| \sum_{n \in \mathbf{Z}^d} a_n \cdot \text{Trans}_{2Rn} f_0 \right\|_{L^p(\mathbf{R}^d)} = \left(\sum |a_n|^p \right)^{1/p}.$$

Assume that at most N of the coefficients a_n are nonzero. We cannot necessarily assume that Tf_0 has compact support but the majority of the mass of Tf_0 can still be concentrated on a compact set. For any $\varepsilon > 0$ we can choose R large enough that

$$\left(\int_{|x| \geq R} |Tf_0(x)|^q \right)^{1/q} \leq \varepsilon.$$

Now for each B_R and $m \in \mathbf{Z}^d$,

$$\left(\int_{x \in 2Rm + B_R} \left| \sum_{n \in \mathbf{Z}^d} a_n \text{Trans}_{2Rn} Tf_0(x) \right|^q dx \right)^{1/q} \geq \left(|a_m|^q - \varepsilon \sum_{n \neq m} |a_n|^q \right)^{1/q}.$$

If, for a fixed sequence $\{a_n\}$, we choose

$$\varepsilon \leq \frac{0.5}{\max_{n \in \mathbf{Z}^d} |a_n| \cdot \left(\sum_{n \in \mathbf{Z}^d} |a_n|^q \right)^{1/q}}.$$

Then we find

$$\left(\int_{x \in 2Rm + B_R} \left| \sum_{n \in \mathbf{Z}^d} a_n \text{Trans}_{2Rn} T f_0(x) \right|^q dx \right)^{1/q} \geq 0.5^{1/q} |a_m|$$

and so summing over all m , we conclude that

$$\left\| \sum_{n \in \mathbf{Z}^d} a_n \text{Trans}_{2Rn} T f_0 \right\|_{L^q(\mathbf{R}^d)} \geq 0.5^{1/q} \left(\sum_{n \in \mathbf{Z}^d} |a_n|^q \right)^{1/q}.$$

Thus we conclude that for *any* sequence $\{a_n\}$ in $l^q(\mathbf{Z}^d)$,

$$\left(\sum_{n \in \mathbf{Z}^d} |a_n|^q \right)^{1/q} \lesssim_q \left(\sum_{n \in \mathbf{Z}^d} |a_n|^p \right)^{1/p}.$$

where the constant is independent of the sequence. For $q < p$ this is impossible. \square

We can also provide a family of functions whose Fourier transforms contradict an extension of the Hausdorff Young inequality for $p > 2$.

Example. Consider the family of functions $f_s(x) = s^{-d/2} e^{-\pi|x|^2/s}$, where $s = 1 + it$ for some $t \in \mathbf{R}$. One can easily calculate using analytic continuation and the Fourier transform for the Gaussian that $\widehat{f_s}(\xi) = e^{-\pi s|\xi|^2}$. We calculate

$$\|f_s\|_{L^p(\mathbf{R}^d)} = |s|^{-d/2} \left(\int e^{-(p/|s|^2)\pi|x|^2} dx \right)^{1/p} = |s|^{d/p-d/2} p^{-d/p}$$

whereas $\|\widehat{f_s}\|_q = q^{-d/2}$. Thus to be able compare the two quantities as $t \rightarrow \infty$, we need $d/p - d/2 \leq 0$, so $p \leq 2$. As $t \rightarrow \infty$, $|f_s(x)| \sim t^{-d/2} e^{-\pi|x/t|^2}$, so the t gives us a decay in f_s . However, when we take the Fourier transform the t only corresponds to oscillatory terms. Thus we need $p \leq 2$ so that the decay in t isn't too important in relation to the overall width of the function. One can obtain analogous examples in \mathbf{T}^d to this example, by applying the Poisson summation formula to the functions f_s and noting that the L^p and L^q norms also follows approximately the same formulas as above.

The Hausdorff-Young inequality shows that the Fourier transforms narrowly supported functions into a function with small magnitude. But the example above shows that the Fourier transform is not so good at transforming functions with small magnitude into functions which are narrowly supported, because the Fourier transform can absorb the small magnitude into an oscillatory property not reflected in the norms. Some kind of way of measuring oscillation needs to be considered to get a tighter control on the function. Of course, in hindsight, we should have never expected too much control of the Fourier transform in terms of the L^p norms, since the Fourier transform measures the oscillatory nature of the input function, and oscillatory properties of a function in phase space are not very well reflected in the L^p norms, except when applying certain orthogonality properties with an L^2 norm, or destroying the oscillation with an L^∞ norm.

4.7 The Poisson Summation Formula

We now show a connection between the Fourier transform on \mathbf{R} , and the Fourier transform on \mathbf{T} . If f is a function on \mathbf{R} , there are two ways of obtaining a ‘periodic’ version of f on \mathbf{T} . Firstly, we can define, for each $x \in \mathbf{T}$,

$$f_1(x) = \sum_{n=-\infty}^{\infty} f(x + 2\pi n),$$

which is a well defined element of $C^\infty(\mathbf{T})$. Secondly, we can define

$$f_2(x) = \sum_{n=-\infty}^{\infty} \hat{f}(n)e_n(x),$$

The Poisson summation formula says that, under an appropriate regularity condition so that we can interpret these formulas correctly, they give the same function.

Theorem 4.19. *Suppose $f \in L^1(\mathbf{R}^d)$. Then the series*

$$\sum_{n \in \mathbf{Z}^d} \text{Trans}_n f$$

converges absolutely in $L^1[0,1]^d$ to a function $g \in L^1[0,1]^d$ with $\hat{g}(n) = \hat{f}(n)$ for each $n \in \mathbf{Z}^d$.

Proof. The fact that the sum converges absolutely in $L^1[0,1]$ follows because

$$\sum_{n \in \mathbf{Z}^d} \|\text{Trans}_n f\|_{L^1[0,1]} = \|f\|_{L^1(\mathbf{R}^d)}.$$

But the absolute convergence in L^1 also justifies the calculation that for each $n \in \mathbf{Z}^d$

$$\begin{aligned} \int_{[0,1]^d} \sum_{m \in \mathbf{Z}^d} (\text{Trans}_n f)(x) e^{2\pi n i x} dx &= \sum_{m \in \mathbf{Z}^d} \int_{[0,1]^d} f(x+m) e^{2\pi n i (x+m)} dx \\ &= \int_{\mathbf{R}^d} f(x) e^{2\pi n i x} dx = \hat{f}(n). \end{aligned} \quad \square$$

We can obtain a much more powerful version of this result if we assume that there is $\delta > 0$ such that

$$|f(x)| \lesssim \frac{1}{1+|x|^{d+\delta}} \quad \text{and} \quad |\hat{f}(\xi)| \lesssim \frac{1}{1+|\xi|^{d+\delta}}.$$

Then we see that the two functions

$$g_1(x) = \sum_{n \in \mathbf{Z}^d} f(x+n) \quad \text{and} \quad g_2(x) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}$$

are continuous functions on \mathbf{T}^d with the same Fourier coefficients. It thus follows that $g_1 = g_2$, i.e. that for each $x \in \mathbf{R}^d$,

$$\sum_{n \in \mathbf{Z}^d} f(x+n) = \sum_{n \in \mathbf{Z}^d} \hat{f}(n) e^{2\pi i n \cdot x}.$$

In particular, this holds if $f \in \mathcal{S}(\mathbf{R})$. From a distributional standpoint, this result says that the tempered distribution $\sum_{n \in \mathbf{Z}^d} \delta_n$ is an eigendistribution of the Fourier transform, i.e. it is its own Fourier transform.

TODO: Also prove this statement under the assumption that f has bounded variation and $f(t) = [f(t+) + f(t-)]/2$ for all $t \in \mathbf{R}$.

4.8 Convergence of Fourier Series via Poisson Summation

TODO

4.9 Radial Functions

Suppose $f \in L^1(\mathbf{R}^d)$ is a radial function. Then \hat{f} is also a radial function. In particular, if we let

$$\|u\|_{L^1([0,\infty),r^{d-1})} = \int_0^\infty r^{d-1}u(r) dr$$

then we have a transform $u \mapsto \tilde{u}$ from $L^1([0,\infty),r^{d-1})$ to $L^\infty[0,\infty)$ where if $f(x) = u(|x|)$, then $\hat{f}(\xi) = \tilde{u}(|\xi|)$. In particular, we calculate quite simply that

$$\tilde{u}(s) = V_d \int_0^\infty r^{d-1}u(r) \left(\int_{S^{d-1}} e^{-2\pi i x_1 s} dx \right).$$

If one recalls the Bessel functions $\{J_s\}$, then we have

$$\tilde{u}(s) = 2\pi s^{1-d/2} \int_0^\infty r^{d/2}u(r)J_{d/2-1}(2\pi sr) dr.$$

If one recalls some Bessel function asymptotics, then one can actually gain some interesting results for the *averaging operator*

$$Af(x) = \oint_{S^{d-1}} f(x-y) d\sigma(y)$$

Example. Suppose $f_R(x) = \mathbf{I}_{|x| \leq R}$. Then

$$\hat{f}(\xi) = 2\pi|\xi|^{1-d/2} \int_0^R r^{d/2}J_{d/2-1}(2\pi sr) dr.$$

The

4.10 Poisson Integrals

s

Chapter 5

Applications of the Fourier Transform

5.1 Applications to Partial Differential Equations

Just as the Fourier series can be used to obtain periodic solutions to certain partial differential equations, the Fourier transform can be used to obtain more general solutions to partial differential equations on \mathbf{R}^d . To begin with, we study the heat equation on \mathbf{R}^d , i.e. we study solutions to the partial differential equation

$$\frac{\partial u}{\partial t} = \Delta u$$

Formally taking Fourier transforms in the spatial variable gives

$$\frac{\partial \hat{u}(\xi, t)}{\partial t} = -4\pi^2 |\xi|^2 \hat{u}(\xi, t)$$

which, if we are given $u(x, 0) = f(x)$, gives that

$$\hat{u}(\xi, t) = \hat{f}(\xi) e^{-4\pi^2 |\xi|^2 t}.$$

Thus, taking the inverse Fourier transform, we might expect the solution to the heat equation to be given by the formula

$$u(x, t) = (H_t * f)(x)$$

where

$$H_t(x) = \frac{1}{(4\pi t)^{d/2}} e^{-|x|^2/4t}.$$

The rapid decay of H_t for large x shows that for any $1 \leq p \leq \infty$ and $f \in L^p(\mathbf{R}^d)$, u is well defined by this formula, lies in $C^\infty(\mathbf{T}^d)$, and solves the heat equation, with the appropriate norm convergence as $t \rightarrow 0$. However, in this case it is not so easy to conclude that u is the unique solution to this equation satisfying the initial conditions, since one cannot necessarily take the Fourier transform of u .

We can get slightly more results if we consider the *steady state* heat equation on the upper half plane \mathbf{H}^d , i.e. we study functions $u(x, t)$, for $x \in \mathbf{R}^d$ and $t > 0$, such that $\Delta u = 0$, subject to the initial condition that $u(x, 0) = f(x)$. Working formally with the Fourier transform leads to the equation

$$\hat{u}(\xi, t) = e^{-2\pi t|\xi|} \hat{f}(\xi)$$

Thus $u(x, t) = (f * P_t)(x)$, where P_t is the Poisson kernel. If $f \in L^1(\mathbf{R}^d)$, it is easy to see that

5.2 Shannon-Nyquist Sampling Theorem

Often, in applications, one deals with band limited function, i.e. functions whose Fourier transforms are compactly supported. For simplicity, we work solely with functions f on \mathbf{R} satisfying a decay condition

$$|f(t)| \lesssim \frac{1}{(1 + |t|)^{1+\delta}}.$$

It follows that $f \in L^p(\mathbf{R}^d)$ for each $1 \leq p \leq \infty$. Suppose that in addition, \hat{f} is supported on $[-1/2, 1/2]$. It follows that, $\hat{f} \in L^p(\mathbf{R}^d)$ for each $1 \leq p \leq \infty$. In particular, it follows that f is smooth, if we alter it on a set of measure zero. Now taking Fourier series on $[-1/2, 1/2]$, noting that $f \in L^1(\mathbf{Z})$ because of its decay, we find that for each $\xi \in \mathbf{R}$,

$$\hat{f}(\xi) = \mathbf{I}(|\xi| \leq 1/2) \sum_{n=-\infty}^{\infty} f(n) e^{-2\pi i n \xi}.$$

But now we conclude by the Fourier inversion formula that

$$\begin{aligned}
f(x) &= \int_{-1/2}^{1/2} \left(\sum_{n=-\infty}^{\infty} f(n) e^{-2\pi i n \xi} \right) e^{2\pi i \xi x} d\xi \\
&= \sum_{n=-\infty}^{\infty} f(n) \int_{-1/2}^{1/2} e^{2\pi i \xi (x-n)} d\xi \\
&= \sum_{n=-\infty}^{\infty} f(n) \cdot \frac{\sin(\pi(x-n))}{\pi(x-n)}.
\end{aligned}$$

In particular, we conclude that the function f is uniquely determined by sampling it's values over the integers. In particular, if N is large, and $|x| \leq N/2$

$$\left| f(x) - \sum_{n=-N}^N f(n) \cdot \frac{\sin(\pi(x-n))}{\pi(x-n)} \right| \lesssim \frac{1}{N},$$

where the implicit constant depends on the decay of f . If we sample on a more fine set of values, then we obtain faster convergence. To do this, we instead take the Fourier series of \hat{f} on $[-\lambda/2, \lambda/2]$, noting that

$$\frac{1}{\lambda} \int_{-\lambda/2}^{\lambda/2} \hat{f}(\xi) e^{2\pi i n \xi / \lambda} d\xi = \frac{f(n/\lambda)}{\lambda}$$

so that

$$\hat{f}(\xi) = \chi(\xi) \sum_{n=-\infty}^{\infty} \frac{f(n/\lambda)}{\lambda} e^{-2\pi i n \xi / \lambda}.$$

where instead of being the indicator on $[-1/2, 1/2]$, χ is the piecewise linear function equal to 1 for $|\xi| \leq 1/2$, and vanishing for $|\xi| \geq \lambda/2$. One can calculate quite easily that

$$\hat{\chi}(x) = \frac{\cos(\pi x) - \cos(\lambda \pi x)}{\pi^2(\lambda - 1)x^2}.$$

Thus it follows from the Fourier inversion formula that

$$\begin{aligned}
f(x) &= \sum_{n=-\infty}^{\infty} \frac{f(n/\lambda)}{\lambda} \int_{-\infty}^{\infty} \chi(\xi) e^{2\pi i \xi (x - n/\lambda)} d\xi \\
&= \sum_{n=-\infty}^{\infty} \frac{f(n/\lambda)}{\lambda} \hat{\chi}(n/\lambda - x) \\
&= \sum_{n=-\infty}^{\infty} f(n/\lambda) \frac{\cos(\pi(n/\lambda - x)) - \cos(\lambda\pi(n/\lambda - x))}{\pi^2 \lambda (\lambda - 1) (n/\lambda - x)^2}.
\end{aligned}$$

It follows that if $|x| \leq N/2\lambda$, then

$$\left| f(x) - \sum_{n=-N}^N f(n/\lambda) \frac{\cos(\pi(n/\lambda - x)) - \cos(\lambda\pi(n/\lambda - x))}{\pi^2 \lambda (\lambda - 1) (n/\lambda - x)^2} \right| \lesssim \left(1 + \frac{1}{\lambda - 1}\right) \frac{1}{N^2}.$$

Thus the rate of convergence of this sum is much better if we *oversample* by a large value λ .

We should not expect f to be obtainable exactly if we undersample, i.e. look at the coefficients $\{f(n/\lambda) : n \in \mathbf{Z}\}$ for some $\lambda < 1$. Thus undersampling often yields artifacts in our reconstruction. For instance, when one takes a video of periodic motion travelling at a much greater frequency than the framerate of a video. To see why this is true, we consider a distributional formulation of the Nyquist sampling theorem.

Theorem 5.1. *For any $\lambda < 1$, there exists $f_1, f_2 \in \mathcal{S}(\mathbf{R})$, with \hat{f}_1 and \hat{f}_2 supported on $[-1/2, 1/2]$, such that $f_1(n/\lambda) = f_2(n/\lambda)$ for any $n \in \mathbf{Z}$.*

Proof. Fix $f_0 \in \mathcal{S}(\mathbf{R})$. Then the Poisson summation formula, appropriately rescaled, tells us that for each $\xi \in \mathbf{R}$,

$$\sum_{n=-\infty}^{\infty} f(n/\lambda) e^{-2\pi i n \xi} = \lambda^d \sum_{n=-\infty}^{\infty} \hat{f}(\xi - \lambda n).$$

One can determine all the coefficients $\{f(n/\lambda)\}$ if one knows the right hand side for all values $\xi \in \mathbf{R}$. Thus if $f_1, f_2 \in \mathcal{S}(\mathbf{R})$ are distinct functions such that \hat{f}_1 and \hat{f}_2 are supported on $[-1/2, 1/2]$, but are equal to one another at a periodization of scale λ , then $f_1(n/\lambda) = f_2(n/\lambda)$ for any $n \in \mathbf{Z}$. This is certainly possible if $\lambda < 1$. \square

We can also get a discretized L^2 identity.

Theorem 5.2. Suppose $f \in L^2(\mathbf{R})$ and \hat{f} is supported on $[-1/2, 1/2]$. Then

$$\sum_{n=-\infty}^{\infty} |f(n)|^2 = \int_{-\infty}^{\infty} |f(x)|^2 dx.$$

Proof. Poisson summation applied to $|f(x)|^2$ implies that

$$\sum_{n=-\infty}^{\infty} |f(n)|^2 = \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} \hat{f}(\xi) \overline{\hat{f}(\xi - n)} dx = \int_{-1/2}^{1/2} |\hat{f}(\xi)|^2 dx = \int_{-\infty}^{\infty} |f(x)|^2 dx.$$

□

5.3 The Uncertainty Principle

The uncertainty principle is a *constraint* placed on the Fourier transform, which prevents a function and its transform from concentrating too tightly in space. Roughly speaking, the uncertainty principle is the heuristic that if f is a function, and its Fourier support is supported on a ball around the origin of radius $1/R$, then f should be ‘locally constant’ at a scale R , because intuitively, f is only composed of waves oscillating at low frequencies. This principle is seen in various scenarios, and we describe some basic situations here. This property is fundamental to the transform. For instance, we see it in the formula for the Fourier transform of the dilation of a function, i.e. that

$$\mathcal{F} \circ \text{Dil}_t = t^d \cdot \text{Dil}_{1/t} \circ \mathcal{F}.$$

Thus dilation which shrinks the support of a function to a scale t , *expand* the support of the Fourier transform of the function by $1/t$. Another, version is the *Paley-Wiener theorem*: an exponentially decaying function must have a Fourier transform which is extendable to a holomorphic function on a neighborhood of the real line. The uncertainty principle appears because holomorphic functions cannot be supported on small sets. Here we consider some other versions of the principle.

5.3.1 The Heisenberg Uncertainty Principle

The first version of the uncertainty principle we give here is the most famous, the *Heisenberg uncertainty principle*, which is an L^2 instance of the heuristic, and has important implications in quantum mechanics.

Theorem 5.3 (Heisenberg). *Suppose $\psi \in \mathcal{S}(\mathbf{R})$. Then for any $x_0, \xi_0 \in \mathbf{R}$,*

$$\left(\int_{-\infty}^{\infty} (x - x_0)^2 |\psi(x)|^2 dx \right) \left(\int_{-\infty}^{\infty} (\xi - \xi_0)^2 |\hat{\psi}(\xi)|^2 d\xi \right) \geq \frac{1}{16\pi^2} \left(\int_{-\infty}^{\infty} |\psi(x)|^2 dx \right)^2.$$

Thus if ψ has L^2 norm A , then for any x_0 and ξ_0 , the L^2 norm of $(x - x_0)\psi$ and $(\xi - \xi_0)\psi$ cannot simultaneously both be $\lesssim A$.

Proof. Normalizing, we may assume that $\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1$ and that $x_0, \xi_0 = 0$. A density argument enables us to assume that $\psi \in \mathcal{S}(\mathbf{R})$. Integration by parts shows that

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} |\psi(x)|^2 dx \\ &= - \int_{-\infty}^{\infty} x \frac{d|\psi(x)|^2}{dx} dx = - \int_{-\infty}^{\infty} (x\psi'(x)\overline{\psi(x)} + x\overline{\psi'(x)}\psi(x)) dx. \end{aligned}$$

Thus

$$\begin{aligned} 1 &\leq 2 \int_{-\infty}^{\infty} |x| |\psi(x)| |\psi'(x)| dx \\ &\leq 2 \left(\int_{-\infty}^{\infty} x^2 |\psi(x)|^2 dx \right)^{1/2} \left(\int_{-\infty}^{\infty} |\psi'(x)|^2 dx \right)^{1/2} \\ &\leq 4\pi \left(\int_{-\infty}^{\infty} x^2 |\psi(x)|^2 dx \right)^{1/2} \left(\int_{-\infty}^{\infty} \xi^2 |\hat{\psi}(\xi)|^2 d\xi \right)^{1/2}. \quad \square \end{aligned}$$

Remark. Taking ψ to be a standard Gaussian function shows the constant $1/16\pi^2$ in the Heisenberg uncertainty principle is tight.

Let us explain the applications of this uncertainty principle in quantum mechanics. Unlike in classical mechanics, the position state of a particle is no longer given by a particular point, but instead given by a state function ψ subject to the normalization condition

$$\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1.$$

It then follows that the position of a particle is nondeterministic, with $|\psi(x)|^2$ giving the probability density function of where the particle is located. If x_0 denotes the expected value of the particle, then the variance of the distribution is given by

$$\sigma_x^2 = \int_{-\infty}^{\infty} |x - x_0|^2 |\psi(x)|^2 dx.$$

On the other hand, the *momentum* of the particle is also random, given in terms of a function ϕ such that the distribution of the momentum is given with density $|\phi|^2$. Thus the variance of the momentum, if ξ_0 is the expectation, is equal to

$$\sigma_\rho^2 = \int_{-\infty}^{\infty} |\rho - \rho_0|^2 |\phi(\rho)|^2 d\xi.$$

The relation between ϕ and ψ is given by the *De-Broglie* relation, which informally says the momentum of a particle is proportional to its frequency, and formally says that

$$\phi(\rho) = h^{-d/2} \int_{\mathbf{R}^d} e^{-2\pi i \rho \cdot x / h} \psi(x) dx = h^{-d/2} \hat{\psi}(\rho/h),$$

where $h > 0$ is Planck's constant. Thus Heisenberg's uncertainty principle tells us precisely that

$$\sigma_x \cdot \sigma_\rho \geq h/4\pi = \hbar/2.$$

Thus we have a fundamental limitation to the minimum possible uncertainty of the position and momentum of a particle.

We can also rephrase the uncertainty principle in terms of the differential operator

$$L = x^2 - \frac{d^2}{dx^2}.$$

This operator is known as the *Hermite operator*. Then for any $f \in \mathcal{S}(\mathbf{R})$,

$$\begin{aligned} (Lf, f) &= \int_{-\infty}^{\infty} x^2 |f(x)|^2 - f''(x) \overline{f(x)} dx \\ &= \int_{-\infty}^{\infty} x^2 |f(x)|^2 dx + \int_{-\infty}^{\infty} |f'(x)|^2 dx \\ &= \int_{-\infty}^{\infty} x^2 |f(x)|^2 dx + 4\pi^2 \int_{-\infty}^{\infty} \xi^2 |\hat{f}(\xi)|^2 d\xi. \end{aligned}$$

The Heisenberg uncertainty principle thus implies that

$$\begin{aligned} (f, f) &\leq 4\pi \left(\int_{-\infty}^{\infty} x^2 |f(x)|^2 dx \right)^{1/2} \left(\int_{-\infty}^{\infty} \xi^2 |\hat{f}(\xi)|^2 d\xi \right)^{1/2} \\ &\leq \int_{-\infty}^{\infty} x^2 |f(x)|^2 dx + 4\pi^2 \int_{-\infty}^{\infty} \xi^2 |\hat{f}(\xi)|^2 d\xi = (Lf, f). \end{aligned}$$

Thus the operator $L - 1$ is formally positive definite. If we consider the operator

$$Af = \frac{d}{dx} + x \quad \text{and} \quad A^* = -\frac{d}{dx} + x$$

then $A^*A = L - 1$. These two operators are called the *annihilation* and *creation* operators respectively.

The uncertainty principle is strongly connected to the *noncommuting* nature of a certain natural pair of operators. Consider the *position* operator

$$Xf(x) = xf(x)$$

and the *momentum* operator

$$Df = (2\pi i)^{-1} f'(x)$$

both continuous, self adjoint operators on $\mathcal{S}(\mathbf{R})$. Given an arbitrary self-adjoint operator A on $\mathcal{S}(\mathbf{R})$, and $f \in \mathcal{S}(\mathbf{R})$ with $\|f\|_{L^2(\mathbf{R})} = 1$, we define the *standard deviation* of the operator to be

$$\sigma_A^2 f = (\langle Af, Af \rangle - \langle Af, f \rangle^2)^{1/2}$$

For two self adjoint operators A and B , the *Robertson uncertainty relation* says that for any $f \in \mathcal{S}(\mathbf{R})$ with $\|f\|_{L^2(\mathbf{R})} = 1$,

$$(\sigma_A^2 f) \cdot (\sigma_B^2 f) \geq \frac{1}{2} \int (AB - BA) f(x) \overline{f(x)} dx.$$

Since $[X, D]$ is the identity map, we see this version of the uncertainty principle implies the Heisenberg uncertainty principle given above as a special case.

5.3.2 Bernstein's Inequality

You can measure how ‘locally constant’ a function is in terms of the smoothness of the function, i.e. if the derivative is small, then the function is roughly speaking, locally constant. Roughly speaking, if a function f is constant at a scale R , then Df should have magnitude on average equal to $1/R$. The Fourier transform of Df is also supported on this ball, so we can iterate this argument, we should expect $D^k f$ to have magnitude on average equal to $1/R^k$. Formulating a precise version of the uncertainty principle here gives Bernstein's inequality. Let's begin with an L^2 version which is easiest to prove, since we have Plancherel.

Lemma 5.4. *If $f \in L^2(\mathbf{R}^d)$ and \hat{f} is supported on a ball of radius $1/R$ about the origin, then for any multi-index α ,*

$$\|D^\alpha f\|_{L^2(\mathbf{R}^d)} \leq (2\pi/R)^{|\alpha|} \|f\|_{L^2(\mathbf{R}^d)}.$$

Proof. Plancherel tells us that

$$\|D^\alpha f\|_{L^2(\mathbf{R}^d)} = \|(2\pi i \xi)^\alpha \hat{f}\|_{L^2(\mathbf{R}^d)}.$$

Since \hat{f} is supported on a ball of radius R , we conclude that on this support, $(2\pi i \xi)^\alpha$ has magnitude at most $(2\pi R)^{|\alpha|}$, and thus

$$\|(2\pi i \xi)^\alpha \hat{f}\|_{L^2(\mathbf{R}^d)} \leq (2\pi R)^{|\alpha|} \|\hat{f}\|_{L^2(\mathbf{R}^d)} = \|f\|_{L^2(\mathbf{R}^d)}.$$

□

There is also a version in more general L^p spaces, but the proof is a little more technical.

Lemma 5.5. *If $f \in L^p(\mathbf{R}^d)$ and \hat{f} is supported on a ball of radius $1/R$ about the origin, then for any multi-index α , and any $q \leq p$,*

$$\|D^\alpha f\|_{L^p(\mathbf{R}^d)} \lesssim R^{d(1/p-1/q)-|\alpha|} \|f\|_{L^q(\mathbf{R}^d)}.$$

Proof. Choose a function $\psi \in \mathcal{S}(\mathbf{R}^d)$ which has a compactly support Fourier transform which is equal to one on the ball of radius 1 about the origin. If $\psi_R(x) = R^{-d}\psi(x/R)$, then the Fourier transform of ψ_R is equal to one on a

ball of radius $1/R$ about the origin. Now we have smoothness bounds on ψ_R , namely

$$|(D^\alpha \psi_R)(x)| \lesssim_{\alpha, N} R^{-d-\alpha} (x/R)^{-N}.$$

Moreover, $f = f * \psi_R$, so the convolution should show that the smoothness of ψ_R transfers to give the smoothness of f . More precisely, Young's inequality implies that if $1/p + 1 = 1/q + 1/r$, then

$$\begin{aligned} \|D^\alpha f\|_{L^p(\mathbf{R}^d)} &= \|f * D^\alpha \psi_R\|_{L^p(\mathbf{R}^d)} \\ &\lesssim \|f\|_{L^q(\mathbf{R}^d)} \|D^\alpha \psi_R\|_{L^r(\mathbf{R}^d)} \\ &= R^{d(1/r-d)-\alpha} \|f\|_{L^q(\mathbf{R}^d)} \\ &= R^{d(1/p-1/q)-\alpha} \|f\|_{L^q(\mathbf{R}^d)}. \end{aligned} \quad \square$$

Here is another version of the inequality. Recall that for discrete quantities, $l^q \leq l^p$ for $p \leq q$. The uncertainty principle says that if the Fourier support of a function is concentrated, then it should be locally constant, and thus, roughly speaking, discrete. Thus we should expect that for such functions, we have $L^q \lesssim L^p$ for $p \leq q$. Making this precise leads to Bernstein's inequality. The result immediately follows from the last argument by taking $\alpha = 0$ in the last argument, but we give an independent argument that involves the l^p and l^q norms directly. First we prove a pointwise version of the uncertainty principle.

Lemma 5.6. *Suppose \hat{f} is supported on a ball of radius $1/R$. Then for any ball B of radius R centered at some $x_0 \in \mathbf{R}^d$, if we define*

$$w^N(x) = (1 + |x|)^{-N},$$

and

$$w_B^N(x) = w_N\left(\frac{x - x_0}{R}\right),$$

then for $x_1 \in B$,

$$|f(x_1)| \lesssim_N \frac{1}{R^d} \int |f(x)| w_B^N(x) dx.$$

Proof. Assume without loss of generality that $x_0 = 0$ and $R = 1$. We have $f = f * \psi$, where $\psi = \psi_1$ is as in the last argument. Thus

$$|f(x_1)| \leq \int |f(x)| |\psi(x_1 - x)| dx \lesssim_N \int |f(x)| w_N(x) dx,$$

where we used the fact that since $|x_1| \leq 1$,

$$|\psi(x_1 - x)| \lesssim_N (1 + |x_1 - x|)^{-N} \lesssim_N (1 + |x|)^{-N} = w_N(x). \quad \square$$

The intuitive idea of the version of the proof using the fact that $l^q \leq l^p$ for $p \leq q$ intuitively precedes as follows. Without loss of generality, suppose that \hat{f} is supported on a ball of radius $1/R$ at the origin. An intuitive proof proceeds as follows: the uncertainty principle tells us that f is locally constant on sidelength $O(R)$ cubes. Thus we can divide \mathbf{R}^d into an almost disjoint union of sidelength cubes $\{I_k : k \in \mathbf{Z}^d\}$, where I_k has centre $R \cdot k$. Roughly speaking, we should be able to find constants $\{c_k\}$ such that

$$f \approx \sum c_k \chi_{I_k}.$$

It then follows that

$$\|f\|_{L^q(\mathbf{R}^d)} \approx R^{d/q} \|c\|_{l^q} \leq R^{d/q} \|c\|_{l^p} \approx R^{-(d/p - d/q)} \|f\|_{L^p(\mathbf{R}^d)}.$$

To make this argument work, we use the bound above. This result really says that for any ball B_R of radius R ,

$$\|f\|_{L^\infty(B)} \lesssim_N R^{-d} \|f\|_{L^1(w_B)}.$$

Interpolation with the trivial inequalities

$$\|f\|_{L^1(B)} \lesssim_N \|f\|_{L^1(w_B)} \quad \text{and} \quad \|f\|_{L^\infty(B)} \lesssim_N \|f\|_{L^\infty(w_B)}$$

gives that for $p \leq q$, $\|f\|_{L^q(B)} \lesssim_N R^{d(1/p - 1/q)} \|f\|_{L^p(w_B)}$. This is not quite a local version of Bernstein's inequality, since w_B is not compactly supported. But w_B has Schwartz tails, so this is a *pseudolocal* version of the inequality, which is often just as good in harmonic analysis. In particular, these bounds lead to another proof of the following lemma, by decomposing \mathbf{R}^d into a family of almost disjoint cubes $\{I_k\}$ and then writing

$$\|f\|_{L^q(\mathbf{R}^d)} \sim \|\|f\|_{L^q(I_k)}\|_{l_k^q} \lesssim_N R^{d(1/p - 1/q)} \|\|f\|_{L^p(w_{I_k})}\|_{l_k^p}.$$

The rapid decay of the weights $\{w_{I_k}\}$ allows us to get a bound of the form

$$\|\|f\|_{L^p(w_{I_k})}\|_{l_k^p} \lesssim \|\|f\|_{L^p(I_k)}\|_{l_k^p} \sim \|f\|_{L^p(\mathbf{R}^d)},$$

which completes the proof. More precisely, we have

$$w_{I_k} \lesssim \sum_{m=0}^{\infty} 2^{-Nm} \chi_{2^m I_k}.$$

The triangle inequality implies that

$$\|f\|_{L^p(w_{I_k})} \lesssim \sum_{m=0}^{\infty} 2^{-Nm} \sum_{m=0}^{\infty} \|f\|_{L^p(2^m I_k)} \lesssim \sum_{m=0}^{\infty} 2^{-Nm} \sum_{|a| \leq 2^m} \|f\|_{L^p(I_{k+a})}.$$

Finally, the triangle inequality then implies that

$$\begin{aligned} \|\|f\|_{L^p(w_{I_k})}\|_{l_k^p} &\leq \sum_{m=0}^{\infty} 2^{-Nm} \sum_{|a| \leq 2^m} \|\|f\|_{L^p(I_{k+a})}\|_{l_k^p} \\ &= \sum_{m=0}^{\infty} 2^{-Nm} 2^{dm} \|\|f\|_{L^p(I_{k+a})}\|_{l_k^p} \\ &\sim \sum_{m=0}^{\infty} 2^{-Nm} 2^{dm} \|f\|_{L^p(\mathbf{R}^d)}. \end{aligned}$$

Taking $N > d$ gives the required bound.

There are extensions of these results to *ellipsoids* rather than balls. One can reformulate the uncertainty principle as follows: if \hat{f} is supported in the *dual* of an ellipsoid E (obtained by inverting the ellipse on each axis), then f is locally constant of translates of the ellipsoid E . It is simple to obtain these results from the results above using balls, by using the way the Fourier transforms under changing coordinates linearly by an element of $GL(d)$. One can also use rectangles, and dual rectangles (we snuck in some applications using cubes instead of balls in the proofs above), since these shapes all have comparable size to their round counterparts.

5.4 Sums of Random Variables

TODO

We now switch to an application of harmonic analysis to studying sums of random variables probability theory. If X is a random vector, it's probabilistic information is given by it's distribution on \mathbf{R}^n , which can be seen

as a measure \mathbf{P}_X on \mathbf{R}^n , with $\mathbf{P}_X(E) = \mathbf{P}(X \in E)$. Given two independent random vectors X and Y , \mathbf{P}_{X+Y} is the convolution $\mathbf{P}_X * \mathbf{P}_Y$ between the measures \mathbf{P}_X and \mathbf{P}_Y , in the sense that

$$\mathbf{P}_{X+Y}(E) = \int \chi_E(x+y) d\mathbf{P}_X(x) d\mathbf{P}_Y(y)$$

If $d\mathbf{P}_X = f_X \cdot dx$ and $d\mathbf{P}_Y = f_Y \cdot dx$, then $d(\mathbf{P}_X * \mathbf{P}_Y) = (f_X * f_Y) \cdot dx$ is just the normal convolution of functions. This is why harmonic analysis becomes so useful when analyzing sums of independent random variables.

It is useful to express the Fourier transform in a probabilistic language. Given a random variable X ,

$$\widehat{\mathbf{P}_X}(\xi) = \int e^{i\xi \cdot x} d\mathbf{P}_X(x)$$

Thus the natural Fourier transform of a random vector X is the characteristic function $\varphi_X(\xi) = \mathbf{E}(e^{i\xi \cdot X})$. It is a continuous function for any random variable X . We can also express the properties of the Fourier transform in a probabilistic language.

Lemma 5.7. *Let X and Y be independent random variables. Then*

- $\varphi_X(0) = 1$, and $|\varphi_X(\xi)| \leq 1$ for all ξ .
- (Symmetry) $\varphi_X(\xi) = \overline{\varphi_X(-\xi)}$.
- (Convolution) $\varphi_{X+Y} = \varphi_X \varphi_Y$.
- (Translation and Dilation) $\varphi_{X+a}(\xi) = e^{ia \cdot \xi} \varphi_X(\xi)$, and $\varphi_{\lambda X}(\xi) = \varphi_X(\lambda \xi)$.
- (Rotations) If $R \in O(n)$ is a rotation, then $\varphi_{R(X)}(\xi) = \varphi_X(R(X))$.

Using the Fourier inversion formula, if φ_X is integrable, then X is a continuous random variable, with density

$$f(x) = \int e^{-i\xi x} \varphi_X(\xi) d\xi$$

In particular, if $\varphi_X = \varphi_Y$, then X and Y are identically distributed. This already gives interesting results.

Theorem 5.8. *If X and Y are independent normal distributions, then $aX + bY$ is normally distributed.*

Proof. Since $\varphi_{aX+bY}(\xi) = \varphi_X(a\xi)\varphi_Y(b\xi)$, it suffices to show that the product of two such characteristic functions is the characteristic function of a normal distribution. If X has mean μ and covariance matrix Σ , then $X \cdot \xi$ has mean $\mu \cdot \xi$ and variance $\xi^T \Sigma \xi$, and one calculates that $\mathbf{E}[e^{i\xi \cdot X}] = e^{-i\mu \cdot \xi - \xi^T \Sigma \xi / 2}$ using similar techniques to the Fourier transform of a Gaussian. One verifies that the class of functions of the form $e^{-i\mu \cdot \xi - \xi^T \Sigma \xi / 2}$ is certainly closed under multiplication and scaling, which completes the proof. \square

Now we can prove the celebrated central limit theorem. Note that if

Theorem 5.9. *Let X_1, \dots, X_N be independent and identically distributed with mean zero and variance σ^2 . If $S_N = X_1 + \dots + X_N$, then*

$$\mathbf{P}(S_N \leq \sigma \sqrt{N}t) \rightarrow \Phi(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-y^2/2} dy$$

Proof. We calculate that

$$\varphi_{S_N/\sigma\sqrt{N}}(\xi) = \varphi_X(\xi/\sigma\sqrt{N})^N$$

Define $R_n(x) = e^{ix} - 1 - (ix) - (ix)^2/2 - \dots - (ix)^n/n!$. Then because of oscillation and the fundamental theorem of calculus,

$$|R_0(x)| = \left| i \int_0^x e^{iy} dy \right| \leq \min(2, |x|)$$

Next, since $R'_{n+1}(x) = iR_n$,

$$R_{n+1}(x) = i \int_0^x R_n(y) dy$$

This gives that $|R_n(x)| \leq \min(2|x|^n/n!, |x|^{n+1}/(n+1)!)$. In particular, we conclude

$$|\varphi_X(\xi) - 1 - \sigma^2 \xi^2/2| = |\mathbf{E}(R_2(\xi X))| \leq \mathbf{E}|R_2(\xi X)| \leq |\xi|^2 \mathbf{E}(\min(|X|^2, |\xi X|^3/6))$$

By the dominated convergence theorem, as $\xi \rightarrow 0$, $\varphi_X(\xi) = 1 - \xi^2 \sigma^2/2 + o(\xi^2)$. But this means that

$$\varphi_{S_N/\sigma\sqrt{N}}(\xi) = (1 - \xi^2/2N + o(\xi^2/\sigma^2 N))^N = \exp(-\xi^2/2)$$

This implies the random variables converge weakly to a normal distribution. \square

5.5 The Wirtinger Inequality on an Interval

Theorem 5.10. Given $f \in C^1[-\pi, \pi]$ with $\int_{-\pi}^{\pi} f(t) dt = 0$,

$$\int_{-\pi}^{\pi} |f(t)|^2 \leq \int_{-\pi}^{\pi} |f'(t)|^2$$

Proof. Consider the fourier series

$$f(t) \sim \sum a_n e_n(t) \quad f'(t) \sim \sum i n a_n e_n(t)$$

Then $a_0 = 0$, and so

$$\int_{-\pi}^{\pi} |f(t)|^2 dt = 2\pi \sum |a_n|^2 \leq 2\pi \sum n^2 |a_n|^2 = \int_{-\pi}^{\pi} |f'(t)|^2 dt$$

equality holds here if and only if $a_i = 0$ for $i > 1$, in which case we find

$$f(t) = A e_n(t) + \bar{A} e_n(-t) = B \cos(t) + C \sin(t)$$

for some constants $A \in \mathbf{C}$, $B, C \in \mathbf{R}$. □

Corollary 5.11. Given $f \in C^1[a, b]$ with $\int_a^b f(t) dt = 0$,

$$\int_a^b |f(t)|^2 dt \leq \left(\frac{b-a}{\pi} \right)^2 \int_a^b |f'(t)|^2 dt$$

5.6 Energy Preservation in the String equation

Solutions to the string equation are

If $u(t, x)$

5.7 Harmonic Functions

The study of a function f defined on the real line can often be understood by extending it's definition holomorphically to the complex plane. Here we will extend this tool, establishing that a large family of functions f defined on \mathbf{R}^n can be understood by looking at a *harmonic* function on the upper half plane \mathbf{H}^{n+1} , which approximates f at it's boundary. This is a

form of the Dirichlet problem, which asks, given a domain and a function on the domain's boundary, to find a function harmonic on the interior of the domain which 'agrees' with the function on the boundary, in one of several senses. As we saw in our study of harmonic functions on the disk in the study of Fourier series, we can study such harmonic functions by convolving f with an appropriate approximation to the identity which makes the function harmonic in the plane. In this case, we shall use the Poisson kernel for the upper half plane.

Theorem 5.12. *If $f \in L^p(\mathbf{R}^n)$, for $1 \leq p \leq \infty$, and $u(x, y) = (f * P_y)(x)$, where*

$$P_y(x) = \frac{\Gamma((n+1)/2)}{\pi^{(n+1)/2}} \frac{1}{(1 + |x|^2)^{(n+1)/2}}$$

then u is harmonic in the upper half plane, $u(x, y) \rightarrow f(x)$ for almost every x , and $u(\cdot, y)$ converges to f in L^p as $y \rightarrow 0$, with $\|u(\cdot, y)\|_{L^p(\mathbf{R}^n)} \leq \|f\|_{L^p(\mathbf{R}^n)}$. If, instead, f is a continuous and bounded function, then $u(\cdot, y)$ converges to f locally uniformly as $y \rightarrow 0$.

Proof. The almost everywhere convergence and convergence in norm follow from the fact that P_y is an approximation to the identity. The fact that u is harmonic follows because

$$u_{xx}(x, y) = (f * P_y'')(x) \quad u_{yy} = (f *)$$

□

Chapter 6

Partial Derivatives and Harmonic Functions

6.1 Conjugate Poisson Kernel

Recall from our discussion of the Poisson kernel that for $f \in L^p(\mathbf{R})$ for $1 \leq p \leq \infty$, if we define

$$u(x, t) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x - t|\xi|} d\xi$$

then we obtain a harmonic function on the upper half plane. TODO If we define

$$\begin{aligned} \frac{\partial v}{\partial t} &= (2\pi i) \int_{-\infty}^{\infty} \hat{f}(\xi) \xi e^{2\pi i \xi x - t|\xi|} d\xi \\ \frac{\partial v}{\partial x} &= -|\xi| \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x - t|\xi|} d\xi. \end{aligned}$$

Chapter 7

Finite Character Theory

Let us review our achievements so far. We have found several important families of functions on the spaces we have studied, and shown they can be used to approximate arbitrary functions. On the circle group \mathbf{T} , the functions take the form of the power maps $\phi_n : z \mapsto z^n$, for $n \in \mathbf{Z}$. The important properties of these functions is that

- The functions are orthogonal to one another.
- A large family of functions can be approximated by linear combinations of the power maps.
- The power maps are multiplicative: $\phi_n(zw) = \phi_n(z)\phi_n(w)$.

The existence of a family with these properties is not dependant on much more than the symmetry properties of \mathbf{T} , and we can therefore generalize the properties of the fourier series to a large number of groups. In this chapter, we consider a generalization to any finite abelian group.

The last property of the power maps should be immediately recognizable to any student of group theory. It implies the exponentials are homomorphisms from the circle group to itself. This is the easiest of the three properties to generalize to arbitrary groups; we shall call a homomorphism from a finite abelian group to \mathbf{T} a character. For any abelian group G , we can put all characters together to form the character group $\Gamma(G)$, which forms an abelian group under pointwise multiplication $(fg)(z) = f(z)g(z)$. It is these functions which are ‘primitive’ in synthesizing functions defined on the group.

Example. If μ_N is the set of N th roots of unity, then $\Gamma(\mu_N)$ consists of the power maps $\phi_n : z \mapsto z^n$, for $n \in \mathbf{Z}$. Because

$$\phi(\omega)^N = \phi(\omega^N) = \phi(1) = 1$$

we see that any character on μ_N is really a homomorphism from μ_N to μ_N . Since the homomorphisms on μ_N are determined by their action on this primitive root, there can only be at most N characters on μ_N , since there are only N elements in μ_N . Our derivation then shows us that the ϕ_N enumerate all such characters, which completes our proof. Note that since $\phi_n \phi_m = \phi_{n+m}$, and $\phi_n = \phi_m$ if and only if $n - m$ is divisible by N , this also shows that $\Gamma(\mu_N) \cong \mu_N$.

Example. The group \mathbf{Z}_N is isomorphic to μ_N under the identification $n \mapsto \omega^n$, where ω is a primitive root of unity. This means that we do not need to distinguish functions ‘defined in terms of n ’ and ‘defined in terms of ω ’, assuming the correspondance $n = \omega^n$. This is exactly the same as the correspondence between functions on \mathbf{T} and periodic functions on \mathbf{R} . The characters of \mathbf{Z}_n are then exactly the maps $n \mapsto \omega^{kn}$. This follows from the general fact that if $f : G \rightarrow H$ is an isomorphism of abelian groups, the map $f^* : \phi \mapsto \phi \circ f$ is an isomorphism from $\Gamma(H)$ to $\Gamma(G)$.

Example. If K is a finite field, then the set K^* of non-zero elements is a group under multiplication. A rather sneaky algebraic proof shows the existence of elements of K , known as primitive elements, which generate the multiplicative group of all numbers. Thus K is cyclic, and therefore isomorphic to μ_N , where $N = |K| - 1$. The characters of K are then easily found under the correspondence.

Example. For a fixed N , the set of invertible elements of \mathbf{Z}_N form a group under multiplication, denoted \mathbf{Z}_N^* . Any character from \mathbf{Z}_N^* is valued on the $\varphi(N)$ ’th roots of unity, because the order of each element in \mathbf{Z}_N^* divides $\varphi(N)$. The groups are in general non-cyclic. For instance, $\mathbf{Z}_8^* \cong \mathbf{Z}_2^3$. However, we can always break down a finite abelian group into cyclic subgroups to calculate the character group; a simple argument shows that $\Gamma(G \times H) \cong \Gamma(G) \times \Gamma(H)$, where we identify (f, g) with the map $(x, y) \mapsto f(x)g(y)$.

7.1 Fourier Analysis on Cyclic Groups

We shall start our study of abstract Fourier analysis by looking at Fourier analysis on μ_N . Geometrically, these points uniformly distribute them-

selves over \mathbf{T} , and therefore μ_N provides a good finite approximation to \mathbf{T} . Functions from μ_N to \mathbf{C} are really just functions from $[n] = \{1, \dots, n\}$ to \mathbf{C} , and since μ_N is isomorphic to \mathbf{Z}_N , we're really computing the Fourier analysis of finite domain functions, in a way which encodes the translational symmetry of the function relative to translational shifts on \mathbf{Z}_N .

There is a trick which we can use to obtain quick results about Fourier analysis on μ_N . Given a function $f : [N] \rightarrow \mathbf{C}$, consider the N -periodic function on the real line defined by

$$g(t) = \sum_{n=1}^N f(n) \chi_{(n-1/2, n+1/2)}(t)$$

Classical Fourier analysis of g tells us that we can expand g as an infinite series in the functions $e(n/N)$, which may be summed up over equivalence classes modulo N to give a finite expansion of the function f . Thus we conclude that every function $f : [N] \rightarrow \mathbf{C}$ has an expansion

$$f(n) = \sum_{m=1}^N \hat{f}(m) e(nm)$$

where $\hat{f}(m)$ are the coefficients of the finite Fourier transform of f . This method certainly works in this case, but does not generalize to understand the expansion of general finite abelian groups.

The correct generalization of Fourier analysis is to analyze the set of complex valued 'square integrable functions' on the domain $[N]$. We consider the space V of all maps $f : [N] \rightarrow \mathbf{C}$, which can be made into an inner product space by defining

$$\langle f, g \rangle = \frac{1}{N} \sum_{n=1}^N f(n) \overline{g(n)}$$

We claim that the characters $\phi_n : z \mapsto z^n$ are orthonormal in this space, since

$$\langle \phi_n, \phi_m \rangle = \frac{1}{N} \sum_{k=1}^N \omega^{k(n-m)}$$

If $n = m$, we may sum up to find $\langle \phi_n, \phi_m \rangle = 1$. Otherwise we use a standard summation formula to find

$$\sum_{k=1}^N \omega^{k(n-m)} = \omega^{n-m} \frac{\omega^{N(n-m)} - 1}{\omega^{n-m} - 1}$$

Since $\omega^{N(n-m)} = 1$, we conclude the sum is zero. This implies that the ϕ_n are orthonormal, hence linearly independent. Since V is N dimensional, this implies that the family of characters forms an orthogonal basic for the space. Thus, for any function $f : [N] \rightarrow \mathbf{C}$, we have, if we set $\hat{f}(m) = \langle f, \phi_m \rangle$, then

$$f(n) = \sum_{m=1}^N \langle f, \phi_m \rangle \phi_m(n) = \sum_{m=1}^N \hat{f}(m) e(mn/N)$$

This calculation can essentially be applied to an arbitrary finite abelian group to obtain an expansion in terms of Fourier coefficients.

7.2 An Arbitrary Finite Abelian Group

It should be easy to guess how we proceed for a general finite abelian group. Given some group G , we study the character group $\Gamma(G)$, and how $\Gamma(G)$ represents general functions from G to \mathbf{C} . We shall let V be the space of all such functions from G to \mathbf{C} , and on it we define the inner product

$$\langle f, g \rangle = \frac{1}{|G|} \sum_{a \in G} f(a) \overline{g(a)}$$

If there's any justice in the world, these characters would also form an orthonormal basis.

Theorem 7.1. *The set $\Gamma(G)$ of characters is an orthonormal set.*

Proof. If e is a character of G , then $|e(a)| = 1$ for each a , and so

$$\langle e, e \rangle = \frac{1}{|G|} \sum_{a \in G} |e(a)| = 1$$

If $e \neq 1$ is a non-trivial character, then $\sum_{a \in G} e(a) = 0$. To see this, note that for any $b \in G$, the map $a \mapsto ba$ is a bijection of G , and so

$$e(b) \sum_{a \in G} e(a) = \sum_{a \in G} e(ba) = \sum_{a \in G} e(a)$$

Implies either $e(b) = 1$, or $\sum_{a \in G} e(a) = 0$. If $e_1 \neq e_2$ are two characters, then

$$\langle e_1, e_2 \rangle = \frac{1}{|G|} \sum_{a \in G} \frac{e_1(a)}{e_2(a)} = 0$$

since e_1/e_2 is a nontrivial character. \square

Because elements of $\Gamma(G)$ are orthonormal, they are linearly independent over the space of functions on G , and we obtain a bound $|\Gamma(G)| \leq |G|$. All that remains is to show equality. This can be shown very simply by applying the structure theorem for finite abelian groups. First, note it is true for all cyclic groups. Second, note that if it is true for two groups G and H , it is true for $G \times H$, because

$$\Gamma(G \times H) \cong \Gamma(G) \times \Gamma(H)$$

since a finite abelian group is a finite product of cyclic groups, this proves the theorem. This seems almost like sweeping the algebra of the situation under the rug, however, so we will prove the statement only using elementary linear algebra. What's more, these linear algebraic techniques generalize to the theory of unitary representations in harmonic analysis over infinite groups.

Theorem 7.2. *Let $\{T_1, \dots, T_n\}$ be a family of commuting unitary matrices. Then there is a basis $v_1, \dots, v_m \in \mathbb{C}^m$ which are eigenvectors for each T_i .*

Proof. For $n = 1$, the theorem is the standard spectral theorem. For induction, suppose that the T_1, \dots, T_{k-1} are simultaneously diagonalizable. Write

$$\mathbb{C}^m = V_{\lambda_1} \oplus \dots \oplus V_{\lambda_l}$$

where λ_i are the eigenvalues of T_k , and V_{λ_i} are the corresponding eigenspaces. Then if $v \in V_{\lambda_i}$, and $j < k$,

$$T_k T_j v = T_j T_k v = \lambda_i T_j v$$

so $T_j(V_{\lambda_i}) = V_{\lambda_i}$. Now on each V_{λ_i} , we may apply the induction hypothesis to diagonalize the T_1, \dots, T_{k-1} . Putting this together, we simultaneously diagonalize T_1, \dots, T_k . \square

This theorem enables us to prove the character theory in a much simpler manner. Let V be the space of complex valued functions on G , and define, for $a \in G$, the map $(T_a f)(b) = f(ab)$. V has an orthonormal basis consisting of the $\chi_a(b) = N[a = b]$, for $a \in G$. In this basis, we compute $T_a \chi_b = \chi_{ba^{-1}}$, hence T_a is a permutation matrix with respect to this basis, hence unitary. The operators T_a commute, since $T_a T_b = T_{ab} = T_{ba} = T_b T_a$. Hence these operators can be simultaneously diagonalized. That is, there is a family $e_1, \dots, e_n \in V$ and $\lambda_{an} \in \mathbf{T}$ such that for each $a \in G$, $T_a e_n = \lambda_{an} f_n$. We may assume $e_n(1) = 1$ for each n by normalizing. Then, for any $a \in G$, we have $f_n(a) = f_n(a \cdot 1) = \lambda_{an} f_n(1) = \lambda_{an}$, so for any $b \in G$, $f_n(ab) = \lambda_{an} f_n(b) = f_n(a) f_n(b)$. This shows each f_n is a character, completing the proof. We summarize our discussion in the following theorem.

Theorem 7.3. *Let G be a finite abelian group. Then $\Gamma(G) \cong G$, and forms an orthonormal basis for the space of complex valued functions on G . For any function $f : G \rightarrow \mathbf{C}$,*

$$f(a) = \sum_{e \in \Gamma(G)} \langle f, e \rangle e(a) = \sum_{e \in \Gamma(G)} \hat{f}(e) e(a) \quad \langle f, g \rangle = \frac{1}{|G|} \sum_{a \in G} f(a) \overline{g(a)}$$

In this context, we also have Parseval's theorem

$$\|f(a)\|^2 = \sum_{e \in \hat{G}} |\hat{f}(e)|^2 \quad \langle f, g \rangle = \sum_{e \in \hat{G}} \hat{f}(e) \overline{\hat{g}(e)}$$

7.3 Convolutions

There is a version of convolutions for finite functions, which is analogous to the convolutions on \mathbf{R} . Given two functions f, g on G , we define a function $f * g$ on G by setting

$$(f * g)(a) = \frac{1}{|G|} \sum_{b \in G} f(b) g(b^{-1}a)$$

The mapping $b \mapsto ab^{-1}$ is a bijection of G , and so we also have

$$(f * g)(a) = \frac{1}{|G|} \sum_{b \in G} f(ab^{-1}) g(b) = (g * f)(a)$$

For $e \in \Gamma(G)$,

$$\begin{aligned}\widehat{f * g}(e) &= \frac{1}{|G|} \sum_{a \in G} (f * g)(a) \overline{e(a)} \\ &= \frac{1}{|G|^2} \sum_{a, b \in G} f(ab) g(b^{-1}) \overline{e(a)}\end{aligned}$$

The bijection $a \mapsto ab^{-1}$ shows that

$$\begin{aligned}\widehat{f * g}(e) &= \frac{1}{|G|^2} \sum_{a, b} f(a) g(b^{-1}) \overline{e(a) e(b^{-1})} \\ &= \frac{1}{|G|} \left(\sum_a f(a) \overline{e(a)} \right) \frac{1}{|G|} \left(\sum_b g(b) \overline{e(b)} \right) \\ &= \widehat{f}(e) \widehat{g}(e)\end{aligned}$$

In the finite case we do not need approximations to the identity, for we have an identity for convolution. Define $D : G \rightarrow \mathbb{C}$ by

$$D(a) = \sum_{e \in \Gamma(G)} e(a)$$

We claim that $D(a) = |G|$ if $a = 1$, and $D(a) = 0$ otherwise. Note that since $|G| = |\Gamma(G)|$, the character space of $\Gamma(G)$ is isomorphic to G . Indeed, for each $a \in G$, we have the maps $\hat{a} : e \mapsto e(a)$, which is a character of $\Gamma(G)$. Suppose $e(a) = 1$ for all characters e . Then $e(a) = e(1)$ for all characters e , and for any function $f : G \rightarrow \mathbb{C}$, we have $f(a) = f(1)$, implying $a = 1$. Thus we obtain $|G|$ distinct maps \hat{a} , which therefore form the space of all characters. It therefore follows from a previous argument that if $a \neq 1$, then

$$\sum_{e \in \Gamma(G)} e(a) = 0$$

Now $f * D = f$, because

$$\widehat{D}(e) = \frac{1}{|G|} \sum_{a \in G} D(a) \overline{e(a)} = \overline{e(1)} = 1$$

D is essentially the finite dimensional version of the Dirac delta function, since it has unit mass, and acts as the identity in convolution.

7.4 The Fast Fourier Transform

The main use of the Fourier series on μ_n in applied mathematics is to approximate the Fourier transform on \mathbf{T} , where we need to compute integrals explicitly. If we have a function $f \in L^1(\mathbf{T})$, then f may be approximated in $L^1(\mathbf{T})$ by step functions of the form

$$f_n(t) = \sum_{k=1}^n a_k \mathbf{I}(x \in (2\pi(k-1)/n, 2\pi k/n))$$

And then $\hat{f}_n \rightarrow \hat{f}$ uniformly. The Fourier transform of f_n is the same as the Fourier transform of the corresponding function $k \mapsto a_k$ on \mathbf{Z}_n , and thus we can approximate the Fourier transform on \mathbf{T} by a discrete computation on \mathbf{Z}_n . Looking at the formula in the definition of the discrete transform, we find that we can compute the Fourier coefficients of a function $f : \mathbf{Z}_n \rightarrow \mathbf{C}$ in $O(n^2)$ addition and multiplication operations. It turns out that there is a much better method of computation which employs a divide and conquer approach, which works when n is a power of 2, reducing the calculation to $O(n \log n)$ multiplications. Before this process was discovered, calculation of Fourier transforms was seen as a computation to avoid wherever possible.

To see this, consider a particular division in the group \mathbf{Z}_{2n} . Given $f : \mathbf{Z}_{2n} \rightarrow \mathbf{C}$, define two functions $g, h : \mathbf{Z}_n \rightarrow \mathbf{C}$, defined by $g(k) = f(2k)$, and $h(k) = f(2k+1)$. Then g and h encode all the information in f , and if $v = e(\pi/n)$ is the canonical generator of \mathbf{Z}_{2n} , we have

$$\hat{f}(m) = \frac{\hat{g}(m) + \hat{h}(m)v^m}{2}$$

Because

$$\begin{aligned} \frac{1}{2n} \sum_{k=1}^n \left(g(k)\omega^{-km} + h(k)\omega^{-km}v^m \right) &= \frac{1}{2n} \sum_{k=1}^n f(2k)v^{-2km} + f(2k+1)v^{-(2k+1)m} \\ &= \frac{1}{2n} \sum_{k=1}^{2n} f(k)v^{-km} \end{aligned}$$

This is essentially a discrete analogue of the Poisson summation formula, which we will generalize later when we study the harmonic analysis of

abelian groups. If $H(m)$ is the number of operations needed to calculate the Fourier transform of a function on μ_{2^n} using the above recursive formula, then the above relation tells us $H(2m) = 2H(m) + 3(2m)$. If $G(n) = H(2^n)$, then $G(n) = 2G(n-1) + 32^n$, and $G(0) = 1$, and it follows that

$$G(n) = 2^n + 3 \sum_{k=1}^n 2^k 2^{n-k} = 2^n(1 + 3n)$$

Hence for $m = 2^n$, we have $H(m) = m(1 + 3 \log(m)) = O(m \log m)$. Similar techniques show that one can compute the inverse Fourier transform in $O(m \log m)$ operations (essentially by swapping the root ν with ν^{-1}).

7.5 Dirichlet's Theorem

We now apply the theory of Fourier series on finite abelian groups to prove Dirichlet's theorem.

Theorem 7.4. *If m and n are relatively prime, then the set*

$$\{m + kn : k \in \mathbf{N}\}$$

contains infinitely many prime numbers.

An exploration of this requires the Riemann-Zeta function, defined by

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

The function is defined on $(1, \infty)$, since for $s > 1$ the map $t \mapsto 1/t^s$ is decreasing, and so

$$\sum_{n=1}^{\infty} \frac{1}{n^s} \leq 1 + \int_1^{\infty} \frac{1}{t^s} = 1 + \lim_{n \rightarrow \infty} \frac{1}{s-1} [1 - 1/n^{s-1}] = 1 + \frac{1}{s-1}$$

The series converges uniformly on $[1 + \varepsilon, N]$ for any $\varepsilon > 0$, so ζ is continuous on $(1, \infty)$. As $t \rightarrow 1$, $\zeta(t) \rightarrow \infty$, because $n^s \rightarrow n$ for each n , and if for a fixed M we make s close enough to 1 such that $|n/n^s - 1| < 1/2$ for $1 \leq n \leq M$, then

$$\sum_{n=1}^{\infty} \frac{1}{n^s} \geq \sum_{n=1}^M \frac{1}{n^s} = \sum_{n=1}^M \frac{1}{n} \frac{n}{n^s} \geq \frac{1}{2} \sum_{n=1}^M \frac{1}{n}$$

Letting $M \rightarrow \infty$, we obtain that $\sum_{n=1}^{\infty} \frac{1}{n^s} \rightarrow \infty$ as $s \rightarrow 1$.

The Riemann-Zeta function is very good at giving us information about the prime integers, because it encodes much of the information about the prime numbers.

Theorem 7.5. *For any $s > 1$,*

$$\zeta(s) = \prod_{p \text{ prime}} \frac{1}{1 - p^s}$$

Proof. The general idea is this – we may write

$$\prod_{p \text{ prime}} \frac{1}{1 - p^s} = \prod_{p \text{ prime}} (1 + 1/p^s + 1/p^{2s} + \dots)$$

If we expand this product out formally, enumerating the primes to be p_1, p_2, \dots , we find

$$\prod_{p \leq n} (1 + 1/p^s + 1/p^{2s} + \dots) = \sum_{n_1, n_2, \dots = 0}^{\infty} \frac{1}{p_1^{n_1}}$$

□

Chapter 8

Complex Methods

In this chapter, we illustrate the intimate connection between the Fourier transform on the real line, and complex analysis. We have already seen some aspects of this for Fourier analysis on the Torus, with the connection between power series of analytic functions on the unit disk. The main theme is that if f is a function initially defined on the real line, then the problem of extending the function to be analytic on a neighbourhood of this line is connected to the Fourier transform of f decaying very rapidly (for instance, exponential decay).

8.1 Fourier Transforms of Holomorphic Functions

For each $a > 0$, let $S_a = \{x + iy : |y| < a\}$ denote the horizontal strip of width $2a$. The next theorem says that functions extendable to be holomorphic on the strip have exponential Fourier decay.

Theorem 8.1. *Let $f : S_a \rightarrow \mathbb{C}$ be holomorphic, integrable on each horizontal line in the strip, such that $f(x + iy) \rightarrow 0$ as $|x| \rightarrow \infty$. Then if \hat{f} is the Fourier transform of the restriction of f to the real line, then for each $b < a$,*

$$|\hat{f}(\xi)| \lesssim_b e^{-2\pi b|\xi|}.$$

Proof. For any $b < a$, R , and $\xi > 0$, consider the contour γ_R on the rectangle with corners $-R$, R , $-R - ib$, and $R - ib$. As $R \rightarrow \infty$, the integral along the

vertical lines of the rectangle tends to zero as $R \rightarrow \infty$, so we conclude that

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) e^{-2\pi i x \xi} dx &= \int_{-\infty}^{\infty} f(x - ib) e^{-2\pi i (x - ib) \xi} dx \\ &= e^{-2\pi i b \xi} \int_{-\infty}^{\infty} f(x - ib) e^{-2\pi i \xi x} dx = e^{-2\pi i b \xi} \hat{f}_b(\xi) \end{aligned}$$

where $f_b(x) = f(x - ib)$. But $|\hat{f}_b(\xi)| \leq \|f_b\|_{L^\infty(\mathbf{R})} \lesssim_b 1$, which implies that

$$|\hat{f}(\xi)| \lesssim_b e^{-2\pi i b \xi}.$$

A similar estimate when $\xi < 0$ completes the argument. \square

It follows that \hat{f} has exponential decay if f satisfies the hypothesis of the theorem. Thus we can always apply the inverse Fourier transform to conclude

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x} d\xi.$$

Conversely, if f is *any* integrable function with $|\hat{f}(\xi)| \lesssim e^{-2\pi a|\xi|}$, then \hat{f} is integrable so the Fourier inversion formula holds. If we define

$$f(x + iy) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{-2\pi \xi y} e^{2\pi i \xi x} d\xi,$$

then this gives a holomorphic extension of f which is well defined on S_a .

Pushing this result to an extreme leads to the Paley-Wiener theorem, which gives precise conditions when a function has a compactly supported Fourier transform.

Theorem 8.2. *A function $f : \mathbf{R} \rightarrow \mathbf{C}$ is bounded, integrable, and continuous. Then f extends to an entire function on the complex plane, such that for all z ,*

$$|f(z)| \lesssim e^{2\pi M|z|},$$

if and only if \hat{f} is supported on $[-M, M]$.

Proof. If \hat{f} is supported on $[-M, M]$, then the Fourier inversion formula comes into play, telling us that for all $x \in \mathbf{R}$,

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x} d\xi = \int_{-M}^M \hat{f}(\xi) e^{2\pi i \xi x} d\xi.$$

But then we can clearly extend f to an entire function by defining

$$f(z) = \int_{-M}^M \hat{f}(\xi) e^{2\pi i \xi z} d\xi,$$

and then $|f(z)| \leq e^{2\pi i M|z|} \|\hat{f}\|_{L^1[-M, M]} \lesssim e^{2\pi i M|z|}$.

Conversely, suppose f is an entire function such that for all $z \in \mathbf{C}$,

$$|f(z)| \leq A g(x) e^{2\pi M|y|},$$

where $g \geq 0$ is integrable on \mathbf{R} . We also assume that $f(x + iy) \rightarrow 0$ uniformly as $x \rightarrow -\infty$, independently of y . Then a contour shift down guarantees that for any y ,

$$\begin{aligned} \hat{f}(\xi) &= \int_{-\infty}^{\infty} f(x) e^{-2\pi i \xi x} dx \\ &= \int_{-\infty}^{\infty} f(x - iy) e^{-2\pi i \xi (x - iy)} dx \\ &\leq A e^{2\pi M y - 2\pi \xi y} \int_{-\infty}^{\infty} g(x) dx \lesssim e^{2\pi(M y - \xi y)}. \end{aligned}$$

If $\xi > M$, then taking $y \rightarrow \infty$ shows $\hat{f}(\xi) = 0$. A contour shift up instead gives $\hat{f}(\xi) = 0$ if $\xi < -M$. Thus the proof is completed in this case.

Now suppose the weaker condition

$$|f(z)| \leq A e^{2\pi M|y|}.$$

For each $\varepsilon > 0$, let

$$f_\varepsilon(z) = \frac{f(z)}{(1 - i\varepsilon z)^2}.$$

Then f_ε is analytic in the lower half plane. Moreover,

$$|f_\varepsilon(x + iy)| \lesssim_\varepsilon \frac{A e^{2\pi M|y|}}{1 + x^2}.$$

Thus we can apply the previous shifting techniques to show that $\hat{f}_\varepsilon(\xi) = 0$ for $\xi > M$. For $x \in \mathbf{R}$, we have $|f_\varepsilon(x)| \leq |f(x)|$, and since $f_\varepsilon \rightarrow f$ pointwise as $\varepsilon \rightarrow 0$, we can apply the dominated convergence theorem to imply $\hat{f}_\varepsilon(\xi) \rightarrow$

$\hat{f}(\xi)$ for each ξ . In particular, we find $\hat{f}(\xi) = 0$ for $\xi > M$. A similar technique with the family of functions

$$f_\varepsilon(z) = \frac{f(z)}{(1 + i\varepsilon z)^2},$$

show that $\hat{f}(\xi) = 0$ for $\xi < -M$.

Finally, it suffices to show that the condition

$$|f(z)| \lesssim e^{2\pi M|z|}$$

implies $|f(x + iy)| \lesssim e^{2\pi M|y|}$. To prove this, we can apply a version of the Phragmén-Lindelöf on the quadrant $\{x + iy : x, y > 0\}$. Let $g(z) = f(z)e^{-2\pi iMy}$. Then we have

$$|g(x)| = |f(x)| \leq \|f\|_{L^\infty(\mathbf{R})},$$

and

$$|g(iy)| = |f(iy)|e^{-2\pi My} \leq A.$$

Since g has at most exponential growth on the quadrant, we can apply the Phragmén-Lindelöf to conclude $|g(z)| \leq \max(A, \|f\|_{L^\infty(\mathbf{R})})$ for all z on the quadrant. A similar argument works for the other quadrants. Thus we conclude that for all $z \in \mathbf{C}$

$$|f(z)| \leq \max(A, \|f\|_{L^\infty(\mathbf{R})})e^{2\pi M|y|},$$

and so we can apply the previous cases to conclude that \hat{f} is supported on $[-M, M]$. \square

Remark. The Paley-Wiener theorem has several variants. For instance, if f is continuous, integrable, and \hat{f} is integrable, and we further assume that $\hat{f}(\xi) = 0$ for all $\xi < 0$, then for $z = x + iy$, we can define

$$f(z) = \int_0^\infty \hat{f}(\xi)e^{2\pi i\xi z} = \int_0^\infty \hat{f}(\xi)e^{-2\pi \xi y}e^{2\pi i\xi x}$$

to extend f to an analytic function in the upper half-plane, i.e. for $y > 0$, which is also continuous and bounded for $y \geq 0$. Conversely, similar techniques to those above enable us to show that if f is continuous, integrable, \hat{f} is integrable, and we can extend f to an analytic function on the open upper half plane, which is continuous and bounded on the closed half plane, then contour shifting shows that $\hat{f}(\xi) = 0$ for $\xi < 0$.

8.2 Classical Theorems by Contours

We now prove some classical theorems of Fourier analysis using techniques of harmonic analysis, given that the functions we study have holomorphic extensions to tubes.

Theorem 8.3. *Let $f : S_b \rightarrow \mathbf{C}$ be holomorphic. Then for any $x \in \mathbf{R}$,*

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i \xi x} d\xi,$$

where \hat{f} is the Fourier transform of f restricted to the real-axis.

Proof. As in the last theorem, the sign of ξ matters. We write

$$\int_{-\infty}^{\infty} \hat{f}(\xi) e^{-2\pi i \xi x} d\xi = \int_0^{\infty} \hat{f}(\xi) e^{-2\pi i \xi x} + \hat{f}(-\xi) e^{2\pi i \xi x} d\xi.$$

Now if $b < a$, we can apply a contour integral argument to conclude that

$$\begin{aligned} \hat{f}(\xi) &= \int_{-\infty}^{\infty} f(x - ib) e^{-2\pi i \xi (x - ib)} dx \\ &= \int_{-\infty}^{\infty} f(x + ib) e^{2\pi i \xi (x + ib)} dx. \end{aligned}$$

Thus by Fubini's theorem, for each $x_0 \in \mathbf{R}$,

$$\begin{aligned} \int_0^{\infty} \hat{f}(\xi) e^{2\pi i \xi x_0} d\xi &= \int_0^{\infty} \int_{-\infty}^{\infty} f(x - ib) e^{2\pi i \xi [x_0 - (x - ib)]} dx d\xi \\ &= \int_{-\infty}^{\infty} f(x - ib) \left(\int_0^{\infty} e^{2\pi i \xi [x_0 - (x - ib)]} d\xi \right) dx \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{f(x - ib)}{(x - ib) - x_0} dx. \end{aligned}$$

Similarly, another application of Fubini's theorem implies

$$\begin{aligned} \int_0^{\infty} \hat{f}(-\xi) e^{-2\pi i \xi x_0} d\xi &= \int_0^{\infty} \int_{-\infty}^{\infty} f(x + ib) e^{-2\pi i \xi [x_0 - (x + ib)]} dx d\xi \\ &= \int_{-\infty}^{\infty} f(x + ib) \int_0^{\infty} e^{-2\pi i \xi [x_0 - (x + ib)]} d\xi dx \\ &= \frac{-1}{2\pi i} \int_{-\infty}^{\infty} \frac{f(x + ib)}{[(x + ib) - x_0]} dx. \end{aligned}$$

In particular, we conclude that

$$\int \hat{f}(\xi) e^{2\pi i \xi x_0} = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - x_0},$$

where γ is the path traces over the two horizontal strips $x + ib$ and $x - ib$. Approximating this integral by rectangles, and then apply Cauchy's theorem, we find this value is equal to $f(x)$. \square

We can also prove the Poisson summation formula.

Theorem 8.4. *Let $f : S_a \rightarrow \mathbf{C}$ be holomorphic. Then*

$$\sum_{n \in \mathbf{Z}} f(n) = \sum_{n \in \mathbf{Z}} \hat{f}(n),$$

where \hat{f} is the Fourier transform of f restricted to the real line.

Proof. The function

$$\frac{f(z)}{e^{2\pi iz} - 1}$$

is meromorphic, with simple poles on \mathbf{Z} , with residue equal to $f(n)$ at each $n \in \mathbf{Z}$. If we apply the Residue theorem to a curve γ_N travelling around the rectangle connecting the points $N + 1/2 - ib$, $N + 1/2 + ib$, $-N - 1/2 + ib$, and $-N - 1/2 - ib$, then we conclude

$$\sum_{|n| \leq N} f(n) = \int_{\gamma_N} \frac{f(z)}{e^{2\pi iz} - 1} dz.$$

These values converge to $\sum_{n \in \mathbf{Z}} f(n)$ as $N \rightarrow \infty$. But this means that

$$\sum_n f(n) = \int_{\gamma} \frac{f(z)}{e^{2\pi iz} - 1} dz,$$

where γ is the two horizontal strips at b and $-b$. Now we use the expansion

$$\frac{1}{z - 1} = \sum_{n=1}^{\infty} z^{-n},$$

for $|z| > 1$, to conclude

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{f(x-ib)}{e^{2\pi i(x-ib)} - 1} dx &= \int_{-\infty}^{\infty} \sum_{n=1}^{\infty} \frac{f(x-ib)}{e^{2\pi ni(x-ib)}} dx \\ &= \sum_{n=1}^{\infty} \int_{-\infty}^{\infty} f(x-ib) e^{-2\pi ni(x-ib)} dx = \sum_{n=1}^{\infty} \hat{f}(n), \end{aligned}$$

where we have performed a contour shift at the end. Similarly, we use the expansion

$$\frac{1}{z-1} = - \sum_{n=0}^{\infty} z^n,$$

to conclude that

$$\begin{aligned} - \int_{-\infty}^{\infty} \frac{f(x+ib)}{e^{2\pi i(x+ib)} - 1} dx &= \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} f(x+ib) e^{2\pi ni(x+ib)} dx \\ &= \sum_{n=0}^{\infty} \hat{f}(-n). \end{aligned}$$

Combining these two calculations completes the proof. \square

8.3 The Laplace Transform

We now look at things from the dual perspective. Instead of looking at whether a function can be extended to a holomorphic function, we look at whether the Fourier transform can be extended to a holomorphic function. For a function $x : \mathbf{R} \rightarrow \mathbf{R}$, this gives rise to the *Laplace transform*

$$X(z) = \int_{-\infty}^{\infty} x(t) e^{-zt} dt,$$

also denoted by $(\mathcal{L}x)(z)$. For $\xi \in \mathbf{R}$, $X(i\xi) = \hat{x}(\xi)$ operates as the usual Fourier transform (slightly rescaled from the version in our notes). But the Laplace transform can also be extended to not-necessarily integrable functions. Given x , we can define $X(z)$ for any $z = x + iy$ such that

$$\int e^{-xt} |f(t)| dt < \infty.$$

It is simple to see this forms a vertical tube in the complex plane, called the *region of convergence* for the Laplace transform. For a particular vertical tube $I \subset \mathbb{C}$, we let $\mathcal{E}(I)$ be the collection of all functions x whose region of convergence for the Dirichlet transform contains I .

Example. Let

$$H(x) = \begin{cases} 0 & : x < 0, \\ 1/2 & : x = 0, \\ 1 & : x > 0. \end{cases}$$

The function H is called the Heavyside Step Function. Its region of convergence consists of the right-most half plane, i.e. all $\omega + i\xi$, where $\omega > 0$. And if $z = \omega + i\xi$, we calculate that

$$\mathcal{L}(H)(z) = \int_0^\infty e^{-zt} dt = z^{-1}.$$

We note that even though the integral formula does not define the Laplace transform of H in the right-most half plane, we can analytically continue $\mathcal{L}(H)$ to a meromorphic function on the entire complex plane.

Example. Similarly, an integration by parts shows that for $z = \omega + i\xi$ with $\omega > 0$, we have

$$\mathcal{L}(tH)(z) = \int_0^\infty te^{-zt} = \int_0^\infty \frac{e^{-zt}}{z} = z^{-2}.$$

Against, $\mathcal{L}(tH)$ extends to a meromorphic function on the entire complex plane.

What distinguishes the Laplace transform from the Fourier transform is the ability to use techniques of complex analysis. If x has region of convergence I , then X is continuous on I , and analytic on I° . We can even calculate an explicit formula for the derivative. As expected from the Fourier transform of the derivative, if $y(t) = tx(t)$, and Y is the Laplace transform of y , then $X'(z) = -Y(z)$. One can verify this quite simply by taking limits of the derivatives of the analytic integrals

$$\int_{-N}^N x(t)e^{-zt} dt,$$

as $N \rightarrow \infty$. Like the Fourier transform, the Laplace transform is symmetric under modulation, translation, and polynomial multiplication:

- If $w \in \mathbf{C}$, and x is a function, set $y(t) = e^{wt}x(t)$. Then if z is in the region of convergence for x , $z - w$ is in the region of convergence for y , and $X(z) = Y(z - w)$.
- If x has region of convergence I , then the region of convergence for $y(t) = tx(t)$ contains I° , and $Y(z) = -X(z)$.
- If x has region of convergence I , $t_0 \in \mathbf{R}$, and we set $y(t) = x(t + t_0)$, then y has region of convergence I , and $Y(z) = e^{zt_0}X(z)$.
- For a function x , define

$$(\Delta_s x)(t) = \frac{x(t+s) - x(t)}{s}.$$

If ω is fixed, if

$$\lim_{s \rightarrow 0} \int |(\Delta_s x)(t) - x'(t)| e^{-\omega t} dt = 0,$$

if $y(t) = x'(t)$, and if $z = \xi + i\omega$ for some $\xi \in \mathbf{R}$, then $Y(z) = zX(z)$.

In particular, this is true if x is supported on $[-N, \infty)$ for some N , has a continuous derivative x' , and there is $\omega_0 < \omega$ such that

$$\lim_{t \rightarrow \infty} x(t)e^{-\omega_0 t} = \lim_{t \rightarrow \infty} x'(t)e^{-\omega_0 t} = 0.$$

Remark. It will be interesting for us to consider functions x supported on $[-N, \infty)$ which have a piecewise continuous derivative x' except at finitely many points t_1, \dots, t_N , such that the left and right-hand limits exist at each t_i . For each $i \in \{1, \dots, N\}$, we let

$$A_i = x(t_i+) - x(t_i-) \quad \text{and} \quad B_i = x'(t_i+) - x'(t_i-).$$

If $y(t) = x'(t)$, we calculate a relation between the Laplace transforms of X and Y at $z = \omega + i\xi$ such that there exists $\omega_0 < \omega$ such that

$$\lim_{t \rightarrow \infty} x(t)e^{-\omega_0 t} = \lim_{t \rightarrow \infty} x'(t)e^{-\omega_0 t} = 0.$$

We consider the function

$$x_1(t) = x(t) - \sum_{i=1}^N A_i H(t - t_i) - \sum_{i=1}^N B_i (t - t_i) H(t - t_i).$$

Then x_1 is continuous everywhere, and moreover, has a continuous derivative. We have

$$x_1'(t) = x'(t) - \sum_{i=1}^N B_i H(t - t_i).$$

Thus if $\omega > 0$, and $z = \omega + i\xi$, if $y_1(t) = x_1'(t)$, we find

$$Y_1(z) = zX_1(z).$$

Now

$$Y_1(z) = Y(z) - \sum_{i=1}^N \frac{B_i e^{-izt_i}}{iz}$$

and

$$X_1(z) = X(z) - \sum_{i=1}^N \frac{A_i e^{-izt_i}}{iz} + \sum_{i=1}^N \frac{B_i e^{-izt_i}}{z^2}.$$

Thus, rearranging, we conclude

$$Y(z) = zX(z) - \sum_{i=1}^N A_i e^{-izt_i}$$

We can carry this through recursively to higher order derivatives. For each k , we set $A_i^k = f^{(k)}(t_i+) - f^{(k)}(t_i-)$. Then if $y(t) = f^{(n)}(t)$, then

$$Y(z) = z^n X(z) - \sum_{k=0}^{n-1} \sum_{i=1}^N z^{n-1-k} A_i^k e^{-izt_i}.$$

This is very useful when wants to solve differential equations, provided the solutions to those differential equations do not grow faster than exponentially.

Example. Suppose we wish to find a formula for the unique real-valued function $x : [0, \infty) \rightarrow \mathbf{R}$ such that $x''(t) - x'(t) - 6x(t) = 5e^{3t}$ for $t \geq 0$, such that $x(0) = 6$ and $x'(0) = 1$. Such a function increases at most exponentially, since it is linear, so we may take the Laplace transform of each sides to conclude that if X is the Laplace transform of x , then

$$\mathcal{L}(x'')(z) = z^2 X(z) - 6z - 1 \quad \text{and} \quad \mathcal{L}(x')(z) = zX(z) - 6.$$

Thus we conclude

$$[z^2X(z) - 6z - 1] - [zX(z) - 6] - (6X) = \frac{5}{z-3}.$$

Thus

$$X(z) = \frac{(3z-4)(2z-5)}{(z-3)^2(z+2)} = \frac{3.6}{z+2} + \frac{2.4}{z-3} + \frac{1}{(z-3)^2}.$$

But this implies that for $t \geq 0$, $x(t) = 3.6e^{-2t} + 2.4e^{3t} + te^{3t}$. In particular, we note that the pole of X determines the large scale behaviour of X , i.e. for large t , and for any $\varepsilon > 0$,

$$e^{(3-\varepsilon)t} \lesssim_\varepsilon x(t) \lesssim_\varepsilon e^{(3+\varepsilon)t}.$$

In the next section, we generalize this situation to give asymptotics of functions whose Laplace transforms extend to meromorphic functions on the complex plane.

8.4 Asymptotics via the Laplace Transform

For simplicity, in this chapter we study integrable functions $x : [0, \infty) \rightarrow \mathbf{R}$, whose Laplace transform is thus well defined on the closed, right half-plane. If the Fourier transform of x is integrable, then we can apply the inversion formula to conclude that for each $t \in \mathbf{R}$,

$$x(t) = \int_{-\infty}^{\infty} X(i\xi) e^{i\xi t} d\xi.$$

Now suppose that X can be analytically continued to a holomorphic function $X(\omega + i\xi)$ for all $\omega \geq -\varepsilon$ which is continuous at the boundary, such that, uniformly for $\omega \in [-\varepsilon, 0]$,

$$\lim_{|\xi| \rightarrow \infty} X(\omega + i\xi) = 0.$$

Then a contour shift argument implies that for each t ,

$$x(t) = \lim_{R \rightarrow \infty} \int_{-R}^R X(-\varepsilon + i\xi) e^{(-\varepsilon + i\xi)t} d\xi = e^{-\varepsilon t} \lim_{R \rightarrow \infty} \int_{-R}^R X(-\varepsilon + i\xi) e^{i\xi t} d\xi.$$

For simplicity, we study functions supported on $[0, \infty)$. The region of convergence for such functions then takes the form of a half plane. For a given $a \in \mathbf{R}$, we let \mathcal{E}_a be the set of functions whose region of convergence contains $\omega + i\xi$ for all $\omega > a$.

Theorem 8.5. Suppose $x : [0, \infty) \rightarrow \mathbf{R}$ is a continuous function such that some ω ,

$$\int |x(t)|e^{-\omega t} dt < \infty.$$

Proof. Since $|X(u + iv)| \rightarrow 0$ uniformly as $v \rightarrow \infty$, we can shift the Fourier inversion formula

$$x(t) = \lim_{R \rightarrow \infty} \frac{1}{2\pi} \int_{-R}^R X(\omega + i\xi) e^{(\omega + i\xi)t} d\xi$$

(where the 2π comes up from our rescaling of the Fourier transform) to conclude that

$$x(t) = \lim_{R \rightarrow \infty} \frac{1}{2\pi}$$

$$X(z) = \lim$$

□

Part II

Distributional Methods

Chapter 9

The Theory of Distributions

Distribution theory is a tool which enables us to justify formal manipulations in harmonic analysis without having to worry about technical analytical issues. For instance, the theory allows us to take the formal derivative of a much more general class of functions than those differentiable pointwise by methods of the classical differential calculus. That such a consistent formal definition of the derivative is possible is hinted at by equations in Fourier analysis such as

$$D^i f = \mathcal{F}^{-1} \left\{ M^i \mathcal{F}\{f\} \right\},$$

where $M^i f(x) = (2\pi i x_i) f(x)$, and $D^i f$ is differentiation in the variable x_i . The quantity $D^i f$ can only be classically interpreted if f is pointwise differentiable in the variable x_i , whereas the right hand side is defined for a much less regular class, namely the family of integrable functions f such that $M^i \widehat{F}\{f\}$ is also integrable. We will find a common domain, a class of generalized functions called tempered distributions, such that both sides of the equation above can be correctly interpreted.

If we consider solutions $u \in C_{\text{loc}}^2(\mathbf{R}^d)$ to elliptic partial differential equations like the Laplace equation $\Delta u = 0$, then the regularity theory of such solutions shows the family of such functions is closed under locally uniform convergence, i.e. in the space $C_{\text{loc}}(\mathbf{R}^d)$. This is no longer true for solutions to certain non-elliptic partial differential equations, such as solutions to the wave equation $\partial_t u = \Delta_x u$ in $C_{\text{loc}}^2(\mathbf{R} \times \mathbf{R}^d)$. For instance, the closure of such a class under uniform convergence would have to contain

all functions of the form

$$f(x+t) + g(x-t)$$

for any $f, g \in C_{\text{loc}}(\mathbf{R} \times \mathbf{R}^d)$. To fix this problem, in PDE it is often of interest to enlarge the class of solutions to consider *weak solutions*, i.e. functions $u \in L^1_{\text{loc}}(\mathbf{R}^d)$ such that for any $\phi \in C_c^\infty(\mathbf{R} \times \mathbf{R}^d)$,

$$\int u(t, x)(\Delta \phi - \partial_t \phi)(t, x) dt dx = 0,$$

an equation true for any classical solution, as verified by integration by parts. This class is then clearly closed under uniform convergence, i.e. in $L^\infty_{\text{loc}}(\mathbf{R}^d)$. The step to distributional solutions to partial differential equations from classical solutions is then not far off, since it allows us to apply differentiable operators to $u \in L^1_{\text{loc}}$, and we find that u will be a weak solution to the wave equation if and only if $\partial_t u = \Delta_x u$. We will also find that the family of *distributional* solutions to the wave equation is closed under a very general topology. These properties are one reason why distributions are a core tool to the formulation of many problems in partial differential equations and modern harmonic analysis.

9.1 Distributions

The path of modern analysis has extended analysis from the study of continuous and differentiable functions to the much larger class of measurable functions. The power of this approach is that we can study a very general class of functions. On the other hand, the more general the class of functions we work with, the more restricted the analytical operations we can perform, e.g. one cannot differentiate elements of $L^1(\mathbf{R}^d)$ in a classical sense. Nonetheless, in pretty much all function spaces we consider on \mathbf{R}^d , the space $C_c^\infty(\mathbf{R}^d)$ forms a dense subclass, and in this subclass we can perform pretty much all possible analytical operations. One approach to studying the general class of measurable functions is to prove results for elements of $C_c^\infty(\mathbf{R}^d)$, in which one can apply all necessary analytical operations, and then apply an approximation result to obtain the result for a wider class of measurable functions. The theory of distributions provides a complimentary approach, using *duality* to formally extend analytical operations on $C_c^\infty(\mathbf{R}^d)$ to larger sets.

From the perspective of set theory, a function $f : X \rightarrow Y$ is a rule for assigning a value in Y to each point in X . However, in analysis this perspective is often not the most useful. This is most clear in measure theory, where we are very used to treating a function as defined ‘up to a set of measure zero’, which has the peculiar feature of a function not actually being defined at any particular point. Distribution theory instead views functions as ‘integrands’, whose properties are understood by integration against a family of ‘test functions’.

As an example of this phenomenon, since the dual space of $L^1(\mathbf{R}^d)$ is $L^\infty(\mathbf{R}^d)$, we can think of elements $f \in L^\infty(\mathbf{R}^d)$ as ‘integrands’, which can be completely understood by ‘testing’ f against an element $\phi \in L^1(\mathbf{R}^d)$, i.e. via a study of the quantities

$$\int f(x)\phi(x) dx.$$

Similarly, if K is a compact topological space, then the dual space to $C(K)$ is the space $M(K)$ of finite Borel measures on K . Thus we can think of the class of measures $M(K)$ as a family of ‘generalized integrands’ which can be understood by testing these measures against continuous functions.

Notice that as we shrink the family of test functions, the resulting family of ‘generalized functions’ becomes larger and larger, and so elements are allowed to behave more erratically. A distribution is a ‘generalized function’ understood by testing against the very regular family of functions in the space $C_c^\infty(\mathbf{R}^d)$, functions which in this situation are classical denoted as $\mathcal{D}(\mathbf{R}^d)$. Because $\mathcal{D}(\mathbf{R}^d)$ is very small, distributions can behave very erratically. But because most operations in analysis can be applied to elements of $\mathcal{D}(\mathbf{R}^d)$ and behave well under duality, we can then use duality to extend these operations to distributions. The class $\mathcal{D}(\mathbf{R}^d)$, or the slightly more general class $\mathcal{S}(\mathbf{R}^d)$ of Schwartz functions to be introduced shortly, is the natural class of functions for most problems studied in harmonic analysis.

One can apply the ideas described in this chapter to many other classes of test functions. Provided that the test functions can be suitably localized, one will likely obtain similar results to that described in this chapter. On the other hand, if one deals with non localizable families of test functions, one is likely to obtain quite a different theory of generalized functions. This is encountered, for instance, if one takes the family of analytic functions as the test functions, which gives the theory of *hyperfunctions*.

Remark. From the perspective of experimental physics, viewing functions as integrands is more natural than viewing functions in the set-theoretic sense. Indeed, points in space are idealizations which do not correspond to real world phenomena. One can never measure the exact value of some quantity of a function at a point, but rather only understand the function by looking at it's averages over a small region around that point. Thus the only physically meaningful properties of a 'function' are those obtained by testing that function against some family of test functions, obtained from some physical measurements. And indeed, physicists had experimented with using distributions in their calculations before a formal definition was introduced by mathematicians.

9.2 Test Functions and Distributions

Fix an open subset Ω of \mathbf{R}^d . There are various families of test functions we can use, which give a theory of distributions:

- We can use the space $\mathcal{D}(\Omega) = C_c^\infty(\Omega)$ of compactly supported smooth functions, under the topology of compact convergence, e.g. the LF space consisting of the inductive limit of the Fréchet spaces $C_c^\infty(K)$ of smooth functions on \mathbf{R}^d , which are compactly supported on a compact set K , equipped with the topology of uniform convergence of the functions and all of their derivatives. This gives the most general space of distributions $\mathcal{D}(\Omega)^*$.
- We can use the space $\mathcal{S}(\mathbf{R}^d)$ of Schwartz functions on \mathbf{R}^d , i.e. the Fréchet space consisting of functions whose derivatives are all rapidly decreasing, i.e. all functions f such that

$$\|f\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)} = \sup_{x \in \mathbf{R}^d} \sup_{|\alpha| \leq m} |D^\alpha f(x)| \langle x \rangle^n$$

is finite for all integers n and m . Schwartz space becomes a Fréchet space if we treat these as continuous seminorms on the space. These test functions then give the space of *tempered distributions* $\mathcal{S}(\mathbf{R}^d)^*$.

- We can use the space $\mathcal{E}(\Omega) = C_{\text{loc}}^\infty(\Omega)$ of all smooth functions on Ω , with the topology of locally uniform convergence of the function and all of it's derivatives. Using these functions as test functions gives the space $\mathcal{E}(\Omega)^*$ of *compactly supported distributions*.

- We can use the space $\mathcal{D}^n(\Omega) = C_c^n(\Omega)$ of compactly supported n -times continuously differentiable functions, under the topology of compact convergence, e.g. with the LF topology analogous to that of $\mathcal{D}(\Omega)$. This gives the space $\mathcal{D}^n(\Omega)^*$ of *distributions of order n* .

We begin by studying the space $\mathcal{D}(\Omega)$. As an LF space, $\mathcal{D}(\Omega)$ is *not metrizable*. This is in some sense necessary; it is *impossible* to make $C_c^\infty(\Omega)$ into a Fréchet space, such that each of the subspaces

$$C_c^\infty(K) = \{f \in C_c^\infty(\Omega) : \text{supp}(f) \subset K\}$$

are closed. The interior of each of these subspaces is empty because no translation of these subspaces is absorbing. The Baire category theorem would then imply that a countable union of such subspaces has nonempty interior, but $C_c^\infty(\Omega)$ is the countable union of such spaces. Nonetheless, as an LF space $\mathcal{D}(\Omega)$ continues to have most of the useful properties of Fréchet spaces. In particular, it follows from the general theory of LF spaces that $\mathcal{D}(\Omega)$ is a complete, locally convex space. The space $\mathcal{D}(\Omega)$ is also a Montel space, since it is the inductive limit of Montel spaces.

We recall some useful properties of the inductive limit. A sequence $\{\phi_n\}$ converges in $\mathcal{D}(\Omega)$ to ϕ if $\bigcup_n \text{supp}(\phi_n)$ is precompact, and if $\{\phi_n\}$ and all derivatives of this sequence converge uniformly to ϕ . The space is Bornological, so that every bounded linear map $T : \mathcal{D}(\Omega) \rightarrow Y$ is continuous, and so in particular, all sequentially continuous linear maps are continuous. Moreover, if X is first countable, then for any bounded linear map $T : X \rightarrow \mathcal{D}(\Omega)$, there exists a compact set $K \subset \Omega$ such that $T(X) \subset K$. The Arzela-Ascoli theorem implies that for any compact set K , each of the spaces $C_c^\infty(K)$ has the Heine-Borel property, and so $\mathcal{D}(\Omega)$ also has the Heine-Borel property, i.e. $\mathcal{D}(\Omega)$ is a Montel space.

TODO: Move to tensor products

Theorem 9.1. *For any two open sets Ω_1 and Ω_2 , $C_{loc}^\infty(\Omega_1 \times \Omega_2)$ is naturally isomorphic to $C_{loc}^\infty(\Omega_2, C_{loc}^\infty(\Omega_1))$.*

Proof. The correspondence is obtained by mapping $f \in C_{loc}^\infty(\Omega_2, C_{loc}^\infty(\Omega_1))$ to $F \in C_{loc}^\infty(\Omega_1 \times \Omega_2)$ given by setting $F(x, y) = f(y)(x)$. The map is clearly one to one and continuous. If F is an arbitrary element of $C_{loc}^\infty(\Omega_1 \times \Omega_2)$, and we define $f[y](x) = F(x, y)$, then the mean-value theorem implies that the quantities

$$\frac{D_x^\alpha F(x, y + he_i) - D_x^\alpha F(x, y)}{h}$$

converge locally uniformly in x to $D_y^i F$, and thus f lies in $C^1(\Omega_2, C^\infty(\Omega_1))$. But iterating this process shows that f is actually C^∞ . Thus the correspondence $f \mapsto F$ is a bijection. and since both spaces are Fréchet spaces, the open mapping theorem implies that the correspondence is an isomorphism. \square

Theorem 9.2. *Finite sums of tensor functions are dense in $\mathcal{D}(\mathbf{R}^d)$.*

Proof. Recall from the theory of multiple Fourier series that if $f \in C^\infty(\mathbf{R}^d)$ is N periodic, in the sense that $f(x+n) = f(x)$ for all $x \in \mathbf{R}^d$ and $n \in (N\mathbf{Z})^d$, then there are coefficients a_m for each $m \in \mathbf{Z}^d$ such that $f = \lim_{M \rightarrow \infty} S_M f$, where the convergence is dominated by the seminorms $\|\cdot\|_{C^n(\mathbf{R}^d)}$, for all $n > 0$, and

$$(S_M f)(x) = \sum_{\substack{m \in \mathbf{Z}^d \\ |m| \leq M}} a_m e^{\frac{2\pi i m \cdot x}{N}}.$$

Note that since

$$e^{\frac{2\pi i m \cdot x}{N}} = \prod_{k=1}^d e^{2\pi i m_i x_i / N}$$

is a tensor product, $S_M f$ is a finite sum of tensor functions. If $\phi \in \mathcal{D}(\mathbf{R}^d)$ is compactly supported on $[-N, N]^d$, we let f be a $10N$ periodic function which is equal to ϕ on $[-N, N]^d$. We then find coefficients $\{a_m\}$ such that $S_M f$ converges to f . If $\psi : \mathbf{R} \rightarrow \mathbf{R}$ is a compactly supported bump function equal to one on $[-N, N]^d$, and vanishing outside of $[-2N, 2N]^d$, then $\psi^{\otimes d} S_M f$ converges to ψ as $M \rightarrow \infty$, and each is a finite sum of tensor functions. \square

Because convergence is so strict in $\mathcal{D}(\Omega)$, almost every operation we want to perform on smooth functions is continuous in this space.

- Since $f \mapsto D^\alpha f$ is a continuous operator from $C_c^\infty(K)$ to itself, it is therefore continuous on the entire space $\mathcal{D}(\Omega)$. More generally, any linear differential operator with coefficients in $\mathcal{D}(\Omega)$ is a continuous operator on $\mathcal{D}(\Omega)$.
- The inclusion $\mathcal{D}(\Omega) \rightarrow L_c^p(\Omega)$ is continuous. To prove this, it suffices to prove for each compact K , the inclusion $C_c^\infty(K) \rightarrow L^p(K)$ is continuous, and this follows because if $f \in C_c^\infty(K)$, then

$$\|f\|_{L^p(K)} \leq |K|^{1/p} \|f\|_{L^\infty(K)}.$$

- Multiplication gives a continuous operator

$$\mathcal{D}(\Omega) \times \mathcal{D}(\Omega) \rightarrow C_b^\infty(\Omega) \times \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega).$$

but the operator

$$C_{\text{loc}}^\infty(\Omega) \times \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)$$

is sequentially continuous, and thus bounded, but not continuous.

- The convolution operator

$$\mathcal{D}(\mathbf{R}^d) \times \mathcal{D}(\mathbf{R}^d) \rightarrow L_c^1(\mathbf{R}^d) \times \mathcal{D}(\mathbf{R}^d) \rightarrow \mathcal{D}(\mathbf{R}^d)$$

is continuous.

Thus $\mathcal{D}(\Omega)$ is an ideal place to study many of the natural operations which occur in analysis.

We now have the tools to explain the idea of a distribution. If $f \in L_{\text{loc}}^1(\Omega)$, then the linear functional $\Lambda[f]$ on $\mathcal{D}(\Omega)$ defined for each $\phi \in \mathcal{D}(\Omega)$ by setting

$$\Lambda[f](\phi) = \int f(x)\phi(x) dx$$

is continuous. Moreover, $\Lambda[f]$ determines f uniquely, and so we can safely identify f with $\Lambda[f]$. We thus speak of ‘the distribution’ f . The idea of the theory of distributions is to treat any continuous linear functional Λ on $\mathcal{D}(\Omega)$ as if it were given by integration against a test function. Thus for such a linear functional Λ , we often abuse notation by denoting the quantity $\Lambda(\phi)$ by

$$\int_{\Omega} \Lambda(x)\phi(x) dx,$$

even if Λ is not given by integration against some function. The space $\mathcal{D}^*(\Omega)$ will be called the space of distributions on Ω . The class of distributions induced by elements of $L_{\text{loc}}^1(\Omega)$ is a fundamental class of distributions, but we will soon see much more erratic distributions.

One huge advantage of this approach is that we can generalize many analytical operations defined on $\mathcal{D}(\Omega)$ *distributionally* to give an operation on $\mathcal{D}^*(\Omega)$, even if the original analytical operations required some degree of smoothness to define. If $A : \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)$ is a continuous operator, then we can consider its adjoint $A^* : \mathcal{D}(\Omega)^* \rightarrow \mathcal{D}(\Omega)^*$. If A^* maps $\mathcal{D}(\Omega)$

continuously into itself, then we can define an extension of A to a map from $\mathcal{D}(\Omega)^* \rightarrow \mathcal{D}(\Omega)^*$ by defining, for a distribution Λ , a distribution $A\Lambda$ by the formula

$$\langle A\Lambda, \phi \rangle = \langle \Lambda, A^*\phi \rangle.$$

This definition has the property that $A(\Lambda[\phi]) = \Lambda[A\phi]$ for any $\phi \in \mathcal{D}(\Omega)$, so that we have really constructed an extension of A so it is defined on all distributions.

For instance, we can use this idea to define the derivative of an arbitrary distribution. For $\phi, \psi \in \mathcal{D}(\mathbf{R})$, integration by parts tells us that

$$\int_{-\infty}^{\infty} \phi'(x)\psi(x) dx = - \int_{-\infty}^{\infty} \phi(x)\psi'(x) dx.$$

Thus if $A\phi = \phi'$ is the derivative operator then its adjoint behaves on distributions by setting $A^*\psi = -\psi'$. Thus, for a distribution Λ on \mathbf{R} , we define its derivative to be the distribution Λ' such that for $\phi \in \mathcal{D}(\mathbf{R})$,

$$\Lambda'(\phi) = \Lambda(A^*\phi) = -\Lambda(\phi').$$

More generally, a similar calculation allows us to set, for a distribution Λ on \mathbf{R}^d , and a multi-index α , we define $D^\alpha \Lambda(\phi) = (-1)^{|\alpha|} \Lambda(D^\alpha \phi)$.

Example. Let $H(x) = \mathbf{I}(x > 0)$ denote the Heaviside step function. Then H is locally integrable, and so for any test function ϕ , we calculate

$$\int_{-\infty}^{\infty} H'(x)\phi(x) dx = - \int_{-\infty}^{\infty} H(x)\phi'(x) dx = - \int_0^{\infty} \phi'(x) dx = \phi(0)$$

Thus the distributional derivative of the Heaviside step function is the Dirac delta function. It is not a function, but if we were to think of it as a ‘generalized function’, it would be zero everywhere except at the origin, where it is infinitely peaked.

Example. Consider the Dirac delta function at the origin, which is the distribution δ such that for any $\phi \in \mathcal{D}(\mathbf{R})$,

$$\int_{-\infty}^{\infty} \delta(x)\phi(x) dx = \phi(0).$$

Then

$$\int_{-\infty}^{\infty} \delta'(x)\phi(x) dx = - \int_{\mathbf{R}^d} \delta(x)\phi'(x) dx = -\phi'(0).$$

This is a distribution that does not arise from integration with respect to a locally integrable function nor integration against a measure, but it is an appropriate model of certain physical situations, i.e. for the distribution of electrical charge in a polarized point mass with positive charge infinitesimally to the left of the origin, and negative charge infinitesimally to the right of the origin.

In general, we define a *distribution* to be a continuous linear functional on the space of test functions $\mathcal{D}(\Omega)$, i.e. an element of $\mathcal{D}(\Omega)^*$. In the last section, our exploration of continuous linear transformations on $\mathcal{D}(\Omega)$ guarantees that a linear functional Λ on $\mathcal{D}(\Omega)$ is continuous if and only if for every compact $K \subset X$ there is an integer n_K such that $|\Lambda\phi| \lesssim_K \|\phi\|_{C^{n_K}(K)}$ for $\phi \in C_c^\infty(K)$. If one integer n works for all K , and n is the smallest integer with such a property, we say that Λ is a distribution of *order* n . If such an n doesn't exist, we say the distribution has infinite order. Applying the Hahn-Banach theorem shows that if $\Lambda \in \mathcal{D}^*(\Omega)$ has order n , then Λ extends uniquely to a continuous linear functional on $\mathcal{D}^n(\Omega)$, since $\mathcal{D}(\Omega)$ is dense in $\mathcal{D}^n(\Omega)$. In fact, we see that we can identify $\mathcal{D}^n(\Omega)^*$ with the space of distributions of order n .

In many other ways, distributions act like functions. For instance, any distribution Λ can be uniquely written as $\Lambda_1 + i\Lambda_2$ for two distributions Λ_1, Λ_2 that are real valued for any real-valued smooth continuous function. However, we cannot write a real-valued distribution as the difference of two positive distributions, i.e. those which are non-negative when evaluated at any non-negative functional. This is because any non-negative distribution is actually given by integration against a Radon measure, and thus has order zero. To see this, given such a non-negative functional Λ (which is automatically continuous), we define Λf for a compactly supported continuous function $f \geq 0$ as

$$\Lambda f = \sup\{\Lambda g : g \in \mathcal{D}(\mathbf{R}^n), g \leq f\}$$

and then in general define $\Lambda(f^+ - f^-) = \Lambda f^+ - \Lambda f^-$. Then Λ is obviously a positive extension of Λ to all continuous functions, and is linear. But then the Riesz representation theorem implies that there is a positive Radon measure such that $\Lambda = \Lambda_\mu$, completing the proof.

Example. If μ is a complex-valued Radon measure, then we can define a distribution $\Lambda[\mu]$ such that for each $\phi \in \mathcal{D}(\mathbf{R}^d)$.

$$\Lambda[\mu](\phi) = \int_{\mathbf{R}^d} \phi(x) d\mu(x)$$

Thus $\Lambda[\mu]$ is a distribution, since if ϕ is supported on K , then

$$|\Lambda[\mu](\phi)| \leq \mu(K) \|\phi\|_{L^\infty(K)}.$$

The fact that this bound does not require information about the derivatives of ϕ implies that $\Lambda[\mu]$ is a distribution of order zero. The Riesz-Markov-Kakutani representation theorem, shows that any distribution of order zero is given by a complex-valued Radon measure.

Example. Let $U \subset \mathbf{R}^d$ be any open set such that ∂U is a C^1 hypersurface. Then we can find a C^1 function $\rho : \mathbf{R}^d \rightarrow \mathbf{R}$ such that $U = \{x : \rho(x) > 0\}$, and $\nabla \rho$ is non-zero on ∂U . Let us calculate the derivatives of the distribution $u = \mathbf{I}_U$. Clearly ∇u is supported on the surface $S = \partial U$. Let us work locally around a point $x_0 \in S$, and without loss of generality, let us rotate so that in a neighborhood of x_0 , if we write $x = (x', x_d)$, then S is described by the equation $x_d = \psi(x')$, where ψ is C^1 . Pick $f \in C^\infty(\mathbf{R})$ such that $f(t) = 0$ for $t < 0$ and $f(t) = 1$ for $t \geq 1$. Then, locally around x_0 , u is the distributional limit of the functions u_ε , where

$$u_\varepsilon(x) = f\left(\frac{x_d - \psi(x')}{\varepsilon}\right).$$

But

$$\nabla u_\varepsilon(x) = (1/\varepsilon) \cdot f'\left(\frac{x_d - \psi(x')}{\varepsilon}\right) \cdot (1, -\nabla \psi(x'))$$

which implies that for any $\phi \in \mathcal{D}(\mathbf{R}^d)$ supported near x_0 , we calculate using a change of variables that

$$\begin{aligned} & \int \phi(x) \nabla u(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int \phi(x) (1/\varepsilon) \cdot f'\left(\frac{x_d - \psi(x')}{\varepsilon}\right) \cdot (1, -\nabla \psi(x')) \\ &= \lim_{\varepsilon \rightarrow 0} \int \phi(x', \psi(x') + \varepsilon t) f'(t) (1, -\nabla \psi(x')) dt dx' \\ &= \int \phi(x', \psi(x')) (1, -\nabla \psi(x')) dx. \end{aligned}$$

Since the surface measure on S is given by $dS = (1 + |\psi'|^2)^{1/2} dx'$, and the normal vector to the surface is given by $n = (1, -\nabla\psi(x'))/(1 + |\psi'|^2)^{1/2}$, it follows that

$$\int \phi(x) \nabla u(x) dx = \int_S \phi(x) n(x) dS.$$

Switching to the global viewpoint, we now see that $\nabla u = n \cdot dS$. In particular, we have proved that for any smooth, compactly supported vector field $X = (X_1, \dots, X_n)$ on \mathbf{R}^d ,

$$\int_S (\nabla \cdot X) = \sum \int u(x) \partial_i X_i(x) dx = - \int_S X \cdot n dS,$$

so our proof amounts exactly to the Gauss-Green divergence theorem. The fact that $n \cdot dS$ is a distribution of order zero implies that the Gauss-Green formula continues to hold for any C^1 compactly supported vector field. Perhaps the most general setting in which the distributional derivative calculation continues to hold for any domain U which is a Caccioppoli set, i.e. such that $u = \mathbf{I}_U$ is a function with bounded variation. We can then define a Radon measure analogous to $n \cdot dS$ such that an analogue of the integration formulas continue to hold. In particular, we can still use the calculation for open sets with Lipschitz boundary, in which case the normal vector is only defined almost everywhere on the boundary.

Example. Consider a functional Λ defined for functions $\phi \in \mathcal{D}(\mathbf{R})$ vanishing in a neighbourhood of the origin by setting

$$\Lambda(\phi) = \int_{-\infty}^{\infty} \frac{\phi(x)}{x} dx.$$

Such functions are not dense in $\mathcal{D}(\mathbf{R})$. But we claim Λ is bounded on its domain, and thus by the Hahn-Banach theorem, extends to at least one continuous functional on the entirety of $\mathcal{D}(\mathbf{R})$. To prove this, fix $\phi \in C_c^\infty[-N, N]$ vanishing on a neighbourhood $(-\varepsilon, \varepsilon)$ of the origin. Then

$$|\Lambda\phi| = \left| \int_{-\infty}^{\infty} \frac{\phi(x)}{x} dx \right| = \left| \int_{\varepsilon \leq |x| \leq N} \frac{\phi(x) - \phi(0)}{x} dx \right|.$$

Applying the mean-value theorem, we find

$$|\Lambda\phi| \leq N \|\phi\|_{C^1[-N, N]}.$$

Since N was arbitrary, it follows that Λ is continuous in the topology induced by that of $\mathcal{D}(\mathbf{R})$, and thus by the Hahn-Banach theorem, extends uniquely to at least one distribution on the entirety of $\mathcal{D}(\mathbf{R})$.

One canonical choice of Λ is the principal value distribution $p.v(1/x)$, defined such that

$$\int_{-\infty}^{\infty} p.v(1/x) \phi(x) dx = \lim_{\delta \rightarrow 0} \int_{|x| \geq \delta} \phi(x)/x dx.$$

We essentially showed that this functional was continuous above. Another choice is the distribution $\lim_{\varepsilon \rightarrow 0} 1/(x + i\varepsilon)$, defined such that

$$\int_{-\infty}^{\infty} \lim_{\varepsilon \rightarrow 0} 1/(x + i\varepsilon) \cdot \phi(x) dx = \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{\infty} \phi(x)/(x + i\varepsilon) dx.$$

If we pick $\delta = \varepsilon^{1/4}$, then we can show using the fact that $1/x$ and $1/(x + i\varepsilon)$ are not too different for large x that

$$\left| \int_{|x| \geq \delta} \frac{\phi(x)}{x} - \int_{-\infty}^{\infty} \frac{\phi(x)}{x + i\varepsilon} \right| \leq \|\phi\|_1 \cdot \varepsilon^{1/2}.$$

A contour integral shift shows that

$$\begin{aligned} \int_{-\delta}^{\delta} \frac{\phi(x)}{x + i\varepsilon} &= \int_{-\delta}^{\delta} \frac{\phi(0)}{x + i\varepsilon} + O(\delta) \\ &= -i\pi\phi(0) + O(\varepsilon/\delta) + O(\delta) \\ &= -i\pi\phi(0) + O(\varepsilon^{1/4}). \end{aligned}$$

Taking $\varepsilon \rightarrow 0$ shows that

$$p.v(1/x) = i\pi\delta + \lim_{\varepsilon \rightarrow 0} 1/(x + i\varepsilon),$$

where δ is the Dirac delta distribution at the origin.

More generally, if Λ_1 and Λ_2 are two distributions which extend the functional Λ , then one can show that for any function ϕ vanishing away from the origin, $\Lambda_1(\phi) - \Lambda_2(\phi) = 0$. We will later define the support of a distribution, and so we have shown here that $\Lambda_1 - \Lambda_2$ is supported at $\{0\}$. It follows from later theorems in this chapter that Λ_1 and Λ_2 , applied to a function $\phi \in \mathcal{D}(\mathbf{R})$,

will differ by a finite linear combination of the values of ϕ and its derivatives at the origin.

The distribution $p.v(1/x)$ can also be described as the distributional derivative of the locally integrable function $\log|x|$, since an integration by parts shows that for each $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\begin{aligned} \int (\log|x|)' \phi(x) dx &= - \int \log|x| \phi'(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \log|x| \phi'(x) \\ &= \lim_{\varepsilon \rightarrow 0} \left(\log(\varepsilon) \cdot (\phi(x) - \phi(-x)) + \int_{|x| \geq \varepsilon} \frac{\phi(x)}{x} \right) \\ &= p.v \int \frac{\phi(x)}{x} dx. \end{aligned}$$

An important analysis of these distributions arises in the theory of the Hilbert transform.

Example. If Ω is an open subset of \mathbf{C} , let us calculate $\partial E_{z_0}/\partial \bar{z}$ in Ω , where $E_{z_0} = p.v\{(z - z_0)^{-1}\}$. To begin with, we note that if X is another open subset of \mathbf{C} containing Ω , and if $\partial\Omega$ (the boundary of Ω relative to X) is C^1 , then for $\phi \in \mathcal{D}(X)^*$, Green's formula gives

$$\int_{\Omega} \frac{\partial \phi}{\partial \bar{z}} = (-i/2) \int_{\partial\Omega} \phi dz.$$

This formula implies the product rule, that for $\phi \in \mathcal{D}(\Omega)^*$,

$$\begin{aligned} \int_{\Omega - B_{\varepsilon}(z_0)} \frac{1}{z - z_0} \frac{\partial \phi}{\partial \bar{z}} &= \int_{\Omega - B_{\varepsilon}(z_0)} \frac{\partial}{\partial \bar{z}} \left\{ \frac{1}{z - z_0} \phi \right\} \\ &= (i/2) \int_{\partial B_{\varepsilon}(z_0)} \frac{1}{z - z_0} \phi dz. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ gives $\int E_{z_0} \partial \bar{z} \phi = -\pi \phi(z_0)$, or in other words, we conclude that in Ω , $\partial \bar{z} E_{z_0} = \pi \delta_{z_0}$.

Example. One reason we could define a distribution agreeing with $1/x$ away from the origin is because there is a lot of cancellation at the origin from either

side of the origin, since $1/x$ switches sign here. One has to rely on other tricks to make sense of a distribution extending $1/x^2$. Indeed, if we write

$$\Lambda(\phi) = \int \frac{\phi(x)}{x^2}$$

for ϕ ranging over all functions vanishing in the neighborhood of the origin, then we can use the mean value theorem to obtain a bound $|\phi(x)| \leq x^2 \|\phi''\|_\infty$, from which it follows that

$$\Lambda(\phi) \lesssim \|\phi''\|_\infty,$$

and so Hahn-Banach extends Λ to a family of distributions. But in this case the principal value

$$\lim_{\delta \rightarrow 0} \int_{|x| \geq \delta} \frac{\phi(x)}{x^2}$$

rarely exists. Indeed, for any fixed $\phi \in \mathcal{D}(\mathbf{R})$ we have

$$\int_{|x| \geq \delta} \frac{\phi(x)}{x^2} = 2\phi(0)/\delta + O(\delta).$$

which will only converge if $\phi(0) = 0$. Thus, to get around this, we define the finite part distribution (or Hadamard regularization) of $1/x^2$, i.e. the distribution $f.p(1/x^2)$ obtained by setting

$$\int f.p(1/x^2) \phi(x) dx = \lim_{\delta \rightarrow 0} \left(\int_{|x| \geq \delta} \phi(x)/x^2 - \frac{2\phi(0)}{\delta} \right),$$

which gets around the result that the distribution might explode near the origin if $\phi(0) \neq 0$. Note that we do not need to account for $\phi'(0)$ because of cancellation on both sides of the integral.

Another approach to extending Λ is to consider the derivative of the distribution $-p.v(1/x)$, since the derivative of this distribution agrees with integration against $1/x^2$ away from the origin. In fact, the derivative of $-p.v(1/x)$ is precisely $f.p(1/x^2)$. We leave it to the reader to use similar tricks to define the finite parts of higher order singularities, such as $1/x^3$.

Example. Let f be a left continuous function on the real line with bounded variation and with $f(-\infty) = 0$. Then f' exists almost everywhere in the classical sense, and $f' \in L^1(\mathbf{R})$. By Fubini's theorem, if we let μ be the measure

defined by $\mu([a, b]) = f(b) - f(a)$, then for any $\phi \in \mathcal{D}(\mathbf{R})$,

$$\begin{aligned}\int_{-\infty}^{\infty} \phi(x) d\mu(x) &= - \int_{-\infty}^{\infty} \int_x^{\infty} \phi'(y) dy d\mu(x) \\ &= - \int_{-\infty}^{\infty} \phi'(y) \int_{-\infty}^y d\mu(x) dy \\ &= - \int_{-\infty}^{\infty} \phi'(y) f(y) dy\end{aligned}$$

Thus if Λ is the distribution corresponding to integration with respect to $f(x) dx$, then Λ' is given by integration with respect to μ . In particular, Λ' is given by integration with respect to $f'(x) dx$ precisely when f is absolutely continuous.

Example. If $f \in C^1(\mathbf{R} - \{0\})$, and if the function $v(x)$ defined to be $f'(x)$ for $x \neq 0$ is integrable, then the limits $f(0-)$ and $f(0+)$ both exist (a simple argument using the fundamental theorem of calculus), and the distributional derivative of f is equal to

$$f' = v + (f(0+) - f(0-))\delta_0.$$

To see this, we calculate that for $\phi \in \mathcal{D}(\mathbf{R})$,

$$\begin{aligned}\int f'(x) \phi(x) dx &= - \int f(x) \phi'(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} - \int_{-\infty}^{-\varepsilon} f(x) \phi'(x) dx - \int_{\varepsilon}^{\infty} f(x) \phi'(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} f(\varepsilon) \phi(\varepsilon) - f(-\varepsilon) \phi(-\varepsilon) + \int_{-\infty}^{-\varepsilon} v(x) \phi(x) dx + \int_{\varepsilon}^{\infty} v(x) \phi(x) dx \\ &= \int v(x) \phi(x) dx + [f(0+) - f(0-)] \phi(0).\end{aligned}$$

As a particular example of this, the distributional derivative of $|x|$ is $\text{sgn}(x)$, and the distributional derivative of the Heaviside step function H given above is evaluated to be δ_0 .

Example. The boundary values of analytic functions are distributions, in the following manner. Let $\Omega \subset \mathbf{R}^n$ be an open set, and let Γ be a convex open cone in \mathbf{R}^n . For $\gamma > 0$, let

$$Z_\gamma = \{z \in \mathbf{C}^n : \text{Re}(z) \in \Omega, \text{Im}(z) \in \Gamma, |\text{Im}(z)| < \gamma\}.$$

If f is analytic in Z_γ , and there is some $N > 0$ such that $|f(x + iy)| \lesssim |y|^{-N}$. We claim then that, if for each $y \in \Gamma$, we consider the analytic function $f_y(x) = f(x + iy)$, then as $y \rightarrow 0$, $\{f_y\}$ converges distributionally to a distribution of order at most $N + 1$ on Ω . In fact, we will obtain an explicit formula for this distribution. For simplicity we deal with the case $n = 1$, where the higher dimensional case is similar (see Hörmander Theorem 3.1.15). Then we might as well assume that $Z_\gamma = \{x + iy : x \in \Omega, 0 < y < \gamma\}$. Given $\phi \in C_c^{N+1}(\Omega)$, write

$$\phi_y(x) = \phi(x + iy) = \sum_{\alpha=0}^N \partial_x^\alpha \phi(x) \frac{(iy)^\alpha}{\alpha!}.$$

Then

$$2 \frac{\partial \phi}{\partial \bar{z}} = \partial^{N+1} \phi(x) \frac{(iy)^N}{N!}.$$

Thus if $0 < y_0 < \gamma$ is fixed, and if $0 < y < \gamma - y_0$, then the Cauchy integral formula implies that

$$\int \phi(x) f(x + iy) dx - \int \phi(x + iy) f(x + iy + iy_0) dx = 2i \int \int_0^{y_0} f(x + iy + it) \frac{\partial \phi}{\partial \bar{z}} dt dx.$$

Thus

$$\int \phi(x) f_y(x) dx = \int \phi_{y_0}(x) f_{y+y_0}(x) dx + \int \int_0^1 f_{y+ty_0}(x) \partial^{N+1} \phi(x) \frac{(iy_0)^{N+1} t^N}{N!} dt dx.$$

Thus we conclude that as $y \rightarrow 0$,

$$\int \phi(x) f_y(x) dx \rightarrow \int \phi_{y_0}(x) f_{y_0}(x) dx + \frac{1}{N!} \int \int_0^1 f_{ty_0}(x) \partial^{N+1} \phi(x) (iy_0)^{N+1} t^N dt dx.$$

The right hand side clearly defines a distribution of order $N + 1$. In higher dimensions, we obtain the similar representation formula

$$\int \phi(x) f_y(x) dx \rightarrow \int \phi_{y_0}(x) f_{y_0}(x) dx + \frac{1}{N!} \int \int_0^1 f_{ty_0}(x) \sum_{|\alpha|=N+1} \partial^\alpha \phi(x) (iy_0)^\alpha t^N dt dx.$$

Thus we have a theory of boundary values of analytic functions.

For $n = 1$, we denote the distribution obtained by $f(x + i0)$. If we, by a similar process, obtained a distribution as a limit of an analytic function defined on the lower half plane, then we would denote that distribution by

$f(x-i0)$. If f is analytic above and below the half plane, and satisfies estimates of the form above, then the distribution u given by

$$\int (u(x+iy)\phi(x,y) dx) dy$$

(the order of integration matters) defines a distribution of order N in the complex plane, the derivative $\partial u/\partial \bar{z}$ is supported on the real line, and on that real line, we have

$$\frac{\partial u}{\partial \bar{z}}(x) = (i/2)(f(x+i0) - f(x-i0)).$$

We can use this, for instance, to calculate the difference between the two distributions $(x+i0)^{-1}$ and $(x-i0)^{-1}$. We have calculated above that the distribution $u(z) = 1/z$ has

$$\frac{\partial u}{\partial \bar{z}} = \pi\delta,$$

and so $(x-i0)^{-1} = (x+i0)^{-1} + 2\pi i\delta$.

Remark. Roughly speaking, this result is tight. If an analytic function f has boundary values defining a distribution of order $N+1$, then it is necessary that $|f(z)| \lesssim |\text{Im}(z)|^{-N-2}$.

There are many other important operations one can apply to distributions. If Ω is a conic subset of \mathbf{R}^d , and $\phi, \psi \in \mathcal{D}(\Omega)$, we find

$$\int_{\Omega} \text{Dil}_{\lambda}\phi(x)\psi(x) dx = \lambda^{-d} \int_{\Omega} \phi(x) \cdot \text{Dil}_{1/\lambda}\psi(x) dx,$$

Thus if Λ is a distribution on Ω , then we define $\text{Dil}_{\lambda}\Lambda$ by setting

$$\text{Dil}_{\lambda}\Lambda(\phi) = \lambda^{-d}\Lambda(\text{Dil}_{1/\lambda}\phi).$$

For $f \in C^{\infty}(\Omega)$, we have an operator $\phi \mapsto f\phi$ on $\mathcal{D}(\Omega)$. The adjoint is clearly $\psi \mapsto f\psi$, so for a distribution Λ on Ω , we define $f\Lambda$ by setting $(f\Lambda)(\phi) = \Lambda(f\phi)$. Thus $\mathcal{D}^*(\Omega)$ is naturally a $C^{\infty}(\Omega)$ module. Similarly, the family $\mathcal{D}^*(\Omega)_k$ consisting of distributions of order k form a $C^k(\Omega)$ module.

Remark. Hörmander developed a sophisticated theory that enables us to define the product of two *distributions* using the Fourier transform. In many basic situations, one can perform a spatial decomposition to define the product. Given a distribution Λ , we define its *singular support* $\text{supp}_{\text{sing}}(\Lambda)$ to be the *complement* of the set of all points x which have a neighborhood U such that $\Lambda|_U \in C^\infty(U)$. For any two distributions Λ and Ψ whose singular supports are disjoint, a decomposition argument enables us to define the product $\Lambda \cdot \Psi$ in a natural way.

9.3 Topologies on the Space of Distributions

As a dual space to an LF space, we can equip the space of distributions $\mathcal{D}(\Omega)^*$ with several topologies. The most important for our purposes is the strong topology, which makes $\mathcal{D}(\Omega)^*$ into a complete locally convex space, and the weak $*$ topology, which is quasi-complete. These topologies are roughly the same for many purposes. For instance, the uniform boundedness theorem implies that a sequence $\{u_n\}$ of distributions converges in the weak $*$ topology if and only if it converges in the strong dual topology, and we call this convergence *distributional convergence*. We note also that $\mathcal{D}(\Omega)^*$, as the strong dual of a Montel space, is also a Montel space, i.e. it is locally convex, barrelled, and satisfies the Heine-Borel property.

Example. If $u \in \mathcal{D}(\mathbf{R}^d)$, and we set $u_\varepsilon(x) = \varepsilon^{-d} \text{Dil}_\varepsilon u(x)$, then for any $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\lim_{\varepsilon \rightarrow 0} \int u_\varepsilon(x) \phi(x) dx \rightarrow \phi(0) \int u(x) dx.$$

Thus u_ε converges distributionally to $(\int u(x) dx) \cdot \delta_0$. Similarly, if $u \in \mathcal{D}(\mathbf{R}^d)$ and for any multi-index α with $|\alpha| \leq k$,

$$\int u(x) x^\alpha dx = 0,$$

and we define $u_\varepsilon(x) = \varepsilon^{-d-k} \text{Dil}_\varepsilon u(x)$, then for any $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\lim_{\varepsilon \rightarrow 0} \int u_\varepsilon(x) \phi(x) dx \rightarrow \frac{1}{k!} \sum_{|\alpha|=k} \left(\int x^\alpha u(x) dx \right) \cdot D^\alpha \phi(0)$$

Thus u_ε converges distributionally to an appropriate linear combination of $D^\alpha \delta_0$.

Example. The distribution $\lim_{\varepsilon \rightarrow 0} 1/(x + i\varepsilon)$ defined above is precisely the limit of the distributions $1/(x + i\varepsilon)$ in the weak * topology. Similarly, $p.v(1/x)$ is the weak * limit of the functions $\mathbf{I}_{|x| \geq \delta}(x) \cdot (1/x)$. The distribution $f.p(1/x^2)$ is the distributional limit of $\mathbf{I}_{|x| \geq \delta}(x) \cdot (1/x^2) - 2\delta_0/\delta$, where δ_0 is the Dirac delta function at the origin.

Example. If n is a positive integer, then integration by parts shows that for any $\phi \in \mathcal{D}(\mathbf{R})$,

$$\int_{-\infty}^{\infty} t^n e^{2\pi i t x} \phi(x) dx = i^{n+1} t^{-1} \int_{-\infty}^{\infty} e^{i t x} \phi^{(n+1)}(x) dx,$$

which converges to zero as $t \rightarrow \infty$. Thus $t^n e^{2\pi i t x}$ converges distributionally to zero as $t \rightarrow \infty$. Another way to see this is to note that the distribution Λ_t given by integration against $t^n e^{i t x}$ can be written as $\Lambda_t(\phi) = t^n \hat{\phi}(-t)$, and the Fourier transform of ϕ decays rapidly. Note that if we tested against functions that were less smooth (say, viewing these distributions as linear functionals on $L^1(\mathbf{R}^d)$, or even C^∞ functions that are only of polynomial decrease as they approach ∞) then this statement would no longer be true.

Example. Let $u_t(x) = t^{1/k} e^{i t x^k}$, where k is an integer bigger than one. Let

$$F(x) = \int_0^x e^{i y^k} dy.$$

When $x > 0$, a contour integration shift shows that

$$F(x) = e^{i\pi/2k} \int_0^x e^{-y^k} dy + O(|x|^{-(k-1)}).$$

If k is even, then for $x < 0$,

$$F(x) = -e^{i\pi/2k} \int_0^x e^{-y^k} dy + O(|x|^{-(k-1)})$$

and for k odd,

$$F(x) = -e^{-i\pi/2k} \int_0^x e^{-y^k} dy + O(|x|^{-(k-1)}).$$

Thus given $\phi \in \mathcal{D}(\mathbf{R})$, we can apply an integration by parts to write

$$\begin{aligned} \int_{-\infty}^{\infty} u_t(x) \phi(x) dx &= \int_{-\infty}^{\infty} t^{1/k} e^{itx^k} \phi(x) dx \\ &= \int_{-\infty}^{\infty} t^{1/k} F'(t^{1/k}x) \phi(x) dx \\ &= - \int_{-\infty}^{\infty} F(t^{1/k}x) \phi'(x) dx, \end{aligned}$$

By decomposing this integral into the region where $|x| \geq t^{1/k}$ and $|x| \leq t^{1/k}$ shows that this quantity converges to

$$-F(\infty) \int_0^{\infty} \phi'(x) dx - F(-\infty) \int_{-\infty}^0 \phi'(x) dx = (F(\infty) - F(-\infty))\phi(0).$$

Thus u_t converges distributionally to $\left(2e^{i\pi/2k} \int_0^{\infty} e^{-y^k} dy\right) \cdot \delta_0$ if k is even, and to $\left(2\cos(\pi/2k) \int_0^{\infty} e^{-y^k} dy\right) \cdot \delta_0$ if k is odd.

It is often of interest to focus on subfamilies of $\mathcal{D}(\Omega)^*$, equipped with a topology which is compatible with the behaviour of distributions. We thus define a *space of distributions* to be a vector subspace X of $\mathcal{D}(\Omega)^*$, equipped with a topology which makes the inclusion map $X \rightarrow \mathcal{D}(\Omega)_b^*$ continuous. Most function spaces are spaces of distributions, for instance, $\mathcal{D}(\Omega)$, L_{loc}^1 , and so on. We will later encounter the space of *tempered distributions* $\mathcal{S}(\Omega)^*$, which will turn out to be a subspace of $\mathcal{D}(\Omega)^*$ equipped with the relative topology.

Since convergence in $\mathcal{D}(\Omega)$ is incredibly strict, a sequence of distributions can very easily converge distributionally. It is therefore surprising that, since a differential operator $L : \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)$ is continuous, its extension to a map $L : \mathcal{D}(\Omega)^* \rightarrow \mathcal{D}(\Omega)^*$, roughly speaking, a rescaling of its adjoint, is also continuous, both in the weak $*$ topologies and in the strong topology.

With either of the two topologies described above, the bilinear multiplication map

$$C_{\text{loc}}^{\infty}(\Omega) \times \mathcal{D}(\Omega)^* \rightarrow \mathcal{D}(\Omega)^*$$

is sequentially continuous, but not jointly continuous.

9.4 Homogeneous Distributions

An important family of distributions are the *homogenous distributions*, which are those distributions Λ on $\mathbf{R}^d - \{0\}$ such that for each $\lambda > 0$, $\text{Dil}_\lambda \Lambda = \lambda^\alpha \Lambda$, where α is the *order* of the homogenous distribution Λ .

Example. If $f \in L_1^{\text{loc}}(\mathbf{R}^d)$ and $f(\lambda x) = \lambda^\alpha f(x)$ for all $x \in \mathbf{R}^d - \{0\}$ then integration against $f(x) dx$ defines a homogenous distribution of order α .

Example. For any complex number a with $\text{Re}(a) > -1$, if we define a distribution on $\mathbf{R} - \{0\}$ by setting

$$x_+^a = \begin{cases} x^a & x > 0 \\ 0 & : x \leq 0 \end{cases},$$

then x_+^a is a homogeneous distribution of order α , and $x \cdot x_+^a = x_+^{a+1}$, and if $\text{Re}(a) > 0$,

$$\frac{d}{dx} (x_+^a) = a x_+^{a-1}.$$

Our goal is to extend this distribution to a larger range of values $a \in \mathbf{C}$, such that the association $a \mapsto x_+^a$ is continuous. For any $\phi \in \mathcal{D}(\mathbf{R})$, the function

$$a \mapsto \langle x_+^a, \phi \rangle$$

is analytic in a for $\text{Re}(a) > -1$. Integration by parts shows that

$$\langle x_+^{a+1}, \phi' \rangle = -(a+1) \langle x_+^a, \phi \rangle.$$

The formula $\langle x_+^a, \phi \rangle = -(a+1)^{-1} \langle x_+^{a+1}, \phi' \rangle$ allows us to extend the definition of x_+^a to all $a \in \mathbf{C}$ with $\text{Re}(a) > -2$, except that we have a pole of order one when $a = -1$. Iterating this allows us to uniquely extend the definition of x_+^a for all $a \in \mathbf{C} - \mathbf{Z}^-$, and these distributions will all be homogeneous.

Marcel Riesz also used some other complex analytic tricks to define x_+^{-k} for all integers k , but then we lose some of the homogeneity. For any $\phi \in \mathcal{D}(\mathbf{R})$, the function $a \mapsto \langle x_+^a, \phi \rangle$ is meromorphic, with simple poles at each integer $-k$ for $k > 0$, and the residue at $-k$ is equal to

$$(-1)^k D^{k-1} \phi(0) / (k-1)!$$

Thus we conclude that, as $a \rightarrow -k$,

$$\langle (a+k)x_+^a, \phi \rangle \rightarrow (-1)^{k-1} D^{k-1} \phi(0)/(k-1)!$$

In fact, expanding things out gives a constant $C_{-k}(\phi)$ such that as $a \rightarrow -k$,

$$\langle x_+^a, \phi \rangle = \frac{(-1)^{k-1} D^{k-1} \phi(0)}{(k-1)!} \cdot \frac{1}{a+k} + C_{-k}(\phi) + O(a+k).$$

We define $\langle x_+^{-k}, \phi \rangle = C_{-k}(\phi)$, i.e. by keeping only the finite part of the integral. Since, for a close to k , we have

$$\begin{aligned} \langle x_+^a, \phi \rangle &= \frac{(-1)^{k-1} D^{k-1} \phi(0)}{(k-1)!} \cdot (a+k)^{-1} \\ &= (-1)^k (a+1)^{-1} \cdots (a+k-1)^{-1} \int_0^\infty \frac{x^{a+k}-1}{a+k} D^k \phi(x) dx \\ &\quad + \frac{(-1)^{k-1}}{a+k} \left((-1-a)^{-1} \cdots (1-a-k) - \frac{1}{(k-1)!} \right) D^{k-1} \phi(0) \\ &\rightarrow \frac{-1}{(k-1)!} \int_0^\infty \log(x) D^k \phi(x) dx + \frac{1}{(k-1)!} \left(\sum_{i=1}^k 1/i \right) D^{k-1} \phi(0). \end{aligned}$$

and thus

$$\langle x_+^{-k}, \phi \rangle = \frac{-1}{(k-1)!} \int_0^\infty \log(x) D^k \phi(x) dx + \frac{1}{(k-1)!} \left(\sum_{i=1}^k 1/i \right) D^{k-1} \phi(0).$$

Our extension of x_+^a for all $a \in \mathbf{C}$ continues to satisfy $x \cdot x_+^a = x_+^{a+1}$ for all $a \in \mathbf{C}$. The derivative formula is not maintained, namely,

$$Dx_+^{-k} = -kx_+^{-k-1} + (-1)^k D^k \delta/k!$$

Moreover, x_+^{-k} is no longer homogeneous of degree $-k$. Plugging into the formula shows that

$$(tx)_+^{-k} = t^{-k} x_+^{-k} + \frac{\log t}{(k-1)!} \cdot D^{k-1} \delta.$$

One can also define x_+^a by first removing the singularity, considering the distributions

$$\langle x_\varepsilon^a, \phi \rangle = \int_\varepsilon^\infty x^a \phi(x) dx.$$

If k is the smallest non-negative integer such that $k + \operatorname{Re}(a) > -1$, then we can integrate by parts to conclude that there are constants C_k such that

$$\langle x_\varepsilon^a, \phi \rangle = \sum_{i=0}^{k-1} C_i \varepsilon^{-i} + (-1)^k \int_0^\infty \frac{1}{(a+1)\dots(a+k)} x^{a+k} D^k \phi(x) dx + o(1).$$

Discarding the singular terms, and letting $\varepsilon \rightarrow 0$ gives the distributions x_+^a above. One can analogously define the distributions x_-^a , and the distributions $|x|^a$, by reflecting and symmetrizing the distributions about the origin.

Example. One can fix the singularities at the integers by normalizing. The appearance of the singularities in the extension of x_+^a occurred because of the a term in the formula $D(x_+^a) = ax_+^{a-1}$. If we consider the normalization $\chi_+^a = x_+^a / \Gamma(a+1)$ when $\operatorname{Re}(a) > -1$, then $D(\chi_+^a) = -\chi_+^{a-1}$, and because the Gamma function has no zeroes, this allows us to extend the definition of χ_+^a to an analytic function for all $a \in \mathbf{C}$, each a homogeneous distribution of order a . But since χ_+^0 is the Heaviside step function, we conclude that $\chi_+^{-k} = (-1)^k D^{k-1} \delta$.

Example. Let δ be the Dirac-delta distribution at the origin in \mathbf{R}^d . Then for $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\int_{\mathbf{R}^d} (\operatorname{Dil}_\lambda \delta)(x) \phi(x) dx = \lambda^{-d} \int_{\mathbf{R}^d} \delta(x) \operatorname{Dil}_{1/\lambda} \phi(x) dx = \lambda^{-d} \phi(0).$$

Thus $\operatorname{Dil}_\lambda \delta = \lambda^{-d} \delta$, which implies δ is a homogenous distribution of order $-d$.

A homogeneous distribution is apriori defined by testing against a distribution in $\mathcal{D}(\mathbf{R}^n - \{0\})$. But in most cases the distribution can be extended so that it can be tested against an arbitrary distribution in $\mathcal{D}(\mathbf{R}^n)$. Before we prove this, we begin with some simple observations. First is a formula due to Euler, which in the distributional setting states that for any distribution u of degree a ,

$$\sum_{i=1}^n x_i \partial_i u = (a+n)u.$$

When $n = 1$, this actually implies that u is a multiple of $|x|^a$ on each coordinate axis. The identity also implies that for any $\psi \in \mathcal{D}(\mathbf{R}^n - \{0\})$ with $\int_0^\infty r^{a+n-1} \psi(rx) dr = 0$ for all x , $\langle u, \phi \rangle = 0$ by rewriting the formula in polar coordinates.

Theorem 9.3. *Let u be a homogeneous distribution on $\mathbf{R}^n - \{0\}$ of order*

$$a \in \mathbf{C} - \{-n, -(n+1), \dots\}.$$

Then u has a unique extension to a distribution $E(u)$ on \mathbf{R}^n , such that for any homogeneous polynomial P , $E(Pu) = PE(u)$, and if u is not a distribution of order $1 - n$, $E(\partial_i u) = \partial_i E(u)$. Moreover, the map $u \mapsto E(u)$ is continuous from $\mathcal{D}(\mathbf{R}^n - \{0\})^$ to $\mathcal{D}(\mathbf{R}^n)^*$.*

Proof. The uniqueness is obvious, because any distribution supported at the origin is a linear combination of derivatives of the Dirac delta function, which are all homogeneous of integer order $\leq -n$. To show existence, we note that if u was locally integrable, then polar coordinates gives

$$\int u(x)\phi(x) = \int_0^\infty \int_{|w|=1} u(w)t^{a+n-1}\phi(tw) d\sigma(w) dt.$$

Thus we need only study the behaviour of u near the unit sphere, which is supported away from the origin. Doing this more formally yields the extension map E . For any $\phi \in \mathcal{D}(\mathbf{R}^n)$, define $R_a\phi(x) = \langle t_+^{a+n-1}, \phi(tx) \rangle$. Then $R_a\phi$ is homogeneous of degree $-n - a$, and is continuous from $\mathcal{D}(K)$ to $C^\infty(\mathbf{R}^n - \{0\})$ for any compact set K . If $\psi \in C_c(\mathbf{R}^n - \{0\})$ and $\int_0^\infty \psi(tx)/t dt = 1$ for all $x \neq 0$, then $\psi R_a\phi \in \mathcal{D}(\mathbf{R}^n - \{0\})$, and $R_a(\psi R_a\phi) = R_a\phi$, so that (because of our observations before the proof) $E(u) = \langle u, \psi R_a\phi \rangle$ is independent of the choice of ψ . Moreover, $\langle u, \psi R_a\phi \rangle = \phi$ for each $\phi \in \mathcal{D}(\mathbf{R}^n - \{0\})$. The continuity of the map $u \mapsto E(u)$ is not too difficult to see, completing the proof. \square

When a is an integer smaller than or equal to $-n$, one can still use the construction in the proof to define an operator $E(u)$, but then it can depend on ψ , may fail to be homogeneous near the origin, and may fail to satisfy the identities $E(Pu) = PE(u)$ and $E(\partial_i u) = \partial_i E(u)$ (See Hörmander, 3.2).

9.5 Localization of Distributions

Just as we can consider the local behaviour of functions around a point, we can consider the local behaviour of a distribution around points, and this local behaviour contains most of the information of the distribution. For

instance, given an open subset U of X , we say two distributions Λ and Ψ are equal on U if $\Lambda\phi = \Psi\phi$ for every test function ϕ compactly supported in U . We recall the notion of a partition of unity, which, for each open cover U_α of Euclidean space, gives a family of C^∞ functions ψ_α which are positive, *locally finite*, in the sense that only finitely many functions are positive on each compact set, and satisfy $\sum \psi_\alpha = 1$ on the union of the U_α .

Theorem 9.4. *If X is covered by a family of open sets U_α , and Λ and Ψ are locally equal on each U_α , then $\Lambda = \Psi$. If we have a family of distributions Λ_α which agree with one another on $U_\alpha \cap U_\beta$, then there is a unique distribution Λ locally equal to each Λ_α .*

Proof. Since we can find a C^∞ partition of unity ψ_α compactly supported on the U_α , upon which we find if ϕ is supported on K , then finitely many of the ψ_α are non-zero on K , and so

$$\Lambda(\phi) = \sum \Lambda(\psi_\alpha \phi) = \sum \Psi(\psi_\alpha \phi) = \Psi(\phi)$$

Thus $\Lambda = \Psi$. Conversely, if we have a family of distributions Λ_α like in the hypothesis, then we can find a partition of unity $\psi_{\alpha\beta}$ subordinate to $U_\alpha \cap U_\beta$, and we can define

$$\Lambda(\phi) = \sum \Lambda_\alpha(\psi_{\alpha\beta} \phi) = \sum \Lambda_\beta(\psi_{\alpha\beta} \phi)$$

The continuity is verified by fixing a compact K , from which there are only finitely many nonzero $\psi_{\alpha\beta}$ on K , and the fact that this definition is independent of the partition of unity follows from the first part of the theorem. \square

In the language of commutative algebra, the association of $\mathcal{D}^*(U)$ to each open subset U of Ω gives the structure of a sheaf of modules on Ω . Given a distribution Λ , we might have $\Lambda(\phi) = 0$ for every ϕ supported on some open set U . The complement of the largest open set U for which this is true is called the *support* of Λ . This agrees with the sheaf theoretic definition.

Theorem 9.5. *If a distribution $\Lambda \in \mathcal{D}^*(\Omega)$ has compact support, then Λ has some finite order n , and extends uniquely to a continuous linear functional on $C^n(\Omega)$.*

Proof. Let Λ be a distribution supported on a compact set. If ψ is a function with compact support with $\psi(x) = 1$ on the support of Λ , then $\psi\Lambda = \Lambda$, because for any ϕ , $\phi - \phi\psi$ is supported on a set disjoint from the support of Λ . But if ψ is supported on some compact set K , then there is n such that for any $\phi \in C_c^\infty(K)$,

$$|\Lambda(\phi)| \lesssim \|\phi\|_{C^n(K)},$$

and so for any other compact set K ,

$$|\Lambda(\phi)| = |\Lambda(\phi\psi)| \lesssim \|\phi\psi\|_{C^n(K)} \lesssim \|\psi\|_{C^n(K)} \|\phi\|_{C^n(K)}.$$

which shows Λ has order N . We have shown that Λ is continuous with respect to the seminorm $\|\cdot\|_{C^N(K)}$ on $C^\infty(X)$, and so by the Hahn Banach theorem, Λ extends uniquely to a continuous functional on $C^\infty(X)$. \square

If $\mathcal{E}(\Omega)$ denotes $C^\infty(\Omega)$, equipped with the topology such that $f_n \rightarrow f$ if $D^\alpha f_n$ converges to $D^\alpha f$ locally uniformly for all α , then the last theorem implies the family of compactly supported distributions embeds itself in $\mathcal{E}(\Omega)^*$. Conversely, *every* element of $\mathcal{E}(\Omega)^*$ is a compactly supported distribution. Indeed, since $\mathcal{E}(\Omega)$ is a Frechét space, if Λ is a continuous linear functional on $\mathcal{E}(\Omega)$, then there exists a compact set K and some $n > 0$ such that

$$|\Lambda(\phi)| \lesssim \|\phi\|_{C^n(K)}.$$

It follows from this that Λ is a distribution with support contained in K .

Remark. For general compact sets K , it is *not* true that if Λ is a distribution supported on a set K , then there exists $n > 0$ such that

$$|\Lambda(\phi)| \lesssim \|\phi\|_{C^n(K)}.$$

Suppose K is not the union of finitely many compact connected sets. Then we can find a family of disjoint compact sets $\{K_i\}$ in K such that $K - (K_1 \cup \dots \cup K_n)$ is compact for any $n > 0$. Fix $x_i \in K_i$, let $x \in K$ be a limit point of this sequence, consider a sequence of numbers $\{a_i\}$ such that $\sum a_i |x_i - x| = 1$, and $\sum a_i = \infty$, and let Λ be the distribution

$$\Lambda(\phi) = \sum_i a_i (\phi(x_i) - \phi(x)).$$

Then

$$|\Lambda(\phi)| \leq \|\phi'\|_{L^\infty(\mathbf{R}^d)},$$

so Λ is a distribution of order at most 1. On the other hand, if we choose a function $\phi \in \mathcal{D}(\Omega)$ which is equal to one on a neighborhood of $K_1 \cup \dots \cup K_n$, and zero on a neighborhood of $K - (K_1 \cup \dots \cup K_n)$, then

$$\Lambda(\phi) = \sum_{i=1}^n a_i,$$

so we cannot have a bound of the form

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq k} \|D^\alpha \phi\|_{L^\infty(K)} \lesssim 1.$$

On the other hand, for any precompact neighborhood U of K , we have a bound

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq 1} \|D^\alpha \phi\|_{L^\infty(U)}.$$

which is almost as good as the bound above.

If Λ is a distribution of order k supported on K , though we do not have a uniform bound $|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq k} \|D^\alpha \phi\|_{L^\infty(K)}$, if the right hand side vanishes, so does the left hand side.

Lemma 9.6. *Suppose Λ is a distribution of order k supported on K , and $\phi \in C^k(\Omega)$ satisfies $D^\alpha \phi(x) = 0$ for all $|\alpha| \leq k$ and $x \in K$, then $\Lambda(\phi) = 0$.*

Proof. By a density argument, we may assume that $\phi \in C^\infty(\Omega)$ without loss of generality. Find $\chi_\varepsilon \in \mathcal{D}(\Omega)$ such that $\chi_\varepsilon(x) = 1$ for $x \in K$, $\chi_\varepsilon(x) = 0$ if $d(x, K) \geq \varepsilon$, and $\|D^\alpha \chi_\varepsilon\|_{L^\infty} \lesssim \varepsilon^{-|\alpha|}$ for all $|\alpha| \leq k$. Then for any $\phi \in \mathcal{D}(\Omega)$,

$$|\Lambda(\phi)| = |\Lambda(\phi \chi_\varepsilon)| \lesssim \sum_{|\alpha| \leq k} \|D^\alpha(\phi \chi_\varepsilon)\|_{L^\infty} \lesssim \sum_{|\alpha| \leq k} \varepsilon^{|\alpha| - k} \|D^\alpha \phi\|_{L^\infty(K_\varepsilon)}.$$

For any $y \in K_\varepsilon$, we can pick $x \in K$ such that $|x - y| \leq \varepsilon$. Taylor's formula at x , together with the fact that all the derivatives of ϕ up to order k vanish at x , implies that

$$|(D^\alpha \phi)(y)| \lesssim \varepsilon^{k+1-|\alpha|}.$$

Thus we conclude that $|\Lambda(\phi)| \lesssim \varepsilon$, and we can then take $\varepsilon \rightarrow 0$. \square

The last lemma implies that the value of *any distribution* Λ of order k supported on a point x_0 depends solely on the values $D^\alpha \phi(x_0)$ for $|\alpha| \leq k$. Thus there exists constants λ_α such that

$$\Lambda(\phi) = \sum_{|\alpha| \leq k} \lambda_\alpha D^\alpha \phi(x_0).$$

This means that Λ is a sum of Dirac delta functions and their derivatives. If we work harder, using the Whitney extension theorem as a black box, we can obtain a similar process for more general supports.

Theorem 9.7 (Whitney). *Let K be a compact set in \mathbf{R}^d , and for each $|\alpha| \leq k$, a function $u_\alpha \in C(K)$. If*

$$U_\alpha(x, y) = \sum_{|\alpha| \leq k} \sup_{x, y \in K} \left| u_\alpha(x) - \sum_{|\beta| \leq k - |\alpha|} u_{\alpha + \beta}(y) \cdot (x - y)^\beta / \beta! \right| \cdot |x - y|^{|\alpha| - k}$$

for $x \neq y$, and $U_\alpha(x, x) = 0$, then provided U_α is continuous on $K \times K$, we can find a function $v \in C^k(\mathbf{R}^d)$ such that $D^\alpha v = u_\alpha$ on K for $|\alpha| \leq k$, and

$$\sum_{|\alpha| \leq k} \|D^\alpha v\|_{L^\infty} \lesssim \sum_{|\alpha| \leq k} \|U_\alpha\|_{L^\infty(K \times K)} + \sum_{|\alpha| \leq k} \|u_\alpha\|_{L^\infty(K)}.$$

A consequence is the following strengthening of the last lemma.

Lemma 9.8. *For any compact set K , and any distribution Λ of order k supported on K , we have*

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq k} \sup_{x, y \in K} \left| D^\alpha \phi(x) - \sum_{|\beta| \leq k - |\alpha|} D^{\alpha + \beta} \phi(y) \cdot (x - y)^\beta / \beta! \right| \cdot |x - y|^{|\alpha| - k} + \sum_{|\alpha| \leq k} \|D^\alpha \phi\|_{L^\infty(K)}.$$

Proof. To do this, we apply the Whitney extension theorem, setting $u_\alpha = D^\alpha \phi|_K$. We then apply the Whitney extension theorem to find $\psi \in C^k(\mathbf{R}^n)$ extending u_α with the required bounds above. Then $D^\alpha(\phi - \psi) = 0$ on K for all $|\alpha| \leq K$, from which it follows that $\Lambda(\phi) = \Lambda(\psi)$. The bound

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq k} \|D^\alpha \psi\|_{L^\infty},$$

which gives the required bound above. □

Recall that a compact set K is *Whitney regular*, which means that K is a finite union of compact, connected components, and for any two points $x, y \in K$ contained in a common component, there exists a rectifiable curve γ from x to y with length $O(|x - y|)$.

Lemma 9.9. *If K is Whitney regular, then for any distribution Λ supported on K , there exists k such that we have a bound*

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq k} \|D^\alpha \phi\|_{L^\infty(K)}.$$

Proof. Fix a rectifiable unit velocity curve $\gamma : [0, L] \rightarrow K$ between two points x and y in K , and let

$$F_\alpha(s) = D^\alpha \phi(\gamma(s)) - \sum_{|\beta| \leq k - |\alpha|} D^{\alpha + \beta} \phi(y) (\gamma(s) - y)^\beta / \beta!$$

Then $|F_\alpha(s)| \lesssim s^{k - |\alpha|} \sum_{|\beta| = k} \|D^\beta \phi\|_{L^\infty(K)}$. This is immediate if $|\alpha| = k$. For $|\alpha| < k$ we prove this result by induction, noting that the case for higher order k implies that

$$\begin{aligned} |dF_\alpha/ds| &\leq \sum_{i=1}^d \left| \left(D^{\alpha + i} \phi(\gamma(s)) - \sum_{|\beta| \leq k - |\alpha|} D^{(\alpha + i) + (\beta - i)} \phi(y) (\gamma(s) - y)^{\beta - i} / (\beta - 1)! \right) \cdot \gamma'_i(s) \right| \\ &\lesssim s^{k - |\alpha| - 1} \sum_{|\beta| = k} \|D^\beta \phi\|_{L^\infty(K)} \end{aligned}$$

Integrating this inequality in s together with the fact that $F_\alpha(0) = 0$ gives the higher order bound. But this means that

$$\begin{aligned} \left| D^\alpha \phi(\gamma(s)) - \sum_{|\beta| \leq k - |\alpha|} D^{\alpha + \beta} \phi(y) (\gamma(s) - y)^\beta / \beta! \right| &= |F_\alpha(L)| \\ &\lesssim L^{k - |\alpha|} \sum_{|\beta| = k} \|D^\beta \phi\|_{L^\infty(K)}. \end{aligned}$$

Choosing γ optimally gives

$$\left| D^\alpha \phi(\gamma(s)) - \sum_{|\beta| \leq k - |\alpha|} D^{\alpha + \beta} \phi(y) (\gamma(s) - y)^\beta / \beta! \right| \lesssim |x - y|^{k - |\alpha|} \sum_{|\beta| = k} \|D^\beta \phi\|_{L^\infty(K)}.$$

The last Lemma, together with this bound, completes the proof. \square

Remark. Similar arguments can be used to show that if Λ is a distribution of order k supported on a compact set K , and there exists $\gamma \leq 1$ such that K is a finite union of connected components, such that for any pair of points x, y in that component, there exists a rectifiable path from x to y with length $O(|x - y|^\gamma)$, and $m \geq k/\gamma$, then

$$|\Lambda(\phi)| \lesssim \sum_{|\alpha| \leq m} \|D^\alpha \phi\|_{L^\infty(K)}.$$

Let us finish by considering a consequence of these results, applied to distributions supported on hyperplanes. For simplicity in notation, we assume this hyperplane is axis oriented.

Theorem 9.10. *Let $x = (x_0, x_1)$, where $x_0 \in \mathbf{R}^{d_1}$, $x_1 \in \mathbf{R}^{d_2}$, and $d = d_1 + d_2$. Let $H = \{(x_0, x_1) \in \mathbf{R}^d : x_1 = 0\}$. If Λ is a distribution of order k compactly supported on H , then there exists distributions Λ_α of order $k - |\alpha|$ on \mathbf{R}^{d_1} for each $|\alpha| \leq k$, where α is a multi-index in the \mathbf{R}^{d_2} variables, and constants γ_α such that*

$$\Lambda(\phi) = \sum \Lambda_\alpha(D^\alpha \phi|_H).$$

Proof. Fix a function $\psi \in \mathcal{D}(\mathbf{R}^{d_1})$ equal to one in a neighborhood of the origin. Given $\phi \in \mathcal{D}(\mathbf{R}^d)$, all derivatives of the function

$$\sum_{|\alpha| \leq k} D^\alpha \phi(x_0, 0) \cdot (x_1^\alpha / \alpha!) \cdot \psi(x_1) = \sum_{|\alpha| \leq k} D^\alpha \phi|_H(x_0) \cdot (x_1^\alpha / \alpha!)$$

agree with ϕ on H , where α ranges over all derivatives in the x_1 direction. It follows that if we define a distribution Λ_α on \mathbf{R}^{d_1} such that for $\psi \in \mathcal{D}(\mathbf{R}^{d_1})$,

$$\Lambda_\alpha(\psi) = \Lambda(\psi \otimes (x_1^\alpha / \alpha!)),$$

then

$$\Lambda(\phi) = \sum_{|\alpha| \leq k} \Lambda_\alpha(D^\alpha \phi|_H).$$

The hard part is showing that Λ_α has order $k - |\alpha|$. If the support of Λ in the x_0 variable is contained in a compact ball B , then, because B is Whitney regular,

$$\begin{aligned} |\Lambda_\alpha(\psi)| &\lesssim \sum_{|\beta_1| + |\beta_2| \leq k} \|D^{\beta_1 + \beta_2} \{\psi \otimes (x_1^\alpha / \alpha!)\}\|_{L^\infty(B \times \{0\})} \\ &= \sum_{|\beta| \leq k - |\alpha|} \|D^\beta \psi\|_{L^\infty(B)}. \end{aligned}$$

This implies Λ_α has order $k - |\alpha|$. □

Remark. This argument does not really need the power of the full extension theorem machinery, since the Whitney extension theorem is relatively trivial in the application we give (we can consider a simple convolution argument to extend a function on a hyperplane to the full space). But the more developed machinery can be applied to characterize distributions on more general sets, which we leave to the reader to experiment with.

9.6 Derivatives of Continuous Functions

One of the main reasons to consider the theory of distributions is so that we can take the derivative of any function we want. We now show that, at least locally, every distribution is the derivative of some continuous function, which means the theory of distributions is essentially the minimal such class of objects which enable us to take derivatives of continuous functions.

Theorem 9.11. *If Λ is a distribution on Ω , and K is a compact set, then there is a continuous function f and α such that for every ϕ ,*

$$\Lambda\phi = (-1)^{|\alpha|} \int_{\Omega} f(x)(D^\alpha\phi)(x) dx$$

Proof. TODO □

Theorem 9.12. *If K is compact, contained in some open subset V , which in turn is a subset of Ω , and Λ has order N , then there exists finitely many continuous functions $f_\beta \in C(\Omega)$ supported on V , for each $|\beta| \leq N + 2$, with supports on V , and with $\Lambda = \sum D^\beta f_\beta$.*

Theorem 9.13. *If Λ is a distribution on Ω , then there exists continuous functions g_α on Ω such that each compact set K intersects the supports of finitely many of the g_α , and $\Lambda = \sum D^\alpha g_\alpha$. If Λ has finite order, then only finitely many of the g_α are nonzero.*

9.7 Convolutions of Distributions

Using the convolution of two functions as inspiration, we will define the convolution of a distribution Λ with a test function ϕ , and under certain conditions, the convolution of two distributions. Recall that if $f, g \in L^1(\mathbf{R}^n)$, then their convolution is the function in $L^1(\mathbf{R}^n)$ defined by

$$(f * g)(x) = \int f(y)g(x - y) dy$$

If we define the translation operators $T_y g(x) = \text{Trans}_y g(x) = g(x - y)$, then $(f * g)(x) = \int f(y)(T_x g^*)(y) dy$, where g^* is the function defined by $g^*(x) = g(-x)$. Thus, if Λ is any distribution on \mathbf{R}^n , and ϕ is a test function on \mathbf{R}^n , we can define a function $\Lambda * \phi$ by setting $(\Lambda * \phi)(x) = \Lambda(T_x \phi^*)$. Notice that since

$$\begin{aligned} \int (T_x f)(y)g(y) dy &= \int f(y - x)g(y) dy = \int f(y)g(x + y) dy \\ &= \int f(y)(T_{-x}g)(y) dy, \end{aligned}$$

so we can also define the translation operators on distributions by setting $(T_x \Lambda)(\phi) = \Lambda(T_{-x} \phi)$. One mechanically verifies that convolution commutes with translations, i.e. $T_x(\Lambda * \phi) = (T_x \Lambda) * \phi = \Lambda * (T_x \phi)$.

Theorem 9.14. $\Lambda * \phi$ is C^∞ , and

$$D^\alpha(\Lambda * \phi) = (D^\alpha \Lambda) * \phi = \Lambda * (D^\alpha \phi).$$

Proof. It is easy to calculate that

$$\begin{aligned} (D^\alpha \Lambda * \phi)(x) &= (D^\alpha \Lambda)(\phi_x^*) = (-1)^{|\alpha|} \Lambda(D^\alpha(T_x \phi^*)) \\ &= \Lambda(T_x(D^\alpha \phi)^*) = (\Lambda * D^\alpha \phi)(x) \end{aligned}$$

If $k \in \{1, \dots, d\}$ and $h \in \mathbf{R}$, we set

$$(\Delta_h f)(x) = \frac{f(x + h e_k) - f(x)}{h}$$

then $\Delta_h \phi$ converges to $D^k \phi$ in $\mathcal{D}(\mathbf{R}^d)$, and as such

$$\begin{aligned} \Delta_h(\Lambda * \phi)(x) &= \frac{(\Lambda * \phi)(x + h e_k) - (\Lambda * \phi)(x)}{h} \\ &= \Lambda \left(\frac{T_{-x - h e_k} \phi^* - T_{-x} \phi^*}{h} \right) \end{aligned}$$

As $h \rightarrow 0$, in $\mathcal{D}(\mathbf{R}^d)$ we have

$$\frac{T_{-x-he_k}\phi^* - T_{-x}\phi^*}{h} \rightarrow -T_{-x}D_k\phi^* = T_{-x}(D_k\phi)^*.$$

Thus, by continuity,

$$\lim_{h \rightarrow 0} \Delta_h(\Lambda * \phi)(x) = \Lambda(T_{-x}(D_k\phi)^*) = (\Lambda * D_k\phi)(x)$$

Iteration gives the general result that $\Lambda * \phi \in C^\infty(\mathbf{R}^d)$. An easy calculation then shows that for each $x \in \mathbf{R}^d$,

$$\begin{aligned} [(D^\alpha \Lambda) * \phi](x) &= (D^\alpha \Lambda)(T_{-x}\phi^*) \\ &= (-1)^{|\alpha|} \Lambda(T_{-x}D^\alpha \phi^*) \\ &= \Lambda(T_{-x}(D^\alpha \phi)^*) \\ &= (\Lambda * D^\alpha \phi)(x). \end{aligned} \quad \square$$

There is a certain duality going on here. Distributions can be viewed as linear functionals on $\mathcal{D}(\mathbf{R}^d)$, but one can also view them as a certain family of linear operators from $\mathcal{D}(\mathbf{R}^d) \rightarrow C^\infty(\mathbf{R}^d)$, and the convolution operator uniquely represents the distribution. In fact, any such operator that is translation invariant and continuous can be represented as convolution by a distribution.

Theorem 9.15. *Let $T : \mathcal{D}(\mathbf{R}^d) \rightarrow C_{loc}^\infty(\mathbf{R}^d)$ be a translation invariant continuous operator. Then there exists a distribution Λ such that $T\phi = \Lambda * \phi$ for all $\phi \in \mathcal{D}(\mathbf{R}^d)$.*

Proof. If we knew $T\phi = \Lambda * \phi$ for some Λ , then we could recover Λ since

$$\int \Lambda(x)\phi(x) dx = T\phi(0).$$

Since T is a continuous operator, the right hand side defines a distribution Λ , and translation invariance allows us to conclude that $T\phi = \Lambda * \phi$ for all $\phi \in \mathcal{D}(\mathbf{R}^d)$. \square

Example. A linear differential operator $P : \mathcal{D}(\mathbf{R}^d) \rightarrow \mathcal{D}(\mathbf{R}^d)$ is translation invariant, from which it follows that there exists a distribution Λ such that $P\phi = \Lambda * \phi$. Of course, $\Lambda(\phi) = P\phi(0)$ is just given by applying the differential operator at the origin.

Theorem 9.16. *If $\phi, \psi \in \mathcal{D}(\mathbf{R}^n)$, then $\Lambda * (\phi * \psi) = (\Lambda * \phi) * \psi$.*

Proof. Let K be a compact set containing the supports of ϕ and ψ . It is simple to verify that for each $x \in \mathbf{R}^d$,

$$(\phi * \psi)^*(x) = \int \phi^*(x + y)\psi(y) dy = \int (T_y \phi^*)(x)\psi(y) dy$$

since the map $y \mapsto (T_y \phi)^* \psi(y)$ is continuous, and vanishes out of the compact set K , we can consider the $C_c^\infty(K)$ valued integral

$$(\phi * \psi)^* = \int_K \psi^*(y) T_y \phi^* ds$$

This means precisely that

$$\begin{aligned} (\Lambda * (\phi * \psi))(0) &= \Lambda((\phi * \psi)^*) = \int_K \psi^*(y) \Lambda(T_y \phi^*) dy \\ &= \int_K \psi^*(y) (\Lambda * \phi)(y) dy = ((\Lambda * \phi) * \psi)(0) \end{aligned}$$

The commutativity in general results from applying the commutativity of the translation operators. \square

A sequence $\{\phi_n\}$ in $\mathcal{D}(\mathbf{R}^n)$ is known as an *approximate identity* in the space of distributions if $\Lambda * \phi_n$ converges to Λ as $n \rightarrow \infty$, for every distribution Λ , and an approximate identity in the space of test functions if $\psi * \phi_n$ converges to ψ in $\mathcal{D}(\mathbf{R}^n)$ for every $\psi \in \mathcal{D}(\mathbf{R}^n)$.

Theorem 9.17. *If $\{\phi_n\}$ is a family of non-negative functions in $\mathcal{D}(\mathbf{R}^n)$ which are eventually supported on every neighbourhood of the origin, and all integrate to one, then $\{\phi_n\}$ is an approximation to the identity in the space of test functions and in the space of distributions.*

Proof. It is easy to verify that if f is a continuous function, then $f * \phi_n$ converges locally uniformly to f as $n \rightarrow \infty$. But now we calculate that if $f \in \mathcal{D}(\mathbf{R}^n)$, then $D^\alpha(f * \phi_n) = (D^\alpha f) * \phi_n$ converges locally uniformly to $D^\alpha f$, which gives that $f * \phi$ converges to f in $\mathcal{D}(\mathbf{R}^n)$. Now if Λ is a distribution, and ψ is a test function, then continuity gives

$$\begin{aligned} \Lambda(\psi^*) &= \lim_{\delta \rightarrow 0} \Lambda(\phi_\delta * \psi) = \lim_{\delta \rightarrow 0} (\Lambda * (\phi_\delta * \psi))(0) \\ &= \lim_{\delta \rightarrow 0} ((\Lambda * \phi_\delta) * \psi)(0) = \lim_{\delta \rightarrow 0} (\Lambda * \phi_\delta)(\psi^*) \end{aligned}$$

and ψ was arbitrary. \square

If Λ is a distribution on \mathbf{R}^n , then the map $\phi \mapsto \Lambda * \phi$ is a linear transformation from $\mathcal{D}(\mathbf{R}^n)$ into $\mathcal{E}(\mathbf{R}^n)$, which commutes with translations. It is also continuous. To see this, we consider a fixed compact K , and consider the map from $C_c^\infty(K)$ to $C_{\text{loc}}^\infty(\mathbf{R}^n)$. Both of these spaces are Fréchet, so we can apply the closed graph theorem. Consider a sequence $\{\phi_n\}$ converging in $C_c^\infty(K)$ to some ϕ , and we suppose that $\{\Lambda * \phi_n\}$ converges in $C_c^\infty(K)$ to a function f . It suffices to show that $\Lambda * \phi = f$. But we calculate that for each $x \in \mathbf{R}^d$,

$$f(x) = \lim_n (\Lambda * \phi_n)(x) = \lim_n \Lambda(T_x \phi_n^*) = \Lambda(\lim_n T_x \phi_n^*) = \Lambda(T_x \phi^*) = (\Lambda * \phi)(x).$$

Here we have used the fact that $T_x \phi_n^*$ converges to $T_x \phi^*$ in $\mathcal{D}(\mathbf{R}^n)$. Thus $\phi \mapsto \Lambda * \phi$ is a continuous, translation invariant operator from $\mathcal{D}(\mathbf{R}^n)$ to $\mathcal{E}(\mathbf{R}^n)$. Surprisingly, the converse is also true.

Theorem 9.18. *If $L : \mathcal{D}(\mathbf{R}^n) \rightarrow \mathcal{E}(\mathbf{R}^n)$ is a continuous linear transformation commuting with translations, then there is a distribution Λ such that $L(\phi) = \Lambda * \phi$.*

Proof. If $L(\phi) = \Lambda * \phi$, then we would have

$$\Lambda(\phi) = (\Lambda * \phi^*)(0) = L(\phi^*)(0)$$

and we take this as the definition of Λ for an arbitrary operator L . Indeed, it then follows that Λ is continuous because all the operations here are continuous, and because L commutes with translations, we conclude

$$(\Lambda * \phi)(x) = \Lambda(T_x \phi^*) = L(T_{-x} \phi)(0) = L(\phi)(x)$$

which gives the theorem. \square

We now move onto the case where a distribution Λ has compact support. Then Λ extends to a continuous linear functional on $\mathcal{E}(\mathbf{R}^n)$, and we can define the convolution $\Lambda * \phi$ if $\phi \in \mathcal{E}(\mathbf{R}^n)$ as above. This is an extension of the convolution operator above to a continuous operator from $\mathcal{E}(\mathbf{R}^n)$ to itself. The same techniques as before, or a density argument, verify that translations and derivatives are carried into the convolution.

Theorem 9.19. *If ϕ and Λ have compact support on \mathbf{R}^d , then $\Lambda * \phi$ has compact support, and moreover,*

$$\text{supp}(\Lambda * \phi) \subset \text{supp}(\Lambda) + \text{supp}(\phi).$$

*The map $\phi \mapsto \Lambda * \phi$ is a continuous operator from $\mathcal{D}(\mathbf{R}^d)$ to itself.*

Proof. Let ϕ be supported on a compact set K , and Λ be supported on K_0 . Then $(\Lambda * \phi)(x) = \Lambda(T_x \phi^*)$. Since $T_x \phi^*$ is supported on $x - K$, $(\Lambda * \phi)(x) = 0$ unless $K_0 \cap (x - K) \neq \emptyset$, i.e. $x \in K_0 + K$.

To obtain continuity of the operator, it suffices to prove that $\phi \mapsto \Lambda * \phi$ is a continuous operator from $C_c^\infty(K)$ to $\mathcal{D}(\mathbf{R}^d)$ for any compact set K . Let Λ be supported on a compact set K_0 . Then $\Lambda * \phi$ is supported on $K + K_0$, and so it suffices to show $\phi \mapsto \Lambda * \phi$ is a continuous operator from $C_c^\infty(K)$ to $C_c^\infty(K + K_0)$. But this follows because the map is continuous into $C_{\text{loc}}^\infty(\mathbf{R}^d)$. \square

Theorem 9.20. *If Λ and ψ have compact support, and $\phi \in C^\infty(\mathbf{R}^n)$, then*

$$\Lambda * (\phi * \psi) = (\Lambda * \phi) * \psi = (\Lambda * \psi) * \phi$$

Proof. Let Λ and ψ be supported on some balanced compact set K . Let V be a bounded, balanced open set containing K . If ϕ_0 is a function with compact support equal to ϕ on $V + K$, then for $x \in V$,

$$(\phi * \psi)(x) = \int \phi(x - y) \psi(y) dy = \int \phi_0(x - y) \psi(y) dy = (\phi_0 * \psi)(x)$$

Thus

$$(\Lambda * (\phi * \psi))(0) = (\Lambda * (\phi_0 * \psi))(0) = ((\Lambda * \psi) * \phi_0)(0)$$

But $\Lambda * \psi$ is supported on $K + K$, so $((\Lambda * \psi) * \phi_0)(0) = ((\Lambda * \psi) * \phi)(0)$. Now we also calculate

$$(\Lambda * (\phi * \psi))(0) = ((\Lambda * \phi_0) * \psi)(0) = ((\Lambda * \phi) * \psi)(0) \int (\Lambda * \phi_0)(-y) \psi(y)$$

where the last fact follows because $\Lambda * \phi_0$ agrees with $\Lambda * \phi$ on K . The general fact follows by applying the translation operators. \square

Now we come to the grand finale, defining the convolution of two distributions. Given two distributions Λ and Ψ , one of which has compact support, we define the linear operator

$$L(\phi) = \Lambda * (\Psi * \phi)$$

This gives a continuous, translation invariant linear operator from $\mathcal{D}(\mathbf{R}^d)$ to $\mathcal{E}(\mathbf{R}^d)$; if Ψ is compactly supported, then $\phi \mapsto \Psi * \phi$ is a continuous operator on $\mathcal{D}(\mathbf{R}^d)$, which gives continuity. If Λ is compactly supported, then

$\Lambda \mapsto \Lambda * \eta$ is a continuous linear operator on $\mathcal{E}(\mathbf{R}^d)$, which gives continuity. But this means that L corresponds to convolution with a distribution, and we define this distribution to be $\Lambda * \Psi$.

Theorem 9.21. *If Λ and Ψ are distributions, one of which has compact support, then $\Lambda * \Psi = \Psi * \Lambda$. Let S_Λ and S_Ψ , and $S_{\Lambda * \Psi}$ denote the supports of Λ , Ψ , and $\Lambda * \Psi$. Then $S_{\Lambda * \Psi} \subset S_\Lambda + S_\Psi$.*

Proof. We calculate that for any two test functions ϕ and ψ ,

$$(\Lambda * \Psi) * (\phi * \psi) = \Lambda * (\Psi * (\phi * \psi)) = \Lambda * ((\Psi * \phi) * \psi)$$

If Λ has compact support, then

$$\Lambda * ((\Psi * \phi) * \psi) = (\Lambda * \psi) * (\Psi * \phi)$$

Conversely, if Ψ has compact support, then

$$\Lambda * ((\Psi * \phi) * \psi) = \Lambda * (\psi * (\Psi * \phi)) = (\Lambda * \psi) * (\Psi * \phi)$$

We also calculate

$$\begin{aligned} \Psi * ((\Lambda * \phi) * \psi) &= \Psi * (\Lambda * (\phi * \psi)) = \Psi * (\Lambda * (\psi * \phi)) \\ &= \Psi * ((\Lambda * \psi) * \phi) = (\Psi * \phi) * (\Lambda * \psi) \end{aligned}$$

But since convolution is commutative, we have

$$((\Lambda * (\Psi * \phi)) * \psi) = \Lambda * ((\Psi * \phi) * \psi) = \Psi * ((\Lambda * \phi) * \psi) = (\Psi * (\Lambda * \phi)) * \psi$$

Since ψ was arbitrary, we conclude

$$(\Lambda * \Psi) * \phi = \Lambda * (\Psi * \phi) = \Psi * (\Lambda * \phi) = (\Psi * \Lambda) * \phi$$

and now since ϕ was arbitrary, we conclude $\Lambda * \Psi = \Psi * \Lambda$. Now we know convolution is commutative, we may assume S_Ψ is compact. The support of $\Psi * \phi^*$ lies in $S_\Psi - S_\phi$. But this means that if $S_\phi - S_\Psi$ is disjoint from S_Λ , which means exactly that S_ϕ is disjoint from $S_\Lambda + S_\Psi$, then

$$(\Lambda * \Psi)(\phi) = (\Lambda * (\Psi * \phi))(0) = 0$$

and this gives the support of $\Lambda * \Psi$. □

This means that the convolution of two distributions with compact support also has compact support. This means that if we have three distributions Λ, Ψ , and Φ , two of which have compact support, then the distributions $\Lambda * (\Psi * \Phi)$ and $(\Lambda * \Psi) * \Phi$ are well defined, so convolution is associative and commutative. We calculate that for any test function ϕ ,

$$(\Lambda * (\Psi * \Phi)) * \phi = \Lambda * (\Psi * (\Phi * \phi))$$

$$((\Lambda * \Psi) * \Phi) * \phi = (\Lambda * \Psi) * (\Phi * \phi)$$

If Φ has compact support, then $\Phi * \phi$ has compact support, and so we can move $(\Lambda * \Psi)$ into the equation to prove equality. If Φ does not have compact support, then Λ and Ψ have compact support, and

$$\Lambda * (\Psi * \Phi) = \Lambda * (\Phi * \Psi)$$

and we can apply the previous case to obtain that this is equal to $(\Lambda * \Phi) * \Psi$. Repeatedly applying the previous case brings this to what we want.

Theorem 9.22. *If Λ and Ψ are distributions, one of which having compact support, then*

$$D^\alpha(\Lambda * \Psi) = (D^\alpha \Lambda) * \Psi = \Lambda * (D^\alpha \Psi).$$

Proof. The Dirac delta function δ satisfies

$$(\delta * \phi)(x) = \int \phi(y) \delta(x - y) dy = \phi(x)$$

so $\delta * \phi = \phi$. Now $D^\alpha \delta$ is also supported at x , since

$$(D^\alpha \delta)(\phi) = (-1)^{|\alpha|} \int \delta(x) (D^\alpha \phi)(x) dx = (-1)^{|\alpha|} (D^\alpha \phi)(0)$$

which means that for any distribution Λ , then $(D^\alpha \delta) * \Lambda$ has compact support,

$$(((D^\alpha \delta) * \Lambda) * \phi)(0) = (D^\alpha \delta)((\Lambda * \phi)^*) = (-1)^{|\alpha|} D^\alpha(\Lambda * \phi)^* = ((D^\alpha \Lambda) * \phi)(0)$$

which verifies that $(D^\alpha \delta) * \Lambda = \delta * (D^\alpha \Lambda)$. But now we find

$$D^\alpha(\Lambda * \Psi) = (D^\alpha \delta) * \Lambda * \Psi = ((D^\alpha \delta) * \Lambda) * \Psi = D^\alpha \Lambda * \Psi$$

$$D^\alpha(\Lambda * \Psi) = D^\alpha(\Psi * \Lambda) = (D^\alpha \Psi) * \Lambda = \Lambda * (D^\alpha \Psi)$$

which verifies the theorem in general. \square

9.8 Schwartz Space and Tempered Distributions

We have already encountered the fact that Fourier transforms are well behaved under differentiation and multiplication by polynomials. If we let $\mathcal{S}(\mathbf{R}^d)$ denote a class of functions under which to study this phenomenon, it must be contained in $L^1(\mathbf{R}^d)$ and $C^\infty(\mathbf{R}^d)$, and closed under multiplication by polynomials, and closed under applications of arbitrary constant-coefficient differential operators. A natural choice is then the family of functions which *decays rapidly*, as well as all of its derivatives; i.e. we let $\mathcal{S}(\mathbf{R}^d)$ be the space of all functions $f \in C^\infty(\mathbf{R}^d)$ such that for any integer n and multi-index α , $|x|^n D^\alpha f \in L^\infty(\mathbf{R}^d)$. The space $\mathcal{S}(\mathbf{R}^d)$ is then locally convex if we consider the family of seminorms

$$\|f\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)} = \sup_{|\alpha| \leq n} \|\langle x \rangle^m D^\alpha f\|_{L^\infty(\mathbf{R}^d)}.$$

Elements of $\mathcal{S}(\mathbf{R}^d)$ are known as *Schwartz functions*, and $\mathcal{S}(\mathbf{R}^d)$ is often known as the *Schwartz space*. The seminorms naturally give $\mathcal{S}(\mathbf{R}^d)$ the structure of a Fréchet space. Sometimes, it is more convenient to use the equivalent family of seminorms $\|f\|_{\mathcal{S}^{\alpha,\beta}(\mathbf{R}^d)} = \|x^\alpha D^\beta f\|_{L^\infty(\mathbf{R}^d)}$, because x^α often behaves more nicely under various Fourier analytic operations. It is obvious that $\mathcal{S}(\mathbf{R}^d)$ is separated by the seminorms defined on it, because $\|\cdot\|_{L^\infty(\mathbf{R}^d)} = \|\cdot\|_{\mathcal{S}^{0,0}(\mathbf{R}^d)}$ is a norm used to define the space. We now show the choice of seminorms make the space complete.

Theorem 9.23. $\mathcal{S}(\mathbf{R}^d)$ is a complete metric space.

Proof. Let $\{f_i\}$ be a Cauchy sequence with respect to the seminorms $\|\cdot\|_{\mathcal{S}^{n,\alpha}(\mathbf{R}^d)}$. This implies that for each integer m , and multi-index α , the sequence of functions $\langle x \rangle^m D^\alpha f_k$ is Cauchy in $L^\infty(\mathbf{R}^d)$. Since $L^\infty(\mathbf{R}^d)$ is complete, there are functions $g_{m,\alpha}$ such that $\langle x \rangle^m D^\alpha f_k$ converges uniformly to $g_{m,\alpha}$. If we set $f = g_{0,0}$, then it is easy to see using the basic real analysis of uniform continuity that f is infinitely differentiable, and $\langle x \rangle^m D^\alpha f = g_{m,\alpha}$. It is then easy to show that f_i converges to f in $\mathcal{S}(\mathbf{R}^d)$. \square

Example. The Gaussian function $\phi : \mathbf{R}^d \rightarrow \mathbf{R}$ defined by $\phi(x) = e^{-|x|^2}$ is Schwartz. For any multi-index α , there is a polynomial P_α of degree at most $|\alpha|$ such that $D^\alpha \phi = P_\alpha \phi$; this can be established by a simple induction. But this

means that for each fixed α , $|P_\alpha(x)| \lesssim_\alpha 1 + |x|^{|\alpha|}$. Since $e^{-|x|^2} \lesssim_{m,\alpha} \langle x \rangle^{-m-|\alpha|}$ for any fixed m and α , we find that for any $x \in \mathbf{R}^d$,

$$|(1 + |x|^m)D^\alpha \phi| \lesssim_{\alpha,m} 1.$$

Since m and α were arbitrary, this shows ϕ is Schwartz.

Example. Any compactly supported smooth function is Schwartz. In particular, the inclusion map

$$\mathcal{D}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$$

is bounded. The proof is left to the reader. An important consequence is that elements of $\mathcal{S}(\mathbf{R}^d)^*$, which we will call tempered distributions, can be viewed as a subspace of the space $\mathcal{D}(\mathbf{R}^d)^*$ of all distributions.

To show that an operator T on $\mathcal{S}(\mathbf{R}^d)$ is bounded, it suffices to show that for each n_0 and m_0 , there is n_1, m_1 such that

$$\|Tf\|_{\mathcal{S}^{n_0,m_0}(\mathbf{R}^d)} \lesssim_{n_0,m_0} \|f\|_{\mathcal{S}^{n_1,m_1}(\mathbf{R}^d)}.$$

For a functional $\Lambda : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathbf{R}$, it suffices to show that there exists n and m such that $|\Lambda f| \lesssim \|f\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)}$. The minimal such choice of n is known as the *order* of the functional Λ . We normally do not care about the constant behind the operators for these norms, since the norms are not translation invariant and therefore highly sensitive to the positions of various functions. We really just care about proving the existence of such a constant. Here are some examples:

- Multiplication gives a continuous bilinear map

$$\mathcal{S}(\mathbf{R}^d) \times \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d).$$

More generally, the bilinear multiplication map

$$C_b^\infty(\mathbf{R}^d) \times \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$$

is continuous.

- For each $x \in \mathbf{R}^d$, the translation operator

$$\text{Trans}_x : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$$

is an isomorphism. For each $\xi \in \mathbf{R}^d$, the modulation operator

$$\text{Mod}_\xi : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$$

is an isomorphism.

- The L^p norms are continuous.
- The Fourier transform is an isomorphism of $\mathcal{S}(\mathbf{R}^d)$.

The last point follows by an application, e.g. of the compatibility of differentiation and polynomial multiplication with the Fourier transform.

Corollary 9.24. *Convolution is a continuous operator*

$$\mathcal{S}(\mathbf{R}^d) \times \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d).$$

Proof. We have

$$f * g = \mathcal{F}^{-1}\{\mathcal{F}\{f\} \cdot \mathcal{F}\{g\}\},$$

and the result then follows from the previous examples. \square

Remark. This result allows us to define the convolution of any two tempered distributions in a natural way extending the theory of convolution of distributions defined before.

Now we get to the interesting part of the theory of Schwartz functions. We have defined a homeomorphic linear transform from $\mathcal{S}(\mathbf{R}^d)$ to itself. The theory of functional analysis then says that we can define a dual map, which is a homeomorphism from the dual space $\mathcal{S}(\mathbf{R}^d)^*$ to itself. Note the inclusion map $\mathcal{D}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$ is continuous, and $\mathcal{D}(\mathbf{R}^d)$ is dense in $\mathcal{S}(\mathbf{R}^d)$. This implies that we have an injective, continuous map from $\mathcal{S}(\mathbf{R}^d)^*$ to $\mathcal{D}(\mathbf{R}^d)^*$, so every functional on the Schwarz space can be identified with a distribution. We call such distributions *tempered*. They are precisely the linear functionals on $\mathcal{D}(\mathbf{R}^d)$ which have a continuous extension to $\mathcal{S}(\mathbf{R}^d)$. Intuitively, this corresponds to having limited growth at infinity.

Example. Recall that for any $f \in L^1_{loc}(\mathbf{R}^d)$, we can consider the distribution $\Lambda[f]$ defined by setting

$$\Lambda[f](\phi) = \int f(x)\phi(x) dx.$$

However, this distribution is not always tempered. If $f \in L^p(\mathbf{R}^d)$ for some p , then, applying Hölder's inequality, we obtain that

$$|\Lambda[f](\phi)| \leq \|f\|_{L^p(\mathbf{R}^d)} \|\phi\|_{L^q(\mathbf{R}^d)}.$$

Since $\|\cdot\|_{L^q(\mathbf{R}^d)}$ is a continuous norm on $\mathcal{S}(\mathbf{R}^d)$, this shows $\Lambda[f]$ is bounded. More generally, if $f \in L^1_{loc}(\mathbf{R}^d)$, and $f(x)(1+|x|)^{-m}$ is in $L^p(\mathbf{R}^d)$ for some m , then $\Lambda[f]$ is a tempered distribution. If $p = \infty$, such a function is known as slowly increasing.

Example. For any Radon measure, μ , we can define a distribution

$$\Lambda[\mu](\phi) = \int \phi(x) d\mu(x)$$

But this distribution is not always tempered. If $|\mu|$ is finite, the inequality $\|\Lambda[\mu](\phi)\| \leq \|\mu\| \|\phi\|_{L^\infty(\mathbf{R}^d)}$ gives boundedness. More generally, if μ is a measure such that for some n ,

$$\int_{\mathbf{R}^d} \frac{d|\mu|(x)}{1+|x|^n} dx < \infty$$

then μ is known as a tempered measure, and acts as a tempered distribution since

$$\begin{aligned} |\Lambda[\mu](\phi)| &\leq \int_{\mathbf{R}^d} |\phi(x)| d|\mu|(x) \\ &\leq \left(\int_{\mathbf{R}^d} \frac{d|\mu|(x)}{1+|x|^n} dx \right) \cdot \|\phi\|_{S^{0,n}(\mathbf{R}^d)}. \end{aligned}$$

Example. Any compactly supported distribution is tempered. Indeed, if Λ is a distribution supported on a compact set K , then it has finite order n for some integer n , and extends to an operator on $C^\infty(\mathbf{R}^d)$. We then find

$$|\Lambda(\phi)| \lesssim \|\phi\|_{C^n(\mathbf{R}^d)} \leq \|\phi\|_{S^{0,n}(\mathbf{R}^d)}.$$

Example. The distribution Λ on \mathbf{R} given by

$$\Lambda(\phi) = p.v \int_{-\infty}^{\infty} \frac{\phi(x)}{x} dx$$

is tempered, since

$$\int_{|x| \geq 1} \frac{\phi(x)}{x} \lesssim \|\phi\|_{S^{1,0}(\mathbf{R}^d)}$$

and

$$p.v \int_{-\infty}^{\infty} \frac{\phi(x)}{x} dx \lesssim \|\phi\|_{C^1(\mathbf{R}^d)} = \|\phi\|_{S^{0,1}(\mathbf{R}^d)}$$

and so Λ is tempered of order 1. This distribution is called the (Cauchy) principal value of $1/x$, often denoted $p.v(1/x)$.

The derivative of a tempered distribution is tempered, and gives a continuous operator on $\mathcal{S}(\mathbf{R}^d)^*$. Multiplication gives a continuous map

$$C_b^\infty(\mathbf{R}^d) \times \mathcal{S}(\mathbf{R}^d)^* \rightarrow \mathcal{S}(\mathbf{R}^d)^*.$$

Furthermore, multiplication by a polynomial is also a continuous operator on $\mathcal{S}(\mathbf{R}^d)$, or more generally, by any smooth function whose derivatives are all slowly increasing.

Let us now apply the distributional method to define the Fourier transform of a tempered distribution. Recall that we heuristically think of Λ as formally corresponding to a regular function f such that

$$\Lambda(\phi) = \int f(x)\phi(x) dx$$

The multiplication formula

$$\int_{\mathbf{R}^d} \widehat{f}(\xi)g(\xi) d\xi = \int_{\mathbf{R}^d} f(x)\widehat{g}(x) dx$$

gives us the perfect opportunity to move the analytical operations on f to analytical operations on g . Thus if Λ is the distribution corresponding to a Schwartz $f \in \mathcal{S}(\mathbf{R}^d)$, the distribution $\widehat{\Lambda}$ corresponding to \widehat{f} , then for any Schwartz $\phi \in \mathcal{S}(\mathbf{R}^d)$,

$$\widehat{\Lambda}(\phi) = \Lambda(\widehat{\phi}).$$

In particular, this motivates us to define the Fourier transform of *any* tempered distribution Λ to be the unique tempered distribution $\widehat{\Lambda}$ such that the equation above holds for all Schwartz ϕ . This distribution exists because the Fourier transform is an isomorphism on the space of Schwartz functions. Clearly, the Fourier transform is a homeomorphism on the space of tempered distributions under the weak topology, and moreover, satisfies all the symmetry properties that the ordinary Fourier transform does, once we interpret scalar, rotation, translation, differentiation, etc, in a natural way on the space of distributions.

Example. Consider the constant function 1, interpreted as a tempered distribution on \mathbf{R}^d . Then for any $\phi \in \mathcal{S}(\mathbf{R}^d)$,

$$1(\phi) = \int \phi(x) dx,$$

Thus for any $\phi \in \mathcal{S}(\mathbf{R}^d)$,

$$\hat{1}(\hat{\phi}) = 1(\phi) = \int \phi(\xi) d\xi = \hat{\phi}(0).$$

Thus $\hat{1}$ is the Dirac delta function at the origin. Similarly, the Fourier inversion formula implies that

$$\hat{\delta}(\hat{\phi}) = \phi(0) = \int \hat{\phi}(\xi) d\xi = 1(\hat{\phi})$$

so the Fourier transform of the Dirac delta function is the constant 1 function.

Example. Let u denote a compactly supported distribution. We claim that $\hat{u} \in C^\infty(\mathbf{R}^d)$ is a smooth function, such that

$$\hat{u}(\xi) = \langle u, e^{-2\pi i \xi \cdot x} \rangle.$$

Indeed, formally speaking,

$$\langle \hat{u}, \phi \rangle = \langle u, \hat{\phi} \rangle = \langle u, \int \phi(x) e^{-2\pi i \xi \cdot x} dx \rangle = \int \phi(x) \langle u, e^{-2\pi i \xi \cdot x} \rangle dx.$$

The proof that \hat{u} is smooth follows because we have control of the derivatives of $e^{-2\pi i \xi \cdot x}$ on the support of u .

Example. The theory of tempered distributions enables us to take the Fourier transform of $f \in L^p(\mathbf{R}^d)$, when $p > 2$ or when $p < 1$. The introduction of distributions is in some sense, essential to this process, because for each $p \notin [1, 2]$, there is $f \in L^p(\mathbf{R}^d)$ such that \hat{f} is not a locally integrable function. Otherwise, we could define an operator $T : L^p(\mathbf{R}^d) \rightarrow L^1(\mathbf{R}^d)$ given by

$$Tf = \hat{f} \mathbf{I}_{|\xi| \leq 1}.$$

If a sequence of functions $\{f_n\}$ converges to f in $L^p(\mathbf{R}^d)$, and Tf_n converges to g in $L^1(\mathbf{R}^d)$, then Tf_n converges distributionally to g , which implies $Tf = g$. The closed graph theorem thus implies that T is a continuous operator from $L^p(\mathbf{R}^d)$ to $L^1(\mathbf{R}^d)$, so there exists $M > 0$ such that

$$\int_{|\xi| \leq 1} |\hat{f}(\xi)| \leq M \|f\|_{L^p(\mathbf{R}^d)}.$$

If $f_\alpha(x) = e^{-\pi\alpha|x|^2}$, then $\widehat{f}_\alpha(\xi) = \alpha^{-d/2}e^{-\pi|\xi|^2/\alpha}$. We have

$$\begin{aligned}\|f_\alpha\|_{L^p(\mathbf{R}^d)} &= \left(\int_{\mathbf{R}^d} e^{-\pi\alpha p|x|^2} dx \right)^{1/p} \\ &= (\alpha p)^{-d/2p} \left(\int_{\mathbf{R}^d} e^{-\pi|x|^2} dx \right)^{1/p} \lesssim_d (\alpha p)^{-1/2p}.\end{aligned}$$

On the other hand, for $|\xi| \leq 1$, $|\widehat{f}_\alpha(\xi)| \geq \alpha^{-d/2}e^{-\pi/\alpha}$, so

$$\int_{|\xi| \leq 1} |\widehat{f}_\alpha(\xi)| \gtrsim_d \alpha^{-d/2}e^{-\pi/\alpha}.$$

Thus we conclude that $\alpha^{-d/2}e^{-\pi/\alpha} \lesssim_d M(\alpha p)^{-d/2p}$, or equivalently,

$$\alpha^{d/2(1/p-1)}e^{-\pi/\alpha} \lesssim_d Mp^{-d/2p}.$$

Taking $\alpha \rightarrow \infty$ gives a contradiction if $p < 1$. For $p > 2$, we give the Gaussian an oscillatory factor that does not affect the L^p norm but boosts the L^1 norm of the Fourier transform. We set

$$g_\delta(x) = \prod_{k=1}^d \frac{e^{-\pi x_k^2/(1+i\delta)}}{(1+i\delta)^{1/2}}.$$

The Fourier transform formula of the Gaussian, when applied using the theory of analytic continuation, shows that

$$\widehat{g}_\delta(\xi) = \prod_{k=1}^d e^{-\pi(1+i\delta)\xi_k^2}.$$

We have

$$\int_{|\xi| \leq 1} |\widehat{g}_\delta(\xi)| = \int_{|\xi| \leq 1} e^{-\pi|\xi|^2} \gtrsim 1.$$

On the other hand, for $\delta \geq 1$,

$$\begin{aligned}\|g_\delta\|_{L^p(\mathbf{R}^d)} &= \left(\int |g_\delta(x)|^p dx \right)^{1/p} \\ &= |1+i\delta|^{-d/2} \left(\int_{-\infty}^{\infty} e^{-p\pi x^2/(1+\delta^2)} dx \right)^{d/p} \\ &\lesssim_d \delta^{-d/2} \delta^{d/p} p^{-d/p} = \delta^{d(1/p-1/2)} p^{-d/p}.\end{aligned}$$

Thus we conclude $1 \lesssim_d M \delta^{d(1/p-1/2)} p^{d/p}$, which gives a contradiction as $\delta \rightarrow \infty$ if $p > 2$.

Example. Consider the Riesz Kernel on \mathbf{R}^d , for each $\alpha \in \mathbf{C}$ with positive real part, as the function

$$K_\alpha(x) = \frac{\Gamma(\alpha/2)}{\pi^{\alpha/2}} |x|^{-\alpha}.$$

Then for $0 < \operatorname{Re}(\alpha) < d$, $\widehat{K}_\alpha = K_{d-\alpha}$. We recall that Γ is defined by the integral formula

$$\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} ds,$$

where $\operatorname{Re}(s) > 0$. We note that if $p = d/\operatorname{Re}(\alpha)$, $K_\alpha \in L^{p,\infty}(\mathbf{R}^d)$. The Marcinkiewicz interpolation theorem implies that if $d/2 < \operatorname{Re}(\alpha) < d$, then K_α can be decomposed as the sum of a $L^1(\mathbf{R}^d)$ function and a $L^2(\mathbf{R}^d)$ function, and so we can interpret the Fourier transform of \widehat{K}_α using techniques in $L^1(\mathbf{R}^d)$ and $L^2(\mathbf{R}^d)$, and moreover, the Marcinkiewicz interpolation theorem implies that

$$\|\widehat{K}_\alpha\|_{L^{q,\infty}(\mathbf{R}^d)} \leq \|K_\alpha\|_{L^{p,\infty}(\mathbf{R}^d)}.$$

where q is the dual of p . In particular, the Fourier transform of K_α is a function. We note that K_α obeys multiple symmetries. First of all, K_α is radial, so \widehat{K}_α is also radial. Moreover, K_α is homogenous of degree $-\alpha$, i.e. for each $x \in \mathbf{R}^d$, $K_\alpha(\varepsilon x) = \varepsilon^{-\alpha} K_\alpha(x)$. This actually uniquely characterizes K_α among all locally integrable functions. Taking the Fourier transform of both sides of the equation for homogeneity, we find

$$\varepsilon^{-d} \widehat{K}_\alpha(\xi/\varepsilon) = \varepsilon^{-\alpha} \widehat{K}_\alpha(x).$$

Thus \widehat{K}_α is homogenous of degree $\alpha - d$. But this uniquely characterizes $\widehat{K}_{d-\alpha}$ out of any distribution, up to multiplicity, so we conclude that for $d/2 < \operatorname{Re}(\alpha) < d$, that \widehat{K}_α is a scalar multiple of $K_{d-\alpha}$. But we know that by a change

into polar coordinates, if A_d is the surface area of a unit sphere in \mathbf{R}^d , then

$$\begin{aligned} \int_{\mathbf{R}^d} K_\alpha(x) e^{-\pi|x|^2} dx &= \frac{\Gamma(\alpha/2)}{\pi^{\alpha/2}} \int_{\mathbf{R}^d} |x|^{-\alpha} e^{-\pi|x|^2} dx \\ &= A_d \frac{\Gamma(\alpha/2)}{\pi^{\alpha/2}} \int_0^\infty r^{d-1-\alpha} e^{-\pi r^2} dr \\ &= A_d \frac{\Gamma(\alpha/2)}{2\pi^{d/2}} \int_0^\infty s^{(d-\alpha)/2-1} e^{-s} ds \\ &= A_d \frac{\Gamma(\alpha/2)\Gamma((d-\alpha)/2)}{\pi^{d/2}}. \end{aligned}$$

But this is also the value of

$$\int_{\mathbf{R}^d} K_{d-\alpha}(x) e^{-\pi|x|^2},$$

so we conclude $\widehat{K}_\alpha = K_{d-\alpha}$ if $d/2 < \operatorname{Re}(\alpha) < d$. We could apply Fourier inversion to obtain the result for $0 < \operatorname{Re}(\alpha) < d/2$, but to obtain the case $\operatorname{Re}(\alpha) = d/2$, we must apply something different. For each $s \in \mathbf{C}$ with $0 < \operatorname{Re}(s) < d$, and for each Schwartz $\phi \in \mathcal{S}(\mathbf{R}^d)$ we define

$$A(s) = \int K_s(\xi) \widehat{\phi}(\xi) d\xi = \frac{\Gamma(s/2)}{\pi^{s/2}} \int |\xi|^{-s/2} \widehat{\phi}(\xi) d\xi.$$

and

$$B(s) = \int K_{d-s}(\xi) \widehat{\phi}(\xi) d\xi = \frac{\Gamma((d-s)/2)}{\pi^{(d-s)/2}} \int |\xi|^{(d-s)/2} \widehat{\phi}(\xi) d\xi.$$

The integrals above converge absolutely for $0 < \operatorname{Re}(s) < d$, and the dominated convergence theorem implies that A and B are both complex differentiable. Since $A(s) = B(s)$ for $d/2 < \operatorname{Re}(s) < d$, analytic continuation implies $A(s) = B(s)$ for all $0 < \operatorname{Re}(s) < d$, completing the proof. For $\operatorname{Re}(\alpha) \geq d$, K_α is no longer locally integrable, and so we must interpret the distribution given by integration by K_α in terms of principal values. The fourier transform of these functions then becomes harder to define.

Example. Let us consider the complex Gaussian defined, for a given invertible symmetric matrix $T : \mathbf{R}^d \rightarrow \mathbf{R}^d$, as $G_T(x) = e^{-i\pi(Tx \cdot x)}$. Then

$$\widehat{G}_T = e^{-i\pi\sigma/4} |\det(T)|^{-1/2} G_{-T^{-1}},$$

where σ is the signature of T , i.e. the number of positive eigenvalues, minus the number of negative eigenvalues, counted up to multiplicity. Thus we need to show that for any Schwartz $\phi \in \mathcal{S}(\mathbf{R}^d)$,

$$e^{-i\pi\sigma/4} |\det(T)|^{-1/2} \int_{\mathbf{R}^d} e^{i\pi(T^{-1}\xi \cdot \xi)} \widehat{\phi}(\xi) d\xi = \int_{\mathbf{R}^d} e^{-i\pi(Tx \cdot x)} \phi(x) dx.$$

Let us begin with the case $d = 1$, in which case we also prove the theorem when T is a complex symmetric matrix. If T is given by multiplication by $-iz$, and if $\sqrt{\cdot}$ denotes the branch of the square root defined for all non-negative numbers and positive on the real-axis, then we note that when $z = \lambda i$,

$$e^{-i\pi\sigma/4} |\det(T)|^{-1/2} = e^{-i\pi \operatorname{sgn}(\lambda)/4} |\lambda|^{-1/2} = \sqrt{z}.$$

Thus it suffices to prove the analytic family of identities

$$\int_{-\infty}^{\infty} e^{-(\pi/z)\xi^2} \widehat{\phi}(\xi) d\xi = \sqrt{z} \int_{-\infty}^{\infty} e^{-\pi z x^2} \phi(x) dx,$$

where both sides are well defined and analytic whenever z has positive real part. But we already know from the Fourier transform of the Gaussian that this identity holds whenever z is positive and real, and so the remaining identities follows by analytic continuation. We note that the higher dimensional identity is invariant under changes of coordinates in $SO(n)$. Thus it suffices to prove the remaining theorem when T is diagonal. But then everything tensorizes and reduces to the one dimensional case. More generally, if $T = T_0 - iT_1$ is a complex symmetric matrix, which is well defined if T_1 is positive semidefinite, then

$$\widehat{G}_T = \frac{1}{\sqrt{i \det(T)}} \cdot G_{-T^{-1}},$$

which follows from analytic continuation of the case for real T .

Example. The Airy function on \mathbf{R} is the tempered distribution defined to be the inverse Fourier transform of $e^{2\pi i \xi^3/3}$, i.e.

$$Ai(x) = \int e^{2\pi i(\xi^3/3 + \xi x)} d\xi.$$

In fact, Ai is an analytic function on \mathbf{R} . For $\varepsilon > 0$, let $\zeta = \xi + i\varepsilon$. Then

$$2\pi i \zeta^3/3 = (2\pi) (i(\xi^3/3 - \varepsilon^2 \xi) - (\varepsilon \xi^2 - \varepsilon^3/3)).$$

This shows that for each $\varepsilon > 0$, $e^{2\pi i(\xi+i\varepsilon)^3/3}$ is a tempered distribution. Since

$$e^{2\pi i(\xi+i\varepsilon)^3/3} - e^{2\pi i\xi^3/3} = e^{2\pi i\xi^3/3} \left(e^{2\pi i\varepsilon^2\xi} e^{-2\pi i\varepsilon\xi^2 + (2\pi/3)\varepsilon^3} - 1 \right).$$

Thus these functions converge locally uniformly to zero, and are uniformly bounded, and thus $e^{2\pi i(\xi+i\varepsilon)^3/3}$ converges in the sense of tempered distributions to $e^{2\pi i\xi^3/3}$. But since $e^{2\pi i(\xi+i\varepsilon)^3/3}$ decreases exponentially, the Fourier transforms of these distributions are holomorphic in a strip of width $O(\varepsilon)$. TODO 7.6.8 of Hörmander Volume 1.

Thus we conclude that

$$|\langle \hat{u}, \phi \rangle| = |\langle u, \hat{\phi} \rangle| \lesssim \|(1 + |x|)^K \phi\|_{L^\infty(\mathbf{R}^d)}.$$

Thus \hat{u} is a distribution of order zero, and thus a measure. But $x^\alpha u$ is a compactly supported distribution for all α , which implies that $D^\alpha \hat{u}$ is a distribution of order zero, and thus a measure.

Example. We know $((-2\pi ix)^\alpha)^\wedge = ((-2\pi ix)^\alpha \cdot 1)^\wedge = D^\alpha \delta$, which essentially provides us a way to compute the Fourier transform of any polynomial, i.e. as a linear combination of dirac deltas and the distribution derivatives of dirac deltas, which are derivatives evaluated at points.

Example. Consider the Hilbert kernel $\Lambda = p.v(1/x)$. We have seen this distribution is tempered, so we can take it's Fourier transform. Now $x\Lambda = 1$, so the derivative of $\hat{\Lambda}$ is $-2\pi i\delta$, where δ is the Dirac delta function at the origin. But this means there exists A such that $\hat{\Lambda}(\xi) = A - 2\pi i \cdot \mathbf{I}(\xi > 0)$. But $\Lambda(-x) = -\Lambda(x)$, implying that $\hat{\Lambda}(-\xi) = -\hat{\Lambda}(\xi)$, and thus $A - 2\pi i = -A$, i.e. $A = i\pi$. Thus

$$\hat{\Lambda}(\xi) = \pi i - 2\pi i \cdot \mathbf{I}(\xi > 0) = -i\pi \cdot \text{sgn}(\xi).$$

Theorem 9.25. If μ is a finite measure, $\hat{\mu}$ is a uniformly continuous bounded function with $\|\hat{\mu}\|_{L^\infty(\mathbf{R}^d)} \leq \|\mu\|$, and

$$\hat{\mu}(\xi) = \int e(-2\pi i x \cdot \xi) d\mu(x)$$

The function $\hat{\mu}$ is also smooth if μ has moments of all orders, i.e. $\int |x|^k d\mu(x) < \infty$ for all $k > 0$.

Proof. Let $\phi \in \mathcal{S}(\mathbf{R}^d)$. We must understand the integral

$$\int_{\mathbf{R}^d} \hat{\phi}(x) d\mu(x).$$

Applying Fubini's theorem, which applies since μ has finite mass, we conclude that

$$\int_{\mathbf{R}^d} \hat{\phi}(x) d\mu(x) = \int_{\mathbf{R}^d} \int_{\mathbf{R}^d} \phi(\xi) e^{-2\pi i \xi \cdot x} d\mu(x) d\xi = \int_{\mathbf{R}^d} \phi(\xi) f(\xi) d\xi,$$

where

$$f(\xi) = \int_{\mathbf{R}^d} e^{-2\pi i \xi \cdot x} d\mu(x).$$

Thus $\hat{\mu}$ is precisely f , and it suffices to show that $\|f\|_{L^\infty(\mathbf{R}^d)} \leq \|\mu\|$, and that f is uniformly continuous. The inequality follows from a simple calculation of the triangle inequality, and the second inequality follows because for some γ ,

$$\begin{aligned} |f(\xi + \eta) - f(\xi)| &= \left| \int_{\mathbf{R}^d} e^{-2\pi i \xi \cdot x} (e^{-2\pi i \eta \cdot x} - 1) d\mu(x) \right| \\ &\leq \int_{\mathbf{R}^d} |e^{-2\pi i \eta \cdot x} - 1| d|\mu|(x). \end{aligned}$$

As $\eta \rightarrow 0$, the dominated convergence theorem implies that this quantity tends to zero, which proves uniform continuity. On the other hand, if $x_i \mu$ is finite for some i , then

$$\frac{f(\xi + \varepsilon e_i) - f(\xi)}{\varepsilon} = \int_{\mathbf{R}^d} e^{-2\pi i \xi \cdot x} \frac{(e^{-2\pi i \varepsilon x_i} - 1)}{\varepsilon} d\mu(x).$$

We can apply the dominated convergence theorem to show that as $\varepsilon \rightarrow 0$, this quantity converges to the classical partial derivative f_i , which has the integral formula

$$f_i(\xi) = (-2\pi i) \int_{\mathbf{R}^d} e^{-2\pi i \xi \cdot x} x_i d\mu(x),$$

which is the Fourier transform of $x_i \mu$. Higher derivatives are similar. \square

Not being compactly supported, we cannot compute the convolution of tempered distributions with all C^∞ functions. Nonetheless, if ϕ is Schwartz, and Λ is tempered, then the definition $(\Lambda * \phi)(x) = \Lambda(T_{-x}\phi^*)$ certainly makes sense, and gives a C^∞ function satisfying $D^\alpha(\Lambda * \phi) = (D^\alpha\Lambda) * \phi = \Lambda * (D^\alpha\phi)$ just as for $\phi \in \mathcal{D}(\mathbf{R}^d)$. Moreover, $\Lambda * \phi$ is a slowly increasing function; to see this, we know there is n such that

$$|\Lambda\phi| \lesssim \|\phi\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)}.$$

Now for $|y| \geq 1$,

$$\|T_y\phi\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)} \leq |x - y|^n \leq 2^n(1 + |y|^n)\|\phi\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)},$$

and so

$$(\Lambda * \phi)(x) = \Lambda(T_{-x}\phi^*) \lesssim_n (1 + |x|^n)\|\phi\|_{\mathcal{S}^{n,m}(\mathbf{R}^d)},$$

which gives that $\Lambda * \phi$ is slowly increasing. In particular, we can take the Fourier transform of $\Lambda * \phi$. Now for any $\psi \in \mathcal{S}(\mathbf{R}^d)$ with $\hat{\psi} \in \mathcal{D}(\mathbf{R}^d)$,

$$\begin{aligned} \int_{\mathbf{R}^d} \widehat{\Lambda * \phi}(\xi) \psi(\xi) d\xi &= \int_{\mathbf{R}^d} (\Lambda * \phi)(x) \hat{\psi}(x) dx \\ &= \int_{\mathbf{R}^d} \Lambda(\hat{\psi}(x) \cdot T_{-x}\phi^*) dx \\ &= \Lambda\left(\int_{\mathbf{R}^d} \hat{\psi}(x) T_{-x}\phi^* dx\right) \\ &= \Lambda\left(\hat{\psi} * \phi^*\right) = \Lambda\left(\hat{\psi} * \hat{\hat{\phi}}\right) \\ &= \Lambda\left(\widehat{\psi\hat{\phi}}\right) = \hat{\Lambda}\left(\psi\hat{\phi}\right) = \hat{\phi}\hat{\Lambda}(\psi). \end{aligned}$$

We therefore conclude that $\widehat{\Lambda * \phi} = \hat{\phi}\hat{\Lambda}$.

Because of the dilation symmetry of the Fourier transform, the family of homogeneous distributions (which are all tempered) is invariant under the Fourier transform. More precisely, the Fourier transform of a distribution on \mathbf{R}^d which is homogeneous of degree σ is a homogeneous distribution of degree $-d - \sigma$.

Lemma 9.26. *If u is homogeneous and $\text{sing-supp}(u) \subset \{0\}$, then \hat{u} is homogeneous, and $\text{sing-supp}(\hat{u}) \subset \{0\}$.*

Proof. Suppose first that u is homogeneous of order a , with $\operatorname{Re}(a) < -n$. Then we can write $u = u_0 + u_1$, where u_0 is supported in a neighborhood of the origin, and u_1 is an integrable function. But the Fourier transform of both of these terms is continuous. Thus \widehat{u} is continuous in this case.

To upgrade this fact, given any homogeneous function u with $\operatorname{sing-supp}(u) \subset \{0\}$, then $D^\alpha u$ is homogeneous with real part less than $-n$ for sufficiently large α , and $\operatorname{sing-supp}(D^\alpha u) \subset \{0\}$, which implies that $|\xi|^\alpha \widehat{u}$ is continuous, and thus \widehat{u} is continuous away from the origin. But since $x^\beta u$ is homogeneous if u is homogeneous, we conclude that $\widehat{x^\beta u} = D^\beta \widehat{u}$ is continuous in $\mathbf{R}^n - \{0\}$ for all β . Thus $\operatorname{sing-supp}(\widehat{u}) \subset \{0\}$. \square

9.9 Kernel Operators

We have seen that all translation-invariant operators $T : \mathcal{D}(\mathbf{R}^d) \rightarrow \mathcal{E}(\mathbf{R}^d)$ are given by convolution by a distribution. Thus convolutions by a distribution are suitably general to represent all continuous translation-invariant operations. Surprisingly, when studying non translation invariant operators T from $\mathcal{D}(\mathbf{R}^d)$ to $\mathcal{E}(\mathbf{R}^d)$, or more generally, from $\mathcal{D}(\Omega_1)$ to $\mathcal{D}(\Omega_2)^*$ for some open subsets Ω_1 and Ω_2 of \mathbf{R}^n and \mathbf{R}^m respectively, we can obtain a similar characterization. Instead of studying operators of the form

$$T\phi(x) = \int \Lambda(x-y)\phi(y) dy$$

we instead study *kernel* operators of the form

$$T\phi(x) = \int K(x,y)\phi(y) dy$$

where K is a distribution on $\Omega_2 \times \Omega_1$ acts as a continuous linear operator from $\mathcal{D}(\Omega_1)$ to $\mathcal{D}(\Omega_2)^*$ by defining, for $\psi \in \mathcal{D}(\Omega_2)^*$,

$$\int \psi(x)(T\phi)(x) = \int \psi(x)K(x,y)\phi(y) dx dy,$$

i.e. by testing K against $\psi \otimes \phi$. The behaviour of the operator T uniquely determines the distribution K , and we call K the *Schwartz kernel* of T . In 1953, Schwartz found the surprising result that kernels define all continuous operators from $\mathcal{D}(\Omega_1)$ to $\mathcal{D}(\Omega_2)^*$. This explains the prevalence of kernel operators in analysis.

Theorem 9.27. Let $T : \mathcal{D}(\Omega_1) \rightarrow \mathcal{D}^*(\Omega_2)$ be a continuous linear operator. Then there exists a unique distribution $K \in \mathcal{D}^*(\Omega_2 \times \Omega_1)$ such that for $\phi \in \mathcal{D}(\Omega_1)$ and $\psi \in \mathcal{D}(\Omega_2)$,

$$\int T\phi(x)\psi(x) dy = \int K(x,y)\psi(x)\phi(y) dx dy.$$

Remark. When $\Omega_1 = \Omega_2 = \mathbf{R}^d$ it is sometimes more elegant to work with a representation of kernel operators in *convolution form*, i.e. writing

$$T\phi(x) = \int k(x,z)\phi(x-z) dz$$

which is just another way to represent the operators above by considering the change of variables $z = x - y$, i.e. we have $k(x,z) = K(x, x-z)$, or $K(x,y) = k(x, x-y)$. We call k the *Schwartz convolution kernel* associated with the operator T . If $k(z)$ is independent of the x -variable, then we have $T\phi = k * \phi$, i.e. the operator T is really given by convolution.

Remark. There are many variants of Schwartz's result. for instance, if $T : \mathcal{S}(\mathbf{R}^n) \rightarrow \mathcal{S}(\mathbf{R}^m)^*$ is a continuous map, then it certainly restricts to a map from $\mathcal{D}(\mathbf{R}^n) \rightarrow \mathcal{D}(\mathbf{R}^m)^*$, and so the vanilla version of Schwartz's theorem shows that there is a distribution kernel K for T . The fact that $\langle K, \phi \otimes \psi \rangle$ is well defined and bilinearly continuous for any $\phi \in \mathcal{S}(\mathbf{R}^n)$ and $\psi \in \mathcal{S}(\mathbf{R}^m)$ implies that K is actually a tempered distribution. Thus every continuous map $T : \mathcal{S}(\mathbf{R}^n) \rightarrow \mathcal{S}(\mathbf{R}^m)^*$ is induced by a kernel in $\mathcal{S}(\mathbf{R}^m \times \mathbf{R}^n)^*$.

Remark. As another variant, if $K \in C^\infty(\Omega_2 \times \Omega_1)$ is a kernel, then we can define

$$Tu(x) = \int K(x,y)u(y) dy$$

for any $u \in \mathcal{E}(\Omega_1)^*$, where we interpret the integral in a distributional sense, and then $Tu \in C^\infty(\Omega_2)$ because u is a continuous linear functional, and K can be viewed as an element of $C^\infty(\Omega_2, C_{\text{loc}}^\infty(\Omega_1))$. Moreover, the operator then extends to a continuous map

$$T : \mathcal{E}(\Omega_1)^* \rightarrow C_{\text{loc}}^\infty(\Omega_2),$$

Conversely, if $T : \mathcal{E}(\Omega_1)^* \rightarrow C_{\text{loc}}^\infty(\Omega_2)$, then the continuity of T implies that

$$K(x,y) = (T\delta_y)(x)$$

lies in $C^\infty(\Omega_2 \times \Omega_1)$. Indeed, it is immediately verified from the continuity of T that $K \in C(\Omega_1, C_{\text{loc}}^\infty(\Omega_2))$ since as $y \rightarrow y_0$, δ_y converges in the strong topology to δ_{y_0} , and so $T\delta_y$ converges to $T\delta_{y_0}$ in $C_{\text{loc}}^\infty(\Omega_2)$. A similar argument shows that K actually lies in $C^\infty(\Omega_1, C_{\text{loc}}^\infty(\Omega_2))$, where

$$D_x^\alpha D_y^\beta K(x, y) = D^\alpha T\{D^\beta \delta_y\}.$$

Thus K is smooth.

Proof. Since Schwartz kernels act via a tensor product, understanding the Schwartz kernel theorem requires an understanding of these tensor products. The key property here is that \mathcal{D}^* is *nuclear*. If X and Y are complete locally convex spaces, X is barrelled, and X_b^* is nuclear, then $L_b(X, Y)$ is isomorphic to $X^* \hat{\otimes} Y$. \square

Remark. The fact that, e.g. the spaces $L^2(\Omega)$ are *not* nuclear spaces (no infinite dimensional norm space can be nuclear) is one reason why, for instance, there are continuous linear operators $T : L^2(\Omega_1) \rightarrow L^2(\Omega_2)$ for which there does not exist a kernel $K \in L^2(\Omega_2 \times \Omega_1)$ such that

$$Tf(x) = \int K(x, y)f(y) dy,$$

i.e. the class of Hilbert-Schmidt operators does not constitute the class of all continuous operators.

Given an operator $T : \mathcal{D}(\Omega_1) \rightarrow \mathcal{D}(\Omega_2)^*$ with associated kernel $K(x, y)$, we can consider the transpose operator $T^t : \mathcal{D}(\Omega_2) \rightarrow \mathcal{D}(\Omega_1)^*$ given by the kernel $K(y, x)$. Then

$$\langle T\phi, \psi \rangle = \int K(x, y)\psi(x)\phi(y) dx dy = \langle \phi, T^*\psi \rangle.$$

Thus T^t is formally adjoint to T , in some sense. In particular, if we consider the adjoint $T^* : \mathcal{D}(\Omega_2)^{**} \rightarrow \mathcal{D}(\Omega_1)^*$, then $T^*\phi = T^t\phi$ for each $\phi \in \mathcal{D}(\Omega_2)$.

Looking at the properties of kernels defining an operator is often a useful technique to gain insight in how an operator behaves, since one can study the singularities and smoothness of the kernel separately from the operator itself. A basic result along this form is that an operator T has Schwartz kernel K , then for any test function ϕ , $\text{supp}(T\phi) \subset \text{supp}(K) \circ \text{supp}(\phi)$. Let us consider some more general properties of operator that can be defined in terms of the kernel K of the operator.

Theorem 9.28. Let $T : \mathcal{D}(\Omega_1) \rightarrow \mathcal{D}(\Omega_2)^*$ be an operator with Schwartz kernel $K \in \mathcal{D}(\Omega_2 \times \Omega_1)^*$:

- (i) T maps $\mathcal{D}(\Omega_1)$ into $\mathcal{E}(\Omega_2)$ if and only if $K \in C^\infty(\Omega_2, \mathcal{D}(\Omega_1)^*)$.
- (ii) T extends to a continuous linear operator from $\mathcal{E}(\Omega_1)^*$ to $\mathcal{D}(\Omega_2)^*$ if and only if $K \in C^\infty(\Omega_1, \mathcal{D}(\Omega_2)^*)$.
- (iii) K is called *separately regular* if it is an element of

$$C^\infty(\Omega_1, \mathcal{D}(\Omega_2)^*) \cap C^\infty(\Omega_2, \mathcal{D}(\Omega_1)^*).$$

Then K is *completely regular* if and only if T maps $\mathcal{D}(\Omega_1)$ into $\mathcal{E}(\Omega_2)$, and extends to a map from $\mathcal{E}(\Omega_1)$ to $\mathcal{D}(\Omega_2)^*$.

- (iv) Suppose $\Omega_1 = \Omega_2$ is the same open set Ω . Then K is called *very regular* if it is *separately regular*, and agrees with a C^∞ function away from the diagonal

$$\Delta = \{(x, x) : x \in \Omega\}.$$

If K is *very regular*, then T is *pseudolocal*, i.e. for any compactly supported distribution u , $\text{sing-supp}(Tu) \subset \text{sing-supp}(u)$.

To make a remark, an operator $T : \mathcal{D}(\Omega)^* \rightarrow \mathcal{E}(\Omega)$ is called *local* if $\text{supp}(Tu) \subset \text{supp}(u)$ for any $u \in \mathcal{D}(\Omega)^*$. It is a result of Peetre that any local operator (continuous or non continuous) is a partial differential operator with coefficients in $C^\infty(\Omega)$.

- (v) We say T is *smoothing*, or *regularizing*, if T extends to a map from $\mathcal{E}(\Omega_2)^*$ to $C^\infty(\Omega_1)$. Then T is *smoothing* if and only if $K \in C^\infty(\Omega_2 \times \Omega_1)$.
- (vi) We say T is *proper* if T maps $\mathcal{D}(\Omega_1)$ into $\mathcal{E}(\Omega_2)^*$, and T^t maps $\mathcal{D}(\Omega_2)$ into $\mathcal{E}(\Omega_1)^*$. This holds if and only if for any $\phi \in \mathcal{D}(\Omega_1)$ and $\psi \in \mathcal{D}(\Omega_2)$, $K\phi$ and ψK are in $\mathcal{E}(\Omega_2 \times \Omega_1)^*$, or equivalently, if for any precompact open set $U \Subset \Omega_1$, there is a precompact open set $V \Subset \Omega_2$ such that if $\text{supp}(\phi) \subset U$, then $\text{supp}(T\phi) \subset V$, and conversely, for each such V , there is U such that if $\text{supp}(\psi) \subset V$, then $\text{supp}(T^t\psi) \subset U$.

Under the assumption that both (ii) and (vi) holds, i.e. $K \in C^\infty(\Omega_1, \mathcal{D}(\Omega_2)^*)$ and T is *proper*, the operator T extends to a map from $\mathcal{D}(\Omega_1)^*$ to $\mathcal{D}(\Omega_2)^*$

Proof. Let us begin with the proof of (i). Suppose $K \in C^\infty(\Omega_2, \mathcal{D}(\Omega_1)^*)$. If $\phi \in \mathcal{D}(\Omega_1)$, then the function

$$u(x) = \int K(x, y) \phi(y) dy$$

then lies in $\mathcal{E}(\Omega_2)$, since one calculates that

$$D^\alpha u(x) = \int (D^\alpha K)(x, y) \phi(y) dy.$$

Conversely, if T maps $\mathcal{D}(\Omega_1)$ into $\mathcal{E}(\Omega_2)$, then for any ϕ ,

$$\int \frac{K(x + \delta e_i, y) - K(x, y)}{\delta} \phi(y) dy = \int \Delta_{y_i, \delta} K(x, y) \phi(y) dy$$

converges locally uniformly in x to $D^i T \phi(x)$. Thus $\{\Delta_{y_i, \delta} K\}$ converges in the weak $*$ topology to $D^i T$, and the uniform boundedness theorem implies that it actually converges in the strong topology to $D^i T$, and that $D^i T : \mathcal{D}(\Omega_1) \rightarrow \mathcal{D}(\Omega_2)^*$ is continuous for all i . The Schwartz kernel of $D^i T$ is clearly the derivative $D^i K$, where K is viewed as a map from $\mathcal{D}(\Omega_1)$ to $\mathcal{D}(\Omega_2)^*$. Iterating this argument shows that $K \in C^\infty(\mathcal{D}(\Omega_1), \mathcal{D}(\Omega_2)^*)$.

Now let us prove (ii). Suppose $K \in C^\infty(\Omega_1, \mathcal{D}(\Omega_2)^*)$. Then (i) implies that T^t maps $\mathcal{D}(\Omega_2)$ into $\mathcal{E}(\Omega_1)$. Taking adjoints shows that the adjoint of T^t is an extension of T to a continuous map from $\mathcal{E}(\Omega_1)^*$ to $\mathcal{D}(\Omega_2)^*$. Conversely, if T extends to a continuous map from $\mathcal{E}(\Omega_1)^*$ to $\mathcal{D}(\Omega_2)^*$, then it is simple to prove that if K is the kernel of T , then $K(x, y) = T \delta_y(x)$, and thus lies in $C^\infty(\Omega_1, \mathcal{D}(\Omega_2)^*)$.

Result (iii) follows immediately from (i) and (ii).

To prove result (iv), we note that if K is very regular, then T maps $\mathcal{D}(\Omega)$ into $\mathcal{E}(\Omega)$. Furthermore, if $\tilde{K} \in C^\infty(X^2 - \Delta)$ is the smooth function agreeing with K away from the origin, then for any $u \in \mathcal{E}(\Omega)^*$, and any $x \notin \text{supp}(u)$,

$$Tu(x) = \int K(x, x') u(x') dy = \int \tilde{K}(x, x') u(x') dy.$$

The right hand side defines a smooth function of x for $x \in \Omega - \text{supp}(u)$. But this gives us the initial estimate $\text{sing-supp}(Tu) \subset \text{supp}(u)$. To prove the full statement, if $u \in \mathcal{E}(\Omega)^*$, and $x_0 \notin \text{sing-supp}(u)$, then there is $\phi \in \mathcal{D}(\Omega)$ which equals one in a neighborhood U of x_0 , such that $\phi u \in \mathcal{D}(\Omega)$. Thus

$T(\phi u) \in \mathcal{E}(\Omega)$. But $\text{supp}((1 - \phi)u)$ does not contain x_0 , which implies that x_0 does not lie in $\text{sing-supp}(T((1 - \phi)u))$. But this means that both $T(\phi u)$ and $T((1 - \phi)u)$ are both smooth in a neighborhood of x_0 , which means the same is true of Tu . Thus $x_0 \notin \text{sing-supp}(Tu)$.

We have already proved (v) in a remark following the statement of the Schwartz kernel theorem.

The first part of the proof of (vi) is left to the reader. The assumptions of (ii) imply T^t maps $\mathcal{D}(\Omega_2)$ to $\mathcal{E}(\Omega_1)$ continuously, and the assumptions of (vi) imply that T^t maps $\mathcal{D}(\Omega_2)$ to $\mathcal{E}(\Omega_1)^*$ continuously. Taking both these properties into account, we conclude that T^t maps $\mathcal{D}(\Omega_2)$ into $\mathcal{D}(\Omega_1)$ continuously. But taking adjoints gives an extension of T as a continuous map from $\mathcal{D}(\Omega_1)^*$ to $\mathcal{D}(\Omega_2)^*$. \square

The pseudodifferential operators we will study later are very regular, so in particular, pseudolocal. In fact, they are *microlocal*, in the sense that they never expand the *wavefront set* of an operator, a subtle refinement of the singular support of a function.

9.10 Test Functions on a Manifold

For a general smooth manifold M , it is unnatural to define the space of test functions $\mathcal{D}(M)$ to be $C_c^\infty(M)$. This is because there is no natural inclusion map from $C^\infty(M)$ to this dual, given that there is no natural definition of integration on a manifold M . Thus one cannot naturally think of the dual of $C_c^\infty(M)$ as being a space of ‘generalized functions’ on M .

We remedy this, we work with *scalar densities* rather than scalar-valued functions. Recall that on any manifold M , we can define a line bundle $\text{Vol}(TM)$, and scalar densities are sections of this line bundle. In coordinates, a scalar density corresponds to a family of functions $\omega_x \in C^\infty(U)$ for each coordinate chart (x, U) , such that for any other coordinate chart (y, V) , on $U \cap V$,

$$\omega_y = \omega_x \cdot \left| \det \left(\frac{\partial y^i}{\partial x^j} \right) \right|.$$

We define $\mathcal{D}(M)$ to be the space of all compactly supported smooth scalar densities, which we can equip with a natural locally convex structure analogous to the topology on the space of test functions in Euclidean space.

Given any $f \in C^\infty(M)$, and any $\omega \in \mathcal{D}(M)$, the quantity

$$\langle f, \omega \rangle = \int_M f \cdot \omega$$

is well defined, where the integral on the right is defined by working in local coordinates. Moreover, the map $\omega \mapsto \langle f, \omega \rangle$ is then a continuous linear functional on $\mathcal{D}(M)$. Thus we have a continuous inclusion $C^\infty(M) \rightarrow \mathcal{D}(M)^*$, which means it makes sense to define $\mathcal{D}(M)^*$ be the space of generalized functions on M .

Remark. If it helps to think more generally, given a smooth line bundle E over M , we define $\mathcal{D}(M, E)$ to be the space of all smooth, compactly supported sections of E^* , the dual line bundle to E . Then the space $\Gamma(E)$ of all smooth sections of E is naturally included in $\mathcal{D}(M, E)^*$, which we call the space of *generalized sections* of E .

As an example of this approach, we can consider the line bundle $\text{Vol}^\alpha(TM)$ for each $0 \leq \alpha \leq 1$, whose sections ω consist of *scalar densities of order α* , which transform in coordinates via the relation

$$\omega_y = \omega_x \cdot \left| \det \left(\frac{\partial y^i}{\partial x^j} \right) \right|^\alpha.$$

Thus $\text{Vol}(TM) = \text{Vol}^1(TM)$, and $TM = \text{Vol}^0(TM)$. The dual bundle to $\text{Vol}^\alpha(TM)$ can naturally be identified with $\text{Vol}^{1-\alpha}(TM)$ since if ω is a scalar density of order α , and η is a scalar density of order $1 - \alpha$, then $\omega\eta$ is a scalar density of order one, and thus can be integrated on M in an invariant manner. Thus we can define the space $\mathcal{D}^\alpha(M)^*$ of generalized scalar densities of order α to be the dual to the space of compactly supported scalar densities of order $1 - \alpha$.

Much of the basic theory of distributions follows for distributions on manifolds. In particular, a version of the Schwartz kernel theorem holds, i.e. for any smooth manifolds M and N , and any continuous linear operator $T : \mathcal{D}(M) \rightarrow \mathcal{D}(N)^*$, there exists a generalized section K of $\text{Vol}(M) \oplus TN$ such that for any smooth, compactly supported function f on M , we formally have

$$Tf(x) = \int K(x, y) f(y) dy,$$

in the sense that for any smooth, compactly supported scalar density ω on N ,

$$\langle Tf, \omega \rangle = \int \omega(x) K(x, y) f(y) dx dy.$$

The idea of the proof is to work locally in coordinates, applying the Schwartz kernel theorem in Euclidean space, and then noting that the kernels patch together if we view them as sections of $\text{Vol}(M) \oplus TN$.

It is more difficult to see how one would build a canonical definition of the space $\mathcal{S}^*(M)$ of tempered distributions on a manifold M , because there is no natural direction to measure the rate of decay of a function. Thus it is often wiser to do the things one does in Schwartz space (e.g. Fourier-type arguments) by working locally in coordinates, or on a compact manifold, where $C^\infty(M)$ should coincide with $\mathcal{S}(M)$.

9.11 Paley-Wiener Theorem

TODO: See Rudin, Functional Analysis, or Treves, Chapter 29, or Hörmander Vol 1, Section 7.3.

Theorem 9.29. *Let $K \subset \mathbf{R}^n$ be a convex, compact subset with supporting function H . If u is a distribution of order m , supported on K , then the Fourier transform of u is pointwise defined by*

$$\hat{u}(\xi) = \langle u, e^{-2\pi i \xi \cdot x} \rangle = \int u(x) e^{-2\pi i \xi \cdot x}.$$

Moreover, this integral formula gives an extension of \hat{u} to an entire function on \mathbf{C}^n , and

$$|\hat{u}(\xi)| \lesssim (1 + |\xi|)^N e^{2\pi H(\text{Im}(\xi))}.$$

Conversely, any entire function $f : \mathbf{C}^n \rightarrow \mathbf{C}$ satisfying estimates of the form

$$|f(\xi)| \lesssim (1 + |\xi|)^N e^{2\pi H(\text{Im}(\xi))}$$

is tempered, and its inverse Fourier transform is supported on K . In particular, if f satisfies estimates of the form

$$|f(\xi)| \lesssim_N (1 + |\xi|)^{-N} e^{2\pi H(\text{Im}(\xi))}$$

for all $N > 0$, then the inverse Fourier transform of f is smooth and compactly supported on K .

Proof. Suppose u is a distribution of order N supported on K . For each $\delta > 0$, define a smooth function χ_δ supported on K_δ , equal to one on $K_{\delta/2}$, and with $\|\partial^\alpha \chi_\delta\| \lesssim_\alpha \delta^{-|\alpha|}$. Then

$$\begin{aligned} |\hat{u}(\xi)| &= \int u(x) \chi_\delta(x) e^{-2\pi i \xi \cdot x} \\ &\lesssim \sup_{|\alpha| \leq N} \|\partial_x^\alpha \{\chi_\delta e^{-2\pi i \xi \cdot x}\}\| \\ &\lesssim e^{2\pi H(\text{Im}(\xi)) + \delta |\text{Im}(\xi)|} \sum_{i=0}^N \delta^{-i} (1 + |\xi|)^{N-i}. \end{aligned}$$

Taking $\delta = (1 + |\xi|)^{-1}$ gives the result.

Conversely, let us begin with the case where f satisfies the decay estimates for all $N > 0$, and show this it's Fourier transform is smooth and compactly supported on K . Then we can define the inverse Fourier transform pointwise via the integral formula

$$u(x) = \int f(\xi) e^{2\pi i \xi \cdot x}.$$

To prove that u is supported on K , we note that we can perform a contour shift, writing

$$u(x) = \int f(\xi + i\eta) e^{2\pi i(\xi + i\eta) \cdot x}.$$

Applying the required decay estimate with $N = n + 1$ gives

$$|u(x)| \leq e^{2\pi(H(\eta) - \eta \cdot x)} \int (1 + |\xi|)^{-n-1} d\xi$$

Thus if, for a given x , there exists η with $H(\eta) \leq \eta \cdot x$, then scaling η gives $u(x) = 0$. But if $H(\eta) \geq \eta \cdot x$ for all $\eta \in \mathbf{R}^d$, it follows that $x \in K$.

Now we consider the general case. Consider an entire function f satisfying estimates of the form above for a fixed N , and let u denote it's Fourier transform. Consider a smooth function ϕ supported in the unit ball, with $\int \phi(x) dx = 1$, and let $\phi_\delta(x) = \delta^{-n} \phi(x/\delta)$. Then the inverse Fourier transform of $f_\delta(\xi) = f(\xi) \hat{\phi}(\delta\xi)$ is $u * \phi_\delta$. Now because ϕ is supported in a unit ball, it's Fourier transform has the estimates proved above, and so in particular,

$$|f_\delta(\xi)| \lesssim_M (1 + |\xi|)^{N-M} e^{2\pi(H(\text{Im}(\eta)) + \delta |\text{Im}(\xi)|)}.$$

Thus by the last paragraph, we conclude that $u * \phi_\delta$ is supported on K_δ , and taking $\delta \rightarrow 0$ completes the proof since $u * \phi_\delta$ converges to u distributionally. \square

9.12 Hyperfunctions

TODO: Treves Chapter 22, and Hörmander 7.3,7.4,7.5,8.4,8.5,8.6,8.7,Chapter 9.

Chapter 10

Microlocal Analysis of Singularities

Suppose u is a distribution on \mathbf{R}^d . The *singular support* of u is the set of points $x_0 \in \mathbf{R}^d$ which *do not* have an open neighbourhood upon which u acts as integration against a C^∞ function. Understanding the singular support of a distribution, and how to control it, is often a useful perspective in harmonic analysis; to reduce the study of u to the study of a C^∞ function one need only smoothen around the singular support of u .

The smoothness of a distribution is linked to the decay of its Fourier transform. In particular, suppose there is a compactly supported bump function $\phi \in C^\infty(\mathbf{R}^d)$ with $\phi(x) = 1$ in a neighbourhood of some point $x_0 \in \mathbf{R}^d$. Since ϕu is compactly supported, the Paley-Wiener theorem implies $\widehat{\phi u}$ is an entire function with polynomial growth at infinity. The Fourier inversion formula implies that $\phi u \in \mathcal{D}(\mathbf{R}^d)$ if and only if for all $N \geq 0$, $|\widehat{\phi u}(\xi)| \lesssim_N |\xi|^{-N}$. Thus we can infer the singular support of u via purely spectral means, provided we localize about a point.

We can therefore gain more detailed information about singularities of a distribution u through the Fourier transform. If x_0 is a singularity of u , then for any bump function $\phi \in C^\infty(\mathbf{R}^d)$ with $\phi(x) = 1$ in a neighbourhood of x_0 , there must exist some direction in frequency space on which $\widehat{\phi u}$ does not decay. However, this does not mean that ϕ is unable to decay in certain directions; there might exist a conical neighbourhood U about the origin containing some frequency ξ_0 such that for all $\xi \in U$ and all $N > 0$,

$$|\widehat{u\phi}(\xi)| \lesssim_N |\xi|^{-N}. \quad (10.1)$$

the set of values ξ_0 which do *not* satisfy (10.1) for any choice of bump function ϕ about x_0 forms a closed conical subset of \mathbf{R}^d , and we call this the *wavefront* of u about the singularity x_0 . The set

$$\text{WF}(u) = \{(x_0, \xi_0) : \xi_0 \text{ is in the wavefront of } u \text{ at } x_0\}$$

is the *wavefront set* of the distribution, and provides a deeper characterization of the singularities of u . For instance, in order to smoothen out a distribution u one need only average along the directions in the wavefront set. This is the beginning of *microlocal analysis*, localization not only in space, but also localization in a conic subset of space / frequency, e.g. the tangent bundle.

Remark. The term *wavefront set* is meant to be in analogy to the theory of Huygens on wave propagation. If one knows the location and tangent plane to a wave, then it is translated along the normal direction to the plane. We will later see that if the principal symbol of a linear partial differential operator is real with constant coefficients, then the wavefront set of linear partial differential equations is invariant under the bicharacteristic flow, which is the case of the wave equation reduces to Huygen's theory.

Let us now discuss the wavefront set a little more precisely. If u is a compactly supported distribution on \mathbf{R}^d , we define $\Gamma(u)$ to be the set of $\xi_0 \in \mathbf{R}^d$ which have no conical neighbourhood U such that for each $N > 0$ and $\xi \in U$,

$$|\hat{u}(\xi)| \lesssim_N |\xi|^{-N}. \quad (10.2)$$

It is simple to verify via a compactness argument that if $\Gamma(u) = \emptyset$, then $u \in C^\infty(\mathbf{R}^d)$.

Lemma 10.1. *If u is a compactly supported distribution and $\phi \in \mathcal{D}(\mathbf{R}^d)$, then*

$$\Gamma(\phi u) \subset \Gamma(u).$$

Proof. Suppose $\xi_0 \notin \Gamma(u)$, so ξ_0 has a conical neighbourhood U such that (10.2) holds. Then there exists $\varepsilon > 0$ such that U contains

$$\left\{ \eta \in \mathbf{R}^d : \frac{\xi_0 \cdot \eta}{|\xi_0||\eta|} \geq 1 - 2\varepsilon \right\}$$

Let V be the conical neighbourhood of ξ_0 defined by setting

$$V = \left\{ \eta \in \mathbf{R}^d : \frac{\xi_0 \cdot \eta}{|\xi_0||\eta|} \geq 1 - \varepsilon \right\}.$$

We claim V satisfies (10.2). Fix $\xi \in V$. Then

$$|\widehat{\phi u}(\xi)| = (\widehat{\phi} * \widehat{u})(\xi) = \int_{\mathbf{R}^d} \widehat{\phi}(\eta) \widehat{u}(\xi - \eta) d\xi.$$

If $|\xi - \eta| \leq 0.25\varepsilon|\xi|$, then it is simple to verify that

$$(\xi_0 \cdot \eta) \geq (1 - 2\varepsilon)|\xi_0||\eta|$$

so $\eta \in U$. Thus for any $N > 0$, $\widehat{u}(\eta) \lesssim_N 1/(1 + |\eta|)^N$. Since $\phi \in L^\infty(\mathbf{R}^d)$, we conclude

$$\begin{aligned} \int_{|\eta| \leq 0.25\varepsilon|\xi|} \widehat{\phi}(\eta) \widehat{u}(\xi - \eta) d\xi &\lesssim_\phi \int_{|\eta| \leq 0.25\varepsilon|\xi|} \frac{1}{1 + |\xi - \eta|^N} \\ &\lesssim_{\varepsilon, d} \frac{|\xi|^d}{(1 + 2|\xi|^N)} \lesssim \frac{1}{1 + |\xi|^{N-d}}. \end{aligned}$$

On the other hand, since u is compactly supported, \widehat{u} is slowly increasing, i.e. there exists $m > 0$ such that

$$|\widehat{u}(\xi)| \leq 1 + |\xi|^m.$$

Since $\phi \in \mathcal{D}(\mathbf{R}^d)$, we have $|\widehat{\phi}(\eta)| \lesssim_M 1/(1 + |\eta|^M)$ for all $M > 0$ and thus we conclude that if $M > m + d$

$$\begin{aligned} \int_{|\eta| \geq 0.25\varepsilon|\xi|} \widehat{\phi}(\eta) \widehat{u}(\xi - \eta) &\lesssim_M \int_{|\eta| \geq 0.25\varepsilon|\xi|} \frac{1 + |\xi - \eta|^m}{1 + |\eta|^M} \\ &\lesssim_{\varepsilon, m} \int_{|\eta| \geq 0.25\varepsilon|\xi|} \frac{1 + |\eta|^m}{1 + |\eta|^M} \\ &\lesssim_{\varepsilon, d} \frac{1}{1 + |\xi|^{M-m-d}}. \end{aligned}$$

Choosing the parameter M and N appropriately, we obtain the required bound which shows that $\xi_0 \notin \Gamma(\phi u)$. \square

This fact means that for each distribution u , and any pair of distributions $\phi_1, \phi_2 \in \mathcal{D}(\mathbf{R}^d)$ such that $\text{supp}(\phi_2)$ is contained in the interior of the support of ϕ_1 , then $\phi_2/\phi_1 \in \mathcal{D}(\mathbf{R}^d)$, and so we conclude that

$$\Gamma(\phi_2 u) = \Gamma((\phi_2/\phi_1)\phi_1 u) \subset \Gamma(\phi_1 u).$$

Thus if u is a distribution, and $x \in \mathbf{R}^d$, then we define $\Gamma_x(U)$ to be equal to

$$\bigcap \left\{ \Gamma(\phi u) : \phi \in \mathcal{D}(\mathbf{R}^d), x \in \text{supp}(\phi) \right\}.$$

If $\{\phi_n\}$ is a sequence in $\mathcal{D}(\mathbf{R}^d)$ such that $\text{supp}(\phi_{n+1})$ is compactly supported in $\text{supp}(\phi_n)$ for each n , and if $\bigcap \text{supp}(\phi_n) = \{x\}$, then $\Gamma_x(u) = \lim_{n \rightarrow \infty} \Gamma(\phi_n u)$. Finally, we define

$$\text{WF}(u) = \{(x, \xi) : \xi \in \Gamma_x(u)\}.$$

This is the *wavefront set* of u .

Remark. The theory of wavefront sets can also be defined via a sheaf theoretic framework. The distributions on a manifold form a sheaf \mathcal{D}^* , since one can restrict and glue distributions defined on open subsets of a manifold. Similarly, the smooth functions on a manifold form a sheaf C^∞ , which is a subsheaf of \mathcal{D}^* . Thus we can consider the quotient sheaf $\mathcal{F} = \mathcal{D}^*/C^\infty$. The support of a distribution in \mathcal{F} is then precisely the singular support of that distribution. Given a manifold M , we can consider a natural sheaf on T^*M . For each open set $U \subset T^*M$, we consider the family of all distributions on M , modulo the space of distributions u with $\text{WF}(u) \cap U = \emptyset$. This is called the *sheaf of microdistributions on M* . The support of a distribution in this sheaf is precisely the wavefront set of the distribution.

Lemma 10.2. *If u is a distribution, then $\pi_x(\text{WF}(u))$ is the singular support of u . If u is compactly supported, then $\pi_\xi(\text{WF}(u)) = \Gamma(u)$.*

Proof. Fix $x_0 \in \mathbf{R}^d$. If $(x_0, \xi_0) \notin \text{WF}(u)$ for all $\xi_0 \in \mathbf{R}^d$, then there exists $\phi \in \mathcal{D}(\mathbf{R}^d)$ such that $\phi(x_0) \neq 0$ and $\Gamma(\phi u) = \emptyset$. But this means $\phi u \in \mathcal{D}(\mathbf{R}^d)$, so x_0 is not in the singular support of u . This shows $\pi_x(\text{WF}(u))$ is a subset of the singular support. The converse is obvious.

On the other hand, let us assume u is compactly supported, and that $\xi_0 \notin \Gamma(u)$. Then $(x_0, \xi_0) \notin \text{WF}(u)$ for any $x_0 \in \mathbf{R}^d$ since $\Gamma(\phi u) \subset \Gamma(u)$ for any

$\phi \in \mathcal{D}(\mathbf{R}^d)$. But if $(x_0, \xi_0) \notin \text{WF}(u)$ for any $x_0 \in \mathbf{R}^d$ we can cover the support of u by a partition of unity $\phi_1, \dots, \phi_N \in \mathcal{D}(\mathbf{R}^d)$ such that $\xi_0 \notin \Gamma(\phi_i u)$ for each i , and summing up shows $\xi_0 \notin \Gamma(u)$. \square

Example. Suppose u is a homogenous distribution which is C^∞ away from the origin. Then \hat{u} is homogenous and C^∞ away from the origin, and we claim that

$$\text{WF}(u) = \{(0, \xi) : \xi \in \text{supp}(\hat{u})\} = \{0\} \times \Gamma(u).$$

Since the singular support of u is $\{0\}$, we know $\text{WF}(u) \subset \{0\} \times \mathbf{R}^d$, and so it suffices to calculate $\Gamma_0(u)$. Fix a non-negative radial function $\phi \in \mathcal{D}(\mathbf{R}^d)$ such that $\hat{\phi}$ is a non-negative Schwartz function with total mass one. Let $\phi_t(x) = \phi(tx)$, let $\psi(\xi) = \hat{\phi}(\xi)$, and let $\psi_t(\xi) = \hat{\phi}_t(\xi) = t^{-d} \psi(\xi/t)$. Set $u_t = \psi_t u$. Then

$$\Gamma_0(u) = \lim_{t \rightarrow \infty} \Gamma(u_t).$$

If v is the homogeneous distribution given by the Fourier transform of u , and v_t is the Fourier transform of u_t , then

$$v_t = \widehat{\phi_t u} = \psi_t * v.$$

Since $\phi_t u$ is no longer homogeneous, neither are the distributions $\{v_t\}$. But they do satisfy the homogeneity relations

$$v_t(\xi) = (t/r)^s \text{Dil}_{t/r} v_r$$

for any $t, r > 0$. In particular, we see from this that $\Gamma(u_t) = \Gamma(u_r)$ for all $t, r > 0$. Thus $\Gamma_0(u) = \Gamma(u_t)$ for any $t > 0$. If v vanishes in a neighborhood of ξ_0 , then it vanishes in a conical neighborhood of ξ_0 . It thus follows that v_t vanishes in a conical neighborhood of ξ_0 if we pick t to be sufficiently small, and thus $\xi_0 \notin \Gamma(u_t) = \Gamma_0(u)$. Conversely, if $\xi_0 \notin \Gamma_0(u)$, then for all $t > 0$, there is an open cone $U_t \subset \mathbf{R}^d$ containing ξ_0 such that $|v_t(\eta)| \lesssim_{t,N} |\eta|^{-N}$ for all $\eta \in U_t$. The homogeneity relation above implies that we actually have

$$|v_t(\eta)| \lesssim_N t^{N+|s|} |\eta|^{-N}.$$

for all $\eta \in U_1$. As $t \rightarrow 0$, $|v_t(\eta)| \rightarrow |v(\eta)|$, and if N is taken larger than $|s|$ in the inequality above, this implies that

$$|v(\eta)| \lesssim_N \limsup_{t \rightarrow 0} t^{N+|s|} |\eta|^{-N} = 0.$$

Thus $v(\eta) = 0$ for all $\eta \in U_1$, so in particular, $\xi_0 \notin \text{supp}(v)$.

Here are some special cases of the above example:

- If δ is the Dirac delta distribution at the origin in \mathbf{R}^d , then $\widehat{\delta}$ is the constant function, and so

$$\text{WF}(\delta) = \{0\} \times \mathbf{R}^d.$$

- If $u(x) = p.v(1/x)$, then $\widehat{u}(\xi) = -i\pi \text{sgn}(\xi)$, and so

$$\text{WF}(u) = \{0\} \times \mathbf{R}.$$

- If $H \subset \mathbf{R}^d$ is a plane, and σ_H is the distribution given by integration against the surface measure on H , then σ_H is homogeneous, and we claim that $\widehat{\sigma}_H = \sigma_{H^\perp}$, from which the argument above implies that

$$\Gamma_0(\sigma_H) = H^\perp.$$

Since σ_H is invariant under translation by elements of H , it follows that for any $x \in H$, $\Gamma_x(\sigma_H) = H^\perp$. But σ_H is C^∞ away from H , so

$$\text{WF}(u) = H \times H^\perp.$$

Example. Let f_0 be a distribution of order $N+1$ defined on an open set $\Omega \subset \mathbf{R}^n$ by taking the boundary values of an analytic function f defined on the set $\Omega \times \Gamma \subset \mathbf{C}^n$, where Γ is a convex open cone in \mathbf{R}^n , and $|f(x+iy)| \lesssim |y|^{-N}$. Let $\Gamma^\circ = \{\xi \in \mathbf{R}^n : y \circ \xi \geq 0 \text{ for all } y \in \Gamma\}$ denote the dual cone of Γ . Then $\text{WF}(f_0) \subset \Omega \times (\Gamma^\circ - \{0\})$. To figure this out, we have that for any $v \geq N$,

$$\begin{aligned} \widehat{f_0 \phi}(\xi) &= \int \phi_{y_0}(x) f_{y_0}(x) e^{-2\pi i(x+iy_0) \cdot \xi} dx \\ &\quad + \frac{1}{v!} \int \int_0^1 f_{ty_0}(x) e^{-2\pi i(x+it y_0) \cdot \xi} \sum_{|\alpha|=v+1} \partial^\alpha \phi(x) (iy_0)^\alpha t^v dx dt. \end{aligned}$$

If $y_0 \cdot \xi < 0$, the first quantity decays exponentially as ξ scales, and the second term is $O((1+|\xi|)^{N-v})$. Taking $v \rightarrow \infty$ yields the claim.

The fact that $(x_0, \xi_0) \notin \text{WF}(u)$ implies precisely that there exists a neighbourhood U_0 of x_0 such that for any $\phi \in \mathcal{D}(U_0)$, and any $N > 0$,

$$\int_{\mathbf{R}^d} u(x) \phi(x) e^{-2\pi i \lambda \xi \cdot x} dx \lesssim_N \langle \xi \rangle^{-N}.$$

It will be useful to consider a nonlinear analogue of this statement, which will be useful for showing the invariance of the wavefront set under changes of variables.

Theorem 10.3. Let u be a distribution, and let $(x_0, \xi_0) \notin WF(u)$. Let U be an open subset of \mathbf{R}^d containing x_0 , let V be an open subset of \mathbf{R}^p containing a_0 , and let $\psi : U \times V \rightarrow \mathbf{R}$ be a C^∞ function with $\nabla_x \psi(x_0, a_0) = \xi_0$. Then there is an open set U_0 of x_0 , an open set V_0 of a_0 such that for any $\phi \in \mathcal{D}(U_0)$, and any $N > 0$,

$$\left| \int u(x) \phi(x) e^{-2\pi i \lambda \psi(x, a)} dx \right| \lesssim_N \lambda^{-N}$$

where the bound is uniform on V_0 .

Proof. Fix $\varepsilon > 0$, to be chosen later, and choose U_0 and V_0 such that $|\nabla_x \psi(x, a) - \xi_0| \leq \varepsilon/2$ for $x \in U_0$ and $a \in V_0$. For any given $\phi \in \mathcal{D}(U_0)$, consider $\tilde{\phi} \in \mathcal{D}(U_0)$ with $\tilde{\phi}\phi = \phi$. Then

$$\begin{aligned} \int u(x) \phi(x) e^{-2\pi i \lambda \psi(x, a)} dx &= \int u(x) \phi(x) \phi_1(x) e^{-2\pi i \lambda \psi(x, a)} dx \\ &= \int \widehat{u\tilde{\phi}}(\xi) \left(\int \phi_1(x) e^{-2\pi i (\lambda \psi(x, a) - \xi)} dx \right) d\xi \\ &= \lambda^d \int \widehat{u\tilde{\phi}}(\xi) \left(\int \phi_1(x) e^{-2\pi \lambda i (\psi(x, a) - \xi)} dx \right) d\xi \\ &= \lambda^d \int \widehat{u\tilde{\phi}}(\lambda \xi) J(\lambda, \xi, a) d\xi. \end{aligned}$$

Let $\eta \in \mathcal{D}(\mathbf{R}^d)$ be a smooth bump function supported on $|\xi| \leq 1$ and with $\eta(\xi) = 1$ for $|\xi| \leq 1/2$. Fix $\varepsilon > 0$, and write $J(\lambda, \xi, a) = J_1(\lambda, \xi, a) + J_2(\lambda, \xi, a)$, where

$$J_1(\lambda, \xi, a) = \eta\left(\frac{\xi - \xi_0}{\varepsilon}\right) \int_{\mathbf{R}^d} \phi_1(x) e^{-2\pi \lambda i (\psi(x, a) - \xi)} dx$$

and

$$J_2(\lambda, \xi, a) = \left(1 - \eta\left(\frac{\xi - \xi_0}{\varepsilon}\right)\right) \int_{\mathbf{R}^d} \phi_1(x) e^{-2\pi \lambda i (\psi(x, a) - \xi)} dx.$$

If ε is chosen appropriately small, then $|\widehat{u\tilde{\phi}}(\lambda \xi)| \lesssim_N \lambda^{-N}$ uniformly for $|\xi - \xi_0| \leq \varepsilon$. Since $|J_1(\lambda, \xi, a)| \lesssim 1$, this implies

$$\left| \int \widehat{u\tilde{\phi}}(\lambda \xi) J_1(\lambda, \xi, a) d\xi \right| \lesssim_N \lambda^{-N}.$$

On the other hand, if $|\xi - \xi_0| \geq \varepsilon$, then $|\nabla_x \phi(x, a) - \xi| = |\xi_0 - \xi| - \varepsilon/2 \geq \varepsilon/2$. Thus the method of stationary phase implies that

$$|J_2(\lambda, \xi, a)| \lesssim_N \lambda^{-N},$$

uniformly in a . Combined with the fact that $\widehat{u\phi}$ is of polynomial growth, this implies that

$$\left| \int \widehat{u\phi}(\lambda\xi) J_1(\lambda, \xi, a) d\xi \right| \lesssim_N \lambda^{-N}.$$

Combining these two estimates completes the proof. \square

For a smooth function $\phi \in \mathcal{D}(V)$ and a smooth diffeomorphism $f : U \rightarrow V$, we can define $f^*\phi \in \mathcal{D}(U)$ by setting $f^*\phi(x) = \phi(f(x))$. Then for $\psi \in \mathcal{D}(V)$,

$$\int f^*\phi(x)\psi(x) = \int \phi(f(x))\psi(x) = \int \phi(y)\psi(f^{-1}(y)) \cdot \frac{1}{|f'(f^{-1}(y))|} dy.$$

Thus for a distribution u on V , to define a distribution f^*u on U such that for $\psi \in \mathcal{D}(\mathbf{R}^d)$,

$$\int (f^*u)(x)\psi(x) = \int u(y)\phi(f^{-1}(y)) \cdot \frac{1}{|f'(f^{-1}(y))|} dy.$$

There is a simple relation between the wavefront set of u and f^*u . We consider $f^* : V \times \mathbf{R}^d \rightarrow U \times \mathbf{R}^d$ by defining $f^*((y, v)) = (f^{-1}(y), f'(y)^T v)$. This agrees with the definition of f^* encountered in differential geometry if we identify $V \times \mathbf{R}^d$ and $U \times \mathbf{R}^d$ with the cotangent bundle T^*V and T^*U .

Theorem 10.4. *For any distribution u on V , $\text{WF}(f^*u) = f^*(\text{WF}(U))$.*

Proof. Assume $(y_0, \eta_0) \notin \text{WF}(u)$, let $(x_0, \xi_0) = f^*((y_0, \eta_0))$, and then define $\psi(y, \xi) = f^{-1}(y) \cdot \xi$. Then

$$\nabla_y \psi(y_0, \xi_0) = (f^{-1}(y_0))'^T(\xi_0) = \eta_0.$$

Thus, applying the previous theorem, since

$$\widehat{f^*(u\phi)}(\lambda\xi) = \int u(y) \frac{\phi(f^{-1}(y))}{|f'(f^{-1}(y))|} e^{-2\pi i \lambda \xi \cdot f^{-1}(y)} = \int u(y) \tilde{\phi}(y) e^{-2\pi i \lambda \psi(y, \xi)} dy,$$

we conclude that $|\widehat{f^*(u\phi)}(\lambda\xi)| \lesssim_N \lambda^{-N}$, which implies $(x_0, \xi_0) \in \text{WF}(f^*(u))$. Thus $\text{WF}(f^*u) \subset f^*(\text{WF}(u))$. The converse statement that $f^*(\text{WF}(u)) \subset \text{WF}(f^*u)$ is obtained by symmetry. \square

Using this change of variables formula, we see that the wavefront set transforms under a change of coordinates like a covector. Since this gives an invariant definition, we can define the wavefront set of distributions on any smooth manifold M , and the wavefront set will then be a closed, conical subset of T^*M .

10.1 Oscillatory Integral Distributions

In this section, we consider distributions on an open subset U of \mathbf{R}^d , formally defined by the formula

$$I_{a,\phi}(x) = \int a(x, \theta) e^{2\pi i \phi(x, \theta)} d\theta.$$

Here a is a *symbol* lying in some class $\mathcal{S}^t(U \times \mathbf{R}^p)$, i.e. a smooth function satisfying bounds of the form

$$|\nabla_x^n \nabla_\theta^m a(x, \theta)| \lesssim_{n,m} \langle \theta \rangle^{t-m},$$

and $\phi \in C^\infty(U \times (\mathbf{R}^p - \{0\}))$ is homogeneous of degree one in θ , and $d\phi$ is nonvanishing on the support of a .

If $t < -d$, then the integrand formally defining $I_{a,\phi}$ is absolutely integrable, and interpreting $I_{a,\phi}$ as a Lebesgue integral gives us a locally integrable function $I_{a,\phi}$. But for $t \geq -d$, the integral defining $I_{a,\phi}$ need not be locally integrable; for instance, our definition will show that the distribution

$$\int_{\mathbf{R}^d} \xi^t e^{2\pi i x \cdot \xi} d\xi$$

acts on functions as a constant multiple of the differential operator D^t .

To define the oscillatory integral distribution rigorously, we fix $\psi \in \mathcal{D}(\mathbf{R}^d)$, and $\rho \in \mathcal{D}(\mathbf{R}^p)$, equal to one in a neighborhood of the origin. The integral

$$\int a(x, \theta) \psi(x) \rho(\theta/R) e^{2\pi i \phi(x, \theta)} d\theta$$

is then well defined, and we claim that the limit

$$\lim_{R \rightarrow \infty} \int a(x, \theta) \psi(x) \rho(\theta/R) e^{2\pi i \phi(x, \theta)} d\theta$$

exists and is independent of the choice of ρ . We can then define this limit to be

$$\int I_{a,\phi}(x) \psi(x) dx$$

and this defines $I_{a,\phi}$ as a distribution. To prove the limit exists, we fix $R_1 \leq R_2$, and let $\tilde{\rho}(\theta) = \rho(\theta/R_2) - \rho(\theta/R_1)$. Then $\tilde{\rho}$ is supported on $R_1 \lesssim |x| \lesssim R_2$. Assume first that $R_2 \leq 2R_1$. Rescaling, we find that if $\eta(x, \theta) = a(x, R_2\theta) \psi(x) \rho(\theta)$, then

$$\begin{aligned} \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i \phi(x, \theta)} a(x, \theta) \psi(x) \tilde{\rho}(\theta) &= R_2^m \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i R_2 \phi(x, \theta)} a(x, R_2\theta) \psi(x) \tilde{\rho}(\theta) \\ &= R_2^p \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i R_2 \phi(x, \theta)} \eta(x, \theta). \end{aligned}$$

Then η is supported on $1/2 \lesssim |\theta| \lesssim 1$ and $|x| \lesssim 1$. Thus the support of η is independent of R_1 and R_2 . It is simple to verify that

$$|\nabla_x^n \nabla_\theta^m \eta(x, \theta)| \lesssim_{n,m} R_2^t \cdot |\nabla_x^n \psi(x)|,$$

where the bound is independent of R_1 and R_2 . Since $\nabla_x \phi$ and $\nabla_\theta \phi$ have no common zeroes on the support of a , we can apply the principle of stationary phase to conclude that

$$\left| R_2^p \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i R_2 \phi(x, \theta)} \eta(x, \theta) \right| \lesssim_N R_2^{p+m-N} \cdot \|\nabla^{\leq N} \psi\|_{L^\infty(\mathbf{R}^d)}.$$

In general, if $R_2 \geq 2R_1$, we consider the largest l such that $2^l R_1 \leq R_2$. If we set $a_k = 2^k R_1$ for $0 \leq k \leq l$, and $a_{l+1} = R_2$, then we write

$$\begin{aligned} &\left| \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i \phi(x, \theta)} a(x, \theta) \phi(x) \tilde{\rho}(\theta) \right| \\ &= \left| \sum_{k=0}^l \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i \phi(x, \theta)} a(x, \theta) \phi(x) (\rho(\theta/a_{k+1}) - \rho(\theta/a_k)) \right| \\ &\lesssim \sum_{k=0}^l a_{k+1}^{p+m-N} \|\nabla^{\leq N} \psi\|_{L^\infty(\mathbf{R}^d)}. \end{aligned}$$

If we choose $N > p + m$, then we conclude that

$$\begin{aligned} \left| \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i \phi(x, \theta)} a(x, \theta) \phi(x) \tilde{\rho}(\theta) \right| &\lesssim (R_1^{p+m-N} + R_2^{p+m-N}) \|\nabla_x^{\leq N} \psi\|_{L^\infty(\mathbf{R}^d)} \\ &\lesssim R_1^{p+t-N} \|\nabla^{\leq N} \psi\|_{L^\infty(\mathbf{R}^d)}. \end{aligned}$$

In particular, this quantity tends to zero as $R_1 \rightarrow \infty$, which gives convergence of the limit, and also gives boundedness, showing $I_{a,\phi}$ is a distribution of order $\leq N$, where N is the smallest integer bigger than $p + m$. A very similar argument shows that if $\rho \in \mathcal{D}(\mathbf{R}^p)$ is equal to zero in a neighborhood of the origin, then

$$\lim_{R \rightarrow \infty} \int_{\mathbf{R}^n} \int_{\mathbf{R}^p} e^{2\pi i \phi(x,\theta)} a(x,\theta) \psi(x) \rho(\theta/R) dx = 0.$$

It follows from the above observation that the definition is independent of the original choice of ρ . It is left as an exercise to show that the map $a \mapsto I_{a,\phi}$ is continuous from $S^t(U \times \mathbf{R}^p)$ to $\mathcal{D}^*(U)$.

Remark. If M is a manifold, and E is a vector bundle over M , then for any homogeneous phase $\phi \in C^\infty(E - \{0\})$ and any symbol $a \in S^t(E^*)$, we can consider the oscillatory integral distribution formally defined by the integral

$$I_{a,\phi}(x) = \int_{E_x} a(x,\theta) e^{2\pi i \phi(x,\theta)} d\theta,$$

which is a distribution on M .

Let us now consider the wavefront set of $I_{a,\phi}$. If ψ is a bump function supported in a neighbourhood of some point x_0 , then rescaling gives

$$\widehat{I_{a,\phi}\psi}(\lambda\xi_0) = \lambda^d \int \int e^{2\pi i \lambda(\phi(x,\theta) - x \cdot \xi_0)} a(x, \lambda\theta) \psi(x) dx d\theta.$$

This is an oscillatory integral, and the phase is non-stationary in the x and θ variables provided that the support of ψ is small enough, and $\nabla_\theta \phi(x_0, \theta_0) \neq 0$, or if $\nabla_\theta \phi(x_0, \theta_0) = 0$, but $\nabla_x \phi(x_0, \theta_0) \neq \xi_0$, provided that ψ is supported on a small enough neighborhood of x_0 . Thus we are lead to conclude that

$$\text{WF}(I_{a,\phi}) \subset \{(x_0, \nabla_x \phi(x_0, \theta_0)) : (x_0, \theta_0) \in \mu\text{-supp}(a) \text{ and } \nabla_\theta \phi(x_0, \theta_0) = 0\}.$$

Here $\mu\text{-supp}(a)$ is the *microsupport* of a , the complement of the set of pairs (x_0, θ_0) which have a conical neighborhood upon which a lies in $S^{-\infty}$.

Theorem 10.5. *If $a \in S^t(U \times \mathbf{R}^p)$, and $\phi \in C^\infty(U \times \mathbf{R}^p)$ is a phase satisfying the standard conditions to define the oscillatory integral distribution $I_{a,\phi}$, then*

$$\text{WF}(I_{a,\phi}) \subset \{(x_0, \nabla_x \phi(x_0, \theta_0)) : (x_0, \theta_0) \in \mu\text{-supp}(a) \text{ and } \nabla_\theta \phi(x_0, \theta_0) = 0\}$$

Proof. Fix $(x_0, \xi_0) \in T^*U$, and suppose that for all $(x_0, \theta_0) \in U \times \mathbf{R}^p$, either $\nabla_\theta \phi(x_0, \theta_0) \neq 0$, $\nabla_x \phi(x_0, \theta_0) \neq \xi_0$, or $(x_0, \theta_0) \notin \mu\text{-supp}(a)$. Without loss of generality, we can assume that this third condition never holds, by decomposing a as the sum of a symbol in $\mathcal{S}^{-\infty}$ and a symbol where a vanishes in a neighborhood of any point (x_0, θ_0) with $\nabla_\theta \phi(x_0, \theta_0) = 0$ and $\nabla_x \phi(x_0, \theta_0) = \xi_0$, and if $a \in \mathcal{S}^{-\infty}$, then $I_{a,\phi} \in C^\infty(U)$. Write

$$\widehat{I_{a,\phi}\psi}(\lambda\xi_0) = \lambda^d \iint e^{2\pi i \lambda \tilde{\phi}(x,\theta;\xi_0)} \tilde{a}_\lambda(x,\theta) dx d\theta$$

where $\tilde{a}_\lambda(x,\theta) = a(x,\lambda\theta)\psi(x)$, and write $\tilde{\phi}(x,\theta;\xi) = \phi(x,\theta) - x \cdot \xi$. If Z is the set of all $\theta \in \mathbf{R}^p$ with $|\theta| = 1$ such that $\nabla_\theta \phi(x_0, \theta) = 0$, then Z is closed, and thus compact. Since $\nabla_x \phi(x_0, \theta) \neq \xi_0$ for all $\theta \in Z$, it follows by compactness that

$$|\nabla_x \phi(x_0, \theta) - t\xi_0| \gtrsim 1$$

for all $\theta \in Z$ and all $t > 0$. By homogeneity, for any $\theta \in \mathbf{R}^p - \{0\}$ such that $\nabla_\theta \phi(x_0, \theta) = 0$,

$$|\nabla_x \phi(x_0, \theta) - \xi_0| \gtrsim |\theta|.$$

By reducing the implicit constant slightly, we may assume that there is $\varepsilon > 0$ such that for any $x \in \mathbf{R}^d$ and $\xi \in \mathbf{R}^d$ with $|x - x_0| \leq \varepsilon$ and $|\xi - \xi_0| \leq \varepsilon$, and any $\theta \in \mathbf{R}^p - \{0\}$ such that $|\nabla_\theta \phi(x_0, \theta)| \leq \varepsilon$, then

$$|\nabla_x \phi(x, \theta) - \xi| \gtrsim |\theta|.$$

Now if ψ has support in a ε neighborhood of x_0 , it follows that for all $x \in \text{supp}(\psi)$ and any $\theta \in \mathbf{R}^p$,

$$|\nabla_{x,\theta} \tilde{\phi}(x, \theta, \xi)| \gtrsim 1.$$

The principle of nonstationary phase thus guarantees that for any $N > 0$,

$$|\widehat{I_{a,\phi}\psi}(\lambda\xi_0)| \lesssim_{\phi,N} \lambda^{-N} \sup_{|\alpha| \leq N} \|D_{x,\theta}^\alpha \tilde{a}_\lambda(x, \theta) |\nabla_{x,\theta} \tilde{\phi}(x, \theta; \xi)|^{|\alpha| - 2N}\|_{L^\infty(U \times \mathbf{R}^p)}.$$

Now

$$|\nabla_{x,\theta} \tilde{\phi}(x, \theta; \xi)| = |\nabla_x \phi(x, \theta) - \xi| \lesssim 1,$$

and if $D_{x,\theta}^\alpha = D_x^{\alpha_1} D_\theta^{\alpha_2}$, then $D_{x,\theta}^\alpha \tilde{a}_\lambda(x, \theta)$ is a finite sum of $O(\alpha)$ terms, each of the form

$$\lambda^{|\alpha_2|} D_x^{\beta_1} D_\theta^{\alpha_2} a(x, \lambda\theta) D_x^{\beta_2} \psi(x),$$

and

$$|\lambda^{|\alpha_2|} D_x^{\beta_1} D_\theta^{\alpha_2} a(x, \lambda\theta) D_x^{\beta_2} \psi(x)| \lesssim_{\beta_1, \beta_2} \lambda^{|\alpha_2|} \langle \lambda \rangle^{t-|\alpha_2|} \lesssim \langle \lambda \rangle^t$$

Thus we conclude that

$$|\widehat{I_{a,\phi}\psi}(\lambda\xi)| \lesssim_{\phi,\psi,N} \langle \lambda \rangle^{t-N}.$$

Thus $\xi_0 \notin \text{WF}(I_{a,\phi})$. □

Example. If we set $\phi(x, \xi) = -2\pi x \cdot \xi$, and $a(x, \xi) = a(\xi)$ is a symbol depending only on the ξ variable, then

$$I_{a,\phi}(x) = \int a(\xi) e^{-2\pi i x \cdot \xi} d\xi$$

is the Fourier transform of a . The result we have just calculated shows that for any symbol a ,

$$\text{WF}(I_{a,\phi}) \subset \{0\} \times \mathbf{R}^n.$$

Thus the Fourier transform of any symbol is smooth away from the origin. This result should be compared to the standard result that the Fourier transform of a homogeneous distribution which is C^∞ away from the origin is also homogeneous and C^∞ away from the origin.

Near the wavefront set of an oscillatory integral distribution, we can compute an asymptotic formula which characterizes the behaviour of the distribution near the wavefront set up to integration against a function in $C^\infty(U)$.

Theorem 10.6. Consider a phase function ϕ_1 . Fix $(x_0, \theta_0) \in U \times \mathbf{R}^p$ such that $\nabla_\theta \phi_1(x_0, \theta_0) = 0$. Let $\xi_0 = \nabla_x \phi_1(x_0, \theta_0)$. Consider any phase function $\phi_2 \in C^\infty(U \times \mathbf{R}^q)$ and $\sigma_0 \in \mathbf{R}^q$ with

$$\nabla_x \phi_1(x_0, \theta_0) = \nabla_x \phi_2(x_0, \sigma_0).$$

Furthermore, assume that the Hessian $H_{x,\theta}(\phi - \psi)$ is nondegenerate at $(x_0, \theta_0, \sigma_0)$. Then there exists a conical neighborhood Γ of (x_0, θ_0) , an open neighborhood V of x_0 , and an open neighborhood Σ of σ_0 , such that if $\psi \in \mathcal{D}(V)$, and a is a symbol on $U \times \mathbf{R}^p$ with $\text{ess sup}(a) \subset \Gamma$, then there exists a family of smooth functions a_k such that as $\lambda \rightarrow \infty$,

$$\begin{aligned} & \int I_{a,\phi_1}(x) \psi(x) e^{-2\pi i \lambda \phi_2(x,\sigma)} dx \\ & \sim e^{-2\pi i \lambda \phi_2(x(\sigma), \sigma)} |\det Q(\sigma)|^{-1/2} e^{(i\pi/4) \text{sgn}(Q(\sigma))} \lambda^{(p-d)/2} \sum_{k=0}^{\infty} a_k(\sigma, \lambda) \cdot \lambda^{-k}, \end{aligned}$$

Here $a_k(\sigma, \lambda)$ is a linear differential operator in a and u at $(x(\sigma), \theta(\sigma), \sigma)$

The phase ϕ of an oscillatory integral distribution is called *nondegenerate* if whenever $\nabla_\theta \phi(x, \theta) = 0$, the matrix $D_{x, \theta}(\nabla_\theta \phi)(x, \theta)$ has full rank p . It follows that

$$\Sigma_\phi = \{(x, \theta) : \nabla_\theta \phi(x, \theta) = 0\}$$

is a d dimensional submanifold of $U \times \mathbf{R}^p$. Moreover, the map f from Σ_ϕ to $U \times \mathbf{R}^d$ given by $(x, \theta) \mapsto (x, \nabla_x \phi(x, \theta))$ is an immersion, the immersed submanifold in the image being denoted by $\Lambda_\phi \subset T^*U$. To verify the map is an immersion, we note that at a point (x, θ) the tangent space of Σ_ϕ consists of vectors $(v, w) \in \mathbf{R}^d \times \mathbf{R}^p$ such that

$$D_x \nabla_\theta \phi(x, \theta) \cdot v + D_\theta \nabla_\theta \phi(x, \theta) \cdot w = 0.$$

Now

$$D_{x, \theta} f(x, \theta)(v, w) = (v, D_x \nabla_x \phi(x, \theta) \cdot v + D_\theta \nabla_x \phi(x, \theta) \cdot w).$$

Thus if (v, w) lies in the tangent space and $Df(x, \theta)(v, w) = 0$, then $v = 0$, which implies

$$D_\theta \nabla_\theta \phi(x, \theta) \cdot w = D_\theta \nabla_x \phi(x, \theta) \cdot w = 0.$$

Since mixed partials commute, this says exactly that $D(\nabla_\theta \phi)^T \cdot w = 0$. The full rank condition thus implies that $w = 0$. Thus $(v, w) = 0$, completing the argument that f is an immersion.

TODO: If $I_{a_1, \phi_1} = I_{a_2, \phi_2}$, prove that $\Lambda_{\phi_1} = \Lambda_{\phi_2}$.

The converse is also true.

Theorem 10.7. Suppose ϕ_1 and ϕ_2 are nondegenerate phase functions on $U \times \mathbf{R}^{p_1}$ and $U \times \mathbf{R}^{p_2}$. Let

The manifold Λ_ϕ of $U \times \mathbf{R}^d$ actually has special geometric structure. Consider the two form

$$\sigma = d\xi^1 \wedge dx^1 + \cdots + d\xi^d \wedge dx^d.$$

The $\sigma = d\omega$, where $\omega = \xi^1 dx^1 + \cdots + \xi^d dx^d$. We claim that for any $p = (x, \theta) \in \Sigma_\phi$, and any $v, w \in T_p \Sigma_\phi$, $\sigma(f_* v, f_* w) = 0$. To see this, we calculate that

$$f^*(\sigma) = f^*(d\omega) = d(f^* \omega),$$

and

$$f^*\omega = \nabla_x \phi \cdot dx = d\phi - \nabla_\theta \phi \cdot d\theta.$$

On Σ_ϕ , $\nabla_\theta \phi = 0$, so $f^*\omega = d\phi$, and so $f^*(\sigma) = d(f^*\omega) = d^2\phi = 0$. Thus Λ_ϕ is a *Lagrangian submanifold* of $T^*\mathbf{R}^d$ with respect to the two form σ .

For any phase function ϕ (possibly degenerate), we can define

$$\Sigma_\phi = \{(x, \theta) : \nabla_\theta \phi(x, \theta) = 0\}$$

$$\Lambda_\phi = \{(x, \nabla_x \phi(x, \theta)) : \nabla_\theta \phi(x, \theta) = 0\}.$$

If Λ_ϕ is an immersed Lagrangian manifold (though not necessarily an immersion through the map $f : \Sigma_\phi \rightarrow \Lambda_\phi$), we say $I_{a,\phi}$ is a *Lagrangian distribution*. The phase ϕ might be degenerate in this case.

Example. A degenerate example of a Lagrangian distribution is given for $p = d + 1$, $\theta = (\theta_0, \theta_1)$ with $\theta_0 \in \mathbf{R}^d$ and $\theta_1 \in \mathbf{R}$, and

$$\phi(x, \theta) = x \cdot \theta_0,$$

then ϕ defines a Lagrangian distribution $I_{a,\phi}$ for any symbol a , provided that $a(x, \theta)$ vanishes for any θ in a cone containing the θ_1 axis. Now $\Sigma_\phi = \{0\} \times \mathbf{R}^{d+1}$, and $\Lambda_\phi = \{0\} \times \mathbf{R}^d$, which is a Lagrangian manifold. Thus the distributions

$$I_{a,\phi}(x) = \int a(x, \theta_0, \theta_1) e^{2\pi i x \cdot \theta_0} d\theta_0 d\theta_1$$

are Lagrangian.

If $f : U \rightarrow V$ is a diffeomorphism between open subsets of \mathbf{R}^d , and we equip T^*U with coordinates (x, ξ) , and T^*V with coordinates (y, η) , then we obtain an isomorphism $g : T^*U \rightarrow T^*V$ mapping (x, ξ) in T^*U to $(y, (Df(x)^T)^{-1}\xi)$ in T^*V . Under this correspondence, if we consider the two-form $\omega_V = \sum \eta_i \wedge dy^i$ on V , then

$$g^*\omega_V = \sum (\eta^i \circ g) \cdot d(y^i \circ g) = \sum ((Df(x)^T)^{-1}\xi)_i df^i(x) = \sum \xi_i dx^i.$$

Thus the Lagrangian form is invariant under coordinate changes, and can thus be well defined on the cotangent bundle of any manifold M . Thus we can discuss Lagrangian submanifolds of T^*M for any manifold M , and Lagrangian distributions on manifolds.

Example. Consider a one-form ψ on M , i.e. a smooth function $\psi : M \rightarrow T^*M$. Working in coordinates (x, ξ) on T^*M , we have

$$\psi^*\omega = \sum \psi^i dx^i = d\psi.$$

Thus we see that $\psi^*\sigma = 0$ if and only if $d\psi = 0$, so ψ defines a Lagrangian submanifold of T^*M if and only if it is closed.

10.2 Singular Operations on Distributions

A subset Γ of $\Omega \times \mathbf{R}^d$ is *conic* if $(x, \xi) \in \Gamma$ implies that $(x, \lambda\xi) \in \Gamma$. Given a closed conic set Γ , let $\mathcal{D}_\Gamma^*(\Omega)$ denote the family of all distributions u with $\text{WF}(u) \subset \Gamma$, with the seminorms

$$u \mapsto \sup_{\substack{\xi \in V \\ \lambda > 0}} \lambda^N |\widehat{\phi u}(\lambda\xi)|$$

where V is a closed conic set disjoint from Γ . Then $\mathcal{D}_\Gamma^*(\Omega)$ is a Fréchet space. To analyze this space, we require a lemma about general distributions.

Lemma 10.8. *Let \mathcal{U} be a family in $\mathcal{D}^*(\Omega)$ such that $\sup_{u \in \mathcal{U}} |u(\phi)| < \infty$ for all $\phi \in \mathcal{D}(\Omega)$. Then for any $\phi \in \mathcal{D}(\Omega)$, there exists $m > 0$ such that for any $\xi \in \mathbf{R}^d$,*

$$|\widehat{\phi u}(\xi)| \lesssim (1 + |\xi|)^m,$$

uniformly in u and ξ . If we have a sequence $\{u_n\}$ converging to some distribution u , then as $n \rightarrow \infty$,

$$|\widehat{\phi u_n}(\xi) - \widehat{\phi u}(\xi)| = o((1 + |\xi|^m)).$$

In particular, $\widehat{\phi u_n}$ converges to $\widehat{\phi u}$ on compact subsets.

Proof. For any compact set $K \subset \Omega$, the assumptions imply there exists m such that for $u \in \mathcal{U}$ and $\phi \in C_c^\infty(K)$,

$$|u(\phi)| \lesssim \|\phi\|_{C^m(K)}.$$

This is a result we have proven earlier following from the uniform boundedness principle. If ϕ is supported on K , then

$$|u(\phi e^{-2\pi i \xi \cdot x})| \lesssim \|\phi e^{-2\pi i \xi \cdot x}\|_{C^m(K)} \lesssim (1 + |\xi|)^m.$$

A similar argument shows the result for a convergent sequence. □

Theorem 10.9. *A sequence of distributions u_n converges to u in $\mathcal{D}_\Gamma^*(U)$ if and only if u_n converges to u distributionally, and for any conic set V disjoint from Γ , and $N > 0$,*

$$\sup_{\xi \in V} \lambda^N |\widehat{\phi u_n}(\lambda \xi)|$$

is bounded independantly of n .

Proof. This conditions are certainly necessary for convergence. Conversely, if these conditions are satisfied, the previous lemma implies that as $n \rightarrow \infty$,

$$\sup_{\lambda \geq 1} \sup_{\xi \in V} \lambda^{-m} |\widehat{\phi u_n}(\lambda \xi) - \widehat{\phi u}(\lambda \xi)| = o(1).$$

We also know that

$$\sup_{\lambda \geq 1} \sup_{\xi \in V} \lambda^{N+1} |\widehat{\phi u_n}(\lambda \xi) - \widehat{\phi u}(\lambda \xi)| < \infty.$$

Let us call this supremum $C > 0$. Given any $\varepsilon > 0$, we find that

$$\sup_{\lambda \geq C/\varepsilon} \lambda^N |\widehat{\phi u_n}(\lambda \xi) - \widehat{\phi u}(\lambda \xi)| \leq \varepsilon.$$

But

$$\sup_{1 \leq \lambda \leq C/\varepsilon} \sup_{\xi \in V} \lambda^N |\widehat{\phi u_n}(\lambda \xi) - \widehat{\phi u}(\lambda \xi)| = o((C/\varepsilon)^{N+m}).$$

Taking n suitably large, depending on C , ε , N , and m , we conclude that

$$\sup_{1 \leq \lambda \leq C/\varepsilon} \sup_{\xi \in V} \lambda^N |\widehat{\phi u_n}(\lambda \xi) - \widehat{\phi u}(\lambda \xi)| \leq \varepsilon.$$

Combining this with the supremum above shows that we have convergence in $\mathcal{D}_\Gamma^*(U)$. \square

Theorem 10.10. *$\mathcal{D}(\Omega)$ is sequentially dense in $\mathcal{D}_\Gamma^*(\Omega)$.*

Proof. Consider a distribution $u \in \mathcal{D}_\Gamma^*(\Omega)$. Without loss of generality we may assume u is compactly supported. Consider an approximation to the identity $\{\phi_\delta\}$. Then $u * \phi_{1/n} \in C^\infty(\Omega)$, and thus an element of $\mathcal{D}_\Gamma^*(\Omega)$. $u * \phi_{1/n}$ converges to u in $\mathcal{D}^*(\Omega)$, so by the last result, it suffices to show that for any admissible choice of ψ , V , and $N > 0$,

$$\sup_{\xi \in V} \sup_n \lambda^N |\widehat{\psi(u * \phi_n)}(\lambda \xi)| < \infty.$$

But this is simple, for we get arbitrarily fast decay if n is suitably large, depending on the distance from V to Γ , and the finitely many smaller choices of n are negligible. \square

We have a continuous map $(\phi, \psi) \rightarrow \phi\psi$ from $\mathcal{D}(\Omega) \times \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)$, which extends to a continuous map $(\phi, u) \rightarrow \phi u$ from $\mathcal{D}(\Omega) \times \mathcal{D}^*(\Omega) \rightarrow \mathcal{D}^*(\Omega)$. However, it is *not* possible to extend this to a continuous map $(u, v) \mapsto uv$ from $\mathcal{D}^*(\Omega) \times \mathcal{D}^*(\Omega) \rightarrow \mathcal{D}^*(\Omega)$. For instance, if $\{\phi_\varepsilon\}$ is an approximation to the identity, then ϕ_ε converges to the Dirac delta distribution δ at the origin, so we would expect ϕ_ε^2 to converge to a distribution representing the product $\delta \cdot \delta$, but this does not happen because if $\psi \in \mathcal{D}(\mathbf{R}^d)$ and $\psi(x) = 1$ for $|x| \leq 1$, then

$$\left| \int \phi_\varepsilon^2(x) \psi(x) dx \right| \gtrsim 1/\varepsilon$$

and thus does not converge. It is a surprising fact that we can use the wavefront set of a distribution to define the product of two distributions, *provided that the wavefront sets satisfy a disjointness relation*.

To see how this is possible, we note that for $\phi, \psi \in \mathcal{D}(\mathbf{R}^d)$, we might expect us to be able to take Fourier transforms, so that

$$\begin{aligned} \int u(x)v(x)\phi(x)\psi(x) dx &= \int (\phi u)(x)(\psi v)(x) dx \\ &= \int \int (\widehat{\phi u} * \widehat{\psi v})(\xi) e^{2\pi i \xi \cdot x} d\xi dx \\ &= \int \int \widehat{\phi u}(\eta) \widehat{\psi v}(\xi - \eta) e^{2\pi i \xi \cdot x} d\xi d\eta. \end{aligned}$$

The only problem with taking this as the *definition* of the product is that the integral we have obtained might not converge in general. However, if at least one of the Fourier transforms decreases rapidly in the right directions.

Theorem 10.11. *Fix conic sets $\Gamma_1, \Gamma_2 \subset \Omega \times \mathbf{R}^d - \{0\}$. If $\Gamma_3 = \Gamma_1 + \Gamma_2$ does not contain any points in $0_\Omega = \Omega \times \{0\}$, then we have a unique continuous map from $\mathcal{D}_{\Gamma_1}^*(\Omega) \times \mathcal{D}_{\Gamma_2}^*(\Omega) \rightarrow \mathcal{D}_{\Gamma_3}^*(\Omega)$ which agrees with multiplication for elements of $C^\infty(\Omega)$.*

Example. Consider the distributions Λ_1 and Λ_2 on \mathbf{R}^2 , given by integration along the x and y axis respectively, i.e.

$$\int \Lambda_1(x, y) \phi(x, y) dx dy = \int \phi(x, 0) dx$$

and

$$\int \Lambda_2(x, y) \phi(x, y) dx dy = \int \phi(0, y) dy.$$

We have seen that $WF(\Lambda_1) = \{(x, 0; 0, \eta) : \eta \neq 0\}$ and $WF(\Lambda_2) = \{(0, \xi; y, 0) : \xi \neq 0\}$. Now

$$WF(\Lambda_1) + WF(\Lambda_2) = \{(x, \xi, y, \eta) : \xi, \eta \neq 0\},$$

which is disjoint from $0_{\mathbf{R}^2}$, and so a product $\Lambda_1 \cdot \Lambda_2$ is well defined. To determine what the product is, we consider a non-negative bump function $\phi \in \mathcal{D}(\mathbf{R}^d)$ equal to one in a neighborhood of the origin, and define

$$\phi_{x,\delta}(x, y) = (1/2\delta) \mathbf{I}(|x| \leq 1/\delta, |y| \leq \delta)$$

$$\phi_{y,\delta}(x, y) = (1/2\delta) \mathbf{I}(|x| \leq \delta, |y| \leq 1/\delta).$$

Then as $\delta \rightarrow 0$, $\phi_{x,\delta} \rightarrow \Lambda_1$ and $\phi_{y,\delta} \rightarrow \Lambda_2$. We find that

$$\phi_{x,\delta} \phi_{y,\delta} = (1/4\delta^2) \mathbf{I}(|x| \leq \delta, |y| \leq \delta).$$

As $\delta \rightarrow 0$, $\phi_{x,\delta} \phi_{y,\delta}$ thus converges to the Dirac delta distribution δ at the origin. Thus $\Lambda_1 \cdot \Lambda_2 = \delta$.

Example. Let $\Lambda = (x + i0^+)^{-1}$, i.e. the distribution

$$\int \Lambda(x) \phi(x) dx = \lim_{y \rightarrow 0^+} \int \frac{\phi(x)}{x + iy} dx = \lim_{y \rightarrow 0^+} \Lambda_y(\phi).$$

The Λ is homogeneous. Moreover, some formal manipulations, plus some contour integrals, show that

$$\hat{\Lambda}(\xi) = -2\pi i \cdot \mathbf{I}(\xi < 0).$$

In particular, $WF(\Lambda) = \{(0, \xi) : \xi < 0\}$. this means $WF(\Lambda) + WF(\Lambda)$ does not contain any zero vectors, so the product $\Lambda \cdot \Lambda$ is well defined. Now Λ is

the limit of the C^∞ functions $\phi_y(x) = 1/(x + iy)$ in $\mathcal{D}^*(\mathbf{R})$, and it requires only a simple calculation to show that Λ is also the limit in $\mathcal{D}_\Gamma^*(\mathbf{R})$, where $\Gamma = \{(0, \xi) : \xi < 0\}$. Since

$$\phi_y(x)^2 = 1/(x + iy)^2,$$

we find by continuity that

$$\Lambda \cdot \Lambda = (x + i0^+)^{-2},$$

i.e.

$$\int \Lambda(x) \Lambda(x) \phi(x) dx = \lim_{y \rightarrow 0} \int \frac{\phi(x)}{(x + iy)^2} dx.$$

To define more sophisticated operations on distributions, we define the generic operations of *pullback*, *pushforward*, and *tensoring*. Intuitively, the pullback of a distribution gives a way to ‘compose’ a distribution with a smooth function in the domain, the push forward enables one to ‘integrate a distribution along fibres’, and tensoring enables us to take the product of distributions.

Let us begin with the pullback. For a smooth map $f : \Omega \rightarrow \Psi$, not necessarily a diffeomorphism, and $\phi \in \mathcal{D}(\Psi)$, we can define $f^*\phi = \phi \circ f \in C^\infty(\Omega)$. This map is continuous in the appropriate topology, and if f is a proper map, f^* is continuous from $\mathcal{D}(\Psi) \rightarrow \mathcal{D}(\Omega)$. To obtain a distributional definition, we apply the Fourier inversion formula; if $\psi \in \mathcal{D}(\mathbf{R}^d)$, then

$$\int (f^*\phi)(x) \psi(x) dx = \int \phi(f(x)) \psi(x) dx = \int \int \hat{\phi}(\eta) \psi(x) e^{2\pi i \eta \cdot f(x)} d\eta dx.$$

For a compactly supported distribution u on Ψ , it is therefore natural to define f^*u on Ω such that

$$\int (f^*u)(x) \psi(x) dx = \int \hat{u}(\eta) \left(\int \psi(x) e^{2\pi i \eta \cdot f(x)} dx \right) d\eta.$$

We can decompose this integral so that ψ is supported on various small sets. If ψ is supported in a neighbourhood of x_0 , then the oscillatory integral on the inside decays fast as $\eta \rightarrow \infty$ provided that $Df(x_0)^T \eta \neq 0$. Thus, provided that $\hat{u}(\eta)$ decays fast whenever $Df(x_0)^T \eta = 0$, the integral above is well defined. Proceeding through this argument more rigorously gives the following result, left as an exercise.

Theorem 10.12. *Given a smooth map $f : \Omega \rightarrow \Psi$, let*

$$N = \{(f(x), \eta) : Df(x)^T \eta = 0\}.$$

Fix a closed cone Γ with $\Gamma \cap N = \emptyset$. Then $f^ : \mathcal{D}(\Psi) \rightarrow C^\infty(\Omega)$ extends to a continuous map from $\mathcal{D}_\Gamma^*(\Psi) \rightarrow \mathcal{D}_{f^*\Gamma}^*(\Omega)$, where*

$$f^*\Gamma = \{(x, Df(x)^T \xi) : (f(x), \xi) \in \Gamma\}.$$

Example. *Given a smooth map $f : \Omega \rightarrow \mathbf{R}^d$, with the property that for any $x \in f^{-1}(0)$, $Df(x)$ has rank d , the set N above is disjoint from $\{0\} \times \mathbf{R}^d$, and so we can define the pullback of the Dirac delta function at the origin, $f^*\delta$, also denoted $\delta(f(x))$.*

As examples of this construction, we can consider the distributions $\delta(x)$ and $\delta(y)$ on \mathbf{R}^2 . This are equal to π_x^δ and $\pi_y^*\delta$, where $\pi_x : \mathbf{R}^2 \rightarrow \mathbf{R}$ and $\pi_y : \mathbf{R}^2 \rightarrow \mathbf{R}$ are the obvious projection maps, then we have*

$$\begin{aligned} \int (\pi_x^*\delta)(x, y) \phi(x, y) dx dy &= \int \hat{\delta}(\xi) \phi(x, y) e^{2\pi i \xi \cdot x} dx dy d\xi \\ &= \int \phi(x, y) e^{2\pi i \xi \cdot x} dx dy d\xi \\ &= \int \phi(0, y) dy. \end{aligned}$$

Thus $\delta(x) = \pi_x^\delta$ is the distribution given by integration on the y -axis. Similarly, one can calculate that $\delta(y) = \pi_y^*\delta$ is the distribution given by integration on the x -axis. It is simple to calculate explicitly, or using the properties of pullback, that*

$$WF(\pi_x^*\delta) \subset \{(0, y; \xi, 0) : \xi \neq 0\}$$

and

$$WF(\pi_y^*\delta) \subset \{(x, 0; 0, \eta) : \eta \neq 0\}.$$

In fact, in these two cases these equations are equalities.

Next, let us define the tensor product. Given a distribution u_1 on Ω_1 and a distribution u_2 on Ω_2 , we define a distribution $u_1 \otimes u_2$ on $\Omega_1 \times \Omega_2$ such that for $\phi \in \mathcal{D}(\Omega_1 \times \Omega_2)$,

$$\int (u_1 \otimes u_2)(x_1, x_2) \phi(x_1, x_2) dx_1 dx_2 = \int u_1(x_1) \left(\int u_2(x_2) \phi(x_1, x_2) dx_2 \right) dx_1,$$

where the function

$$\tilde{\phi}(x_1) = \int u_2(x_2) \phi(x_1, x_2) dx_2$$

is smooth, where one can easily verify that

$$D^\alpha \tilde{\phi}(x_1) = \int u_2(x_2) D^\alpha \phi(x_1, x_2) dx_2.$$

Thus the tensor product of any two distributions is well defined. It is simple to check that

$$\text{WF}(u_1 \otimes u_2) \subset \text{WF}(u_1) \times \text{WF}(u_2) \cup \text{WF}(u_1) \times \{0\} \cup \{0\} \times \text{WF}(u_1).$$

obtained by isolating each variable separately with a bump function and then tensoring the Fourier transform.

Example. Given $\phi, \psi \in \mathcal{D}(\Omega)$, we have

$$\phi \cdot \psi = i^*(\phi \otimes \psi),$$

where $i(x) = (x, x)$, which gives us another way to define the product of distributions by a tensoring, combined with a pullback.

Finally, we define the pushforward of a distribution. This is most naturally defined distributionally. Given a smooth map $f : \Omega \rightarrow \Psi$, $\phi \in \mathcal{D}(\Omega)$, and $\psi \in \mathcal{D}(\Psi)$, we define

$$\int f_* \phi(y) \psi(y) dy = \int \phi(x) \psi(f(x)) dx.$$

Thus f_* is just the adjoint of f^* . One problem which prevents us from directly using this definition to extend the definition to distributions is that $\psi \circ f$ need not be compactly supported if ψ is compactly supported. One way to resolve this is to consider only the pushforwards of compactly supported distributions. Another way to solve the problem is to assume f is a *proper map*, i.e. inverse images of compact sets are compact. It is then simple to define

$$\int f_* u(y) \psi(y) dy = \int u(x) \psi(f(x)) dx.$$

for a distribution u on Ω and $\psi \in \mathcal{D}(\Psi)$. To understand the wavefront set of u , we consider a bump function ϕ supported in a neighbourhood on Ω and consider

$$\int f_*(u\phi)(y)e^{-2\pi i\eta \cdot y} dy = \int u(x)\phi(x)e^{-2\pi i\eta \cdot f(x)} dx.$$

We have already show that for such an oscillatory integral, provided that $(f(x_0), Df(x_0)^T \eta) \notin \text{WF}(u)$, this integral converges. Thus

$$\text{WF}(f_*u) \subset \{(y, \eta) : \text{There is } (x, \xi) \in \text{WF}(u) \text{ and } Df(x)^T \eta = \xi\}.$$

Now we have defined pushforward, pullback, and tensoring, let us see how they can be used to define useful operations on distributions.

Example. Let M be a manifold with a C^∞ immersed submanifold M_0 . Then we have a smooth inclusion map $f : M_0 \rightarrow M$. That f is proper is equivalent to the assumption that M_0 is a closed submanifold of M . If $u \in \mathcal{D}(M_0)^*$, then f_*u is precisely the distribution on M given by $\langle f_*u, \phi \rangle = \langle u, \phi|_{M_0} \rangle$. The wavefront set $\text{WF}(f_*u)$ is equal to the unique of the conormal bundle of M_0 , i.e. the set

$$N^*(M_0) = \{(x, \xi) \in T^*M : x \in M_0 \text{ and } f^*(x)(\xi) = 0\},$$

and $\text{WF}(u)$.

Let us consider an important example which occurs in the theory of kernel operators. Recall that the Schwartz kernel theorem says that if $T : \mathcal{D}(Y) \rightarrow \mathcal{D}^*(X)$ is any continuous linear map, then there exists a distribution $K \in \mathcal{D}^*(X \times Y)$ such that, formally speaking, if $\phi \in \mathcal{D}(Y)$ and $\psi \in \mathcal{D}(X)$,

$$\int T\phi(x)\psi(x) dx = \int \psi(x)K(x, y)\phi(y) dy.$$

In other words, if $\pi(x, y) = x$ and $\Delta(x, y) = (x, y, y)$, then, unwinding the definition, we find

$$T\phi = \pi_*(\Delta^*(K \otimes \phi)).$$

Now define the *wavefront* or *canonical relation*

$$\text{WF}(K)' = \{(x, \xi; y, \eta) : (x, \xi; y, -\eta) \in \text{WF}(K)\},$$

and

$$\text{WF}_X(K)' = \text{WF}(K)' \circ 0_Y = \{(x, \xi) : (x, \xi; y, 0) \in \text{WF}(K) \text{ for some } y \in Y\}.$$

Working through the definition shows that $\text{WF}(T\phi) \subset \text{WF}_X(K)'$. We can also use the pushforward equation to extend the domain of T to certain compactly supported distributions. Going through the definition shows that for a compactly supported distribution u , the expression $\pi_*(\Delta^*(K \otimes u))$ is well defined provided that $\text{WF}(u)$ is disjoint from $\text{WF}_Y(K)'$, where

$$\text{WF}_Y(K)' = \{(y, \eta) : (x, 0; y, -\eta) \in \text{WF}(K) \text{ for some } x \in X\}.$$

In this case, we define $Tu = \pi_*(\Delta^*(\phi \otimes K))$. This gives a sequentially continuous map from the subspace of compactly supported distributions in $\mathcal{D}_\Gamma^*(\Omega)$ to $\mathcal{D}^*(\Omega)$ for any Γ with $\text{WF}_Y(K)' \cap \Gamma = \emptyset$. If, in addition, the projection map $\pi(x, y) = x$ is proper on $\text{supp}(K)$, then this can be extended to a sequentially continuous map from $\mathcal{D}_\Gamma^*(\Omega)$ to $\mathcal{D}^*(\Omega)$. Again, working through the definitions shows that

$$\text{WF}(Tu) \subset \text{WF}(K)' \circ \{\text{WF}(u) \cup 0_Y\}.$$

A simple way to remember the results of this construction is that K can be applied to any compactly supported distribution u such that

$$\text{WF}(K)' \circ (\text{WF}(u) \cup 0_\Omega)$$

contains no zero vector, and then $\text{WF}(Ku)$ is equal to this composition.

Example. Consider a pseudodifferential operator T given by a symbol a , i.e.

$$T\phi(x) = \int a(x, \xi) \hat{\phi}(\xi) e^{2\pi i \xi \cdot x} d\xi = \int a(x, \xi) \phi(y) e^{2\pi i \xi \cdot (x-y)} d\xi dy.$$

We can also think of T as a kernel operator with kernel

$$K(x, y) = \int a(x, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi.$$

The kernel is a distribution defined by an oscillatory integral distribution, and our calculations for such distributions show that

$$\text{WF}(K) \subset \{(x, -\xi; x, \xi) : x \in \Omega, \xi \in \mathbf{R}^n - \{0\}\}.$$

Thus

$$\text{WF}(K)' \subset \{(x, \xi; x, \xi) : x \in \Omega, \xi \in \mathbf{R}^n - \{0\}\}.$$

In particular, $\text{WF}(K)'$, viewed as a relation, contains no zero vectors, and so T extends to a continuous operator from $\mathcal{E}(\mathbf{R}^d)^*$ to $\mathcal{D}(\mathbf{R}^d)^*$, and if a is compactly supported in the x variable, T extends to a continuous operator from $\mathcal{D}(\mathbf{R}^d)^*$ to itself. For any distribution u , we find that $\text{WF}(Tu) \subset \text{WF}(u)$. This is part of the pseudolocal nature of pseudodifferential operators; when T is applied to some distribution u supported near (x_0, ξ_0) in phase space, we should expect the same will be true of Tu .

Example. Given a distribution u , the operator given by convolution by u has kernel

$$K(x, y) = u(x - y).$$

In other words, $K = f^*u$, where $f : \mathbf{R}^{2d} \rightarrow \mathbf{R}^d$ is given by $f(x, y) = x - y$. Since f is surjective, the pullback f^*u is always well defined, and moreover,

$$\text{WF}(K) \subset f^*\text{WF}(u) = \{(x_1, -\xi, x_2, \xi) : (x_2 - x_1, \xi) \in \text{WF}(u)\},$$

and therefore

$$\text{WF}(K)' \subset \{(x + a, \xi; x, \xi) : (a, \xi) \in \text{WF}(u)\}.$$

We actually have equality here. To see this, for $a \in \mathbf{R}^d$, and let $g : \mathbf{R}^d \rightarrow \mathbf{R}^{2d}$ such that $g(x) = (x + a, a)$. Then $u = g^*K$, and so it follows that

$$\text{WF}(u) \subset g^*\text{WF}(K) = \{(x, \xi) : (x + a, \xi; a, \eta) \in \text{WF}(K)\}.$$

It follows from this that we have equality. Thus the operation $v \mapsto u * v$ is a continuous operator from $\mathcal{E}^*(\mathbf{R}^d)$ to $\mathcal{D}(\mathbf{R}^d)^*$, and

$$\text{WF}(u * v) \subset \{(x + a, \xi) : (a, \xi) \in \text{WF}(u), (x, \xi) \in \text{WF}(v)\}.$$

If u is a distribution with singular support only at the origin, then $\text{WF}(K)'$ is a subset of the diagonal in $T^*\mathbf{R}^d \times T^*\mathbf{R}^d$, and so $\text{WF}(u * v) \subset \text{WF}(v)$ for any distribution v for which the convolution can be computed.

How about the composition of kernel operators? Intuitively, if $A : \mathcal{D}(Y) \rightarrow \mathcal{D}^*(X)$ and $B : \mathcal{D}(Z) \rightarrow \mathcal{D}^*(Y)$, with kernels $K_A(x, y)$ and $K_B(y, z)$, then, if we could define a kernel operator $C = A \circ B : \mathcal{D}(Z) \rightarrow \mathcal{D}^*(X)$, it should have kernel

$$K_C(x, z) = \int K_A(x, y) K_B(y, z) dy.$$

This would be possible and well defined, for instance, if the kernel of B lay in $\mathcal{D}(Y \times Z)$. The integral of K_C would be well defined, the composition of operators would be well defined, for if $\phi \in \mathcal{D}(Z)$, then $B\phi \in \mathcal{D}(Y)$, and so $A(B\phi)$ is well defined, and one finds that K_C is the kernel of this operator. To generalize this composition, we might want to define $K_C = \pi_* \Delta^*(K_A \otimes K_B)$, for the appropriate maps Δ and π . Now this operation is well defined provided that $(x, 0; y, \eta) \in \text{WF}(K_A)'$ and $(y, \eta; z, 0) \in \text{WF}(K_B)'$ for some $x \in X$, $y \in Y$, $z \in Z$, and $\eta \in \mathbf{R}^d$, and the projection map $\pi(x, y, z) = (x, z)$ is proper on the support of $\Delta^*(K_A \otimes K_B)$, which would hold, for instance, if either $\pi(x, y) = x$ was proper on the support of K_A , or $\pi(y, z) = z$ was proper on the support of K_B . In this situation, we have

$$\text{WF}(K_C)' \subset \{\text{WF}(K_A)' \circ \text{WF}(K_B)'\} \cup \{\text{WF}_X(K_A)' \times 0_Z\} \cup \{0_X \times \text{WF}_Z(K_B)'\}.$$

Thus we have defined a fairly general composition operator on distributions, which is the unique bilinear continuous extension of the composition of two operators with kernels in $\mathcal{D}(X \times Y)$ and $\mathcal{D}(Y \times Z)$ to the pair $\mathcal{D}_{\Gamma_A}^*(X \times Y)$ and $\mathcal{D}_{\Gamma_B}^*(Y \times Z)$ for any pair of conic sets Γ_A and Γ_B for which there does not exist any $(x, 0; y, -\eta) \in \Gamma_A$ and $(y, \eta; z, 0) \in \Gamma_B$.

Example. Let us use this construction to define the composition of two pseudodifferential operators P and Q on some domain U , such that one of these operators is proper in the sense above. Indeed, the wave front sets of P and Q are never incompatible, so we can always define $P \circ Q$ using the theory above, and we find that

$$\text{WF}(P \circ Q)' = \text{WF}(P) \circ \text{WF}(Q).$$

In particular, since $\text{WF}(P)$ and $\text{WF}(Q)$ are both subsets of the diagonal in $T^*U \times T^*U$, it follows that $P \circ Q$ is also pseudolocal in the sense that $\text{WF}((P \circ Q)(u)) \subset \text{WF}(u)$. Of course, in the theory of pseudodifferential operators one shows that $P \circ Q$ is also a pseudolocal operator for any two pseudolocal operators P and Q , and then this result follows from that theory.

Example. Suppose $T : \mathcal{D}(Y) \rightarrow \mathcal{D}^*(X)$ is continuous, and thus has a Schwartz kernel $K \in \mathcal{D}^*(X \times Y)$. Then if we consider the operator $T^* : \mathcal{D}(X) \rightarrow \mathcal{D}(Y)^*$ induced by the kernel $K^*(y, x) = \overline{K(x, y)}$, then we find that

$$\langle T\phi, \psi \rangle = \langle \phi, T^*\psi \rangle$$

for any $\phi \in \mathcal{D}(Y)$ and $\psi \in \mathcal{D}(X)$. Thus T^* is the formal adjoint of T . It is simple to verify that

$$WF(K^*)' = \{(y, \eta; x, \xi) : (x, \xi; y, \eta) \in WF(K)'\}.$$

Thus if $\pi(x, y) = y$ is a proper map on $\text{supp}(K)$ and $WF_X(K)' = \emptyset$, we can define $T^* \circ T$, and we have

$$WF(T^* \circ T)' \subset \{(y_1, \eta_1; y_2, \eta_2) : (x, \xi; y_1, \eta), (x, \xi; y_2, \eta_2) \in WF(K)'\}.$$

In particular, if $WF(K)$ is contained in the graph of a function, i.e. there exists a function f such that

$$WF(K) \subset \{(x, \xi; y, \eta) : (y, \eta) = f(x, \xi)\},$$

then $WF(K^* \circ K)$ is a subset of the diagonal of $T^*Y \times T^*Y$, and thus $T^* \circ T$ has various pseudolocal properties. This idea comes up in the theory of Fourier integral operators, for then if T is a Fourier integral operator defined by a Lagrangian distribution that is locally a graph of a function, then T^*T will be a pseudodifferential operator.

10.3 Propagation of Singularities

One important relation between u and $WF(u)$ is the *propagation of singularities theorem*. If u is a solution to a linear partial differential equation

$$\sum_{|\alpha| \leq K} a_\alpha(x) (\partial_\alpha u)(x) = v$$

where v is a distribution, then for any $(x, \xi) \in WF(u) - WF(v)$,

$$q(x, \xi) = \sum_{|\alpha| \leq K} a_\alpha(x) \xi^\alpha = 0,$$

and $WF(u) - WF(v)$ is invariant under the flow generated by the Hamiltonian vector field

$$H_{x, \xi} = \sum_{j=1}^d \frac{\partial q}{\partial x^j} \frac{\partial}{\partial \xi^j} - \frac{\partial q}{\partial \xi_j} \frac{\partial}{\partial x^j}.$$

As a particular example, if $u(t, x, y)$ is a distributional solution to the wave equation $u_{tt} = \Delta u$ and we let $v_t(x, y) = u(t, x, y)$, then $\Delta v_t = u_{tt}$, and so by the propagation of singularities theorem $\text{WF}(v_t) \subset \text{WF}(u_{tt})$.

Then the Paley-Wiener theorem implies that \hat{u} is an analytic function on \mathbf{R}^d . If \hat{u} decays rapidly, then u is also a smooth function. However, even if u is not smooth, \hat{u} may still decrease rapidly in certain directions, which implies that the singularities of u ‘propagate’ in certain directions and understanding these directions is often useful to understanding the distribution u . We can also get even more information about the distribution u by looking at the singular frequencies.

To begin with, let

To begin with, a distribution u is *nonsingular* at a point $x \in \mathbf{R}^d$ if u is locally a C^∞ function in a neighbourhood of x , i.e. there exists a bump function $\phi \in C^\infty(\mathbf{R}^d)$ with $\phi(x) \neq 0$ such that $\phi u \in C^\infty(\mathbf{R}^d)$. The *singular support* of a compactly supported distribution u to be the set of all points $x \in \mathbf{R}^d$ upon which u is not nonsingular.

A degree m constant coefficient linear differential operator $P(D)$ on \mathbf{R}^n is said to be of *real principal type* if the principal symbol P_m is a real-coefficient polynomial, and ∇P_m is non-vanishing on $\mathbf{R}^n - \{0\}$.

Lemma 10.13. *If $P(D)$ is a differential operator of real principal type, then there exists distributions E_+ and E_- which are parametrices for P ,*

$$WF(E_+) \subset (\{0\} \times \mathbf{R}^n) \cup \{(t \cdot \nabla P_m(\xi), \xi) : t > 0, \xi \in Char(P), P_m(\xi) = 0\},$$

and

$$WF(E_-) \subset (\{0\} \times \mathbf{R}^n) \cup \{(t \cdot \nabla P_m(\xi), \xi) : t < 0, \xi \in Char(P), P_m(\xi) = 0\}.$$

Chapter 11

Distributional Methods to PDEs

Distribution theory was originally invented to provide a more amenable setting to the theory of existence for linear partial differential equations. Let us use the theory we have now established to solve some differential equations in the language of distributions. We begin with the most basic differential equation, namely solutions to the transport equation $D^i u = 0$.

Theorem 11.1. *If $u \in \mathcal{D}^*(\mathbf{R}^d)$ and there exists an index i such that $D^i u = 0$, then there exists $v \in \mathcal{D}^*(\mathbf{R}^{d-1})$ such that*

$$\int_{\mathbf{R}^d} u(x) \phi(x) dx = \int_{\mathbf{R}^{d-1}} v(x) \left(\int_{-\infty}^{\infty} \phi(x) dx^i \right) dx,$$

i.e. u is ‘constant’ in the direction i . In particular, if $d = 1$, and $Du = 0$, then u is a constant.

Proof. Suppose without loss of generality that $i = d$. Suppose $\phi \in \mathcal{D}(\mathbf{R}^d)$ and for each $x \in \mathbf{R}^{d-1}$,

$$\int_{-\infty}^{\infty} \phi(x, t) dt = 0.$$

Then the function

$$\psi(x, t) = \int_{-\infty}^t \phi(x, s) ds = 0$$

has compact support and $D^i \psi = \phi$. Thus

$$\begin{aligned} \int_{-\infty}^{\infty} u(x, t) \phi(x, t) \, dx \, dt &= \int_{-\infty}^{\infty} u(x, t) D^i \psi(x, t) \, dx \, dt \\ &= - \int_{-\infty}^{\infty} D^i u(x, t) \psi(x, t) \, dx \, dt = 0. \end{aligned}$$

Now fix $\phi_0 \in \mathcal{D}(\mathbf{R})$ with $\int_{-\infty}^{\infty} \phi_0(x) \, dx = 1$. Then given any $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\int_{-\infty}^{\infty} u(x, t) \phi(x, t) \, dx \, dt = \int_{-\infty}^{\infty} u(x, t) \phi_0(t) \left(\int_{-\infty}^{\infty} \phi(x, s) \, ds \right) \, dx \, dt.$$

Thus it suffices to set

$$v(x) = \int_{-\infty}^{\infty} u(x, t) \phi_0(t) \, dt. \quad \square$$

Remark. Applying this result repeatedly shows that if $u \in \mathcal{D}^*(\mathbf{R}^d)$ is a distribution, and $\nabla u = 0$, then u is a constant.

A change of variables gives versions of this result for any transport equation of the form $w \cdot \nabla u = 0$ for a fixed vector $w \in \mathbf{R}^d$. A corollary is a regularity result for distributional solutions to the equation $w \cdot \nabla u(x) + a \cdot u = f$, where $f \in C(\mathbf{R})$, and $a \in C^\infty(\mathbf{R})$. We begin with the case $d = 1$.

Lemma 11.2. *If $u \in \mathcal{D}(\mathbf{R})^*$ and $Du + au = f$, for $f \in C(\mathbf{R})$, and $a \in C^\infty(\mathbf{R})$, then $u \in C^1(\mathbf{R})$, and so u is a classical solution to the equation.*

Proof. We just apply classical techniques distributionally. First assume $a = 0$. If F is an antiderivative of f in the i th direction, then $F \in C^1(\mathbf{R})$, and $D(u - F) = 0$, so u differs from F by a constant, and is therefore also in $C^1(\mathbf{R})$. For $a \neq 0$, let A be an antiderivative of a , and set

$$E(x) = e^{A(x)}.$$

Then $E \in C^\infty(\mathbf{R})$. Thus if u is a distribution solving $Du + au = f(x)$, and if $v = Eu$ then the product rule shows that $Dv = Ef$. The $a = 0$ case implies that $v \in C^1(\mathbf{R})$, and so $u \in C^1(\mathbf{R})$. \square

Remark. The idea of this result generalizes to a system of differential equations given by a matrix a with C^∞ entries, and where f is a vector with

continuous entries, by finding an invertible matrix $E(x)$ such that $E'(x) = E(x) \cdot a(x)$. In particular, since higher order ordinary differential equations can be reduced to one dimensional systems of ordinary differential equations, we conclude that if u is a distribution satisfying the equation

$$D^m u + a_{m-1} D^{m-1} u + \cdots + au = f,$$

for $a_i \in C^\infty(\mathbf{R})$, and $f \in C(\mathbf{R})$, then u actually lies in $C^m(\mathbf{R})$, and satisfies this equation pointwise in the classical sense.

Higher dimensional analogues of these results are not as strong. Indeed, we have already seen that distributional solutions to $D^i u = 0$ may fail to be classical solutions ‘normal to the direction i ’. On the other hand, we can ‘almost’ obtain such a result if we assume apriori that u is a continuous function.

Lemma 11.3. *Suppose u and f are continuous functions in $C(\mathbf{R}^d)$, and u , viewed as a distribution, satisfies the equation $D^i u = f$. Then $D^i u$ exists pointwise, in the classical sense, for all $x \in \mathbf{R}^d$, and $D^i u(x) = f(x)$ for all $x \in \mathbf{R}^d$.*

Proof. Assume $i = d$ without loss of generality, and write $x = (x_0, t)$, for $x_0 \in \mathbf{R}^{d-1}$ and $t \in \mathbf{R}$. Set

$$v(x_0, t) = \int_0^t f(x_0, s) ds.$$

Then v is a distributional solution to the equation $D^i v = f$, and so $D^i(u - v) = 0$. It follows that there exists a distribution $w \in \mathcal{D}(\mathbf{R}^{d-1})^*$ such that $u(x, t) - v(x, t) = w(x)$. The proof of the existence of w actually implies that w is continuous, since u and v are continuous. But then $u(x, t) = v(x, t) + w(x)$ is differentiable in the t -variable by the fundamental theorem of calculus. \square

Theorem 11.4. *Fix $a_1, \dots, a_n \in C^1(\mathbf{R}^d)$, and $b \in C^0(\mathbf{R}^d)$, and suppose that $u \in C(\mathbf{R}^d)$ is pointwise differentiable, and in a pointwise sense, the equation $a_1 D_1 u + \cdots + a_n D_n u + bu = f$, where $f \in C(\mathbf{R}^d)$. Then the same equation holds in a distributional sense (which makes sense because $D_i u$ is a distribution of order one, and thus can be multiplied against C^1 functions).*

Proof. We adapt the proof of Cauchy’s theorem due to Goursat. Our goal is to prove, given the assumptions in the theorem, that for any $\phi \in C_c^\infty(\mathbf{R}^d)$,

$$\int (b\phi - D_1(a_1\phi) + \cdots + D_n(a_n\phi)) \cdot u = \int f\phi.$$

For any cube I , let

$$A_I = \int_I (f\phi - (b\phi - D_1(a_1\phi) - \cdots - D_n(a_n\phi)) \cdot u) + \int_{\partial I} u(a \cdot n) dS.$$

For any fixed $x_0 \in \mathbf{R}^d$, we claim that

$$\lim_{\substack{x_0 \in I \\ |I| \rightarrow 0}} \frac{|A_I|}{|I|} = 0.$$

To prove this, we may replace u in the formula above with its first order Taylor expansion, and f with the resultant formula $a_1 D_1 u + \cdots + a_n D_n u + bu$, without loss of generality, since the error term here disappears in the limit. But then the formula is easy to verify. But now it follows that we must have $A_I = 0$ for all I , because otherwise we could find a nested family of cubes A_I with $|A_I| \gtrsim |I|$. But if we now take I large enough that it contains the support of ϕ , the theorem is proved. \square

Using techniques from the next section, we obtain a higher dimensional variant of the ODE result above.

Theorem 11.5. *Let $U = V_x \times I_t$, where $V \subset \mathbf{R}^n$ is open, and $I \subset \mathbf{R}$ is an open interval. If $u \in \mathcal{D}(U)^*$ satisfies*

$$\partial_t^m u + L_{m-1}\{\partial_t^{m-1} u\} + \cdots + L_0\{u\} = f,$$

where $f \in C(I, \mathcal{D}(V)^)$, and L_0, \dots, L_{m-1} are differential operators with coefficients in $C^\infty(U)$, then $u \in C^m(I, \mathcal{D}(V)^*)$.*

Proof. Assuming we are working with vector-valued inputs reduces us to the study of $m = 1$, i.e. an equation of the form $\partial_t u + Lu = f$, and it suffices to show that $u \in C^1(I, \mathcal{D}(V)^*)$. The case $n = 0$ has already been considered above. Localizing if necessary, write $u = \sum_{\beta_1, \beta_2} \partial_t^{\beta_1} D_x^{\beta_2} u_\beta$, where $u_\alpha \in C(U)$, and write $f = \sum_{|\alpha| \leq N} D_x^\alpha f_\alpha$, where $f_\alpha \in C(U)$. If $u_\beta = 0$ for any β with $\beta_1 > 0$, then $u \in C(I, \mathcal{D}(V)^*)$. Otherwise, let M be the smallest integer such that $u_\beta = 0$ for any β with $\beta_1 > M$. Then

$$\partial_t u = f - Lu = \sum D^\alpha f_\alpha - \sum L D^\alpha u_\alpha.$$

But, antidifferentiating, we can write

$$\sum D^\alpha f_\alpha - \sum L D^\alpha u_\alpha = \partial_t \left\{ \sum_{\beta} \partial_t^{\beta_1} D^{\beta_2} g_\beta \right\},$$

where $g_\beta = 0$ for $\beta_1 > M - 1$. But we have already seen from the fact that these quantities are equal, that

$$u - \sum_{\beta} \partial_t^{\beta_1} D^{\beta_2} g_\beta$$

lies in $C(I, \mathcal{D}(V)^*)$. Applying induction on M , we conclude $u \in C(I, \mathcal{D}(V)^*)$. But $\partial_t u = u - Lf$ also lies in $C(I, \mathcal{D}(V)^*)$, and so $u \in C^1(I, \mathcal{D}(V)^*)$. \square

Remark. If f extends to an element of $C(\bar{I}, \mathcal{D}(V)^*)$, and u extends to a distribution defined in an open neighborhood of $\bar{I} \times V$, then the proof actually shows that $u \in C^m(\bar{I}, \mathcal{D}(V)^*)$.

11.1 Fundamental Solutions

Let us now discuss the idea of a *fundamental solution* to a partial differential equation, a technique very useful to the study of the existence and uniqueness of such equations. For a differential operator $L = \sum c_\alpha D^\alpha$ with constant coefficients on \mathbf{R}^n , a *fundamental solution* for L is a distribution $\Phi \in \mathcal{D}(\mathbf{R}^n)^*$ such that $L\Phi = \delta$ is the Dirac delta function at the origin. The reason that fundamental solutions are so useful to the study of constant coefficient partial differential equations is that for any compactly supported distribution $v \in \mathcal{D}(\mathbf{R}^d)^*$, the distribution $u = \Phi * v$ is a solution to the equation $Lu = v$, since $Lu = (L\Phi) * v = \delta * v = v$. Thus fundamental solutions give rise to a natural right inverse to a differential operator. It is a general result that *all* constant coefficient differential equations have fundamental solutions, though here we only consider particular examples.

Example. Consider the differential operator Δ , defining Poisson's equation $\Delta u = v$. To guess a fundamental solution for Δ , we first note that since Δ is invariant under rotations, as is the Dirac delta, the equation $\Delta u = \delta$ is radially symmetric. Thus we might expect to find a radially symmetric fundamental solution. Since Δ is elliptic, we expect fundamental solutions to be

smooth away from the origin. Thus to make a guess on the fundamental solution, it will be a useful calculation to determine all smooth, radially symmetric functions $u : \mathbf{R}^n - \{0\} \rightarrow \mathbf{C}$ such that $\Delta u = 0$. If $u(x) = f(|x|)$ for $x \neq 0$ and $f \in C^\infty((0, \infty))$, then for $r > 0$,

$$f''(r) + \frac{n-1}{r}f'(r) = 0.$$

For $n > 2$, this implies that $f(r) = a_1 r^{2-n} + a_2$, and for $n = 2$, $f(r) = a_1 \log(r) + a_2$. We might expect that some choice of constants gives the fundamental solution if we extend these functions to distributions on \mathbf{R}^n . We may without loss of generality pick $a_2 = 0$, since constants do not factor into the output of the operator Δ . The correct choice of the constant a_1 gives the Poisson kernel

$$\Phi(x) = \begin{cases} -\frac{1}{2\pi} \log |x| & : \text{if } n = 2 \\ -\frac{1}{c_n} \frac{1}{n-2} \frac{1}{|x|^{n-2}} & : \text{if } n > 2. \end{cases}$$

Here c_n is the surface area of the unit sphere in \mathbf{R}^n . These distributions are all locally integrable near the origin and thus extend uniquely to distributions on \mathbf{R}^n . To prove that Φ is a fundamental solution, it suffices to show for any $\phi \in \mathcal{D}(\mathbf{R}^d)$,

$$\phi(0) = \int \Phi(x) \Delta \phi(x) dx.$$

Since Φ is locally integrable, Gauss' formula implies that as $\varepsilon \rightarrow 0$,

$$\int_{|x| \geq \varepsilon} \Phi(x) \Delta \phi(x) dx = \int_{|x|=\varepsilon} \Phi(x) [\nabla \phi(x) \cdot n(x)] dS - \int \nabla \Phi(x) \cdot \nabla \phi(x) dx.$$

Now a simple estimate gives that as $\varepsilon \rightarrow 0$,

$$\int_{|x|=\varepsilon} \Phi(x) [\nabla \phi(x) \cdot n(x)] dS \rightarrow 0.$$

Thus we conclude that

$$\int \Phi(x) \Delta \phi(x) dx = - \int \nabla \Phi(x) \cdot \nabla \phi(x) dx.$$

Again, we approximate away from the origin by a ε ball, integrate by parts, and calculate that

$$\begin{aligned} \int_{|x| \geq \varepsilon} \nabla \Phi(x) \cdot \nabla \phi(x) dx &= \int_{|x|=\varepsilon} \phi(x) [\nabla \Phi(x) \cdot n(x)] dS - \int_{|x| \geq \varepsilon} \Delta \Phi(x) \phi(x) dx \\ &= \int_{|x|=\varepsilon} \phi(x) [\nabla \Phi(x) \cdot n(x)] dS. \end{aligned}$$

For $n = 2$, we have $\nabla\Phi(x) = (2\pi)^{-1}(x/|x|^2)$, hence

$$\begin{aligned}\int_{|x|=\varepsilon} \phi(x)[\nabla\Phi(x) \cdot n(x)]dS &= \frac{1}{2\pi\varepsilon} \int_{|x|=\varepsilon} \phi(x) dx \\ &= \phi(0) + O(\varepsilon).\end{aligned}$$

For $n > 2$, $\nabla\Phi(x) = c_d^{-1}(x/|x|^d)$, so

$$\int_{|x|=\varepsilon} \phi(x)[\nabla\Phi(x) \cdot n(x)]dS = \frac{1}{c_d\varepsilon^{d-1}} \int_{|x|=\varepsilon} \phi(x) dS = \phi(0) + O(\varepsilon).$$

Taking $\varepsilon \rightarrow 0$ gives the result.

An alternate approach to obtaining this fundamental solution is to take Fourier transforms, assuming that we can find a tempered fundamental solution Φ . If $\Psi = \hat{\Phi}$, then we conclude that Φ is a fundamental solution if and only if Ψ is tempered and

$$-4\pi^2|\xi|^2 \cdot \Psi(\xi) = 1.$$

For $n > 2$, we can interpret the formula $\Psi(\xi) = (-1/4\pi^2)|\xi|^{-2}$ as defining a distribution by integration against a locally integrable function. For $n = 2$ a version of this equation remains true provided that we interpret the distribution $1/|\xi|^2$ at the origin in the right way. Thus we have the alternate expression for the fundamental solution above.

Example. Next, consider the operator $L = \partial_t - \Delta$ on \mathbf{R}^{n+1} , which gives rise to the heat equation. Set

$$\Phi_+(x, t) = \frac{1}{(4\pi t)^{n/2}} \exp\left(-\frac{|x|^2}{4t}\right)$$

for $t > 0$, and $\Phi(x, t) = 0$ for $t \leq 0$. Then Φ is locally integrable, and a fundamental solution for L . Indeed, Φ is tempered in the x -variable, and the Fourier transform of Φ in the x variable is verified to be

$$U(\xi, t) = e^{-4\pi^2 t |\xi|^2}$$

which satisfies $\partial_t U = -4\pi^2 |\xi|^2 U$, which is equivalent to the equation $L\Phi = \delta$. We see immediately that $\Phi(0+) = \lim_{t \rightarrow 0} \Phi(t)$ is the Dirac delta function at the origin.

More generally, similar calculations enable us to find fundamental solutions to any operator of the form $L = \partial_t - S$, where $S = \sum A_{ij} \partial_i \partial_j$. If we assume that there exists a fundamental solution Φ_+ to L , tempered in the x -variable and supported on $t \geq 0$, then taking Fourier transforms of the equation $L\Phi = \delta$ in the x -variable, and setting Ψ_+ to be this Fourier transform, we are lead to conclude that $\partial_t \Psi + 4\pi^2(\xi^T A \xi) \Psi = \delta(t)$. Let us assume that Φ is supported on $t \geq 0$. Thus we can write

$$\Psi(\xi, t) = H(t)A(\xi)e^{-4\pi^2(\xi^T A \xi)t}$$

for some distribution $A \in \mathcal{D}(\mathbf{R}^n)^*$. But we then calculate that

$$L\Psi = \delta(t)A(\xi)$$

and so we must have $A(\xi) = 1$, i.e.

$$\Psi(\xi, t) = H(t)e^{-4\pi^2(\xi^T A \xi)t}.$$

But this means that, in order to get a tempered fundamental solution we must assumed $\text{Re}(A)$ is positive semidefinite. If we, in addition, assume that A is invertible, then we have the Fourier transform for Ψ , namely, we conclude that

$$\Phi(x, t) = \frac{H(t)}{(4\pi t)^{n/2}(\det(A))^{1/2}} e^{-x^T A^{-1} x/4t}.$$

If the real part of A is positive definite, then the same is true for A^{-1} , and so Φ can be interpreted as locally integrable on $\mathbf{R} \times \mathbf{R}^n$. On the other hand, if A is positive semidefinite this is not the case, for instance, the Schrödinger equation $L = \partial_t - i\Delta_x$ has fundamental solution

$$\Phi(x, t) = \frac{H(t)}{(4\pi it)^{n/2}} e^{i|x|^2/4t},$$

which is not locally integrable, but we must treat this solution as a principal value in the t -variable, using the principle of stationary phase to show the limit is well defined, which allows us to conclude Φ is a distribution of order $n + 1$.

Before we consider other quadratic partial differential operators, let us perform a few calculations. Take an operator of the form

$$L_A = \sum_{i,j=1}^n A_{ij} D_x^{ij}$$

for some symmetric $n \times n$ matrix A . If T is an invertible linear transformation, then the chain rule implies that

$$D^i\{f \circ T\} = \sum_{k=1}^n T_{ki} \cdot (D^k\{f\} \circ T),$$

i.e. $\nabla\{f \circ T\} = T^t\{(\nabla f) \circ T\}$. Iterating the chain rule, we find that

$$D^{ij}\{f \circ T\} = \sum_{k=1}^n \sum_{l=1}^n T_{lj} T_{ki} \cdot (D^{kl}\{f\} \circ T).$$

Thus the Hessian is $H\{f \circ T\} = T^t \cdot \{Hf \circ T\} \cdot T$. This means that if T^*L_A is the operator given by the equation $T^*L_A\{f\} = L_A\{f \circ T\} \circ T^{-1}$, then

$$T^*L_A f = (L_{TAT^t} f) \circ T.$$

The use of these pullbacks is that if $f \in C^\infty(\mathbf{R}^n)$, and if $T^*f = f \circ T^{-1}$, then

$$T^*\{L_A f\} = L_A f \circ T^{-1} = L_A\{T^*f \circ T\} \circ T^{-1} = (T^*L_A)\{T^*f\}.$$

We can take several important points away from this result:

- $T^*L = L$ precisely when $TAT^t = A$, which is equivalent to the condition that if $B_A(x, y) = x^t A y$ is the symmetric bilinear form specified by A , then $B_A(Tx, Ty) = B(x, y)$. The family of such linear maps from a *generalized orthogonal group*.
- Sylvester's law of inertia implies we can write any symmetric $n \times n$ matrix A as $T^{-1}BT^{-t}$, where B is a diagonal matrix consisting of zeroes, ones, and negative ones. But this means that

$$T^*L = \sum B_{ii}(D^{ii}\{f\} \circ T).$$

Thus the theory of a real-coefficient order two differential equation with constant coefficients is equivalent to one of the form

$$Lf = \sum_{1 \leq i < n_1} \frac{\partial^2 f}{\partial x_i^2} - \sum_{n_1 \leq i < n_2} \frac{\partial^2 f}{\partial x_i^2},$$

where $n_1 + n_2 \leq n$. This simplifies our calculations considerably, e.g. the theory of an operator L given by a positive definite matrix

A , i.e. one which has signature $(n, 0)$, is equivalent to the theory of the Laplacian Δ . The theory of L given by a $(n+1) \times (n+1)$ matrix with signature $(1, n)$ is equivalent to the theory of the wave operator $\square = \partial_t^2 - \Delta_x$, and so on and so forth.

Inspired by these calculations, we now find a fundamental solution to any quadratic partial differential operator $L = \sum B_{ij} \partial^{ij}$, such that B is invertible.

Lemma 11.6. *Let A be a real, invertible matrix with signature (n_+, n_-) . If c_n is the surface area of the unit sphere in \mathbf{R}^n , then for $n > 2$, if we let $B = A^{-1}$, and*

$$L_B = \sum B_{ij} \partial^{ij},$$

then $L_B\{(A \pm i0)^{1-n/2}\} = -(n-2)c_n |\det(A)|^{-1/2} e^{\mp i\pi n_-/2} \delta_0$.

Proof. By conjugation, it suffices to show that

$$L_B\{(A + i0)^{1-n/2}\} = -(n-2)c_n |\det A|^{-1/2} e^{-i\pi n_-/2} \delta_0.$$

We will begin by showing that if A is a symmetric matrix with complex coefficients such that $\operatorname{Re}(A)$ is positive definite, and if B is the inverse of A , then

$$L_B\{A^{1-n/2}\} = -(n-2)c_n (\det A)^{-1/2} \delta$$

where the determinant power is defined to be the unique analytic branch such that $(\det A)^{-1/2} > 0$ if A is positive definite. Analytic continuation means we only need to prove this formula if A , and thus B , is positive definite. But then we can find a matrix S such $SBS^t = I$, and thus $S^{-t}AS^{-1} = I$. But then

$$\begin{aligned} S^*\{L_B\{A^{1-n/2}\}\} &= (S^*L_B)\{S^*A^{1-n/2}\} \\ &= L_I\{|x|^{2-n}\} \\ &= \Delta\{|x|^{2-n}\} \\ &= -(n-2)c_n \delta_0. \end{aligned}$$

The result then follows because

$$(S^{-1})^* \delta_0 = \det(S)^{-1} \delta_0 = \det(A)^{-1/2}.$$

To obtain the general result, we now note that if

$$A_\varepsilon = \varepsilon - iA \quad \text{and} \quad B_\varepsilon = (\varepsilon - iA)^{-1}.$$

Then

$$\det(A_\varepsilon)^{-1/2} = |\det(A)|^{-1/2} e^{i\pi \operatorname{sgn}(A)/4},$$

and $B_\varepsilon \rightarrow iB$ and $A_\varepsilon^{1-n/2} \rightarrow i^{-1} e^{i\pi n/4} (A + i0)^{1-n/2}$, so that this gives the result in the limit. \square

We would hope to define a fundamental solution to the equation L_B as the pull back $A^* \chi_+^{1-n/2}$. This is not quite possible using the spectral analysis of singularities since A is singular at the origin. But provided A is nonsingular, we can define the pullback viewing A as a map from $\mathbf{R}^n - \{0\} \rightarrow \mathbf{R}$. Since $A^* \chi_+^{1-n/2}$ is then a homogeneous distribution of degree $2 - n$ on $\mathbf{R}^n - \{0\}$, this distribution then uniquely extends to a distribution on \mathbf{R}^n , and we claim this is a constant multiple of a fundamental solution.

Corollary 11.7. *Using the setup to the previous lemma,*

$$L_B \{A^* \chi_\pm^{1-n/2}\} = \pm 4\pi^{n/2-1} \sin(\pi n_\pm/2) |\det A|^{-1/2} \delta_0.$$

Proof. The result follows from the previous result given that

$$\chi_+^a(s) = \frac{i}{2\pi} \Gamma(-a) ((s - i0)^a e^{i\pi a} - (s + i0)^a e^{-i\pi a})$$

for all a that is not a positive integer. To obtain this, we use the fact that $\Gamma(a+1)\Gamma(-a) = -\pi/\sin(\pi a)$, which means that

$$\begin{aligned} \chi_+^a(s) &= \frac{1}{\Gamma(a+1)} s_+^a \\ &= -\frac{1}{\pi} \Gamma(-a) \sin(\pi a) \cdot s_+^a \\ &= \frac{i}{2\pi} \Gamma(-a) (e^{i\pi a} - e^{-i\pi a}) s_+^a. \end{aligned}$$

For $\operatorname{Re}(a) > 0$, s_+^a is locally integrable, and

$$(s \pm i0)^a = \begin{cases} s^a & : s \geq 0, \\ |s| e^{\pm i\pi a} & : s < 0. \end{cases}$$

Thus we conclude that

$$(e^{i\pi a} - e^{-i\pi a})s_+^a = e^{i\pi a}(s - i0)^a - e^{-i\pi a}(s + i0)^+.$$

The general case follows by analytic continuation. \square

Example. Consider the d'Alembertian, or wave operator $\square f = \partial_t^2 f - \Delta_x f$. The fundamental solutions for this equation are significantly richer than for the Laplace equation. There is no single preferred fundamental solution in this setting. Physical intuition tells us that waves travel at a finite speed of propagation, in this case, at a unit velocity. Thus we expect to find a fundamental solution supported on the interior of the light cone

$$\Sigma = \{(x, t) \in \mathbf{R}^n \times \mathbf{R} : Q(x, t) = 0\},$$

i.e. supported on $\Sigma_+ = \{(x, t) \in \mathbf{R}^n \times \mathbf{R} : Q(x, t) \geq 0\}$, where $Q(x, t) = t^2 - |x|^2$. In fact, for $n \geq 2$, we have already done the calculations to find such a fundamental solution, namely we have a fundamental solution of the form

$$\Phi = \frac{1}{4\pi^{\frac{n-1}{2}}} Q^* \left(\chi_+^{-\frac{n-1}{2}} \right).$$

For $n = 1$, this formula actually continues to hold, namely, we have

$$\Phi(x, t) = (1/2)\mathbf{I}((x, t) \in \Sigma_+) = (1/2)H(t - x)H(t + x),$$

though we have to make do with more rudimentary calculations. These choices give fundamental solutions to the wave equation. To verify the equation gives a fundamental solution to the wave operator for $n = 1$, we calculate that

$$\begin{aligned} \partial_t \Phi(x, t) &= (1/2)\delta(t - x)H(t + x) + (1/2)\delta(t + x)H(t - x) \\ &= (1/2)(\delta(t - x) + \delta(t + x))H(t), \end{aligned}$$

and so

$$\begin{aligned} \partial_t^2 \Phi(x, t) &= (1/2)(\delta(t - x) + \delta(t + x))\delta(t) + (1/2)(\delta'(t - x) + \delta'(t + x))H(t) \\ &= \delta(x, t) + (1/2)(\delta'(t - x) + \delta'(t + x))H(t). \end{aligned}$$

Next, we calculate that

$$\begin{aligned} \partial_x \Phi(x, t) &= (1/2)H(t - x)\delta(t + x) - (1/2)H(t + x)\delta(t - x) \\ &= (1/2)(\delta(t + x) - \delta(t - x))H(t), \end{aligned}$$

and thus

$$\begin{aligned}\partial_x^2 \Phi(x, t) &= (1/2) (\delta(t+x) - \delta(t-x)) \delta(t) + (1/2) (\delta'(t+x) + \delta'(t-x)) H(t) \\ &= (1/2) (\delta'(t+x) + \delta'(t-x)) H(t).\end{aligned}$$

Thus $\square \Phi(x, t) = \delta(t, x)$. In particular, we note that when n is odd, then the support of

$$\chi_+^{-\frac{n-1}{2}}$$

is equal to $\{0\}$. This means that when $n \geq 3$ is odd, then Φ is actually supported on Σ , which hints at Huygen's principle, i.e. that in odd dimensions, if u is a solution to the Cauchy problem $\square u = 0$ with initial conditions u_0 , then the behaviour of u at a point (t, x) is determined by the values of u_0 on a sphere of radius t around x .

There are two important alternate fundamental solutions, the forward and backward, or advanced and retarded fundamental solution

$$\Phi_+(x, t) = 2H(t)\Phi(x, t) \quad \text{and} \quad \Phi_-(x, t) = 2H(-t)\Phi(x, t).$$

That these two distributions are also fundamental solution follows from the symmetry of all objects involved under time reflection. These are actually the unique fundamental solutions supported on the forward and backward light cone and their interior. Indeed, if $u \in \mathcal{D}^*(\mathbf{R}^n \times \mathbf{R})$ is supported on the forward light cone and $\square u = 0$, then

$$u = (\square \Phi_+) * u = \Phi_+ * \square u = 0,$$

where the convolution is well defined because the projection

$$((x, t), (y, s)) \mapsto (x + y, t + s)$$

is then proper on $\text{supp}(u) \times \text{supp}(\Phi_+)$.

The wave equation plays nicely with the Fourier transform. If Φ is any fundamental solution, and we assume that it is tempered in the x -variable (which is the case for the distributions we consider above), supported on Σ , then applying the Fourier transform in this variable, i.e. setting $\Psi = \mathcal{F}_x \Phi$, then

$$\partial_t^2 \Psi(\xi, t) + 4\pi^2 |\xi|^2 \Psi(\xi, t) = \delta(t)$$

This implies that, for $t > 0$, we have

$$\Psi(\xi, t) = A_+(\xi) e^{2\pi i |\xi| t} + B_+(\xi) e^{-2\pi i |\xi| t},$$

and a similar formula, allowing different functions, for $t < 0$. Because Ψ is smooth away from the boundary of the light cone, this implies that A_+ and B_+ are both smooth functions. We calculate that

$$\partial_t \Psi(\xi, 0) = 2\pi i |\xi| [A_+(\xi) - B_+(\xi)]$$

If we assume that Ψ is supported on $t \geq 0$, and we denote it by Ψ_+ , then for any $\phi \in \mathcal{D}(\mathbf{R}^n \times \mathbf{R})$, an integration by parts shows that

$$\begin{aligned} & \int_0^\infty \int_{\mathbf{R}^n} (\partial_t^2 \phi(\xi, t) + 4\pi^2 |\xi|^2 \phi(\xi, t)) \Psi_+(\xi, t) \\ &= -[A_+(\xi) + B_+(\xi)] \partial_t \phi(\xi, 0) + 2\pi i |\xi| [A_+(\xi) - B_+(\xi)] \phi(\xi, 0). \end{aligned}$$

Thus to get a fundamental solution, we should have $A_+(\xi) + B_+(\xi) = 0$, and $A_+(\xi) - B_+(\xi) = 1/2\pi i |\xi|$. Thus we conclude that

$$\Psi_+(\xi, t) = \frac{\sin(2\pi |\xi| t)}{4\pi |\xi|}.$$

This implies that there is a unique fundamental solution supported on $t > 0$ and tempered in the x -variable, i.e. the fundamental solution Φ_+ we define above, and moreover, we have the Fourier representation

$$\Phi_+(x, t) = \frac{H(t)}{2} \int \frac{\sin(2\pi |\xi| t)}{2\pi |\xi|} e^{2\pi i \xi \cdot x} d\xi.$$

A similar formula holds for the other fundamental solutions we constructed above, namely

$$\Phi_-(x, t) = \frac{H(-t)}{2} \int \frac{\sin(-2\pi |\xi| t)}{2\pi |\xi|} e^{2\pi i \xi \cdot x} d\xi,$$

and

$$\Phi(x, t) = \frac{1}{4} \int \frac{\sin(2\pi |\xi| t)}{2\pi |\xi|} e^{2\pi i \xi \cdot x} d\xi.$$

However, the Fourier calculation above yields some other interesting fundamental solutions, such as the Feynman fundamental solution

$$\Phi_F(x, t) = \frac{1}{4i} \int \frac{1}{2\pi |\xi|} e^{2\pi i (\xi \cdot x + |\xi| t)} d\xi$$

One quirk of this fundamental solution is that it is not contained in the light cone, despite the finite speed of propagation of the d'Alembertian. In fact, $\text{supp}(\Phi_F) = \mathbf{R} \times \mathbf{R}^n$. To see this, let $u(t) = \Phi_F(0, t)$ and $v(x) = \Phi_F(x, 0)$. That $\text{supp}(\Phi_F) = \mathbf{R} \times \mathbf{R}^n$ will follow from the symmetry of Φ_F under the Lorentz transformation, and that $\text{supp}(u) = \mathbf{R}$, $\text{supp}(v) = \mathbf{R}^n$. To obtain this, we can explicitly calculate that for $n = 1$, $v(x)$ is a multiple of $\text{sgn}(x)$, and for $n > 1$, $v(x)$ is a multiple of $|x|^{1-n}$. In both cases, $\text{supp}(v) = \mathbf{R}^n$. Next, we calculate that

$$u(t) = \frac{1}{4i} \int \frac{e^{2\pi i |\xi| t}}{2\pi |\xi|} d\xi = \frac{A_{n-1}}{4i} \int_0^\infty \tau^{n-2} e^{2\pi i \tau t} d\tau.$$

Thus u is the Fourier transform of a homogeneous function of order $n-2$ in the τ variable, and is thus a homogeneous function of order $1-n$ in the t variable. Time reversal symmetry, and the fact that $u(t)$ is not supported at the origin (since its Fourier transform is not a polynomial) imply that $\text{supp}(u) = \mathbf{R}$.

TODO: Move to Cauchy Problem For the study of the Cauchy problem for the wave operator, it is useful to study the behaviour of Φ_+ for $t > 0$. Since $\partial_t^2 \Phi - \Delta_x \Phi = 0$, it follows from the general theory of distributional solutions to ODEs that $\Phi \in C^2([0, \infty), \mathcal{D}(\mathbf{R}^n)^*)$. In particular, if we consider the distributions $\Phi(t)$ on $\mathcal{D}(\mathbf{R}^n)^*$, then both $\Phi(0+) = \lim_{t \rightarrow 0+} \Phi(t)$ and $\partial_t \Phi(0+) = \lim_{t \rightarrow 0+} \partial_t \Phi(t)$ are well defined. Moreover, for $\phi \in \mathcal{D}(\mathbf{R}^{n+1})$ we have the representation formula

$$\langle \Phi_+, \phi \rangle = \int_0^\infty \langle \Phi_+(t), \phi(t) \rangle dt,$$

since both sides of the equation define homogeneous distributions of order $1-n$ which agree for $\phi \in \mathcal{D}(\mathbf{R}^{n+1} - \{0\})$, and thus agree for all $\phi \in \mathcal{D}(\mathbf{R}^{n+1})$. But integration by parts implies that

$$\begin{aligned} \phi(0) &= \langle \square \Phi_+, \phi \rangle \\ &= \int_0^\infty \langle \Phi_+(t), \square \phi(t) \rangle dt \\ &= \int_0^\infty \langle \Phi_+(t), \partial_t^2 \phi(t) \rangle - \langle \Phi_+(t), \Delta \phi(t) \rangle dt \\ &= \langle \partial_t \Phi_+(0+), \phi(0) \rangle - \langle \Phi_+(0+), \partial_t \phi(0) \rangle. \end{aligned}$$

Since this is true for arbitrary $\phi \in \mathcal{D}(\mathbf{R}^{n+1})$, we conclude that $\Phi_+(0+) = 0$ and $\partial_t \Phi_+(0+) = \delta$. For $t > 0$, we can compute a formula for the distribution Φ_+ .

The map $(t, x) \mapsto (t^2 - |x|^2, x)$ then has an inverse

$$H(s, x) = ((s + |x|^2)^{1/2}, x).$$

Thus $|\det(DH)| = (1/2)(s + |x|^2)^{-1/2}$. If $\phi \in \mathcal{D}(\mathbf{R}^n \times \mathbf{R})$ is supported on $t > 0$, and we set

$$\langle \Phi_+, \phi \rangle = \frac{1}{4\pi^{\frac{n-1}{2}}} \langle \chi_+^{-\frac{n-1}{2}}, \psi \rangle,$$

where

$$\psi(s) = \int \phi(H(s, x))(s + |x|^2)^{-1/2} dx.$$

If we set

$$\tilde{\phi}(t, r) = \frac{1}{r} \int_{|x|=r} \phi(t, x) dx,$$

then

$$\psi(s) = \int_{s^{1/2}}^{\infty} \tilde{\phi}(t, (t^2 - s)^{1/2}) dt$$

If we consider the compactly supported distribution $\Phi_+(t)$ on \mathbf{R}^n , then for $\eta \in C^\infty(\mathbf{R}^n)$,

$$\langle \Phi_+(t), \eta \rangle = \frac{1}{4\pi^{\frac{n-1}{2}}} \int_0^{t^2} \chi_+^{-\frac{n-1}{2}}(s) \tilde{\eta}((t^2 - s)^{1/2}) ds,$$

where

$$\tilde{\eta}(r) = \frac{1}{r} \int_{|x|=r} \eta(x) dx.$$

For $n = 1$, we have $\chi_+^0(s) = H(s)$, and

$$\Phi_+(t) = (1/2)\mathbf{I}_{[-t, t]}.$$

For $n = 2$, we have $\chi_+^{-1/2} = \chi_+^{-1/2} \pi^{-1/2}$, and so

$$\Phi_+(t) = (1/2\pi) \cdot \max(t^2 - |x|^2, 0)^{-1/2}.$$

For odd values $n = 2m + 1$, we have $\chi_+^{-\frac{n-1}{2}} = \chi_+^{-m} = D^{m-1} \delta_0$, and this leads to the conclusion that

$$\langle \Phi_+(t), \eta \rangle = \frac{1}{4\pi^m} \frac{d^{m-1}}{ds^{m-1}} \left(\tilde{\eta}(s^{1/2}) \right) \Big|_{s=t^2}.$$

Thus this value depends on the values of η , and their normal derivatives up to order $m - 1$, on a sphere of radius t . In particular, for $n = 3$, we have

$$\langle \Phi_+(t), \eta \rangle = \frac{1}{4\pi} \tilde{\eta}(t) = \frac{1}{4\pi t} \int_{|x|=t} \eta(x) dx.$$

TODO: Move to Cauchy Problem The existence and uniqueness of solutions to the Cauchy problem for the wave equation follow immediately from this calculation. Namely, if $\phi_0, \phi_1 \in C^\infty(\mathbf{R}^n)$, and $f \in C^\infty(\mathbf{R}^n \times [0, \infty))$, then there exists a unique $u \in C^\infty(\mathbf{R}^n \times [0, \infty))$ such that $\square u = f$ on $\mathbf{R}^n \times [0, \infty)$, and we have

$$u(t) = \Phi_+(t) * \phi_1 + (\partial_t \Phi_+)(t) * \phi_0 + \int_0^t (\Phi_+(t-s) * f(s)) ds.$$

Uniqueness follows the fact the fact that if u is a distribution supported on the forward light cone and $\square u = 0$, then $u = 0$. Conversely, the equation above certainly defines an element of $C^\infty([0, \infty) \times \mathbf{R}^n)$ and gives a solution since

$$\square(\Phi_+ * \phi_1) = \delta * \phi_1 = \phi_1$$

$$\square(\partial_t \Phi_+ * \phi_0) = \partial_t(\square \Phi_+ * \phi_0) = \partial_t \phi_0 = 0$$

and since $\Phi_+(0) = 0$, and $\partial_t \Phi_+(0) = \delta$, we find that

$$\partial_t \left\{ \int_0^t (\Phi_+(t-s) * f(s)) ds \right\} = \int_0^t (\partial_t \Phi_+(t-s) * f(s)) ds$$

and so

$$\partial_t^2 \left\{ \int_0^t (\Phi_+(t-s) * f(s)) ds \right\} = f(t) + \int_0^t (\partial_t^2 \Phi_+(t-s) * f(s)) ds.$$

Since $\square \Phi_+$ vanishes away from the origin,

$$\square \left\{ \int_0^t (\Phi_+(t-s) * f(s)) ds \right\} = f(t),$$

which completes the proof.

Example. Consider the Cauchy-Riemann equations on \mathbf{R}^2 , i.e. the operators $\partial_z = (1/2)(\partial_x - i\partial_y)$ and $\partial_{\bar{z}} = (1/2)(\partial_x + i\partial_y)$. The divergence theorem tells us that

$$\int_{\Omega} \frac{\partial \phi}{\partial z} dx dy = \frac{1}{2} \int_{\partial \Omega} \phi d\bar{z}$$

and

$$\int_{\Omega} \frac{\partial \phi}{\partial \bar{z}} dx dy = \frac{1}{2} \int_{\partial \Omega} \phi dz$$

The variant of Green's formula then tells us that

$$\int_{\Omega} \left(u \frac{\partial v}{\partial z} + \frac{\partial u}{\partial \bar{z}} v \right) dx dy = \frac{1}{2} \int_{\partial \Omega} uv d\bar{z}.$$

Thus the operators ∂_z and $\partial_{\bar{z}}$ are self adjoint. In particular, applying this formula gives that for any $\phi \in \mathcal{D}(\mathbf{R}^2)$,

$$\begin{aligned} \int_{|z| \geq \varepsilon} \frac{\partial \phi}{\partial \bar{z}} \left(\frac{1}{\pi i} \frac{1}{z} \right) dx dy &= \frac{1}{2\pi i} \int_{|z|=\varepsilon} \frac{\phi}{z} dz \\ &= \frac{1}{2\pi} \int_0^{2\pi} \phi(\varepsilon e^{i\theta}) d\theta \\ &= \phi(0) + O(\varepsilon). \end{aligned}$$

Taking $\varepsilon \rightarrow 0$ allows us to conclude that

$$\Phi(x, y) = \frac{1}{i\pi} \frac{1}{x + iy}$$

is a fundamental solution to the operator $\partial_{\bar{z}}$. Similarly,

$$\Phi(x, y) = \frac{1}{i\pi} \frac{1}{x - iy}$$

is a fundamental solution to the operator ∂_z . The first fundamental solution is called the Cauchy Kernel.

A simple consequence of the fundamental solution technique is the intriguing statement that *any* distribution is a successive derivative of a family of continuous functions.

Theorem 11.8. For any open set $U \subset \mathbf{R}^n$, and any distribution $u \in \mathcal{D}(U)^*$, there exists $f_\alpha \in C(U)$, such that the cover $\{\text{supp}(f_\alpha)\}$ is locally finite on U , and such that $u = \sum_\alpha D^\alpha f_\alpha$.

Proof. Suppose first that $U = \mathbf{R}^n$. For any m , the function

$$\Phi(x) = \max(x_1, 0)^m \dots \max(x_n, 0)^m / (m!)^n$$

is a fundamental solution to the operator $L = (\partial_1 \dots \partial_n)^{m+1}$, and lies in $C^{m-1}(\mathbf{R}^d)$. If u is a distribution of order $m-1$, then it follows that $v = \Phi * u$ lies in $C(\mathbf{R}^d)$, and $Lv = u$, which completes the proof in this case. In general, we just localize. \square

The fundamental solutions to the heat equation and Cauchy-Riemann equation are smooth away from the origin. It will turn out that this is true for any constant coefficient partial differential equation which is *hypoelliptic*. Here are several consequences.

Lemma 11.9. Suppose Φ is a fundamental solution to a constant coefficient partial differential operator L on \mathbf{R}^n , and suppose that $\text{sing-supp}(\Phi) = \{0\}$. Then for any open set $U \subset \mathbf{R}^n$, and any distribution u on U , $\text{sing-supp}(Lu) = \text{sing-supp}(u)$. Thus any distribution u on U with $Lu = 0$ lies in $C^\infty(U)$. If $\{u_n\}$ is a sequence of solutions to the equation $Lu = 0$, with $u_n \rightarrow u$ distributionally, then $u_n \rightarrow u$ in $C^\infty(U)$.

Proof. It is simple to see that $\text{sing-supp}(Lu) \subset \text{sing-supp}(u)$. Conversely, fix a compact set $K \subset U$, and consider $\psi \in \mathcal{D}(U)$ equal to one on a neighbourhood of K . Then for any distribution u on U with $Lu = 0$,

$$\begin{aligned} K \cap \text{sing-supp}(Lu) &= K \cap \text{sing-supp}(L(\psi u)) \\ &= K \cap \text{sing-supp}(\Phi * (\psi u)) \\ &= K \cap [\text{sing-supp}(\Phi) + \text{sing-supp}(\psi u)] \\ &= K \cap \text{sing-supp}(u). \end{aligned}$$

The first part is therefore completed, since K was arbitrary. Now given the Fréchet space $X = \{u \in \mathcal{D}(U)^* : Lu = 0\}$, we have a natural inclusion map $i : X \rightarrow C^\infty(U)$. It's graph is obvious closed, so by the closed graph theorem, this map is continuous, and this implies the sequence property above. \square

The fundamental solutions above are actually better than just being smooth away from the origin, they are *analytic* away from the origin. This is true of any *elliptic* partial differential equation with constant coefficients. The following lemma then applies.

Lemma 11.10. *Suppose Φ is a fundamental solution to a constant coefficient partial differential operator L on \mathbf{R}^n , and suppose that Φ is real analytic away from the origin. Then for any open set $U \subset \mathbf{R}^n$, there is an open set $V \subset \mathbf{C}^n$ such that if u is a distribution on U , and $Lu = 0$, then u is an analytic function extending to a complex analytic function on V . In particular, we conclude that if u vanishes in a neighborhood of a point $x_0 \in U$, then u vanishes on the connected component of U containing x_0 . Moreover, if u_n is a sequence in $C^\infty(U)$ with $Lu_n = 0$, and with $u_n \rightarrow u$, then the extension of the functions u_n to analytic functions on V converge locally uniformly to the extension of u .*

Proof. Choose any open set $\mathbf{R}^n - \{0\} \subset V_0 \subset \mathbf{C}^n$ such that Φ extends to a complex analytic function on V_0 . Since the question is local, consider a compact set K , and $\psi \in \mathcal{D}(U)$ equal to one on a neighborhood of K . Then for $x \in K$, we have

$$u(x) = (\psi u)(x) = (\Phi * L(\psi u))(x) = \int \Phi(x - y) L(\psi u)(y) dy,$$

and this formula continues to define a complex analytic function if we replace x with a complex parameter z provided that for any $y \in \text{supp}(\nabla(\psi u))$, $z - y \in V_0$. This is a neighborhood of K . We can then perform this extension for many different choices of ψ , and thus of K , to show that u extends to an analytic function on some set V depending only on U . To get the latter part of the Lemma, we just apply the closed graph theorem. \square

Let us now extend the approximation theorem of Runge to the case of general elliptic partial differential equations.

Lemma 11.11. *Suppose Φ is a fundamental solution to a constant coefficient partial differential operator L on \mathbf{R}^n , and suppose that Φ is real analytic away from the origin. Consider two open sets $V \subset U \subset \mathbf{R}^n$, such that $U - V$ does not have a compact connected component. Then every solution $v \in C^\infty(V)$ to the equation $Lv = 0$ is a limit in $C^\infty(V)$ of the restriction of solutions $u \in C^\infty(U)$ to the equation $Lu = 0$.*

Proof. By duality, it suffices to show that if w is a compactly-supported distribution in V , and

$$\int_V w(x)u(x) dx = 0$$

for any $u \in C^\infty(U)$ such that $Lu = 0$, then for any $v \in C^\infty(V)$ with $Lv = 0$,

$$\int_V w(x)v(x) dx = 0.$$

If L^* is the adjoint of the operator L , and we set $\Psi(x) = \Phi(-x)$, then Ψ is an analytic fundamental solution to L^* . If we can show that $\Psi * w$ is a compactly supported distribution in V , then for any $v \in C^\infty(V)$ with $Lv = 0$,

$$\int w(x)v(x) dx = \int L^*(\Psi * w)(x)v(x) dx = \int (\Psi * w)(x)Lv(x) dx = 0.$$

This would therefore complete the proof.

Let K denote the support of w . The distribution $\Psi * w$ is analytic function outside of K . Moreover, the support of $\Psi * w$ is certainly contained in U , because if $x \notin U$, then the function $u(y) = \Psi(x - y)$ satisfies $Lu = 0$, and therefore by assumption on w , $(\Psi * w)(x) = 0$. Applying analyticity, it follows that $\Psi * w$ vanishes in every component of K^c containing a point in U^c . We claim this is sufficient to prove the result. If W is a bounded connected component of K^c contained in U , then the set $W \cap V^c$ is a compact connected component of $U - V$. This contradicts the hypothesis of the Lemma, and therefore W cannot exist. Because K^c has at most one unbounded connected component, we conclude that the theorem is proved, except in the case where U contains the unbounded connected component of K^c . We will address this case in two paragraphs. For now, note that this proves the theorem when U is a open ball.

Next, we address the case where $U = \mathbf{R}^d$, and V is a ball B , without loss of generality, centred at the origin. This is because if $v \in C^\infty(B)$ solves $Lv = 0$, then for each k , we can recursively find a sequence $u_{k,i} \in C^\infty(kB)$, such that $Lu_{k,i} = 0$, and such that $u_{1,i}$ converges to v in $C^\infty(B)$ as $i \rightarrow \infty$, such that $\|u_{k+1,i} - u_{k,i}\|_{L^\infty(kB)} \leq 1/2^{k+i}$ for any $|\alpha| \leq k+i$. It follows that we can define $u_i \in C^\infty(\mathbf{R}^d)$ as the pointwise limit of the functions $u_{k,i}$ (smooth because of analyticity). Then $Lu_i = 0$, and $u_i \rightarrow v$ in $C^\infty(B)$.

Now returning to the general case of the proof, if K is contained in a ball B of radius R , then for $|x| > R$, the function $u_x(y) = \Phi(y - x)$ solves

the equation $Lu = 0$ in B , and thus is the limit in $C^\infty(B)$ of functions u_i solving $Lu_i = 0$ in $C^\infty(\mathbf{R}^n)$. But then

$$(\Phi * w)(x) = \int w(x)u(x) dx = \lim_{i \rightarrow \infty} \int w(x)u_i(x) dx = 0,$$

which completes the proof. \square

Remark. This result is *only* possible given the topological hypothesis. Suppose the result is true, and K is a compact connected component of $U - V$. If $u_i \in C^\infty(U)$ is a sequence of solutions to the equation $Lu_i = 0$ such that u_i converges to a function $u \in C^\infty(V)$, then we claim that u_i converges to some $u \in C^\infty(V \cup K)$. In particular, we conclude every solution on $C^\infty(V)$ extends to a solution on $C^\infty(V \cup K)$. This causes a contradiction, since if $x_0 \in K$, then the function $u(y) = \Phi(y - x_0)$ solves the equation on V , but cannot be extended to a solution on $V \cup K$ (e.g. because any such solution would be analytic).

Let us now prove that u can be extended under our assumptions. If we consider $\phi \in \mathcal{D}(V \cup K)$ equal to one on a neighborhood W of K , then $L(\phi u_i)$ vanishes on W , and hence is supported in V . Since u_i converges to u in V , so $L(\phi u_i)$ converges to Lu in $V \cup K$, and this means that $u_i = \Phi * L(\phi u_i)$ converges in $C^\infty(V \cup K)$ to some function u in $C^\infty(V \cup K)$, which completes the proof of our claim.

Using Runge's approximation theorem, we can now finally prove our first major existence theorem for partial differential equations on proper subsets U of \mathbf{R}^n , assuming the existence of a fundamental solution (i.e. the existence of solutions to the equation $Lu = f$ for compactly supported distributions f on \mathbf{R}^n). To obtain the idea of the proof of the existence of solutions to the equation $Lu = f$, for a distribution f on U (not necessarily compactly supported), imagine that such a solution u existed. For any $\phi \in \mathcal{D}(U)$ equal to one on some open set W , we can find a distribution u_W such that $L(u_W) = \phi f$. Then $L(u - u_W) = 0$ on W , which implies that $u - u_W$ is analytic on W . If $u_V - u_W = 0$ on $V \cap W$ for any $V, W \subset U$, these solutions would combine to give us a choice of u on the whole domain. Instead, we only have $u_V - u_W$ *analytic* on $V \cap W$. The Runge approximation theorem will enable us to adjust these functions so that $u_V - u_W$ is *almost* equal to zero, and then we can take limits.

Theorem 11.12. *Suppose Φ is a fundamental solution to a constant coefficient partial differential operator L on \mathbf{R}^n , and suppose that Φ is real analytic away from the origin. Then for any open set U in \mathbf{R}^n , and any distribution $f \in \mathcal{D}(U)^*$, then there exists $u \in \mathcal{D}(U)^*$ such that $Lu = f$.*

Proof. Let $U_R = \{x \in U : |x| < R \text{ and } d(x, U^c) > 1/R\}$. Then we can apply Runge's approximation theorem for $U_R \subset U$. We leave this proof of this to the end for it is fairly technical and unenlightening. Now if $\phi_i \in \mathcal{D}(U)$ equals one on U_i , we set $u_i = \Phi * (\phi_i f)$. Then $Lu_i = \phi_i f$, in particular, $Lu_i = f$ on U_i . It suffices to modify u_i so that it converges distributionally to some function u , since it will then follow that $Lu = f$ on U . we note that $L(u_{i+1} - u_i) = 0$ in U_i . Thus Runge's theorem implies that there exists an analytic function w_i defined on U such that $\|(u_{i+1} - u_i) - w_i\|_{L^\infty(U_i)} \leq 1/2^i$. But then if we define $v_i = u_i - w_1 - \cdots - w_{i-1}$, then $Lv_i = \phi_i f$ on U , and

$$\|v_{i+1} - v_i\|_{L^\infty(U_i)} = \|(u_{i+1} - u_i) - w_i\|_{L^\infty(U_i)} \leq 1/2^i,$$

which implies the required convergence property.

To show that $U - U_R$ has no compact connected components, consider a potential compact connected component K of $U - U_R$. If B_R is the ball of radius R centred at the origin, $K \cap B_R^c$ is compact. It is also open in B_R^c , because we can write $K = V \cap U \cap U_R^c$ for some open set V , and then $K \cap B_R^c = V \cap U \cap U_R^c \cap B_R^c = V \cap U \cap B_R^c$ since $U_R \subset B_R$, and U and V are both open. But this means that $K \cap B_R^c = \emptyset$, i.e. $K \subset B_R$. Next, we claim $d(x, y) > 1/R$ for any $y \in U^c$. Otherwise, take y with $d(x, y) \leq 1/R$, and consider the line segment L between x and y . Since U^c is closed, we may assume without loss of generality (by picking the closest point in U^c to x on the line and replacing y with it), that $L \cap U^c = \{y\}$. Since $x \notin U_R$, and $d(x_1, y) < 1/R$ for any other point x_1 on the line, $L \cap U_R = \emptyset$. Thus $L - \{y\} \subset U - U_R$. But $L - \{y\}$ is connected and contains $x \in K$, so $L - \{y\} \subset K$. But this is impossible for it implies that ∂K contains y , which is not an element of K , so K cannot be closed. Put together, this means that for any $x \in K$, $d(x, U^c) = 1/R$. But then it is impossible for K to be an open subset of $U - U_R$, so K cannot exist. \square

Example. Suppose $L = \sum_{|\alpha|} c_\alpha D^\alpha = P(D)$ is a homogeneous elliptic partial differential equation of degree m on \mathbf{R}^n . Then $1/P(\xi)$ is a homogeneous function defined on $\mathbf{R}^n - \{0\}$, of degree $-m$, analytic in $\mathbf{R}^n - \{0\}$. Thus we have a fundamental solution Φ defined such that $\hat{\Phi} = 1/P$.

- If $n > m$, then L has a fundamental solution $\Phi = \Phi_0$ is homogeneous of degree $m - n$, and C^∞ in $\mathbf{R}^n - \{0\}$.
- If $n \leq m$, then L has a fundamental solution $\Phi = \Phi_0 - Q(x) \log |x|$, where Φ_0 is homogeneous of degree $m - n$ and

$$Q(x) = \frac{1}{(2\pi)^m (m - n)!} \int_{|\xi|=1} \frac{(2\pi i x \cdot \xi)^{m-n}}{P(\xi)} d\xi.$$

TODO: Technical FT Calculation in Hörmander Volume 1, Theorem 7.1.20.

In either case, Φ_0 is analytic in $\mathbf{R}^n - \{0\}$.

11.2 The Cauchy Problem

Recall the Cauchy problem: Given a linear differential operator $L = \sum c_\alpha D^\alpha$ of order m , defined on an open set $\Omega_0 \subset \mathbf{R}^{n+1}$, with coefficients in $C^\infty(\Omega_0)$. We suppose that Ω_0 is divided into two parts by a smooth hypersurface S . Thus Ω_0 is divided into the union of three disjoint set $\Omega_0^- \cup S \cup \Omega_0^+$, where Ω_0^- and Ω_0^+ are connected, open sets.

Classically, the goal of the Cauchy problem, given some data f on Ω_0 , and $\phi_0, \dots, \phi_{m-1}$ on S , is to find a solution $Lu = f$, such that the first $m - 1$ derivatives of u in the direction tangent to S agrees with the functions $\{\phi_i\}$. For smooth data this is often a fine way to pose the problem; but for more general data. Instead, we consider $f \in \mathcal{D}(\Omega_0)$, and

Consider a linear differential operator $L = \sum c_\alpha D^\alpha$ of order m , defined on an open set. Finally, we consider some $f \in \mathcal{D}(\Omega_0)^*$.

For rough data, the best way to formulate the Cauchy problem is as follows: We consider some initial data $\phi \in \mathcal{D}$

Cauchy problem for this data is, given some set $X \subset \mathcal{D}(\Omega_0)^*$, to find $u \in X$ such that $Lu = f$.

11.2.1 First Order Operators

TODO

11.2.2 Calderon's Uniqueness Theorem

Let us discuss the phenomenon of *uniqueness* for the Cauchy problem. Thus our goal is to determine, for a given set $X \subset \mathcal{D}(\Omega_0)^*$, whether $L : X \rightarrow \mathcal{D}(\Omega_0)^*$ is injective. If X is a linear subspace of $\mathcal{D}(\Omega_0)^*$, it suffices to show that if $u \in X$, and $Lu = 0$, then $u = 0$. This simplifies the notation in the problem. To make the problem possible to study microlocally, we work locally. Thus we focus on the case where X solely contains functions which vanish on Ω_0^- , and will try and find conditions which guarantee that if $Lu = 0$, then u vanishes on a neighborhood of S .

Let $P(x, \xi)$ denote the principal symbol of L . We begin by assuming that S is *non characteristic* at any of its points, i.e. that if $x_0 \in S$, and $\eta_0 \in T_{x_0}^* \mathbf{R}^{n+1} - \{0\}$ has $T_{x_0} S$ as its kernel (conormal to S), then $P(x_0, \eta_0) \neq 0$. We actually assume S is *strongly non characteristic*, i.e. that for any $\eta_0 \in T_{x_0}^* \mathbf{R}^{n+1}$ conormal to S , and any $\eta_1 \in T_{x_0}^* \mathbf{R}^{n+1}$ not conormal to S , the polynomial $P(x_0, \eta_0 + z\eta_1)$ has m simple roots in the z -variable. For $m = 1$, this is equivalent to being noncharacteristic. For $m > 1$, the surface S can be noncharacteristic but not satisfy this property, for instance, if S is the x -axis, and $L = (D_x + iD_y)^2 = D_x^2 + 2iD_xD_y - D_y^2$.

Being strongly noncharacteristic is a *local* conditions on L and of S ; if we replace S with any other surface tangent to S at x_0 , the condition will continue to hold in a neighborhood of x_0 . Thus, working locally, we may switch to a coordinate system in which it suffices to study a more familiar version of the Cauchy problem, namely, we consider a set $\Omega_0 = (-T, T) \times \Omega$, where Ω is a connected, open subset of \mathbf{R}^n , $T > 0$, where $S = \{0\} \times \Omega$, and where

$$\Omega^- = \{(t, x) \in (-T, T) \times \Omega : t < 0\} \quad \text{and} \quad \Omega^+ = \{(t, x) \in (-T, T) \times \Omega : t > 0\}.$$

The principal symbol will then be written in the (t, x) variables, and their dual variables (τ, ξ) variables, i.e. as the polynomial $P(t, x, \tau, \xi)$. The assumption slightly stronger than being noncharacteristic then locally amounts to the fact that for any $\xi \in \mathbf{R}^n - \{0\}$, and any $z \in \mathbf{C}$, if $P(t, x, z, \xi) = 0$, then $\partial_z P(t, x, z, \xi) \neq 0$. It follows that locally on Ω we can write the solutions to $P(t, x, \tau, \xi) = 0$ in the τ variable for $1 \leq i \leq m$ as the smooth functions $\tau_1, \dots, \tau_m : (-T, T) \times \Omega_x \times (\mathbf{R}_\xi^n - \{0\}) \rightarrow \mathbf{C}$. Given that P is homogeneous in the ξ variable of degree m , τ_1, \dots, τ_m will be homogeneous of degree one.

Example. Let $L = \partial_t^2 + \Delta_x$. Then L has principal symbol $-4\pi^2(\tau^2 + |\xi|^2)$, and has roots $\pm i|\xi|$. The principal symbol factors as $-4\pi^2(\tau - i|\xi|)(\tau + i|\xi|)$.

Since P is a homogeneous polynomial of degree m , the coefficient of τ^n in the expansion of P must be a non-vanishing, C^∞ function of P . If it vanished at a point (t_0, x_0) , then we would have $P(t_0, x_0, \tau, 0) = 0$ for all $\tau \in \mathbf{R}$, which contradicts the fact that L is strongly noncharacteristic since then $\partial_\tau P(t_0, x_0, \tau, 0) = 0$. If $a \in C^\infty((-T, T) \times \Omega)$ is the coefficient of τ^n , it is therefore clear that the solutions of the operator $Lu = 0$ are the same as solutions of the operator $a^{-1} \cdot Lu = 0$. Thus in the sequel, we will assume without loss of generality that $a(t, x) = 1$ for all $t \in (-T, T)$ and all $x \in \Omega$.

Lemma 11.13. *Suppose that L is a differential operator of order m on $(-T, T) \times \Omega$, such that $S = \{(t, x) : t = 0\}$ is strongly noncharacteristic with respect to L . Then there exists m classical pseudodifferential operators Z_1, \dots, Z_m , each of order one, with principal symbols z_1, \dots, z_m such that*

$$L = (D_t - Z_1) \dots (D_t - Z_m) + R,$$

where R is a smoothing operator.

Proof. The composition calculus shows that $L - (D_t - z_1(t, x, D_x)) \dots (D_t - z_m(t, x, D_x))$ is a pseudodifferential operator R_1 of order $m - 1$. As with many regularizing calculi. We proceed asymptotically, like in the construction of parametrices. Given a choice of operators $\{Z_i^N\}$ such that $R_N = L - (D_t - Z_1^N) \dots (D_t - Z_m^N)$ is order $m - N$, we must consider operators W_1, \dots, W_m of order $1 - N$ such that $R_{N+1} = L - (D_t - Z_1^N - W_1) \dots (D_t - Z_m^N - W_m)$ is an operator of order $m - N - 1$. If w_1, \dots, w_n are the symbols of W_1, \dots, W_m , then it clearly suffices to choose these symbols such that if r_N is the principal symbol of R_N , then

$$r_N(t, x, \tau, \xi) = \sum_{i=1}^n w_i \left[\prod_{j \neq i} (\tau - z_j) \right].$$

If we pick

$$w_i = \frac{r_N}{\partial_z P(t, x, z_i, \xi)} = \frac{r_N}{\prod_{j \neq i} (z_i - z_j)},$$

then w_k is homogeneous of degree $m - N - (m - 1) = 1 - N$, and since the rational function

$$f(\tau) = \sum_{i=1}^n \prod_{j \neq i} \frac{\tau - z_j}{z_i - z_j}$$

is a degree $m - 1$ polynomial with $f(z_1) = \cdots = f(z_m) = 1$, and so $f(\tau) = 1$ for all $\tau \in \mathbf{R}$, which implies the required formula for r_N . Now we just take $Z_i = \lim Z_i^N$, and we get the required result. \square

The next remark shows that we can assume that there exists a compact set K' such that $\text{supp}_x(u) \subset K'$, such that the coefficients of L vanish outside of K' , and the (x, y) support of the kernels of Z_1, \dots, Z_m , and of R vanish outside of K' , uniformly in t .

Remark. Switch to a coordinate system in which $S = \{(t, x) \in (-T, T) \times \Omega : t = |x|^2\}$, after possibly shrinking Ω . In this coordinate system, if u vanishes on Ω_0^- , then $u(t, x)$ vanishes for $t < |x|^2$. Thus, after shrinking T , we may assume that Ω contains a ball of radius T^2 , and then $\text{supp}_x(u)$ is contained in that ball of radius T^2 . Switching back to our original coordinate system, we may assume without loss of generality that there exists a compact set $K \subset \Omega$ such that $\text{supp}_x(u) \subset K$. If $g \in C_c^\infty(\Omega)$ is equal to one on a neighborhood of K , then $0 = Lu = g \cdot Lu = (D_t - gZ_1) \cdots (D_t - gZ_m) + R^\# u$, which is a similar expansion to the original except that we can now assume that the (x, y) supports of the kernels of all operators involved are contained in $K' \times K'$ for some compact set K' .

Define $u_1 = u$, and for $1 \leq i \leq m - 1$, iteratively define

$$u_{i+1} = (D_t - Z_i)u_i.$$

Then since $L = (D_1 - Z_1) \cdots (D_n - Z_m)u + R$, it follows that

$$(D_t - Z_m)u_m = Lu - Ru = -Ru.$$

We can then consider the diagonal matrix valued pseudodifferential operator

$$A = i \cdot \text{diag}(Z_1, \dots, Z_m).$$

Let N be the standard nilpotent matrix, with ones directly above the diagonal. Let $\tilde{R}(t)$ be the $m \times m$ equal to $R(t)$ in the lower left corner. Then if $U = (u_1, \dots, u_m)$, then

$$\mathcal{L}U = \frac{\partial U}{\partial t} - A(t)U - i[N - \tilde{R}(t)]U = 0.$$

Let us now assume that $u \in C^i((-T, T), H_c^{m-j}(\Omega))$. Then $u_k \in C^j((-T, T), H_c^{m-k+1-j})$. Thus for $j = 0$ and $j = 1$, $U \in C^j((-T, T), H_c^{1-j}(\Omega) \otimes \mathbf{C}^m)$.

The uniqueness will now follow from an *inequality of Carleman type*. Namely, if $V(t, x) = \chi(t)U(t, x)$, where $\chi(t) = 1$ for $t < 8T/10$, and vanishes for $t > 9T/10$, then for all suitably large positive values $\rho > 0$,

$$\rho \int_0^T \int_{\Omega} e^{\rho(T-t)^2} |V(t, x)|^2 dx dt \lesssim \int e^{\rho(T-t)^2} |\mathcal{L}V(t, x)|^2 dx dt.$$

We then claim that U vanishes for $t < T/2$. Indeed, we can restrict the integration on the right hand side to the region $8T/10 \leq t \leq T$, because $\mathcal{L}U$ because it satisfies the ODE. But if we limit the integration on the left hand side to $0 \leq t \leq t/2$, we get that

$$\rho \cdot e^{\rho(24/25)T^2} \int_0^{t/2} \int_{\Omega} |V(t, x)|^2 dx dt \lesssim \int_{8T/10}^T \int_{\Omega} |\mathcal{L}V|^2 dx dt.$$

Unless the right hand side is equal to zero, it will become arbitrarily large as $\rho \rightarrow \infty$, which eventually gives a contradiction. Thus V must vanish on $0 \leq t \leq T/2$.

Remark. Note that we did not use ρ in the proof. It's utility is that if ρ is suitably large, then we can replace \mathcal{L} by the diagonal part $\mathcal{L}_0 = I_m(\partial/\partial t) - A(t)$, and thus discard the other terms as extraneous.

Calderon proved this result (Uniqueness in the Cauchy problem of partial differential equations, 1958) under two assumptions:

- The roots z_1, \dots, z_m are all valued in $\mathbf{C} - \mathbf{R}$, i.e. so that P is elliptic.
- The roots z_1, \dots, z_m are all valued in \mathbf{R} . Under this condition, we say that z_1, \dots, z_m are *strongly hyperbolic*.

TODO: Rest of argument (Treves, Vol 1. Starting on page 112).

Chapter 12

Symbol Classes

In various settings in harmonic analysis, especially generalizations of settings where *homogeneous functions* are the classical objects of study, it is useful to study various *symbol classes*. For instance, pseudodifferential operators historically dealt with operators $a(x, D)$, where $a : \mathbf{R}_x^d \times \mathbf{R}_\xi^d \rightarrow \mathbf{C}$ is a smooth function defined by a finite, or asymptotic sum of homogeneous functions of various orders in the ξ variable. If the highest degree of the terms in the sum was t , then for any n and m , $\nabla_x^n \nabla_\xi^m a$ is also a sum of homogeneous functions, with highest degree $t - m$. Thus we have bounds of the form

$$|\nabla_x^n \nabla_\theta^m a(x, \xi)| \lesssim \langle \xi \rangle^{t-m}.$$

Given a quantity t , a non-negative integer p , and an open subset $\Omega \subset \mathbf{R}^d$, we define the symbol class $\mathcal{S}^t(\Omega \times \mathbf{R}^p)$, consisting of *symbols of order t* , as the family of all functions $a \in C^\infty(\Omega_x \times \mathbf{R}_\theta^p)$ such that

$$|\nabla_x^n \nabla_\theta^m a(x, \theta)| \lesssim_{n,m} \langle \theta \rangle^{t-m}.$$

where the implicit constant is uniform in x . We take the optimal constants in these inequalities as a family of seminorms which gives $\mathcal{S}^t(\Omega \times \mathbf{R}^p)$ the structure of a Frechét space. Similar to other function spaces, we can also consider the local symbol classes $\mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^p)$.

The classes $\mathcal{S}^t(\Omega \times \mathbf{R}^p)$ are decreasing as $t \rightarrow -\infty$, and we define $\mathcal{S}^{-\infty}(\Omega \times \mathbf{R}^p)$ to be the intersection of all these classes of symbols. Operators defined by such functions are often highly regular. For instance, a pseudodifferential operator defined by such a symbol is called a *smoothing operator*, and maps any compactly supported distribution to a smooth function. The

class $\mathcal{S}^{-\infty}(\Omega \times \mathbf{R}^p)$ is dense in any of the classes $\mathcal{S}^t(\Omega \times \mathbf{R}^p)$, since it contains any symbol compactly supported in θ , and we can take cutoffs as $\theta \rightarrow \infty$.

A useful strategy to understand a symbol is to break it down into an asymptotic series of simpler symbols. Suppose $\{a_n\}$ is a sequence of symbols, then we write

$$a \sim \sum_{n=0}^{\infty} a_n$$

for some symbol a , if for any $t \in \mathbf{R}$, there exists N_0 such that for $N \geq N_0$, $a - \sum_{n=0}^N a_n$ is a symbol of order t . If a_n is a symbol of order t_n , and $\lim_{n \rightarrow \infty} t_n = -\infty$, then a symbol a always exists satisfying these asymptotics.

Theorem 12.1. *Consider a sequence of symbols $\{a_n\}$, with $a_n \in \mathcal{S}^{t_n}(\Omega \times \mathbf{R}^p)$, where $\lim_{n \rightarrow \infty} a_n = -\infty$, and let $t = \max t_n$. Then there exists a symbol $a \in \mathcal{S}^t(\Omega \times \mathbf{R}^p)$ such that $a \sim \sum a_n$.*

Proof. Fix a bump function $\phi \in \mathcal{D}(\mathbf{R}^p)$ equal to 0 when $|x| \leq 1/2$, and equal to one when $|x| \geq 1$. Find a rapidly increasing sequence $\{r_n\}$ such that

$$|\nabla_x^j \nabla_\lambda^k \{\phi(\theta/r_n) a_n(x, \theta)\}| \leq 2^{-n} \langle \theta \rangle^{t_n+1-k}$$

for $x \in \Omega$, where $i, j \leq n$. We define

$$a(x, \theta) = \sum_{n=0}^{\infty} \phi(\theta/r_n) \cdot a_n(x, \theta),$$

which is smooth, since it is a locally finite sum. For any N , if we set

$$R_N(x, \theta) = \sum_{n=N}^{\infty} \phi(\theta/r_n) \cdot a_n(x, \theta),$$

then

$$a - \sum_{n=0}^{N-1} a_n = \sum_{n=0}^{N-1} (\phi(\theta/r_n) - 1) a_n(x, \theta) + R_N(x, \theta)$$

If $x \in \Omega$, we find that

$$|\nabla_x^j \nabla_\lambda^k R_N(x, \theta)| \lesssim_{N,i,j} \langle \theta \rangle^{\max_{n \geq N} t_n + 1 - k}.$$

Thus $R_N \in \mathcal{S}^{\beta_N}(\Omega \times \mathbf{R}^p)$, where $\beta_N = \max_{n \geq N} t_n + 1$. On the other hand,

$$E_N(x, \theta) = \sum_{n=0}^{N-1} (\phi(\theta/r_n) - 1) a_n(x, \theta)$$

vanishes for $|\theta| \geq r_n$, and is thus compactly supported in θ , which implies that $E_N \in \mathcal{S}^{-\infty}(\Omega \times \mathbf{R}^p)$. \square

Remark. A similar formula holds for local families of symbols.

To verify asymptotic formulae, the following Lemma is often helpful.

Lemma 12.2. *Suppose $a \in C^\infty(\Omega \times \mathbf{R}^p)$, and for any $n, m > 0$, there exists t_{nm} such that*

$$|\nabla_x^n \nabla_\theta^m a(x, \theta)| \lesssim_{n,m} \langle \theta \rangle^{t_{nm}}.$$

If, for any $t \in \mathbf{R}$,

$$|a(x, \theta)| \lesssim_t \langle \theta \rangle^t,$$

then $a \in \mathcal{S}^{-\infty}(\Omega \times \mathbf{R}^p)$.

Proof. We begin by showing that if $f \in C^2(\mathbf{R})$, $\|f\|_{L^\infty(\mathbf{R})} \leq A$, and $\|f''\|_{L^\infty(\mathbf{R})} \leq B$, then $\|f'\|_{L^\infty(\mathbf{R})} \leq \sqrt{2AB}$. this follows because for any x , and $\varepsilon > 0$, there exists θ_1 lying between x and $x - \varepsilon$ such that

$$f(x) - f(x - \varepsilon) = \varepsilon f'(\theta_1) + \varepsilon^2 f''(\theta_1)/2$$

and θ_2 lying between x and $x + \varepsilon$ such that

$$f(x + \varepsilon) - f(x) = \varepsilon f'(\theta_2) + \varepsilon^2 f''(\theta_2)/2.$$

Thus

$$f(x + \varepsilon) - f(x - \varepsilon) = 2\varepsilon f'(\theta) + \varepsilon^2/2(f''(\theta_1) + f''(\theta_2)).$$

Rearranging gives

$$f'(\theta) = (f(x + \varepsilon) - f(x - \varepsilon))/2\varepsilon - (\varepsilon/4)(f''(\theta_1) + f''(\theta_2)),$$

and thus

$$|f'(\theta)| \leq A/\varepsilon + B\varepsilon/2.$$

Taking $\varepsilon = \sqrt{2A/B}$ completes the proof.

It follows from this that if K and K' are compact sets, with K contained in the interior of K' , then

$$\|\nabla_\theta \phi\|_{L^\infty(K)} \lesssim_K \sqrt{\|\phi\|_{L^\infty(K')} \|\nabla_\theta^2 \phi\|_{L^\infty(K')}}.$$

The theorem then follows by successively differentiating in θ . \square

Corollary 12.3. Suppose $\{a_n\}$ are a family of symbols, with $a_n \in \mathcal{S}^{t_n}(\Omega \times \mathbf{R}^p)$ for each n , and $\lim_{n \rightarrow \infty} t_n = -\infty$. Then if $a \in C^\infty(\Omega \times \mathbf{R}^p)$, and for each N and M , there exists t_{NM} such that

$$|\nabla_x^N \nabla_\theta^M a(x, \theta)| \lesssim \langle \theta \rangle^{t_{NM}}.$$

If for each n , there exists β_n such that

$$|a(x, \theta) - \sum_{k=0}^n a_k(x, \theta)| \lesssim_n \langle \theta \rangle^{\beta_n},$$

and $\lim_{n \rightarrow \infty} \beta_n = -\infty$, then $a \sim \sum a_n$.

Sometimes one has to use more powerful notions of homogeneity than the simple decay estimates above. In this case, it is useful to focus on *classical symbols*, i.e. symbols which satisfy an asymptotic formula of the form

$$a(x, \theta) \sim \sum_{n=0}^{\infty} a_{t-n}(x, \theta),$$

where $a_{t-n} \in S^{\text{Re}(t)-n}(\Omega \times \mathbf{R}^p)$ is homogeneous of degree $t - n$ in the θ variables for suitably large θ (the symbol must be smooth, and so cannot be homogeneous for small θ unless it is a polynomial). We denote the class of such symbols of order t by $\mathcal{S}_{\text{cl}}^t(\Omega \times \mathbf{R}^p)$. Sometimes this class is also called the class of *polyhomogeneous symbols*, and denoted $\mathcal{S}_{\text{phg}(\Omega \times \mathbf{R}^p)}^t$. We can also consider polyhomogeneous symbols with non integer step sizes, i.e. the class $\mathcal{S}_{\text{phg}}^{t,h}(\Omega \times \mathbf{R}^p)$, i.e. those symbols that satisfy an asymptotic expansion of the form

$$a(x, \theta) \sim \sum_{n=0}^{\infty} a_{t-hn}(x, \theta)$$

where $a_{t-hn} \in S^{\text{Re}(t)-hn}(\Omega \times \mathbf{R}^p)$ is homogeneous of degree $t - hn$.

Remark. Let M be a manifold, and let E be a vector bundle over M . We can define the space $\mathcal{S}_{\text{loc}}^t(E)$ of symbols of order t on E to be the family of all scalar functions a on E which are symbols of order t in local coordinates. These have a very similar theory of the theory we have expounded above. In particular, one can consider asymptotic developments of symbols.

Chapter 13

Pseudodifferential Operators

The goal of this chapter is to define the calculus of pseudodifferential operators, a general family of operators which allows us to manipulate the spatial and frequential properties of functions simultaneously. It is impossible to do this completely pointwise because of the uncertainty principle, but one can do things *pseudolocally*, i.e. the position of the support in time and space is approximately preserved, up to a rapidly decaying error. Roughly speaking, we will define a family of operators $a(x, D)$, associated with functions $a(x, \xi)$, such that if the support of a function f is concentrated near a point x_0 , and the support of \hat{a} is concentrated near ξ_0 , then we will find $a(x, D)f \approx a(x_0, \xi_0)f$. Before we begin, let us consider some basic examples that allow us to control space or time exclusively, to get an idea of what we want out of such a theory.

The most basic spatial multipliers in analysis are the *position operators*, which are the family of operators $X^\alpha : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$, defined by setting

$$X^\alpha f(x) = x^\alpha f(x),$$

and the *momentum operators* $D^\alpha : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$, which is the Fourier multiplier

$$\widehat{D^\alpha f}(\xi) = \xi^\alpha \hat{f}(\xi).$$

Note that in this chapter, the operators $\{D^\alpha\}$ will be normalized as such, and thus differ from the usual differential operators, which we will here denote by ∂^α , by the constant $(2\pi i)^{-|\alpha|}$. For each $m \in C^\infty(\mathbf{R}^d)$ such that m and all of its derivatives are slowly increasing, we can define a bounded

operator $m(X) : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$ by setting

$$m(X)f(x) = m(x)f(x).$$

We can also define an operator $m(D) : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$ by setting

$$\widehat{m(D)f}(\xi) = m(\xi)\widehat{f}(\xi).$$

Thus we have found two homomorphisms from a ring of smooth functions on \mathbf{R}^d to the ring of bounded operators on $\mathcal{S}(\mathbf{R}^d)$.

The family of such operators is very useful in analysis, since families of functions are more amenable to intuition than families of operators, and so we can try and understand what features of the function m tell us about the resulting operators $m(X)$ and $m(D)$. For instance, a very important application of operators of the form $m(D)$ is to the study of *elliptic* differential operators with constant coefficients. Recall that a partial differential operator $L = \sum c_\alpha D^\alpha$ of degree k is *elliptic* if the homogeneous polynomial $\sum_{|\alpha|=k} c_\alpha \xi^\alpha$ is non-vanishing away from the origin.

Theorem 13.1. *If L is an elliptic partial differential operator on \mathbf{R}^d , with constant coefficients, then L has a fundamental solution, i.e. there exists a distribution $\Phi \in \mathcal{D}(\mathbf{R}^d)^*$ such that $L(\Phi) = \delta$.*

Proof. Suppose L has degree k , and write $L = \sum c_\alpha D^\alpha$ for some constants $\{c_\alpha\}$. Since L is elliptic, there exists $R > 0$ such that the polynomial $P(\xi) = \sum_\alpha c_\alpha \xi^\alpha$ satisfies $|P(\xi)| \sim |\xi|^k$ for $|\xi| \geq R$. If $\chi \in \mathcal{D}(\mathbf{R}^d)$ is chosen such that $\chi(\xi) = 1$ for $|\xi| \leq R$, and we define a distribution Φ_0 such that

$$\widehat{\Phi_0}(\xi) = \frac{(1 - \chi(\xi))}{P(\xi)}.$$

Then $\widehat{\Phi_0}$ is a smooth function with $|\widehat{\Phi_0}(\xi)| \lesssim \langle \xi \rangle^{-k}$ for all $\xi \in \mathbf{R}^d$, which means that $\widehat{\Phi_0}$ is a well defined tempered distribution, and thus Φ_0 is also a well defined tempered distribution. But then

$$\widehat{L\Phi_0} = 1 - \chi(\xi).$$

Since $\chi \in \mathcal{D}(\mathbf{R}^d)$, it follows by the Paley-Wiener theorem, taking the inverse Fourier transform that $L\Phi_0 = \delta - w$, where w is an entire analytic

function of at most polynomial increase. The Cauchy-Kovalevskaya theorem (i.e. solving the equation by expanding out power series) allows us to find an entire analytic function u of at most exponential increase such that $Lu = w$. Then $\Phi = \Phi_0 + u$ is a fundamental solution for L . \square

The theory of pseudodifferential operators was introduced primarily to generalize these kinds of constructions to elliptic linear partial differential equations with *non constant* coefficients. A (*left*) *parametrix* for a linear differential operator L with smooth coefficients on a domain Ω is an operator $S : \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)^*$ such that $1 - S \circ L$ is a *smoothing operator*. We think of S as given an ‘approximate inverse’ for the operator T . The existence of a regular parametrix for an elliptic linear differential operator, which will be justified by the theory of pseudodifferential operators, is quite important in the theory of differential equations. In particular, it proves that certain differential operators are *hypoelliptic*, i.e. that if $u \in \mathcal{D}(\Omega)$, then $\text{sing-supp}(Lu) = \text{sing-supp}(u)$; it suffices to show $\text{sing-supp}(u) \subset \text{sing-supp}(Lu)$, since the inclusion $\text{sing-supp}(Lu) \subset \text{sing-supp}(u)$ is true for any differential operator L with smooth coefficients.

Theorem 13.2. *Let L be a differential operator with smooth coefficients. If L has a very regular left parametrix S , then L is hypoelliptic.*

Proof. For any very regular operator S , $\text{sing-supp}(Su) \subset \text{sing-supp}(u)$. It suffices to prove that for any *compactly supported* distribution u , we have $\text{sing-supp}(u) \subset \text{sing-supp}(Lu)$, since the general case follows by localization. Since $1 - S \circ L$ is a smoothing operator, we have

$$\begin{aligned} \text{sing-supp}(u) &\subset \text{sing-supp}((1 - S \circ L)u) \cup \text{sing-supp}((S \circ L)u) \\ &\subset \text{sing-supp}((S \circ L)u) \\ &\subset \text{sing-supp}(Lu). \end{aligned} \quad \square$$

The family of pseudodifferential operators we will study are *microlocal*, i.e. not only do they not expand the singular support of distributions, but they also do not expand the wavefront set of distributions. It will therefore follow from the theory that any elliptic differential operator has a pseudodifferential parametrix, and an analogous argument to that given above gives the strong equation $\text{WF}(Lu) = \text{WF}(u)$ for any distribution u .

Returning to the general question of constructing a functional calculus which includes both the position and momentum operators, we recall the

spectral calculus, whose goal, for a suitable algebra of normal operators A , is to produce an isomorphism of A with an algebra of functions on some space X , called the *spectrum* of A . A natural hope would be to find such a calculus for an algebra A of operators which includes the position and momentum operators. This would, in particular, enable us to analyze linear differential operators with non-constant coefficients. However, we quickly see that such a theory would not quite work in as standard a way as the spectral calculus provides, because the families of operators $\{X^\alpha\}$ and $\{D^\alpha\}$ do *not* commute with one another, i.e. the chain rule implies that

$$[D^i, X^i] = D^i X^i - X^i D^i = 1.$$

The key thing we should notice from this equation, however, is that this equation indicates that position and momentum operators commute ‘up to lower order terms’. In other words, if we think of X^α and D^α as being operators of *order* $|\alpha|$, then $[D^\alpha, X^\beta]$ is equal to zero, *modulo terms of order* $|\alpha| + |\beta| - 1$. This fact will enable us to obtain an ‘approximate’ functional calculus for the desired algebra of operators. This is precisely the *calculus of pseudodifferential operators*.

We will associate, with each suitably regular function $a(x, \xi)$, an operator $a(x, D)$, which is a homomorphism ‘modulo lower order terms’. This association will have the property that if $a(x, \xi) = \sum c_\alpha(x) \xi^\alpha$, then $a(x, D)$ will be the differential operator $\sum c_\alpha(x) D^\alpha$. Indeed, this is where the notation $a(x, D)$ comes from. The association will also generalize the two families of multiplier operators; if $a(x, \xi) = m(x)$, then $a(x, D)$ is equal to $m(X)$, and if $a(x, \xi) = m(\xi)$, then $a(x, D)$ is equal to $m(D)$. To get an idea for what this operator should look like, we calculate that if $a(x, \xi) = \sum_\alpha c_\alpha(x) \xi^\alpha$ is the symbol of a differential operator with nonconstant coefficients, then the corresponding differential operator satisfies

$$\begin{aligned} a(x, D)f &= \sum c_\alpha(x) D^\alpha f(x) \\ &= \int_{\mathbf{R}^d} \sum_\alpha c_\alpha(x) \xi^\alpha \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi \\ &= \int_{\mathbf{R}^d} a(x, \xi) e^{2\pi i \xi \cdot x} \widehat{f}(\xi) d\xi. \end{aligned}$$

We use this integral formula to define $a(x, D)$ for a much more general family of functions $a(x, \xi)$.

Fix an open set $\Omega \subset \mathbf{R}^d$. Given any distribution $a(x, \xi)$ which is tempered in the ξ variable, i.e. any continuous, bilinear map $a : \mathcal{D}(\Omega_x) \times \mathcal{S}(\mathbf{R}_\xi^d) \rightarrow \mathbf{C}$, or more technically, any element of $\mathcal{D}(\Omega_x)^* \hat{\otimes} \mathcal{S}(\mathbf{R}_\xi^d)^*$, we can associate an operator $a(x, D) : \mathcal{D}(\Omega) \rightarrow \mathcal{D}(\Omega)^*$, such that for any $f, g \in \mathcal{D}(\Omega)$,

$$\langle a(x, D)f, g \rangle = \int a(x, \xi) \hat{f}(\xi) e^{2\pi i \xi \cdot x} g(x) dx d\xi.$$

We call any operator T which can be given in the form $a(x, D)$ a *pseudodifferential operator*. The symbol a is uniquely determined by the operator T , since the action of $a(x, D)$ on $\mathcal{D}(\Omega)$ determines the behaviour of a , viewed as a bilinear map, on an arbitrary element of $\mathcal{D}(\Omega_x) \times \mathcal{S}(\mathbf{R}_\xi^d)$. For any set $S \subset \mathcal{D}(\Omega_x)^* \hat{\otimes} \mathcal{S}(\mathbf{R}_\xi^d)^*$, the notation $\text{Op}(S)$ is often used to denote the family of all pseudodifferential operators defined by an element of S .

Example. Consider the Laplacian Δ on \mathbf{R}^n . Then Δ is the pseudodifferential operator on \mathbf{R}^n of order two, corresponding to the symbol $a(x, \xi) = -4\pi^2 |\xi|^2$. Since Δ is a constant coefficient operator, it just acts as a Fourier multiplier. If $\Delta u = f$, then, modulo a harmonic function, which is arbitrarily smooth, $u = \Phi * f$, where Φ is a fundamental solution to the Laplacian. The operator $f \mapsto \Phi * f$ is a pseudodifferential operator with symbol $b(x, \xi) = f.p(1/4\pi^2 |\xi|^2)$.

Example. The Cauchy-Riemann operator on \mathbf{R}^2 given by

$$\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$$

is a pseudodifferential operator of order one with symbol $i\pi(\xi + i\eta)$.

Any pseudodifferential operator $T = a(x, D)$ is continuous from $\mathcal{D}(\Omega)$ to $\mathcal{D}(\Omega)^*$, and has Schwartz kernel

$$K_a(x, y) = \int a(x, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi,$$

where in general the oscillatory integral must be interpreted formally. It is also useful to write this kernel in the convolution form $k_a(x, z) = K_a(x, x - z)$, because we then have

$$T\phi(x) = \int k_a(x, z) f(x - z) dz,$$

which reflects the fact that when $a(x, \xi)$ is independent of x , k_a is a function of z , and then $Tf = k_a * f$ is a convolution operator. In fact, all pseudodifferential operators are infinite sums of convolution operators, in the following sense: if $a \in \mathcal{D}(\mathbf{R}^d)^* \hat{\otimes} \mathcal{S}(\mathbf{R}^d)^*$, then we can write a as a sum of the form

$$\sum_{i=1}^{\infty} u_i \otimes v_i,$$

where $\{u_i\}$ are in $\mathcal{D}(\mathbf{R}^d)^*$, and $\{v_i\} \in \mathcal{S}(\mathbf{R}^d)^*$, and the convergence occurs unconditionally, in the distributional topology. It then follows that for $\phi \in \mathcal{D}(\Omega)$,

$$a(x, D)\phi = \sum_{i=1}^{\infty} u_i \cdot (v_i * \phi).$$

Thus $a(x, D)$ is, in some sense, a non-constant coefficient sum of convolution operators.

As we increase the regularity of a , we no longer need to treat the definition of a pseudodifferential operator quite as formally, and so we can define the operator on a more general family of functions. Here are some non-comprehensive examples of this phenomenon:

- If $a \in \mathcal{S}(\mathbf{R}^d \times \mathbf{R}^d)^*$, then $a(x, D)$ extends to a continuous linear operator from $\mathcal{S}(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^d)^*$. The Schwartz kernel theorem implies that any continuous linear operator from $\mathcal{S}(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^d)^*$ is of this form, which probably indicates that the family of such operators is too general to obtain interesting results.
- If $a \in \mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^d)$, then we will see later on that $a(x, D)$ extends to a continuous operator from $\mathcal{E}(\Omega)^*$ to $\mathcal{D}(\Omega)^*$ and from $\mathcal{D}(\Omega)$ to $\mathcal{E}(\Omega)$.
- If $a \in \mathcal{S}^t(\mathbf{R}^d \times \mathbf{R}^d)$, then $a(x, D)$ extends to a continuous operator from $\mathcal{S}(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^d)$.
- If $a \in \mathcal{S}_{\text{loc}}^{-\infty}(\Omega \times \mathbf{R}^d)$, then we will see later in this chapter that $a(x, D)$ has a kernel in $C^\infty(\mathbf{R}^d \times \mathbf{R}^d)$, and is therefore a smoothing operator, thus extending to a continuous operator from $\mathcal{E}(\mathbf{R}^d)^*$ to $\mathcal{E}(\mathbf{R}^d)$.

Conversely, we will also see that if T is *any* ‘pseudolocal’ smoothing operator, in the sense that it has a kernel $K \in C^\infty(\Omega \times \Omega)$ satisfying

bounds of the form

$$|\nabla_x^n \nabla_y^m K(x, y)| \lesssim_N \langle x - y \rangle^{-N},$$

then $T \in \text{Op}(\mathcal{S}_{\text{loc}}^{-\infty}(\Omega \times \mathbf{R}^d))$. In particular, any proper smoothing operator is of this form.

- If $a(x, D)$ is a proper operator, then it maps $\mathcal{D}(\Omega)$ into $\mathcal{E}(\Omega)^*$ and from $\mathcal{E}(\Omega)$ into $\mathcal{D}(\Omega)^*$. In combination with the previous results, we conclude that if $a \in \mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^d)$, then $a(x, D)$ is an operator from $\mathcal{D}(\Omega)^*$ to itself, $\mathcal{E}(\Omega)^*$ into itself, and from $\mathcal{D}(\Omega)$ into itself. If $a \in \mathcal{S}_{\text{loc}}^{-\infty}(\Omega \times \mathbf{R}^d)$, then $a(x, D)$ maps $\mathcal{D}(\Omega)$ to itself and maps $\mathcal{D}(\Omega)^*$ into $\mathcal{E}(\Omega)$.

Conversely, let $T : \mathcal{D}(\Omega) \rightarrow \mathcal{E}(\Omega)^*$ be *any* proper operator, and let K be it's kernel. Then K lies in $\mathcal{D}(\Omega_x)^* \hat{\otimes} \mathcal{E}(\Omega_y)^*$. Thus we can define a symbol $a \in \mathcal{D}(\Omega_x)^* \hat{\otimes} \mathcal{S}(\mathbf{R}_\xi^d)^*$ by setting

$$a(x, \xi) = \int K(x, y) e^{2\pi i \xi \cdot (x-y)} dy.$$

We verify using the Fourier multiplication formula that

$$T_a \phi(x) = \int a(x, \xi) \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi = \int K(x, y) f(y) dy = T \phi(x).$$

Thus *any* proper operator is a pseudodifferential operator.

That every proper operator, and that every operator on Schwartz space, is a pseudodifferential operator, indicates that the theory of pseudodifferential operators is too general to study more detailed in the form above. We will mostly focus on pseudodifferential operators defined by various symbol classes, since most practical operator occuring in PDE and analysis are of this form, and we can still get a sophisticated calculus.

13.1 Basic Definitions

There are two varieties of the theory of pseudodifferential operators, whose basic results are roughly analogous to one another. The first, which works

best when considering pseudodifferential operators on \mathbf{R}^d , works with operators specified by symbols $a : \mathbf{R}^d \times \mathbf{R}^d \rightarrow \mathbf{C}$ in $\mathcal{S}^t(\mathbf{R}^d \times \mathbf{R}^d)$, i.e. satisfying estimates of the form

$$|\nabla_x^n \nabla_\xi^m a(x, \xi)| \lesssim_{n,m} \langle \xi \rangle^{t-m}$$

where the bound holds uniformly in both x and ξ , for any integers n and m . As mentioned above, $a(x, D)$ then extends to a continuous operator from $\mathcal{S}(\mathbf{R}^d)$ to itself, which leads to an elegant theory. However, this theory is less easy to work with locally, e.g. working in various different coordinate systems, or obtaining a definition of pseudodifferential operator that applies to operators on manifolds. The approach here is best taken by describing a theory described by symbols $a \in \mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^d)$, i.e. satisfying an inequality of the form above uniformly in ξ , but only *locally uniformly* in x .

Fix an open set $\Omega \subset \mathbf{R}^d$, and consider a symbol $a \in \mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^d)$. From this symbol, we can define a continuous operator $T_a : \mathcal{D}(\Omega) \rightarrow \mathcal{E}(\Omega)$ by setting

$$T_a f(x) = \int a(x, \xi) e^{2\pi i \xi \cdot x} \hat{f}(\xi) d\xi,$$

where the integral can now be interpreted in the usual Riemann / Lebesgue sense, and $T_a f$ is smooth since $a \in C^\infty(\Omega_x, \mathcal{S}(\mathbf{R}_\xi^d)^*)$. We then say T_a is a *pseudodifferential operator of order t* . The family of all such operators is denoted $\Psi_{\text{loc}}^t(\Omega)$; we reserve the notation $\Psi^t(\Omega)$ to denote those operators defined by elements of $\mathcal{S}^t(\Omega \times \mathbf{R}^d)$.

The kernel of a pseudodifferential operator given by a symbol $a(x, \xi)$ of the class above is of the form

$$K_a(x, y) = \int a(x, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi,$$

where the integral is now an oscillatory integral distribution. In particular, we know from our discussion of oscillatory integral distributions that

$$\text{WF}(K_a) \subset \{(x, x; \xi, \xi) : x, \xi \in \mathbf{R}^d\}.$$

The microlocal analysis of distributions implies the existence of a continuous extension $T_a : \mathcal{E}(\Omega) \rightarrow \mathcal{D}(\Omega)^*$, and we find that $\text{WF}(T_a u) \subset \text{WF}(u)$. Thus the operator T_a preserves the location of singularities of a distribution

in both position and frequency. This is the first instance of the *microlocal nature* of pseudodifferential operators; these operators roughly preserve the location of the mass and frequency support of a function, but with some additional ‘fuzz’ that is usually negligible to the problem, but must be managed.

The symbol a is uniquely determined from the operator T_a . To actually recover the symbol from an operator, we have several methods. Formally, we can calculate that

$$a(x, \xi) = e^{-2\pi i \xi \cdot x} T_a(e^{2\pi i \xi \cdot y}).$$

The wavefront calculation above shows that the convolution kernel $k_a(x, z)$ of a pseudodifferential operator agrees with a smooth function away from the line $z = 0$. We will see very shortly that it decays rapidly away from this line, and therefore k_a is tempered in the z -variable. The above formal equation then implies the less formal equation

$$a(x, \xi) = \int k_a(x, z) e^{2\pi i \xi \cdot z} dy.$$

Thus the symbol a is obtained by taking the Fourier transform of the convolution kernel k_a in the z variable.

Here is a quantitative estimate on the kernel of a pseudodifferential operator, which show another instance of its pseudolocal nature, i.e. localization on the spatial side of things. In particular, the result implies that if $\Omega = \mathbf{R}^d$, then T_a extends to a continuous operator from $\mathcal{S}(\mathbf{R}^d)$ to itself. As mentioned above, $K_a \in \mathcal{D}(\Omega \times \Omega)^*$ agrees with a C^∞ function away from the diagonal Δ_Ω . Thus if $f \in \mathcal{D}(\mathbf{R}^d)$, and $x \notin \text{supp}(f)$, then the multiplication formula for tempered distributions implies that

$$T_a f(x) = \int a(x, \xi) \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi = \int K(x, y) f(y) dy,$$

where, since the integral on the right hand side vanishes in a neighborhood of x , we can actually interpret the right hand integral as a Lebesgue integral, rather than a formal integral. Moreover, we have even better estimates for the behaviour of K_a away from the origin, which reflects the pseudolocal behaviour of the operator. To discuss these estimates, we introduce the differential operators $\partial_z^i = \partial_x^i - \partial_y^i$, and the induced operators ∇_z^m , which measures the derivatives measured in the direction normal to the diagonal in $\Omega \times \Omega$.

Theorem 13.3. *Let $a \in \mathcal{S}_{loc}^t(\Omega)$. Then for any pair of integers $n, m \geq 0$, and any $N \geq 0$ such that $t + d + m + N \geq 0$,*

$$|\nabla_x^n \nabla_z^m K_a(x, y)| \lesssim_{n,m,N} |x - y|^{-t-d-m-N},$$

where the implicit constant is locally uniform in x . If $a \in \mathcal{S}^t(\Omega)$, then we can choose the implicit constant to be uniform in x .

Proof. If $\text{supp}_\xi(a)$ is compact, then $a \in \mathcal{S}_{loc}^{-\infty}(\Omega)$, and by the compactness, we conclude that for any $N \geq 0$,

$$|\nabla_x^n \nabla_z^m K_a(x, y)| = |\mathcal{F}_\xi \{a\}(x, x - y)| \lesssim_N 1/|x - y|^N,$$

where the uniform estimate happens because a is compactly supported in the ξ variable, uniformly in x , and smooth, locally uniformly in x . Thus, without loss of generality, in the remainder of the proof we may assume $a(x, \xi) = 0$ for $|\xi| \leq 1$. We can then perform a Littlewood-Paley decomposition, i.e. writing

$$a(x, \xi) = \sum_{i=0}^{\infty} a_i(x, y, \xi),$$

where $a_i(x, \xi) = \rho(\xi/2^i)a(x, \xi)$ is supported on $|\xi| \sim 2^i$. Let K_i be the kernel of the pseudodifferential operator $a_i(x, D)$. Then

$$K(x, z) = \sum_{i=0}^{\infty} K_i(x, z),$$

where the convergence is distributional. We claim that for any i , and any $n, m, N \geq 0$,

$$|\nabla_x^n \nabla_z^m K_i(x, y)| \lesssim_{n,m,N} |x - y|^{-N} 2^{i(t+d+m-N)},$$

where the implicit constant is locally uniform in x . This follows from a simple integration by parts, applied to the integral

$$K_i(x, y) = \int \rho(\xi/2^i) a(x, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi.$$

These bounds, if N is taken large enough, imply that the sum $K = \sum K_i$ converges uniformly on any set of the form

$$\{(x, y) \in \Omega \times \Omega : x \in K, |y - x| > \varepsilon\},$$

where $K \subset \Omega$ is compact. Summing up these bounds for sufficiently large N gives the required inequality for $|x - y| \geq 1$. For $0 < |x - y| \leq 1$, we break the sum into two parts, i.e. writing

$$K(x, y) = \sum_{2^i \leq 1/|x-y|} K_i(x, y) + \sum_{2^i > 1/|x-y|} K_i(x, y).$$

For the first sum, we take $N = 0$, and for the second sum, we take $N > t + d + m$, which gives the required bounds. \square

These singularity conditions characterize those kernels K_a induced by pseudodifferential operators of some order $t < 0$. For $t \geq 0$, we must also assume an additional cancellation condition.

Lemma 13.4. *Let $K \in \mathcal{D}(\Omega \times \Omega)^*$ be a Schwartz kernel with*

$$\text{sing-supp}(K) \subset \{(x, x) : x \in \Omega\}.$$

Suppose that for some t , and any non-negative integers n, m , and N , the kernel K satisfies the growth condition

$$|\nabla_x^n \nabla_z^m K(x, y)| \lesssim |x - y|^{-t-d-m-N}$$

locally uniformly in x . Then:

- *If $t < 0$, and if, for each $x \in \Omega$, and any $\phi \in \mathcal{D}(\Omega)$,*

$$\int K(x, y) \phi(y) dy = \lim_{\varepsilon \rightarrow 0} \int_{|x-y| > \varepsilon} K(x, y) \phi(y) dy,$$

where the right hand side exists, and defines a distribution induced by a locally integrable function on Ω by virtue of the growth condition with m and N equal to zero, then K is the Schwartz kernel of a pseudodifferential operator given by a local symbol of order t .

- *If $t \geq 0$, and any $\phi \in \mathcal{D}(\Omega)$, the kernel K satisfies the cancellation conditions*

$$\left| \int D_x^\alpha K(x, y) \phi((x - y)/R) dy \right| \lesssim_{\phi, \alpha} R^t,$$

where the implicit constant is independent of R , locally uniform in x , and a continuous seminorm on $\mathcal{D}(\Omega)$ for each multi-index α , then K is the kernel of some pseudodifferential operator given by a local symbol of order t .

If we replace the inequalities that locally uniformly depend on x with inequalities that are uniform in x , then the symbols we find can also be chosen to be uniform.

Proof. TODO: Ask Andreas about this. \square

In addition to studying the behaviour of Ψ DOs away from the diagonal, which reflects the pseudolocal behaviour of the distribution, it is also of interest to determine the behaviour of the operator under highly oscillatory, but non-stationary, phenomena, which is related to its microlocal nature. Consider a symbol $a(x, \xi)$, a smooth function $f(y)$, and a smooth phase $\phi(y)$ with $\nabla\phi(y)$ nonvanishing on $\text{supp}_x(a)$. Our goal is to try to determine the asymptotic behaviour of the function $T_a(fe^{2\pi i\lambda\phi})$ as $\lambda \rightarrow \infty$. Since T_a is pseudolocal, the value at a point x should be determined to a large degree by the behaviour of $fe^{2\pi i\lambda\phi}$ near x , which, roughly speaking, oscillates near the frequency $\lambda\nabla\phi(x)$. Thus we might expect that

$$T_a\{fe^{2\pi i\lambda\phi}\}(x) \approx a(x, \lambda\nabla\phi(x))f(x)e^{2\pi i\lambda\phi(x)}.$$

This is correct up to first order in λ , and in fact, we can obtain a complete asymptotic development as $\lambda \rightarrow \infty$. For simplicity, we assume $\text{supp}_x(a)$ is compact.

Theorem 13.5. Fix a symbol $a \in \mathcal{S}^t(\Omega \times \mathbf{R}^d)$, compactly supported in the x -variable, a smooth function $f \in \mathcal{D}(\Omega)$, and a smooth, real-valued function $\phi \in C^\infty(\Omega)$ with $\nabla\phi$ nonvanishing on $\text{supp}_x(a)$. Let

$$r_x(y) = \nabla\phi(x) \cdot (x - y) - (\phi(x) - \phi(y)).$$

Then for any $N > 0$ and $\lambda > 0$, we can write $e^{-2\pi i\lambda\phi(x)}T_a\{fe^{2\pi i\lambda\phi}\}(x)$ as

$$\sum_{|\beta| < N} \frac{1}{\beta! \cdot (2\pi i)^\beta} \cdot \partial_\xi^\beta a(x, \lambda\nabla\phi(x)) \partial_y^\beta \{e^{2\pi i\lambda r_x} f\} \Big|_{y=x} + R_N(x, \lambda),$$

where $\lambda^{t-[N/2]}R_N \in L^\infty(\mathbf{R}^d \times (0, \infty))$. In particular, for $N = 3$, we find that

$e^{-2\pi i \lambda \phi(x)} T_a \{f e^{2\pi i \lambda \phi}\}(x)$ is equal to

$$\begin{aligned} & a(x, \lambda \nabla \phi(x)) f(x) \\ & + \frac{1}{2\pi i} \sum_{k=1}^d (\partial_\xi^k a)(x, \lambda \nabla \phi(x)) \cdot (\partial_x^k f)(x) \\ & - \frac{i\lambda}{4\pi} \sum_{|\beta|=2} (\partial_\xi^\beta a)(x, \lambda \nabla \phi(x)) (\partial_x^\beta \phi)(x) f(x) \\ & + O(\lambda^{t-2}). \end{aligned}$$

If $\phi(x) = \xi \cdot x$ for some ξ , then we find

$$T_a \{f e^{2\pi i \lambda \phi}\}(x) \sim \sum_{\beta} \frac{1}{\beta! (2\pi i)^\beta} \partial_\xi^\beta a(x, \lambda \xi) \cdot \partial_x^\beta f(x).$$

Proof. We write

$$e^{-2\pi i \lambda \phi(x)} T_a \{f e^{2\pi i \lambda \phi}\}(x) = \lambda^d \int a(x, \lambda \xi) f(y) e^{2\pi i \lambda [(\xi - \nabla \phi(x)) \cdot (x-y) + r_x(y)]} dy d\xi.$$

This integral is oscillatory, with phase

$$\Phi_x(\xi, y) = (\xi - \nabla \phi(x)) \cdot (x - y) + r_x(y).$$

Now $\nabla_\xi \Phi_x(\xi, y) = 0$ precisely when $y = x$, and $\nabla_y \Phi_x(\xi, x) = 0$ precisely when $\xi = \nabla \phi(x)$. If we consider a smooth cutoff $\psi \in C_c^\infty(\mathbf{R}_\xi^d)$ supported on

$$(1/2) \cdot \inf_{x \in \text{supp}_x(a)} |\nabla \phi(x)| \leq |\xi| \leq 2 \cdot \sup_{x \in \text{supp}_x(a)} |\nabla \phi(x)|$$

then we can write $e^{-2\pi i \lambda \phi(x)} T_a \{f e^{2\pi i \lambda \phi}\}(x) = I_1 + I_2$, where I_1 is obtained by substituting ψ into the integrand, and I_2 is obtained by substituting $1 - \psi$ into the integrand. Nonstationary phase tells us that

$$|I_2| \lesssim_N \lambda^{-N}$$

for all $N > 0$, where the implicit constant is independent of λ . Thus it suffices to concentrate on I_1 . In the sequel, we will therefore assume without loss of generality that $a(x, \lambda \xi) = a(x, \lambda \xi) \psi(\xi)$, i.e. $a(x, \lambda \xi)$ is supported

near $|\xi| \sim 1$. A change of variables $\xi \mapsto \xi + \nabla\phi(x)$ allows us to rewrite I_1 as

$$\lambda^d \int e^{2\pi i \lambda(\xi \cdot (x-y) + r_x(y))} a(x, \lambda[\nabla\phi(x) + \xi]) f(y) d\xi dy.$$

Using Taylors formula, we write

$$a(x, \lambda\nabla\phi(x) + \lambda\xi) = \sum_{|\beta| < N} \frac{\lambda^\beta}{\beta!} \partial_\xi^\beta a(x, \lambda\nabla\phi(x)) \xi^\beta + R_{N,\lambda}(x, \xi),$$

where

$$R_{N,\lambda}(x, \xi) = \sum_{|\beta|=N} \xi^\beta \frac{N}{\beta!} \int_0^1 (1-t)^{N-1} \lambda^N \partial_\xi^\beta a(x, \lambda\nabla\phi(x) + t\lambda\xi) dt.$$

We only care about this formula when $|\xi| \sim 1$, and differentiating this formula gives that, for such ξ ,

$$|\partial_\xi^\alpha R_{N,\lambda}(x, \xi)| \lesssim_\alpha \lambda^{t-|\alpha|},$$

and $\partial_\xi^\alpha R_{N,\lambda}(x, 0) = 0$ for all $|\alpha| < N$. Stationary phase thus implies that

$$\left| \lambda^d \int \int e^{2\pi i \lambda((x-y) \cdot \xi + r_x(y))} R_{N,\lambda}(x, \xi) f(y) d\xi dy \right| \lesssim \lambda^{t-[N/2]}.$$

Finally, we note that via an integration by parts,

$$\begin{aligned} & \int e^{2\pi i \lambda(\xi \cdot (x-y) + r_x(y))} \xi^\beta f(y) d\xi dy \\ &= \lambda^{-(d+\beta)} (2\pi i)^{-\beta} \partial_y^\beta \{ e^{2\pi i \lambda r_x(y)} f(y) \} \Big|_{y=x} \end{aligned}$$

and substituting them into the formula completes the proof. \square

Remark. If T is a pseudodifferential operator on $\Omega \subset \mathbf{R}^d$ defined by a symbol a with $\text{supp}_x(a)$ compact, and $\kappa : \Psi \rightarrow \Omega$ is a diffeomorphism, then we can consider the operator $S : \mathcal{D}(\Psi) \rightarrow \mathcal{D}(\Psi)^*$ given by ‘changing the coordinates of T ’, i.e.

$$S\phi(x) = T(\phi \circ \kappa^{-1})(\kappa(x)).$$

We can thus write, after a change of coordinates, with $\tilde{a}(x, \xi) = a(\kappa(x), \xi)$,

$$S\phi(x) = \int \tilde{a}(x, \xi) e^{2\pi i \xi \cdot (\kappa(x) - \kappa(y))} |\det(\kappa(y))| \phi(y) dy d\xi.$$

This leads to an analysis of quantities similar to that obtained in the above proof, which will lead us to conclude that S itself is a pseudodifferential operator. Note, however, that purely from the spectral analysis of singularities, we see immediately from the integral kernel that S is a microlocal operator, i.e. it preserves the wavefront set of distributional inputs.

This calculation will also prove handy when understanding the composition of a pseudodifferential operator with a *Fourier integral operator*.

As the order of the symbol a decreases, we expect the behaviour of the corresponding pseudodifferential operator to become more and more regular. In particular, for $t < -d$, the symbol is actually *integrable* in ξ . A Ψ DO of order $-\infty$ has a kernel K lying in $C^\infty(\Omega \times \Omega)$, and satisfying estimates of the form

$$|\nabla_x^n \nabla_y^m K(x, y)| \lesssim_{n,m,N} \langle x - y \rangle^{-N}$$

for any $N \geq 0$. Thus a Ψ DO of order $-\infty$ is smoothing. Conversely, if $K \in C^\infty(\Omega \times \Omega)$ satisfies estimates of the form above, then it is the Ψ DO corresponding to the symbol

$$a(x, \xi) = \int K(x, y) e^{2\pi i \xi \cdot (x - y)} dy,$$

which is a symbol of order $-\infty$. In particular, all properly supported smoothing operators are Ψ DOs of order $-\infty$.

Much of the theory of pseudodifferential operators to come is most elegantly explained *modulo smoothing operators*. Working modulo smoothing operators is usually fine in harmonic analysis provided that we are trying to establish localized estimates for certain objects, since smoothing operators, once localized, satisfy all the inequalities we might be interested in. In particular, working modulo smoothing operators often makes the theory more flexible. For instance, one interesting operator to study is the square root of the Laplacian, which is the pseudodifferential operator with symbol $a(x, \xi) = 2\pi|\xi|$. The fact that a is not smooth near the origin means that $a(x, D)$ does not quite fit the theory of pseudodifferential operators

given above. Nonetheless, for any bump function $\rho \in \mathcal{D}(\mathbf{R}_\xi^d)$ equal to one in a neighborhood of the origin, the symbol $\tilde{a}(x, \xi) = 2\pi|\xi|(1 - \rho(\xi))$ lies in $\mathcal{S}^1(\mathbf{R}^d \times \mathbf{R}^d)$. Since $\text{supp}_\xi(a - \tilde{a})$ is compact, it follows that $a(x, D) - \tilde{a}(x, D)$ is a smoothing operator. Thus, modulo smoothing operators, it makes sense to identify a with a symbol of order one. Thus we define the symbol class $\dot{\mathcal{S}}^t(\Omega \times \mathbf{R}^d)$ to be the space of all distributional symbols which differ from an element of $\mathcal{S}_{\text{loc}}^t(\Omega \times \mathbf{R}^d)$ by a distributional symbol which induces a smoothing operator. Then $a \in \dot{\mathcal{S}}^1$.

One can already see from this example, that this theory works well with pseudodifferential operators defined in terms of homogeneous functions, of which the square root of the Laplacian is a special example, since homogeneous functions often fail to be smooth at the origin. Another case where this works well is that it only suffices to specify an element of $\dot{\mathcal{S}}^t$ by a symbol $a(x, \xi)$ only defined for suitably large ξ , since any two extensions of the symbols near the origin will differ by a symbol compactly supported in the ξ variable, and thus smoothing.

Another virtue of working modulo smoothing operators is that asymptotics can be used to specify symbols. In other words, for any family of symbols $a_k \in \mathcal{S}_{\text{loc}}^{t_k}(\Omega \times \mathbf{R}^d)$, provided that $t_k \rightarrow -\infty$ and with $t = \max(t_k)$, we can define a symbol $a \in \dot{\mathcal{S}}^t(\Omega \times \mathbf{R}^d)$ by an asymptotic formula of the form

$$a(x, \xi) \sim \sum_{k=0}^{\infty} a_k(x, \xi),$$

since the difference of any two symbols satisfying this asymptotic formula lies in $\mathcal{S}_{\text{loc}}^{-\infty}(\Omega \times \mathbf{R}^d)$, and thus induces a smoothing operator.

13.2 Compound Operators and Quantization

It is natural to wish to study a more general family of operators with a *compound symbol* of the form $a(x, y, \xi)$, i.e. an operator of the form

$$T_a f(x) = \int a(x, y, \xi) e^{2\pi i \xi \cdot (x-y)} f(y) d\xi dy.$$

However, any such operator is already a pseudodifferential operator, and we can calculate an explicit asymptotic expansion for the symbol of this operator.

Lemma 13.6. For any $a \in \mathcal{S}_{loc}^t(\Omega_x \times \Omega_y \times \mathbf{R}^d)$, the operator T_a lies in $\Psi_{loc}^t(\Omega)$, and the symbol of T_a has the asymptotic expansion

$$\sum_{\beta} \frac{1}{\beta!} \frac{1}{(2\pi i)^{|\beta|}} \partial_{\xi}^{\beta} \partial_y^{\beta} a(x, x, \xi).$$

Proof. We begin by noting that, by very similar techniques to those above, the kernel K of an operator defined by a compound symbol of order $-\infty$ is smooth, and satisfies estimates of the form

$$|\nabla_x^n \nabla_z^m K(x, y)| \lesssim_{n,m,N} \langle x - y \rangle^{-N}$$

locally uniformly in x and y , which means that K is the kernel of an operator in $\Psi_{loc}^{-\infty}(\Omega)$. Thus we obtain the result for $t = -\infty$. In general, we perform a Taylor expansion, writing

$$a(x, y, \xi) = \sum_{|\beta| \leq N} \frac{1}{\beta!} \partial_y^{\beta} a(x, x, \xi) \cdot (y - x)^{\beta} + R_N(x, y, \xi),$$

where $\partial_y^{\beta} R_N(x, x, \xi) = 0$ for all $|\beta| \leq N$. We can find C^{∞} functions $b_{\beta}(x, y, \xi)$, for $|\beta| = N + 1$, such that

$$R_N(x, y, \xi) = \sum_{|\beta|=N+1} (2\pi i)^{N+1} (y - x)^{\beta} b_{\beta}(x, y, \xi).$$

Now integration by parts shows that

$$\begin{aligned} & \int R_N(x, y, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi \\ &= (-1)^{N+1} \sum_{|\alpha|=N+1} \int (D_{\xi}^{\alpha} b_{\alpha})(x, y, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi. \end{aligned}$$

The functions b_{β} are symbols of order t , so $D_{\xi}^{\alpha} b_{\beta}$ are symbols of order $t - (N + 1)$. Thus the operators specified by the compound symbol R_N have order at most $t - (N + 1)$. On the other hand, another integration by parts again shows that

$$\begin{aligned} & \int \partial_y^{\beta} a(x, x, \xi) \cdot (y - x)^{\beta} \cdot e^{2\pi i \xi \cdot (x-y)} d\xi \\ &= \frac{1}{(2\pi i)^{|\beta|}} \int \partial_{\xi}^{\beta} \partial_y^{\beta} a(x, x, \xi) e^{2\pi i \xi \cdot (x-y)} d\xi. \end{aligned}$$

Thus the operator corresponding with symbol $\partial_y^\beta a(x, x, \xi) \cdot (y - x)^\beta$ also corresponds to the symbol $1/(2\pi i)^{|\beta|} \cdot \partial_\xi^\beta \partial_y^\beta a(x, x, \xi)$. Thus if we consider a symbol $\tilde{a}(x, \xi)$ satisfying the asymptotic equation defined in the theorem, then we see that $\tilde{a}(x, D)$ differs from T_a by a compound symbol of order $-\infty$, which, as previously discussed, is an element of $\Psi_{\text{loc}}^{-\infty}(\Omega)$. \square

Remark. If $b(x, \xi) = a(x, x, \xi)$, then the formula above can be written more formally as $a(x, y, \xi) \sim e^{2\pi i D_x \cdot D_\xi} b$. This makes sense, since if K is the kernel of T_a , and T_a corresponds to a pseudodifferential operator, then it's symbol would correspond precisely to

$$\begin{aligned} a(x, \xi) &= \int K(x, x - y) e^{-2\pi i \xi \cdot y} dy \\ &= \int \int a(x, x - y, \xi - \eta) e^{-2\pi i \eta \cdot y} d\eta dy. \end{aligned}$$

If we define $a_x(y, \xi) = a(x, y, \xi)$, and $c(x, \xi) = e^{-2\pi i \xi \cdot x}$, then $a(x, \xi) = (a_x * c)(x, \xi)$. But c is a Gaussian, and thus if (ξ', x') are the dual variables to (x, ξ) , then $\hat{c}(\xi', x') = e^{2\pi i \xi' \cdot x'}$. Thus

$$a(x, \xi) = e^{2\pi i D_x \cdot D_\xi} a_x(x, \xi),$$

where $e^{2\pi i D_x \cdot D_\xi}$ is the Fourier multiplier operator with symbol $e^{2\pi i \xi' \cdot x'}$. Taking the power series expansion of the exponential gives the expansion above.

For any pseudodifferential operator T with kernel $K_a(x, y)$, we can consider it's formal adjoint T^* with kernel $K(x, y) = \overline{K_a(y, x)}$. We now use the calculation above to show the adjoint of a pseudodifferential operator $a(x, \xi)$; it is simple to calculate that the adjoint of any ΨDO $a(x, D)$ is a pseudodifferential operator with compound symbol $(x, y, \xi) \mapsto \overline{a(y, \xi)}$. Nonetheless, the above theorem implies that the adjoint can be given by a symbol $a^*(x, \xi)$ where $a^*(x, \xi) = e^{2\pi i D_x \cdot D_\xi} \overline{a}(x, \xi)$, which we can write explicitly as an asymptotic expansion as

$$a^*(x, \xi) \sim \sum_{\beta} \frac{1}{\beta!} \frac{1}{(2\pi i)^{|\beta|}} \overline{\partial_\xi^\beta \partial_x^\beta a(x, \xi)}.$$

In particular, if a is a symbol of order t , then $a^*(x, \xi) - \overline{a(x, \xi)}$ is a symbol of order $t - 1$, which we might write as saying that $a^* \approx \overline{a}$, up to lower

order terms. In particular, if a is a symbol correspond to a *self adjoint* pseudodifferential operator, then $a \approx \text{Re}(a)$, up to lower order terms.

The choice of (x, ξ) variables is common, but certainly not standard. The association of the pseudodifferential operator $a(x, D)$ with any symbol $a(x, \xi)$ is called the *Kohn-Nirenberg quantization*. We could also use the *adjoint Kohn-Nirenberg quantization* to associate an operator with every symbol a in two variables, using the (y, ξ) variables instead of the (x, ξ) variables. We find, using the expansion above, that modulo smoothing operators, any symbol in the (y, ξ) variables can be written in the (x, ξ) variables, and moreover,

$$a(y, \xi) \sim \sum_{\beta} \frac{1}{\beta!} \frac{1}{(2\pi i)^{|\beta|}} \partial_{\xi}^{\beta} \partial_x^{\beta} a(x, \xi).$$

In particular, the symbol $(x, y, \xi) \mapsto a(y, \xi) - a(x, \xi)$ is a pseudodifferential operator of order $t - 1$. Thus the two quantizations describe exactly the same family of operators, and the association of an operator with a symbol only matters up to lower order terms.

The family of operators one can describe via the adjoint Kohn-Nirenberg quantization is the same as the Kohn-Nirenberg quantization. Thus, in the sequel, there is no harm in sticking with the Kohn-Nirenberg quantization. On the other hand, the symbols representing various operators change. For instance, we previously found that under the Kohn-Nirenberg quantization, the symbol $a(x, \xi) = \sum c_{\alpha}(x) \xi^{\alpha}$ corresponded to the differential operator $Lf = \sum c_{\alpha} D^{\alpha} f$. Under the adjoint Kohn-Nirenberg quantization, the symbol $a(y, \xi) = \sum c_{\alpha}(y) \xi^{\alpha}$ corresponds to the differential operator $Lf = \sum D^{\alpha} (c_{\alpha} f)$. If t is the order of these operators, then the difference of these operators is a differential operator of order $t - 1$, which reflects the equivalence described above.

Thus we see that, roughly speaking, the operators differ in the order in which they apply spatial and frequency modulation. It is sometimes useful to deal with a quantization that does both in a ‘symmetric’ manner. To do this, we introduce the *Weyl quantization*, which associates with each symbol $a(x, \xi)$ gives the Pseudodifferential operator T with compound symbol $(x, y, \xi) \mapsto a((x + y)/2, \xi)$. This approach has the advantage that T will be self-adjoint if and only if a is real-valued, rather than just this only being true up to lower order terms. The Weyl quantization is the approach that works best in a generalization of a functional calculus for any

finite family of noncommuting operators (there are notes by Tao which describes this process in detail, but it is beyond the scope of these notes).

13.3 Compositions of Ψ DOs

Let $a(x, D)$ and $b(x, D)$ be pseudodifferential operators defined by local symbols of order t and s . It is not always possible to define the composition $a(x, D) \circ b(x, D)$. This is because the image of an element of $\mathcal{D}(\Omega)$ under the operator $b(x, D)$ in general lies in $\mathcal{E}(\Omega)$, and one cannot necessarily apply $a(x, D)$ to elements of $\mathcal{E}(\Omega)$. If we were working on \mathbf{R}^d and working with uniform symbols this wouldn't be a problem, since if $a \in \mathcal{S}^t(\mathbf{R}^d \times \mathbf{R}^d)$ and $b \in \mathcal{S}^s(\mathbf{R}^d \times \mathbf{R}^d)$, then $a(x, D)$ and $b(x, D)$ are both continuous operators from $\mathcal{S}(\mathbf{R}^d)$ to itself, and so the composition is well defined. In our approach, we must make a slightly technical assumption: we assume that either $a(x, D)$ or $b(x, D)$ maps $\mathcal{D}(\mathbf{R}^d)$ to itself, which is true in particular if $a(x, D)$ or $b(x, D)$ is a *proper* pseudodifferential operator.

Regardless of which method we use, the composition of a Ψ DO of order t and a Ψ DO of order s will then be a Ψ DO of order $t + s$, and we have an asymptotic formula for the symbol of such an expansion, reflecting the lack of commutivity between the spatial and frequential variables. In particular, the symbol of the composition is, to first order, the product of the symbols of the two operators.

Theorem 13.7. *Let $a(x, \xi)$ and $b(x, \xi)$ be symbols of order t and s , corresponding to operators T_a and T_b . Then $T_a \circ T_b$ is a Ψ DO of order $t + s$, and has symbol*

$$(a \circ b)(x, \xi) = e^{2\pi i D_\xi \cdot D_y} \{a(x, \xi) b(y, \eta)\} \Big|_{y=x, \eta=\xi}.$$

In particular, we have the asymptotic expansion

$$(a \circ b)(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} \frac{1}{(2\pi i)^{|\alpha|}} \partial_{\xi}^{\alpha} a(x, \xi) \cdot \partial_x^{\alpha} b(x, \xi).$$

Thus $(a \circ b)(x, \xi) - a(x, \xi)b(x, \xi)$ is a symbol of order $t + s - 1$.

Remark. If we consider the two form

$$\omega = dx \wedge d\xi - d\xi \wedge dx$$

on $T^*\mathbf{R}^n$, then we can define the *Poisson bracket* of two functions on $T^*\mathbf{R}^n$ by setting

$$\{a, b\} = \omega(\nabla a, \nabla b) = \sum_{i=1}^d \frac{\partial a}{\partial \xi^i} \frac{\partial b}{\partial x^i} - \frac{\partial a}{\partial x^i} \frac{\partial b}{\partial \xi^i}.$$

For any two symbols a and b , the result above implies that the commutator $[a(x, D), b(x, D)]$ of the two operators is a pseudodifferential operator of order $t + s$ with some symbol $[a, b]$, such that the symbol $[a, b] - (4\pi i)^{-1} \{a, b\}$ has order $t + s - 2$.

Proof. We can write

$$\begin{aligned} (T_a \circ T_b)f(x) &= \int a(x, \eta) e^{2\pi i \eta \cdot (x-z)} T_b f(z) dz d\eta \\ &= \int a(x, \eta) b(z, \xi) e^{2\pi i (\eta - \xi) \cdot (x-z)} e^{2\pi i \xi \cdot (x-y)} f(y) dy dz d\xi d\eta. \end{aligned}$$

Thus we see that we can view the composition as a Ψ DO with kernel

$$c(x, \xi) = \int \int a(x, \eta) b(z, \xi) e^{2\pi i (\eta - \xi) \cdot (x-z)} d\eta dz.$$

This is an oscillatory integral, with stationary point when $z = x$ and $\eta = \xi$. Thus we expand power series near this point, i.e. writing

$$a(x, \eta) = \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} a(x, \xi) (\eta - \xi)^{\alpha}$$

and

$$b(z, \xi) = \sum_{\beta} \frac{1}{\beta!} \partial_x^{\beta} b(x, \xi) (z - x)^{\beta}.$$

Using the Fourier inversion formula, we calculate that

$$\begin{aligned} &\int (\eta - \xi)^{\alpha} (z - x)^{\beta} e^{2\pi i (\eta - \xi) \cdot (x-z)} d\eta dz \\ &= \int \tau^{\alpha} y^{\beta} e^{-2\pi i \tau \cdot y} d\tau dy \\ &= \begin{cases} 0 & \alpha \neq \beta, \\ \alpha! / (2\pi i)^{\alpha} & \alpha = \beta. \end{cases} \end{aligned}$$

Working like in our analysis of compound symbols, it suffices to show that if g_1 and g_2 are symbols of order t and s , then

$$f(x, \xi) = \int \int (\eta - \xi)^\alpha (z - x)^\beta g_1(x, \eta) g_2(z, \xi) e^{2\pi i(\eta - \xi) \cdot (x - z)} d\eta dz$$

is a symbol of order $t + s - M - 1$. Applying sufficiently many integration by parts, it actually suffices to show integrals of the form

$$f(x, \xi) = \int \int g_1(x, \eta) g_2(z, \xi) e^{2\pi i(\eta - \xi) \cdot (x - z)} d\eta dz,$$

have order $t + s$, where g_1 has order t , and g_2 has order s . We write $\lambda = |\xi|$, and $\xi = \lambda \tilde{\xi}$, and write

$$f(x, \xi) = \lambda^d \int \int g_1(x, \lambda \eta) g_2(z, \xi) e^{2\pi i \lambda (\eta - \tilde{\xi}) \cdot (x - z)} d\eta$$

We can decompose the domain dyadically. For $|\eta| \leq 1/2$ and $|x - z| \leq 1$, an integration by parts in z gives rapid decay in t . Similarly, we can dyadically sum over the regions where $|\eta| \leq 1/2$ and $|x - z| \sim 2^k$ by first integrating in η using integration by parts, then integration in parts in z . This also gives rapid decay in t . Similar arguments give rapid decay in ξ for $|\eta| \sim 2^l$, in fact giving estimates which are summable in l . Thus we are left with giving decay for an integral of the form

$$t^d \int \int g_1(x, t\eta) g_2(z, \xi) \rho(|x - z|) \rho(|\eta| - 1) e^{2\pi i t(\eta - \tilde{\xi}) \cdot (x - z)} d\eta dz.$$

This domain has a stationary point when $\eta = \xi$ and $z = x$. However, the stationary point is nondegenerate. Thus the integral is $O(\lambda^{t+s} \lambda^{-d})$ and so $|f(x, \xi)| \lesssim \langle \xi \rangle^{t+s}$. Replacing g_1 and g_2 with appropriate derivatives gives a full argument that f is a symbol of order $t + s$. \square

Remark. This result shows that $\Phi = \bigcup_t \Phi^t$ and the subfamily $\Phi_{\text{loc,prop}}$ of proper pseudodifferential operators in $\Phi_{\text{loc}} = \bigcup_t \Phi_{\text{loc}}^t$ form graded algebras. We note for any pseudodifferential operator $T \in \Phi_{\text{loc}}^t$, there is a properly supported pseudodifferential operator $\tilde{T} \in \Phi_{\text{loc,prop}}^t$ such that $T - \tilde{T}$ is a smoothing operator. If we define $\dot{\Phi}^t$ to be the space of all pseudodifferential operators which differ from an element of Φ_{loc}^t by a pseudodifferential smoothing operator, modulo the set of smoothing operators, then

$\dot{\Phi} = \bigcup_t \dot{\Phi}^t$ is a graded algebra under addition and composition, which induces, by bijection, an alternate graded algebra structure on \dot{S} , agreeing up to first order with the standard algebra structure on this space given by multiplication.

13.4 Parametrices for Elliptic Operators

A *pseudodifferential parametrix* for a pseudodifferential operator T is a pseudodifferential operator S such that $S \circ T$ and $T \circ S$ are the identity operator, modulo smoothing operators. One useful result of our calculations is that we can easily construct *parametrices* for suitable pseudodifferential operators. A symbol $a \in \dot{S}^t(\Omega \times \mathbf{R}^d)$ is called *elliptic* if, locally in the x variable, we can find $R > 0$ such that for $|\xi| \geq R$,

$$|a(x, \xi)| \sim \langle \xi \rangle^t,$$

where the implicit constant is also locally uniform. In this case, we can interpret $b_0(x, \xi) = 1/a(x, \xi)$ as an element of \dot{S}^{-t} , since the reciprocal is well defined for large ξ , and satisfies the required symbol estimates. By the composition calculus, $1 - (a \circ b_0)(x, \xi)$ is a symbol $r_1 \in \dot{S}^{-1}$. For $i \geq 1$, given $r_i \in \dot{S}^{-i}$, if we define $b_i = r_i/a$ in \dot{S}^{t-i} , then the composition calculus tells us that $r_{i+1} = 1 - (a \circ (b_0 + \cdots + b_i))$ lies in \dot{S}^{-i-1} , and we can continue the iteration. If we consider $b \in \dot{S}^{-t}$ defined by the asymptotic development

$$b \sim \sum_{i=0}^{\infty} b_i$$

then $b(x, D)$ is a right pseudodifferential parametrix for $a(x, D)$. Similarly, we can construct $c \in \dot{S}^{-t}$ such that $c(x, D)$ is a left pseudodifferential parametrix for $a(x, D)$. But this means that, in \dot{S}^{-t} ,

$$b = b \circ (a \circ c) = (b \circ a) \circ c = c$$

so $b = c$, modulo smoothing operators. Thus a left parameterix for an elliptic operator is automatically a right parameterix, and we have constructed such a parametrix.

Remark. The condition that $|a(x, \xi)| \sim \langle \xi \rangle^t$ for large ξ is necessary in order to construct a parametrix of order $-t$. Without loss of generality, it suffices to analyze the case $t = 0$, since in general we can replace $a(x, \xi)$ with $a(x, \xi)\langle \xi \rangle^{-t}$ and $b(x, \xi)$ with $b(x, \xi)\langle \xi \rangle^t$. If there is $b \in \dot{S}^0$ such that $b(x, D)$ is a parametrix for $a(x, D)$, then the composition calculus tells us that $1 - a(x, \xi)b(x, \xi)$ is a local symbol of order -1 . Thus we find that $|a(x, \xi)b(x, \xi) - 1| \lesssim \langle \xi \rangle^{-1}$ locally uniformly in x , which implies that, locally in x , there exists $R > 0$ such that for $|\xi| \geq R$,

$$|a(x, \xi)b(x, \xi) - 1| \leq 1/2.$$

Thus

$$|a(x, \xi)| \geq \frac{0.5}{|b(x, \xi)|},$$

and combined with the fact that $|b(x, \xi)| \lesssim 1$, we conclude that $|a(x, \xi)| \gtrsim 1$.

It follows from our theory that if T is an elliptic pseudodifferential operator on Ω , then for any $u \in \mathcal{E}(\Omega)^*$, $\text{sing-supp}(Tu) = \text{sing-supp}(u)$. This is very similar to hypoellipticity, except that for differential operators, this condition can be applied to arbitrary distributions, not just the compactly supported distributions. More generally, we actually find that $\text{WF}(Tu) = \text{WF}(u)$ for $u \in \mathcal{E}(\Omega)^*$ in virtue of the microlocal nature of pseudodifferential operators.

We can use similar asymptotic tools to construct formal fractional powers of an elliptic pseudodifferential operator. For simplicity, we assume we are constructing the fractional powers of an elliptic symbol $a(x, \xi)$ such that $a(x, \xi)$ is never a negative, real number, so that $a(x, \xi)^{p/q}$ is well defined as a principal branch of $z^{p/q}$. Of course, one can consider a similar development for any other choice of branch, assuming an appropriate constraint on the range of a .

Theorem 13.8. *Let $a \in \dot{S}^t$ be an elliptic symbol such that $a(x, \xi)$ is negative a negative, real number, and suppose that*

$$|\arg(a(x, \xi)) - \pi| \gtrsim 1$$

for $\xi \gtrsim 1$, where the implicit constants are locally uniform in x . Then for any pair of positive integers p and q , there exists a unique $b \in \dot{S}^{t(p/q)}$ with principal symbol $a(x, \xi)^{p/q}$, such that $a(x, D)^p = b(x, D)^q$, modulo smoothing operators.

Proof. Let $b_0(x, \xi) = a(x, \xi)^{p/q}$. Then b_0 is a symbol of order $t(p/q)$, and the composition calculus implies that $a(x, D)^p - b_0(x, D)^q$ is a pseudodifferential operator of order $tp - 1$, say, with symbol r_1 . Given that we have chosen b_0, \dots, b_N such that $a(x, D)^p - (b_0 + \dots + b_N)(x, D)^q$ is a pseudodifferential operator of order $tp - (N + 1)$, say, with symbol r_N . The composition calculus means if we choose $b_{N+1} \in \dot{S}^{t(p/q) - (N+1)}$ by setting

$$b_{N+1} = q^{-1} r_N / (b_0 + \dots + b_N)^{q-1}.$$

Here we rely on the fact that a is elliptic, so that the denominator is non-vanishing for large ξ . Then

$$a(x, D)^p - (b_0 + \dots + b_N + b_{N+1})^q$$

is a pseudodifferential operator of order $tp - (N + 2)$, allowing us to continue the construction. If we pick $b \sim \sum_{n=0}^{\infty} b_n$, then $a(x, D)^p - b(x, D)^q$ will be a smoothing operator. Moreover, it is clear that the choice of such symbols at each step is essentially unique, which shows that b is unique up to a smoothing operator. \square

Given the assumptions of the theorem, we denote the unique operator b by $a^{p/q}$. Using a^{-1} instead of a , one can also construct negative fractional powers of the symbol a . It is simple to see that for $r_1, r_2 \in \mathbf{Q}$, we have $a^{r_1} \circ a^{r_2} = a^{r_1 + r_2}$.

13.5 Regularity Theory

Let us now discuss the boundedness of certain pseudodifferential operators with respect to various norm spaces. We first note that a differential operator of degree m given by

$$L = \sum c_\alpha(x) D^\alpha,$$

where c_α is bounded, maps $H^s(\mathbf{R}^d)$ to $H^{s-m}(\mathbf{R}^d)$ for each s . This feature remains true for a general pseudodifferential operator.

Theorem 13.9. *For any $a \in \mathcal{S}_{loc}^t(\mathbf{R}^d)$, $a(x, D)$ extends uniquely to a continuous operator from $H_c^s(\mathbf{R}^d)$ to $H_{loc}^{s-m}(\mathbf{R}^d)$.*

Proof. Let T have symbol $a(x, \xi)$. Without loss of generality, since we need only prove local estimates in the output we may assume that a is compactly supported in the x -variable. Then a is uniformly integrable in the x -variable, and we let

$$A(\lambda, \xi) = \int a(x, \xi) e^{-2\pi i \lambda \cdot x} dx$$

denote the Fourier transform of a in the x -variable. For any input $\phi \in \mathcal{S}(\mathbf{R}^d)$, $T\phi \in \mathcal{S}(\mathbf{R}^d)$, and we may calculate that

$$\widehat{T\phi}(\lambda) = \int A(\lambda - \xi, \xi) \widehat{\phi}(\xi) d\xi.$$

The assumptions on the symbol a imply that

$$|A(\lambda, \xi)| \lesssim_N \langle \xi \rangle^t \langle \lambda \rangle^{-N}.$$

Without loss of generality, assume that $s = t$. Applying Schur's lemma, the operator

$$Sf(\lambda) = \int \langle \lambda - \xi \rangle^{-N} f(\xi) d\xi,$$

somewhat analogous to the Hardy-Littlewood-Sobolev fractional integration operator, is bounded from $L^2(\mathbf{R}^d)$ to itself provided that $N > d$. But this means that if we pick $N > d$, then

$$\begin{aligned} \|T\phi\|_{L^2(\mathbf{R}^d)} &\lesssim_N \left\| \int \langle \lambda - \xi \rangle^{-N} \langle \xi \rangle^t \widehat{\phi}(\xi) d\xi \right\|_{L^2(\mathbf{R}^d)} \\ &\lesssim \|\langle \xi \rangle^t \widehat{\phi}\|_{L^2(\mathbf{R}^d)} \\ &\lesssim \|\phi\|_{H^t(\mathbf{R}^d)}. \end{aligned}$$

Thus T is bounded from $H^t(\mathbf{R}^d)$ to $L^2(\mathbf{R}^d)$. But for general $s \in \mathbf{R}^d$, T will be bounded from $H^s(\mathbf{R}^d)$ to $H^{s-t}(\mathbf{R}^d)$ if and only if $(1 - \Delta)^{s-t} T (1 - \Delta)^{t-s}$ is bounded from $H^t(\mathbf{R}^d)$ to $L^2(\mathbf{R}^d)$, and this follows because $(1 - \Delta)^{s-t} T (1 - \Delta)^{t-s}$ is also a pseudodifferential operator of order t . The bounds we have proven show that there is a unique extension of T to $H^s(\mathbf{R}^d)$ for all $s \in \mathbf{R}^d$ such that $\|Tf\|_{H^{s-t}(\mathbf{R}^d)} \lesssim \|f\|_{H^s(\mathbf{R}^d)}$ for all $f \in H^s(\mathbf{R}^d)$. The closed graph theorem shows that this extension agrees with the definition of Tf given for any compactly supported f , which we can view as an element of $\mathcal{E}(\mathbf{R}^d)^*$. \square

Remark. For symbols of order zero, a simpler proof follows by writing

$$T^\lambda \phi(x) = \int A(\lambda, \xi) \hat{\phi}(\xi) e^{2\pi i(\lambda + \xi) \cdot x} d\xi.$$

then the Fourier inversion formula shows that

$$T\phi(x) = \int T^\lambda \phi(x) d\lambda.$$

Now the operators $\{T^\lambda\}$ are just Fourier multiplier operators with symbols $\{m_\lambda\}$, where $m_\lambda(\xi) = A(\lambda, \xi) e^{2\pi i \lambda \cdot x}$, the bounds on A imply that $\|m_\lambda\|_{L^\infty} \lesssim \langle \lambda \rangle^{-N}$, and this immediately gives boundedness from $L^2(\mathbf{R}^d)$ to itself. The general result follows from the same trick as in the end of the last proof using the composition calculus.

Remark. If a is *properly supported* pseudodifferential operator of order t , then $a(x, D)$ extends to a continuous operator from $H_{\text{loc}}^s(\Omega)$ to $H_{\text{loc}}^{s-t}(\Omega)$.

Using Calderon-Zygmund theory, we can obtain better estimates. Let us restrict ourselves at first to pseudodifferential operators of order 0. The kernel of a Ψ DO of order zero satisfies estimates of the form

$$|K(x, y)| \lesssim \frac{1}{|x - y|^d}.$$

Thus we focus on obtaining $L^2 \rightarrow L^2$ estimates, so that the standard theory of singular integrals gives $L^p \rightarrow L^p$ estimates for all $1 < p < \infty$.

Theorem 13.10. *If $a \in \mathcal{S}^0(\mathbf{R}^d \times \mathbf{R}^d)$, then for any $f \in \mathcal{S}$,*

$$\|T_a f\|_{L^2(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^d)}.$$

Proof. If $\text{supp}_x(a)$ is compact, then we have already proven this result. To prove the result for more general symbols, we work with a kernel representation of T_a . Thus we write

$$T_a f(x) = \int_{\mathbf{R}^d} K(x, y) f(y) dy,$$

where

$$K(x, y) = \int_{\mathbf{R}^d} a(x, \xi) e^{2\pi i \xi \cdot (x - y)} d\xi.$$

We have already shown that the kernel K is C^∞ away from the diagonal, and decays rapidly away from that diagonal. This is one instance of the pseudolocal nature of these operators. Another quantitative result reflecting this nature is that for each $N > 0$ and $x_0 \in \mathbf{R}^d$,

$$\int_{|x-x_0| \leq 1} |T_a f(x)|^2 dx \lesssim_N \int_{\mathbf{R}^d} \frac{|f(x)|^2}{\langle x - x_0 \rangle^N} dx.$$

Thus we can ‘almost’ bound the magnitude of $T_a f$ in a neighbourhood of x_0 by the magnitude of f in a neighbourhood of x_0 . We focus on the case $x_0 = 0$, the other cases treated in much the same way. Write $f = f_1 + f_2$, where f_1 is supported on $|x| \leq 3$, f_2 is supported on $|x| \geq 2$, and $|f_1|, |f_2| \leq |f|$. If $\eta(x)$ is a smooth cutoff supported on $|x| \leq 3$, then the symbol $\eta(x)a(x, \xi)$ is compactly supported, and so

$$\begin{aligned} \int_{|x| \leq 1} |T_a f_1(x)|^2 dx &= \int_{|x| \leq 1} |T_{\eta a} f_1(x)|^2 \lesssim \|f_1\|_{L^2(\mathbf{R}^d)}^2 \\ &\lesssim_N \int_{\mathbf{R}^d} \frac{|f_1(x)|^2}{\langle x \rangle^N} dx \leq \int_{\mathbf{R}^d} \frac{|f(x)|^2}{\langle x \rangle^N} dx. \end{aligned}$$

On the other hand, since $f_2(x)$ is supported on $|x| \geq 2$, we find that

$$\begin{aligned} \int_{|x| \leq 1} |T_a f_2(x)|^2 dx &= \int_{|x| \leq 1} \left| \int K(x, y) f_2(y) dy \right|^2 dx \\ &\leq \int \int_{|x| \leq 1} |K(x, y)|^2 |f_2(y)|^2 dy dx \\ &\lesssim_N \int \int_{|x| \leq 1} \frac{|f_2(y)|^2}{|x - y|^N} dy dx \\ &\lesssim \int \int_{|x| \leq 1} \frac{|f_2(y)|^2}{\langle y \rangle^N} dy \\ &\lesssim \int \frac{|f_2(y)|^2}{\langle y \rangle^N} dy. \end{aligned}$$

But we now find that if $N > d$, then

$$\begin{aligned} \int |T_a f(x)|^2 dx &\lesssim \int \int_{|x-y| \leq 1} |T_a f(y)|^2 dy dx \\ &\lesssim_N \int \int \frac{|f(y)|^2}{\langle x-y \rangle^N} dy dx \\ &\lesssim \int |f(y)|^2 dy, \end{aligned}$$

which gives L^2 boundedness. \square

Sobolev norms follow simply from these bounds. Namely, it follows simply from this that if $a(x, \xi)$ is a symbol of order t , then for $1 < p < \infty$, and any s , we have bounds of the form

$$\|T_a f\|_{L_s^p(\mathbf{R}^d)} \lesssim_{p,s} \|f\|_{L_{t+s}^p(\mathbf{R}^d)}.$$

If T_a is an elliptic pseudodifferential operator of order t , then it has a parametrix S of order $-t$, and so for any $r > 0$

$$\begin{aligned} \|f\|_{L_{t+s}^p(\mathbf{R}^d)} &= \|ST_a f + (I - ST_a)f\|_{L_{t+s}^p(\mathbf{R}^d)} \\ &\leq \|ST_a f\|_{L_{t+s}^p(\mathbf{R}^d)} + \|(I - ST_a)f\|_{L_{t+s}^p(\mathbf{R}^d)} \\ &\lesssim_{p,s,r} \|T_a f\|_{L_s^p(\mathbf{R}^d)} + \|f\|_{L_{-r}^p(\mathbf{R}^d)} \end{aligned}$$

Thus T is *almost* invertible as a map from $L_{t+s}^p(\mathbf{R}^d)$ to $L_t^p(\mathbf{R}^d)$, except that we cannot quite obtain the bound $\|Tf\|_{L_s^p(\mathbf{R}^d)} \sim_{p,s} \|f\|_{L_{t+s}^p(\mathbf{R}^d)}$, for instance, by virtue of the fact that T might not even be invertible. But we can obtain a less quantitative result.

Theorem 13.11. *Let T be an elliptic pseudodifferential operator of order t . If $f \in \mathcal{E}(\mathbf{R}^d)^*$, and Tf lies in $L_s^p(\mathbf{R}^d)$, then f lies in $L_{s+t}^p(\mathbf{R}^d)$.*

Proof. Since T is elliptic, we can find a parametrix S , which is a pseudodifferential operator of order $-t$. Thus there exists a smoothing operator U such that $1 = ST + U$. Since $Tf \in L_s^p(\mathbf{R}^d)$, $STf \in L_{s+t}^p(\mathbf{R}^d)$. Since f is compactly supported, $Uf \in C^\infty(\mathbf{R}^d)$, and thus in $L_{s+t, \text{loc}}^p(\mathbf{R}^d)$. This means $f \in L_{s+t, \text{loc}}^p(\mathbf{R}^d)$, and since f is compactly supported, this means $f \in L_{s+t}^p(\mathbf{R}^d)$. \square

13.6 Pseudodifferential Operators on Manifolds

It is an important fact that the class of pseudodifferential operators whose kernels are compactly supported is invariant under a change of coordinates, modulo smoothing operators. By an analysis of how these pseudodifferential operators change under a change of coordinates, we will be able to obtain a theory of pseudodifferential operators on manifolds.

Let $\kappa : U \rightarrow V$ be a diffeomorphism, where U and V are open sets in \mathbf{R}^n , and consider $a \in S_{\text{loc}}^t(U)$ which gives a pseudodifferential operator $T : \mathcal{D}(U) \rightarrow \mathcal{E}(U)$ of order t . Define an operator $S : \mathcal{D}(V) \rightarrow \mathcal{E}(V)$ by setting

$$Sf(\kappa(x)) = T(f \circ \kappa)(x) = \int a(x, \xi) e^{2\pi i \xi \cdot (x-y)} f(\kappa(y)) dy d\xi,$$

i.e. S is defined such that $\kappa_* \circ S = T \circ \kappa_*$. One can verify that S is pseudolocal, so one might expect S to be a pseudodifferential operator, of order t as well. By localizing, we may assume that $\text{supp}_x(a)$ is compact, so that T extends to a continuous operator from $\mathcal{D}(U)^*$ to $\mathcal{D}(U)^*$, and S to a continuous operator from $\mathcal{D}(V)^*$ to $\mathcal{D}(V)^*$. If a symbol $b \in S_{\text{loc}}^t(V)$ existed such that $S = b(y, D)$, then we would find

$$b(\kappa(x_0), \eta) = \left\{ S e^{2\pi i \eta \cdot z} \right\} (\kappa(x_0)) = \int a(x_0, \xi) e^{2\pi i [\xi \cdot (x_0 - x) + \eta \cdot \kappa(x)]} dx d\xi$$

Provided we can show that the right hand side defines a symbol $b \in S_{\text{loc}}^t(V)$, then $b(y, D)$ will be a continuous operator whose kernel is compactly supported in $V \times V$, and the fact that $S e^{2\pi i \eta \cdot z} = b(y, D) e^{2\pi i \eta \cdot z}$ for all η implies, by continuity and the fact that linear combinations of exponentials are dense in $\mathcal{D}(V)^*$, that $S = b(y, D)$. Thus, given $a \in S_{\text{loc}}^t(U)$, such that $\text{supp}_x(a)$ is compact, our goal is an analysis of the function

$$b(\kappa(x_0), \eta) = \int a(x_0, \xi) e^{2\pi i [\xi \cdot (x_0 - x) + \eta \cdot \kappa(x)]} dx d\xi.$$

If $\phi \in \mathcal{D}(V)$ is a bump function equal to one in a neighborhood of the projections of the support of the kernel of S onto each coordinate, then

$$b(\kappa(x_0), \eta) = \left\{ \phi S \left\{ \phi \cdot e^{2\pi i \eta \cdot z} \right\} \right\} (\kappa(x_0)),$$

which implies that $b \in C^\infty(T^*V)$. If we set

$$\Phi(\xi, \eta) = \int \phi(x) e^{2\pi i [\eta \cdot \kappa(x) - \xi \cdot x]} dx,$$

then

$$b(\kappa(x_0), \eta) = \int a(x_0, \xi) \Phi(\xi, \eta) e^{2\pi i \xi \cdot x_0} d\xi.$$

Now Φ is a standard oscillatory integral, whose phase $\phi(x) = \eta \cdot \kappa(x) - \xi \cdot x$ has a stationary point for values of x such that $D\kappa(x)^T \eta = \xi$. Since $D\kappa(x)^T$ is invertible, this can only happen when $|\eta| \sim |\xi|$. More precisely, suppose $|D\kappa(x)|, |D\kappa(x)^{-1}| \leq C$. Then $|D\kappa(x)^T \eta - \xi| \geq |\xi|/2 \gtrsim_C |\xi| + |\eta|$, and $|D\kappa(x)^T \eta - \xi| \geq |\eta|/2 \gtrsim_C |\xi| + |\eta|$ for $|\xi| \leq |\eta|/2C$. Thus unless $2C|\eta| \geq |\xi| \geq |\eta|/2C$, we conclude by the principle of nonstationary phase that for each $N > 0$,

$$|\Phi(\xi, \eta)| \lesssim_N [1 + |\xi| + |\eta|]^{-N},$$

where the implicit constant will be uniformly bounded over a family of diffeomorphisms κ and a family of pseudodifferential operators T if the kernels of the resulting operators S are uniformly supported on a common compact subset of $V \times V$, and if we have uniform upper and lower bounds on the derivative of κ on the support of these kernels. If we write $1 = \chi_1 + \chi_2$, for $\chi_1, \chi_2 \in C^\infty(\mathbf{R}^d)$ with $\chi_1(\alpha)$ supported on $1/2C \leq |\alpha| \leq 2C$ and equal to one when $1/C \leq |\alpha| \leq C$, then this induces a decomposition $b(\kappa(x_0), \eta) = b_1(\kappa(x_0), \eta) + b_2(\kappa(x_0), \eta)$, where

$$b_i(\kappa(x_0), \eta) = \int \chi_i(\xi/|\eta|) a(x_0, \xi) \Phi(\xi, \eta) e^{2\pi i \xi \cdot x_0} d\xi.$$

The bound on $\Phi(\xi, \eta)$ above implies that $b_1 \in S^{-\infty}$. To analyze b_2 , we apply the method of stationary phase. To simplify our formulas, let us assume without loss of generality that $x_0 = \kappa(x_0) = 0$ and that $B = D\kappa(x_0)$. If we fix $|\eta| = 1$, consider $\lambda > 1$, and do a change of variables, replacing ξ with $\lambda(\xi + B^T \eta)$, we find that

$$b_2(\kappa(x_0), \lambda\eta) = \lambda^d \int \psi(x, \xi) a(x, \lambda B^T \eta + \xi) e^{2\pi i \lambda \phi(x, \xi)} d\xi dx,$$

where $\psi(\xi, x) = \chi_2(\xi + B^T \eta) \phi(x_0 + x)$ and $\phi(x, \xi) = \eta \cdot (\kappa(x) - Bx) - \xi \cdot x$. This is a continuous family of non-degenerate stationary phase integrals, each

with a unique stationary point when $(x, \xi) = (0, 0)$. Since ψ is equal to one in a neighborhood of this point, it will not show up in the corresponding asymptotics. The Hessian at the origin is precisely $A + B$, where

$$A = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix}$$

and

$$C = \begin{pmatrix} \tilde{C} & 0 \\ 0 & 0 \end{pmatrix}$$

where $\tilde{C} = \tilde{C}(\eta)$ is the Hessian of the map $x \mapsto \eta \cdot \kappa(x)$ at the origin. If we set

$$r(x) = \eta \cdot (\kappa(x) - Bx) - (1/2)(x^T A x)$$

which satisfies $\partial^\alpha r(0) = 0$ for all $|\alpha| \leq 2$, then it follows that if $a_\lambda(x_0, \xi) = a(x_0, \lambda\xi)$, then

$$b_2(\kappa(x_0), \lambda\eta) \sim \sum_{2\nu \geq 3\mu} \frac{1}{(-2)^\nu} \frac{1}{\mu! \nu!} \frac{1}{(2\pi i \lambda)^{\nu-\mu}} \langle (A+B)^{-1} \nabla, \nabla \rangle^\nu \{r^\mu a_\lambda\}(x, D\kappa(x)^T).$$

The next Lemma implies that we can write this asymptotic development as

$$b_2(\kappa(x_0), \lambda\eta) \sim \sum_{\nu=0}^{\infty} \langle i \cdot \nabla_x / 2\pi, \nabla_\xi / \lambda \rangle^\nu \left\{ e^{2\pi i \lambda r(x)} a_\lambda(0, \xi) \right\}_{x=0, \xi=B^T \eta}$$

where if we sum over $\nu \leq N$, then the error term will be $O(\lambda^{(d-N)/2})$.

Lemma 13.12. *Let A be a symmetric, invertible matrix, let B be a symmetric matrix, and suppose $\det(A + tB)$ is independent of t . Then there exists k such that $(A^{-1}B)^k = 0$ for some k . For this k , and for any $N > 0$,*

$$\sum_{j < N} \frac{1}{j!} (2i\lambda)^{-j} \langle (A+B)^{-1} \nabla, \nabla \rangle^j u(0) = \sum_{j < N} \frac{1}{j!} (2i\lambda)^{-j} \langle A^{-1} \nabla, \nabla \rangle^j \left\{ e^{2\pi i \lambda x^T B x / 2} u \right\}(0) + O(\lambda^{-N/k}).$$

Proof. See Hormander, 7.7. □

Thus we have proved the following representation formula for pseudodifferential operators under changes of coordinates.

Theorem 13.13. Let $\kappa : U \rightarrow V$ be a diffeomorphism, where U and V are open sets in \mathbf{R}^n , and suppose $T = a(x, D)$ is a pseudodifferential operator of order t on U . If S is an operator defined on V defined such that $\kappa_* \circ S = T \circ \kappa_*$, then S is a pseudodifferential operator of order t , whose symbol a_κ has the asymptotic development

$$a_\kappa(\kappa(x), \eta) \sim \sum \frac{1}{(2\pi i)^\alpha} \frac{1}{\alpha!} (\partial_\xi^\alpha a)(x, D\kappa(x)^T \eta) \left\{ \partial_y^\alpha e^{2\pi i \eta \cdot r_x(y)} \right\} \Big|_{y=x},$$

where $r_x(y) = \kappa(y) - \kappa(x) - D\kappa(x)(y - x)$. The term in the sum corresponding to a multi-index α is a symbol of order $t - [|\alpha|/2]$, and thus this really is an asymptotic expansion.

Pseudodifferential operators defined by a local symbol of order t are therefore invariant under a change of coordinates, modulo smoothing operators. Moreover, if $\kappa_* \circ S = T \circ \kappa_*$, where T is a ΨDO with symbol $a \in S_{\text{loc}}^t(U)$, and S is a ΨDO with symbol $b \in S_{\text{loc}}^t(V)$, then

$$b(y, \eta) - a(\kappa^{-1}(y), D\kappa(x)^{-T} \cdot \eta)$$

is a pseudodifferential operator of order $t - 1$.

Given a manifold M , a continuous operator $T : \mathcal{D}(M) \rightarrow \mathcal{E}(M)$ is called a *pseudodifferential operator of order t* if whenever (x, U) is a coordinate chart on M , the operator $T_x : \mathcal{D}(x(U)) \rightarrow \mathcal{E}(x(U))$ given by T in coordinates is a pseudodifferential operator of order t on $x(U)$. We let $\Psi_{\text{loc}}^t(M)$ denote the family of operators of this form. The next Lemma shows that when $M = U$ is an open subset of \mathbf{R}^n , this really is the family of pseudodifferential operators defined by symbols in $S_{\text{loc}}^t(T^*U)$.

Lemma 13.14. Suppose $T : \mathcal{D}(U) \rightarrow \mathcal{E}(U)$ is a pseudodifferential operator on U of order t , where U is viewed as a manifold as above. Then we can find a symbol $a(x, \xi) \in S_{\text{loc}}^t(U \times \mathbf{R}^d)$ such that $T - a(x, D)$ is a smoothing operator. The symbol a is uniquely determined up to a symbol in $S^{-\infty}(\mathbf{R}^d \times \mathbf{R}^d)$.

Proof. The idea is to work on a partition of unity, which we can sum up appropriately to get a sum over local estimates. The complete proof is supplied in Hormander, Proposition 18.1.19. \square

Remark. Under the assumption that the operator has a kernel which is smooth away from the diagonal, an operator $T : \mathcal{D}(M) \rightarrow \mathcal{E}(M)$ is a pseudodifferential operator if and only if it is pseudodifferential operator when

transferred in coordinates on a family of coordinates charts that form an atlas for M . The assumption on the kernel is needed, for instance, since if we take $M = \mathbf{R}$, we consider an atlas of the form $\{(n, n+1)\}$, and we consider $Tf(x) = f(x-2)$, then T vanishes on each of these coordinates charts, and so looks to be a pseudodifferential operator in these charts, whereas T clearly is not a pseudodifferential operator on \mathbf{R} since it is not pseudolocal.

It is often to discuss operators that can be asymptotically expanded in terms of homogeneous symbols. That is, a symbol $a \in \dot{S}^t$ of order t is classical if there exist a sequence of symbols $\{a_k\}$, where $a_k(x, \xi)$ is smooth away from $\xi = 0$, homogeneous of order $t - k$, and

$$a \sim \sum_{k=0}^{\infty} a_k.$$

Such symbols are called *classical*, and denoted $\mathcal{S}_{\text{cl}}^t$, since this was the family of pseudodifferential operators initially studied by Kohn and Nirenberg. Given a manifold M , we write $\Psi_{\text{cl}}^t(M) = \text{Op}(\mathcal{S}_{\text{cl}}^t)(M)$ for the class of all pseudodifferential operators T such that T_x is classical for any coordinate system (x, U) on M . It suffices to check this on a cover because of the asymptotic expansion for the change of variables formula. The homogeneous function which agrees with the leading term in the expansion is invariant under coordinate changes if we interpret it as a function on $T^*M - 0_M$, so for a classical pseudodifferential operator T in $\Psi_{\text{cl}}^t(M)$, we can define the *principal symbol* $a \in C^\infty(T^*M - 0_M)$ to be that homogeneous function of order t which agrees with the leading term in the asymptotic expansions above. For nonclassical operators, there is no canonical choice of a principal symbol, though one can consider the principal symbol as an element of $\mathcal{S}_{\text{loc}}^t(M)/\mathcal{S}_{\text{loc}}^{t-1}(M)$. A pseudodifferential operator is then called *elliptic* if its principal symbol is nonvanishing at a suitably distance away from the origin, or equivalently, if it is elliptic in coordinates ranging over the entirety of M .

Example. Let $\mathbf{T} = \mathbf{R}/\mathbf{Z}$. For any ‘symbol of order t ’ on \mathbf{T}^d , i.e. any function $a : \mathbf{T}_x^d \times \mathbf{Z}_\xi^d \rightarrow \mathbf{C}$ with extends to an element of $\mathcal{S}^t(\mathbf{T}_\xi^d \times \mathbf{R}_\xi^d)$, we can consider the operator $T : \mathcal{E}(\mathbf{T}^d) \rightarrow \mathcal{E}(\mathbf{T}^d)$ defined by setting

$$T\phi(x) = \sum_{\xi \in \mathbf{Z}^d} a(x, \xi) \hat{\phi}(\xi) e^{2\pi i \xi \cdot x},$$

where $\hat{\phi} : \mathbf{Z}^d \rightarrow \mathbf{C}$ is the Fourier transform of ϕ on \mathbf{T}^d . We claim that T is a pseudodifferential operator. By translation invariance, it will suffice to find a neighborhood Ω of the origin upon and a coordinate system on Ω upon which the transfer of T is a pseudodifferential operator. Write

$$\begin{aligned} T\phi(x) &= \sum_{\xi} a(x, \xi) \hat{\phi}(\xi) e^{2\pi i \xi \cdot x} \\ &= \int_{\mathbf{T}^d} \left(\sum_{\xi} a(x, \xi) e^{2\pi i \xi \cdot (x-y)} \right) \phi(y) dy \\ &= \int_{\mathbf{T}^d} K(x, y) \phi(y) dz, \end{aligned}$$

where we treat

$$K(x, y) = \sum_{\xi} a(x, \xi) e^{2\pi i \xi \cdot (x-y)}.$$

as the distribution on $\mathcal{D}(\mathbf{T}^d \times \mathbf{T}^d)^*$ obtained as the distributional limit of the partial sums. Consider the periodic kernel $\tilde{K} \in \mathcal{D}(\mathbf{R}^d \times \mathbf{R}^d)^*$ induced by K . Then the Poisson summation formula implies that

$$\tilde{K}(x, y) = \sum_{\xi} a(x, \xi) e^{2\pi i \xi \cdot (x-y)} = \sum_n (\mathcal{F}_{\xi}^{-1} a)(x, (x-y) + n) = \sum_n K_a(x, y + n),$$

where K_a is the kernel of the pseudodifferential operator $a(x, D)$. If $\tilde{\Omega} = (-1/4, 1/4)^d$, then for $x, y \in \tilde{\Omega}$, and $n \neq 0$,

$$|\partial_x^{\alpha} \partial_y^{\beta} K_a(x, y + n)| \lesssim_N |n|^{-N}.$$

This implies that

$$(x, y) \mapsto \sum_{n \neq 0} K_a(x, y + n)$$

lies in $C^{\infty}(\tilde{\Omega} \times \tilde{\Omega})$, and so induces a smoothing operator on $\tilde{\Omega}$. Thus \tilde{T} , restricted to $\tilde{\Omega}$, differs by a smoothing operator from the pseudodifferential operator $a(x, D)$. But this means that $\tilde{T} \in \Psi^t(\tilde{\Omega})$. If Ω is the subset of \mathbf{T}^d corresponding to $\tilde{\Omega}$, then this means T behaves like a pseudodifferential operator on $\tilde{\Omega}$. Thus T is actually a pseudodifferential operator of order t on \mathbf{T}^d . Modulo smoothing operators, working backwards through this argument also clearly

shows that all pseudodifferential operators of order t can be written in the form introduced at the beginning of this example.

As a particular example of this kind of construction, consider the operator $T : \mathcal{E}(\mathbf{T}) \rightarrow \mathcal{E}(\mathbf{T})$ given by

$$P_+ \phi(t) = \sum_{n>0} \hat{\phi}(n) e^{2\pi n i t},$$

then P_+ is a classical pseudodifferential operator on \mathbf{T} of order zero, and its principal symbol is $\mathbf{I}(\xi > 0)$. Similarly, if

$$P_- \phi(t) = \sum_{n<0} \hat{\phi}(n) e^{2\pi n i t},$$

then P_- is a classical pseudodifferential operator on \mathbf{T} of order zero with principal symbol $\mathbf{I}(\xi < 0)$. For any non-zero complex values $a_+, a_- \in \mathbb{C} - \{0\}$, $a_+ P_+ + a_- P_-$ is an elliptic classical pseudodifferential operator on \mathbf{T} of order zero.

Example. On a compact Riemannian manifold M , the operator $-\Delta$ is a positive-semidefinite differential operator of order two with symbol given in local coordinates as

$$a(x, \xi) = \sum g^{ij}(x) \xi_i \xi_j.$$

TODO: Change argument by arguing it is elliptic, and so we can take fractional powers.

The spectral theory of this operator decomposes it as a sum $\sum \lambda_i^2 E_i$, where E_i is an orthogonal projection onto a one dimensional eigenspace, and $\lambda_0 \leq \lambda_1 \leq \dots$ are ordered by multiplicity. Then we can define the operator $\sqrt{-\Delta}$ to be the operator $\sum \lambda_i E_i$. Let us argue that $\sqrt{-\Delta}$ is a classical pseudodifferential operator of order one on M . More precisely, we will construct a positive definite pseudodifferential operator Q of order one such that $Q - \sqrt{-\Delta}$ is smoothing.

To construct Q , we first consider the modified Laplacian $Lu = E_0 u + \sum_{i \geq 1} \lambda_i^2 E_i$. Then L is invertible since it is positive definite, and differs from $-\Delta$ by the smoothing operator

$$E_0 u = \frac{1}{\text{Vol}(M)} \int_M u(x) dV(x).$$

We will not exploit the fact that L is invertible until the end of the argument, but keep this invertibility in mind for later.

To begin with, we construct a positive definite $Q \in \Psi_{cl}^1(M)$ such that $L - Q^2$ is a smoothing operator. To do this we may assume without loss of generality that we are working in a small coordinate patch in local coordinates, and that we are patching these locally defined parts of the operator globally using a partition of unity. We then define

$$\tilde{b}_1(x, \xi) = \chi_1(\xi) \left(\sum g^{i,j}(x) \xi_i \xi_j \right)^{1/2}$$

where χ_1 is smooth and vanishes for $|\xi| \leq 1$. Now define $Q_1 = b_1(x, D)$ to be equal to the pseudodifferential operator $\text{Re}(\tilde{b}_1(x, D)) = (1/2)(\tilde{b}_1(x, D) + \tilde{b}_1(x, D)^*)$. The calculus of pseudodifferential operators tells us that b_1 is a classical symbol of order one, and the principal part of b_1 agrees with the principal part of \tilde{b}_1 . We therefore conclude that $R_1 = L - Q_1^2$ is a classical pseudodifferential operator of order one with principal symbol $r_1(x, \xi)$.

We now proceed inductively, assuming that we have found classical symbols b_i of order $2 - i$ for each $1 \leq i \leq n$, such that if $Q_i = b_i(x, D)$, then Q_i is self-adjoint, and $R_n = L - (Q_1 + \cdots + Q_n)^2$ is a classical pseudodifferential operator of order $2 - n$ with some principal symbol $r_n(x, \xi)$ of order $2 - n$. Our goal is now to find a non-negative symbol $\tilde{b}_{n+1}(x, \xi)$, homogeneous of order $1 - n$ for large ξ , defining a self-adjoint pseudodifferential operator Q_{n+1} such that $L - (Q_1 + \cdots + Q_{n+1})^2$ is a classical pseudodifferential operator of order $1 - n$. Expanding out the order $2 - n$ part of $L - (Q_1 + \cdots + Q_{n+1})^2$, this is possible provided that for large ξ ,

$$r_n(x, \xi) = 2b_1(x, \xi)b_{n+1}(x, \xi)$$

Thus we define

$$\tilde{b}_{n+1}(x, \xi) = \chi_{n+1}(\xi) \left(\frac{r_n(x, \xi)}{2b_1(x, \xi)} \right)$$

where χ_{n+1} is smooth, equal to one for large ξ , and vanishes for values of ξ such that $b_1(x, \xi) = 0$, and then set b_{n+1} to be the classical symbol of order $1 - n$ such that $Q_{n+1}b_{n+1}(x, D) = \text{Re}(\tilde{b}_{n+1}(x, D))$. Then $R_{n+1} = L - (Q_1 + \cdots + Q_{n+1})^2$ is a classical pseudodifferential operator of order $1 - n$, allowing us to continue our induction.

Thus we find now that we have constructed an infinite sequence of classical pseudodifferential operators b_1, b_2, \dots , where b_i is order $2 - i$. Using symbol asymptotics, we can therefore pick any representative symbol \tilde{b} , such that

$$\tilde{b} \sim \sum_{i=1}^{\infty} b_i,$$

and then \tilde{b} will be a classical symbol of order 1. The fact that each of the operators Q_i is self-adjoint implies that the difference between $\tilde{b}(x, D)$ and $\text{Re}(\tilde{b}(x, D))$ is a smoothing operator. Thus if we let b be the symbol of $\text{Re}(\tilde{b}(x, D))$, then we find

$$b \sim \sum_{i=1}^{\infty} b_i,$$

the operator $Q = b(x, D)$ is self-adjoint, and $L - Q^2$ is a smoothing operator.

We claim that $\sqrt{L} - Q$ is also a smoothing operator. To see this, we employ heuristics from the holomorphic functional calculus. Since the spectrum of L is contained in (ε, ∞) for some $\varepsilon > 0$, and the function \sqrt{z} is holomorphic in a neighborhood of this region, if we consider the rectifiable curve $\gamma(t) = |t| + i \text{sgn}(t) \cdot t$, then we should expect to have

$$L^{-1/2} = \frac{1}{2\pi i} \int_{\gamma} z^{-1/2} (z - L)^{-1} dz.$$

The operator $(z - L)^{-1}$ is bounded, with operator norm $O(1/z)$ for large z , and operator norm $O(1)$ for small z , which implies that the operator-valued integral is absolutely convergent. Thus the left hand and right hand side of this equation both define bounded operators on $L^2(M)$. And for each of the eigenvectors e_i for L^{-1} ,

$$\frac{1}{2\pi i} \int_{\gamma} z^{-1/2} (z - L)^{-1} e_i dz = \left(\frac{1}{2\pi i} \int_{\gamma} z^{-1/2} (z - \lambda_i)^{-1} \right) e_i = \lambda_i^{-1/2} e_i,$$

so $L^{-1/2}$ and the integral formula have the same eigenvectors and eigenvalues, and are therefore equal. Similarly,

$$Q^{-1} = \frac{1}{2\pi i} \int_{\gamma} z^{-1/2} (z - Q^2)^{-1},$$

since Q^{-2} is bounded and positive definite, hence diagonalizable, and so one can apply analogous arguments to those above. If we set $R = L - Q^2$, then

$$\begin{aligned} L^{-1/2} - Q^{-1} &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/2} ((z - L)^{-1} - (z - Q^2)^{-1}) \\ &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/2} ((z - L)^{-1} - (z - L + R)^{-1}) \\ &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/2} (z - (L + R))^{-1} R (z - L)^{-1}. \end{aligned}$$

Because the operator norm of $(z - (L + R))^{-1}$ and $(z - L)^{-1}$ are each individually $O(1/z)$, and R is a smoothing operator, if we let T_z be the integrand of this operator, then for any $k > 0$,

$$\|T_z u\|_{L_k^2(M)} \lesssim_k \frac{1}{1 + z^2} \cdot \|u\|_{L^2(M)},$$

and from this we see that $L^{-1/2} - Q^{-1}$ is a smoothing operator. But now we can write

$$Q - L^{1/2} = Q(L^{-1/2} - Q^{-1})L^{-1/2},$$

and so we see that $Q - L^{1/2}$ is a smoothing operator. But this means that $Q - \sqrt{-\Delta}$ is smoothing.

13.7 Ψ DOs and Microsupport

Fix an open set $\Omega \subset \mathbf{R}^d$. The theory of microlocal analysis asks us to think of the singularities of a distribution $u \in \mathcal{D}(\Omega)^*$ as having both position *and* direction, i.e. existing on the wavefront set $\text{WF}(u) \subset T^*M$, and indicated by the directions that the Fourier transform of u does not decay rapidly, once sufficiently localized. The heuristics of pseudodifferential operators also ask us to view a distribution u as living on T^*M , such that a pseudodifferential operator S given by a symbol a acts on u by multiplying that part of u that ‘lives at’ the point (x, ξ) by the quantity $a(x, \xi)$. In particular, if a is made to decay rapidly in the directions that define the wavefront set u , then we should expect Su to be smooth, i.e. application of S annihilates the singularities of a . In this section, we elaborate on these ideas, as well as introducing further notions of the microlocal properties of pseudodifferential operators.

If $T : \mathcal{D}(\Omega) \rightarrow \mathcal{E}(\Omega)$ is a pseudodifferential operator with symbol a , then we define the microsupport $\mu\text{-supp}(T)$ of T to be equal to the microsupport $\mu\text{-supp}(a)$ of it’s symbol, defined in the parts of these notes on oscillatory integral distributions.

Lemma 13.15. *Let T be a pseudodifferential operator on Ω . Then it’s canonical relation is equal to*

$$\mathcal{C}_T = \{(x, x; \xi, \xi) : (x, \xi) \in \mu\text{-supp}(T)\}.$$

This follows because an open conic set $\Gamma \subset \Omega \times \mathbf{R}^d$ is disjoint from $\mu\text{-supp}(T)$ if and only if Γ is disjoint from $\text{WF}(Tu)$ for any $u \in \mathcal{E}(\Omega)^$.*

Proof. Let $T = a(x, D)$ for some symbol $a(x, \xi)$. It follows from the general theory of oscillatory integral distributions that the canonical relation of T is equal to

$$\mathcal{C}_T \subset \{(x, \xi; x, \xi) : (x, \xi) \in \mu\text{-supp}(T)\}.$$

Thus if Γ is disjoint from $\mu\text{-supp}(T)$, then it follows from the general theory that Γ is disjoint from $\text{WF}(Tu)$ for any $u \in \mathcal{E}(\Omega)^*$. If we assume the converse, then for any $(x_0, \xi_0) \in \Gamma$, we can find a pseudodifferential operator $S : \mathcal{D}(\Omega) \rightarrow \mathcal{E}(\Omega)$ with a symbol $b(x, \xi)$ supported on Γ and equal to one in a conic neighborhood of (x_0, ξ_0) . It follows that ST is a smoothing operator. It follows from the composition calculus that $a \cdot b \in \mathcal{S}^{-\infty}(\Omega \times \mathbf{R}^d)$. Thus $(x_0, \xi_0) \notin \mu\text{-supp}(a) = \mu\text{-supp}(T)$. \square

Similar to the argument above, the composition calculus implies that for any two properly supported pseudodifferential operators T and S ,

$$\mu\text{-supp}(TS) \subset \mu\text{-supp}(T) \cap \mu\text{-supp}(S),$$

and for any pseudodifferential operator T ,

$$\mu\text{-supp}(T^t) = \{(x, -\xi) : (x, \xi) \in \mu\text{-supp}(T)\} \quad \mu\text{-supp}(T^*) = \mu\text{-supp}(T).$$

One can prove these either using the microsupport of the symbols defining T and S , or the properties of the canonical relation of T and S .

Theorem 13.16. *Let Γ be a closed conic set, and suppose T is a properly pseudodifferential operator with $\mu\text{-supp}(T) \cap \Gamma = \emptyset$, then T maps $\mathcal{D}_\Gamma^*(\Omega)$ continuously into $\mathcal{E}(\Omega)$.*

Proof. TODO: For any open set $U \subset \Omega$, control the derivatives of $T\phi$ on U by decomposing T into inputs outside of U , a Ψ DO with a symbol in $\mathcal{S}^{-\infty}$, and a Ψ DO localized near $\mu\text{-supp}(T)$, and so on. \square

In fact, a sequence $\{u_n\}$ converges in $\mathcal{D}_\Gamma^*(\Omega)$ to some $u \in \mathcal{D}_\Gamma^*(\Omega)$ if and only if it converges distributionally to u , and $Tu_n \rightarrow Tu$, where T is an arbitrary properly supported Ψ DO with $\Gamma \cap \mu\text{-supp}(T) = \emptyset$. The proof is left to the reader.

Theorem 13.17. *Fix $u \in \mathcal{D}(\Omega)^*$. Then $(x_0, \xi_0) \notin \text{WF}(u)$ if and only if there exists a conic neighborhood Γ of (x_0, ξ_0) such that for any properly supported pseudodifferential operator T on Ω with $(x_0, \xi_0) \in \mu\text{-supp}(T)$, $Tu \in \mathcal{E}(\Omega)$.*

Remark. It follows from this theorem, and the fact that the class of pseudodifferential operators and the microsupport of such operators are invariant under diffeomorphism, that the wavefront set of a distribution is invariant under diffeomorphisms.

We have already seen the regularity theory of pseudodifferential operators. Recall that if T is a properly supported pseudodifferential operator of order t , then T maps $H_{\text{loc}}^s(\Omega) \rightarrow H_{\text{loc}}^{s-t}(\Omega)$. By virtue of the fact that this is an isomorphism if T is elliptic, one can define $H_{\text{loc}}^s(\Omega)$ to be the space of all distributions $u \in \mathcal{D}(\Omega)^*$ such that $Tu \in L_{\text{loc}}^2(\Omega)$, where T can be an arbitrary properly supported ΨDO of order m .

Let us also now look at the *microlocal regularity* of T , i.e. the regularity of T where we only care about regularity in certain conical subsets of $T^*\Omega$. For a conic open set $\Gamma \subset \Omega \times \mathbf{R}^d$, we define $H_{\Gamma, \text{loc}}^s(\Omega)$ to be the family of all distributions $u \in \mathcal{D}(\Omega)^*$ such that for any properly supported pseudodifferential operator T of order zero with conically compact microsupport $\mu\text{-supp}(T) \subset \Gamma$, $Tu \in H_{\text{loc}}^s(\Omega)$. By the results above, and the Sobolev embedding theorem, $\lim_{s \rightarrow \infty} H_{\Gamma, \text{loc}}^s(\Omega) = \mathcal{D}_{\Gamma}^*(\Omega)$.

Theorem 13.18. *If T is a properly supported pseudodifferential operator of order t on Ω , then T maps $H_{\Gamma, \text{loc}}^s(\Omega)$ into $H_{\Gamma, \text{loc}}^{s-t}(\Omega)$.*

Proof. Fix a conically compact set $\Gamma_1 \subset \Gamma$. If S is a properly supported pseudodifferential operator of order zero with $\mu\text{-supp}(S) \subset \Gamma$, it suffices to show that ST maps $H_{\Gamma, \text{loc}}^s(\Omega)$ into $H_{\Gamma, \text{loc}}^{s-t}(\Omega)$. But $ST = STU_1 + STU_2$, where U_1 and U_2 are properly supported pseudodifferential operators of order zero, such that $\mu\text{-supp}(U_1)$ is conically compact and contained in Γ , and $\mu\text{-supp}(U_2)$ is disjoint from Γ_1 . But this means that if $v \in H_{\Gamma, \text{loc}}^s(\Omega)$, then $U_1v \in H_{\text{loc}}^s(\Omega)$, and since ST is a properly support pseudodifferential operator of order t , $STU_1v \in H_{\text{loc}}^{s-t}(\Omega)$. On the other hand, $\text{WF}(U_2v)$, and thus $\text{WF}(TU_2v)$, is disjoint from Γ_1 , which implies that STU_2 is smooth, and thus lies in $H_{\text{loc}}^{s-t}(\Omega)$ trivially. \square

Ellipticity can also be microlocalized. If T was an elliptic pseudodifferential operator, we found a pseudodifferential parametrix S for T , i.e. such that $S \circ T$ and $T \circ S$ are smoothing operators. We say a symbol $a \in \dot{S}^t(\Omega)$ is *elliptic* on a conic open set $\Gamma \subset \Omega \times \mathbf{R}^d$ if for $(x, \xi) \in \Gamma$,

$$|a(x, \xi)| \sim \langle \xi \rangle^t$$

with an implicit constant uniform in ξ , and locally uniform in x . In this case, we can find a ‘parametrix’ on *conically compact* subcones of Γ .

Theorem 13.19. *If T is a pseudodifferential operator with a symbol $a \in \dot{S}^t(\Omega)$ which is elliptic on a conic set Γ , then there exists a pseudodifferential operator $S \in \dot{S}^{-t}(\Omega)$ such that $T \circ S$ and $S \circ T$ are regularizing on Γ .*

Proof. If Γ_1 is a conic open subset of Γ , we can find a pseudodifferential operator S with a symbol b by a recursive formula similar to that of the construction of a parametrix of an elliptic symbol, though applying cutoffs outside of Γ_1 so that the symbol vanishes outside of Γ and remains smooth and well defined. \square

As a consequence, if a Ψ DO T is elliptic on Γ , then for any compactly supported distribution u , it follows that

$$\text{WF}(Tu) \cap \Gamma = \text{WF}(u) \cap \Gamma,$$

i.e. the wavefront set is preserved on Γ .

If T is a properly supported classical pseudodifferential operator of order t with principal symbol $p(x, \xi)$, then the *characteristic set* of T is

$$\text{Char}(T) = \{(x, \xi) : p(x, \xi) = 0\}.$$

The characteristic set is closed and conic, and T is elliptic on the complement of $\text{Char}(T)$. Thus it follows that for $u \in \mathcal{D}(\Omega)^*$, $\text{WF}(u) \subset \text{WF}(Tu) \cup \text{Char}(T)$. Therefore, if $Tu \in \mathcal{E}(\Omega)$, then $\text{WF}(u) \subset \text{Char}(T)$.

13.8 Vector-Valued Pseudodifferential Operators

For certain applications of pseudodifferential operators, e.g. to systems of partial differential equations, it is useful to have a theory of vector-valued pseudodifferential operators acting on systems of distributions. The theory is almost entirely analogous to the scalar-valued theory, except that products of pseudodifferential operators may not commute.

We consider only finite dimensional vector-valued quantities here, though the generalization to Banach spaces is not too difficult to imagine. If V is a finite dimensional vector space, and $\Omega \subset \mathbf{R}^d$, we can define a family $\mathcal{D}(\Omega; V)$ of V -valued test functions, and thus obtain a family $\mathcal{D}(\Omega, V)^*$ of

V -valued distributions on Ω . These spaces are really just $\mathcal{D}(\Omega) \hat{\otimes} V$ and $\mathcal{D}(\Omega)^* \hat{\otimes} V$. Thus if V has a basis $\{e_1, \dots, e_n\}$, then elements of $\mathcal{D}(\Omega, V)$ can be uniquely expanded as $\phi_1 e_1 + \dots + \phi_n e_n$ for test functions $\{\phi_i\}$ in $\mathcal{D}(\Omega)$, and elements of $\mathcal{D}(\Omega, V)^*$ can be uniquely expanded as $u_1 e_1 + \dots + u_n e_n$ for distributions $\{u_i\}$ in $\mathcal{D}(\Omega)^*$. Similarly, we can define $\mathcal{E}(\Omega, V)$, $\mathcal{E}(\Omega, V)^*$, $H^s(\Omega, V)$, and virtually all of the other function spaces of interest to us in analysis, and these also just turn out to be tensor products.

If V and W are both finite dimensional linear spaces, then we have a natural isomorphism

$$L(X \otimes V, Y \otimes W) \cong L(X, Y) \otimes L(V, W).$$

In particular, after fixing bases $\{e_1, \dots, e_n\}$ and $\{e'_1, \dots, e'_m\}$ for V and W , an arbitrary operator T in $L(X \otimes V, Y \otimes W)$ can be written uniquely as

$$T \left(\sum_{i=1}^n x_i \otimes e_i \right) = \sum_{i=1}^n \sum_{j=1}^m T_{ij}(x_i) e_j$$

where T_{ij} lie in $L(X, Y)$. Thus every vector-valued continuous linear map is given by an $n \times m$ matrix of scalar-valued continuous linear maps.

It is trivial that we have a Schwartz kernel for such operators, i.e. for any $T : \mathcal{D}(\Omega_1, V) \rightarrow \mathcal{D}(\Omega_2, W)^*$, there exists a matrix-valued distribution $K \in \mathcal{D}(\Omega_2 \times \Omega_1, V \otimes W)^*$ such that

$$\langle T\phi, \psi \rangle = \int K(x, y) (\phi(x) \otimes \psi(y)) dx dy.$$

Let us now specialize to the study of vector valued pseudodifferential operators

We define a pseudodifferential operator T of order t on Ω , *valued in* $L(V, W)$, to be an operator from $\mathcal{D}(\Omega, V)$ to $\mathcal{E}(\Omega, W)^*$ induced by an element of $\Psi_{\text{loc}}^t(\Omega) \hat{\otimes} L(V, W)$. With any such operator we can associate $a : \Omega \times \mathbf{R}^d \rightarrow L(V, W)$, a *vector-valued symbol* of order t , such that

$$T\phi(x) = \int a(x, \xi) \phi(y) e^{2\pi i \xi \cdot (x-y)} dy.$$

The symbolic calculus remains unperturbed, except for a few modifications:

- The operators T^t and T^* are $L(W^*, V^*)$ -valued.
- If T is an $L(V, W)$ valued proper pseudodifferential operator of order t , and S an $L(W, U)$ valued operator of order s , then the operator $T \circ S$ is an $L(V, U)$ valued pseudodifferential operator of order $t + s$ with an analogous asymptotic expansion to the scalar-valued case, but the non-commutativity implies that the commutator $[T, S] = T \circ S - S \circ T$ is *not necessarily* a pseudodifferential operator of $t + s - 1$.
- If T is a classical $L(V, W)$ valued pseudodifferential operator with principal symbol $p(x, \xi)$, then the characteristic set of T is

$$\text{Char}(T) = \{(x, \xi) : p(x, \xi) : V \rightarrow W \text{ is not injective}\}.$$

Then T is elliptic if $\text{Char}(T) = \emptyset$. One can consider the analogous microlocal variants.

We can also consider pseudodifferential operators valued in linear maps between two vector bundles E and F on a space Ω , i.e. an operator $T : C^\infty(\Omega; E) \rightarrow C^\infty(\Omega; F)$ which, when taken in coordinates given by trivializations of E and F , look like matrix valued pseudodifferential operators. We can then associate such an operator T with a symbol, i.e. a family of sections $\mathbf{R}^n \rightarrow \Gamma(\Omega, L(E, F))$ with appropriate smoothness and decay. One can then define $\text{Char}(T)$ as above, and the notion of an elliptic operator.

Example. If M is a smooth manifold, then the exterior derivative $\Omega^n(M) \rightarrow \Omega^{n+1}(M)$ is a pseudodifferential operator. Let us focus on the case where M is an open submanifold of \mathbf{R}^d so that it has a natural coordinate system. Since

$$d(f dx^S) = \sum_{j \notin S} \frac{\partial f}{\partial x^j} dx^j \wedge dx^S = \sum_{j \notin S} (2\pi i) (D_x^j f) dx^j \wedge dx^S,$$

it follows that the symbol of the exterior derivative is

$$a(\xi) = \sum_{\substack{S \subset \{1, \dots, d\} \\ \#(S)=n}} \sum_{j \notin S} \sigma(j, S) \cdot 2\pi i \xi_j \cdot E_{S, S \cup \{j\}},$$

where $E_{S, T} : L(\Lambda^n, \Lambda^m)$ is simply the bundle map that maps dx^S to dx^T , and everything else to zero, and where $\sigma(j, S) \in \{-1, 1\}$ is chosen such that $dx^j \wedge$

$dx^S = \sigma(j, S) dx^{S \cup \{j\}}$. For $n = 0$ in particular, we have

$$a(\xi) = \sum_{j=1}^d 2\pi i \xi_j E_j,$$

thus we see that the exterior differential in this case is always injective, and so $d : \Omega^0(M) \rightarrow \Omega^1(M)$ is elliptic. For $n > 0$ on the other hand, is not elliptic. For instance, in the case $n = 1$, we have

$$a(\xi) = \sum_{i < j} 2\pi i (\xi_j E_{i, \{i, j\}} - \xi_i E_{j, i \cup \{i, j\}}).$$

But then we notice that for each fixed $\xi^0 \in \mathbf{R}^d$, $a(\xi^0)$ maps $\sum \xi_i^0 dx^i$, and thus $\text{Char}(T) = \Omega \times \mathbf{R}_{\xi}^d$.

Example. We recall that the principal symbol of a classical pseudodifferential operator on a manifold M is invariantly defined on T^*M . Given a manifold M , we can consider the line bundle $\text{Vol}^{1/2}(M)$ of half scalar densities. We claim there is a more rich invariant that can be associated with pseudodifferential operators from $\text{Vol}^{1/2}(M)$ to itself, namely, the subprincipal symbol, given in coordinates such that if T has symbol $a \sim \sum_{i=0}^{\infty} a_{m-i}$, then

$$a^{sp} = a_{m-1} + C \cdot \partial_j^x \partial_j^{\xi} a_m$$

TODO: GET COEFFICIENT C CORRECT. is invariantly defined as a $\text{Hom}(\text{Vol}^{1/2}(M))$ valued symbol of order $m-1$ on T^*M . In particular, if a_0 vanishes up to second order, the a_{m-1} is invariantly defined on T^*M .

13.9 The Index Theorem

Let $T \in \Psi_{\text{loc}}^0(\Omega)$. Then T is a continuous operator from $L_c^2(\Omega)$ to $L_{\text{loc}}^2(\Omega)$. We begin this section by determining what conditions ensure that this operator is compact, i.e. what conditions ensure that T maps bounded subsets of $L_c^2(\Omega)$ to bounded subsets of $L_{\text{loc}}^2(\Omega)$. This is equivalent to the induced maps $L^2(K) \rightarrow L_{\text{loc}}^2(\Omega)$ being compact for any compact set $K \subset \Omega$, and our job is simplified considerably by the following lemma.

Lemma 13.20. *Let X be a Banach space, and let Y be a Fréchet space. Then a bounded linear operator $T : X \rightarrow Y$ is compact if and only if for any sequence $\{x_i\}$ converging weakly to zero in X , $\{Tx_i\}$ converges in the standard topology of Y .*

Proof. TODO: Maybe move to functional analysis notes?

Since Y is a Fréchet space, T is compact if and only if for any bounded sequence $\{x_i\}$ in X , $\{Tx_i\}$ has a convergent subsequence in Y .

If T is compact, and a sequence $\{x_i\}$ converges weakly to zero, then that sequence is bounded, hence $\{Tx_i\}$ is precompact. But $\{Tx_i\}$ converges weakly to zero, since T is continuous from the weak topology on X to the weak topology on Y . But this means $\{Tx_i\}$ converges in norm to zero, since any subsequence has a further subsequence converging in norm to zero.

Conversely, suppose T maps a sequence converging weakly to zero to a sequence converging in norm. If $\{x_i\}$ is a bounded sequence in X , then by Banach-Alaoglu, there exists a subsequence of $\{x_i\}$ which is Cauchy in the weak topology, which without loss of generality we will assume to be $\{x_i\}$ itself. To show this implies $\{Tx_i\}$ converges in Y , we note that if d_Y is a translation invariant metric defining Y , and $\{Tx_i\}$ did not converge, then there would be $\varepsilon > 0$ and a strictly increasing pair of sequences $\{i_j\}$ and $\{i'_j\}$ such that for all j , $d_Y(Tx_{i_j}, Tx_{i'_j}) \geq \varepsilon$. But $x_{i_j} - x_{i'_j}$ converges to zero in the weak topology, which gives a contradiction. \square

Thus an operator $T : L_c^2(\Omega) \rightarrow L_{\text{loc}}^2(\Omega)$ is compact if and only if, for any compact set $K \subset \Omega$, and any sequence $\{f_i\}$ in $L^2(K)$ converging weakly to zero, $\{Tf_i\}$ converges to zero in $L_{\text{loc}}^2(\Omega)$, which means that for any other compact set $K_1 \subset \Omega$,

$$\lim_{i \rightarrow \infty} \int_{K_1} |Tf_i(x)|^2 dx = 0.$$

Any operator TODO-

Any smoothing operator $T : L_c^2(\Omega) \rightarrow L_{\text{loc}}^2(\Omega)$ is compact, since if $K \in C^\infty(\Omega \times \Omega)$, then it is certainly true that $K \in$

The dual of $L_c^2(\Omega)$ can be identified with $L_{\text{loc}}^2(\Omega)$. Thus a family $\{f_i\}$ converges to zero weakly in $L_c^2(\Omega)$ if and only if

$$\int f_i(x)g(x) dx \rightarrow 0$$

A family $\{f_i\}$ in $L_c^2(\Omega)$ converges weakly to zero precisely when it converges in $L_{\text{loc}}^2(\Omega)$ to zero.

Thus an operator $T : L_c^2(\Omega) \rightarrow L_{\text{loc}}^2(\Omega)$ is compact if and only if

13.10 Self-Adjoint Elliptic Pseudo-Differential Operators on a Compact Manifold

Given a scalar density $d\omega$ on a manifold M , we can consider $L^2(M)$ as a Hilbert space induced by the inner product

$$\langle u, v \rangle = \int u(x) \overline{v(x)} d\omega.$$

Thus given a scalar density we can consider the adjoint of a pseudodifferential operator on M by considering the operator as an unbounded operator on $L^2(M)$ with domain $H_c^k(M)$, and in particular, we can consider *self-adjoint unbounded* operators on M ; we are being somewhat lax with terminology here, by a self-adjoint operator we mean one which is self adjoint on $\mathcal{D}(\mathbf{R}^n)$ with the natural pairing, i.e. a formally self-adjoint operator. A self-adjoint unbounded operator is then an unbounded operator self-adjoint as defined in the theory of such operators. If T is a classical self-adjoint pseudodifferential operator of order t , then it follows from the calculus of pseudodifferential operators that its principal symbol is real-valued. If T is self-adjoint and also *elliptic*, and M is connected and has dimension bigger than one, it also follows that the principal symbol is either positive everywhere in $T^*\mathbf{R}^n - 0$, or negative everywhere for large ξ . Thus we might as well assume in what follows that the principal symbol is positive. We begin with a calculation which tells us we are safe to use the spectral calculus for unbounded operators for the class of operators considered here provided we are working on a compact manifold, so that $H_c^k(M) = H^k(M)$.

Theorem 13.21. *Suppose T is a self-adjoint pseudodifferential operator of order t on a compact manifold M . Then T is a closed, self-adjoint unbounded operator.*

Proof. Let us begin by showing T is closed. We begin by noting that T is a continuous operator from $\mathcal{D}(M)^*$ to itself. We desire to show

$$\{(u, v) \in H^t(M) \times L^2(M) : Tu = v\}$$

is closed in $L^2(M) \times L^2(M)$. If $\{u_i\}$ is a sequence in $H^t(M)$ converging in the $L^2(M)$ norm to some $u \in L^2(M)$, and $\{Tu_i\}$ converges in the $L^2(M)$ norm to some $v \in L^2(M)$, then $\{u_i\}$ converges distributionally to u , and so $Tu = v$. Applying elliptic regularity, since $v \in L^2(M)$, we conclude that $u \in H^t(M)$. Thus we have shown T is closed.

Next, we show that T is a self-adjoint unbounded operator. To do this, since T is already self-adjoint, it suffices to show that if $v \in L^2(M)$ induces an inequality of the form

$$|\langle Tu, v \rangle| \lesssim \|u\|_{L^2(M)}$$

for all $u \in H^t(M)$, then $v \in H^t(M)$. But this again follows by elliptic regularity. Indeed, the inequality then holds for $u \in C^\infty(M)$, and the distributional theory thus implies that for such u ,

$$\langle Tu, v \rangle = \langle u, Tv \rangle.$$

Thus we have an inequality of the form

$$|\langle u, Tv \rangle| \lesssim \|u\|_{L^2(M)}$$

for all $u \in C^\infty(M)$, and a density argument thus shows that $Tv \in L^2(M)$, so that $v \in H^t(M)$. \square

We begin our analysis by focusing on the case where T has *order one*, though we will later find many question about elliptic self-adjoint operators of other orders reduce to this question. If the principal symbol of this elliptic, self-adjoint classical operator T is positive away from the origin, one might expect T to be almost positive semidefinite, in some sense.

Lemma 13.22. *Let T be a classical self-adjoint elliptic pseudodifferential operator of order 1. Then there exists a classical self-adjoint elliptic pseudodifferential operator S of order 1/2 such that $T - S^*S$ is a smoothing operator.*

Proof. We proceed as in the proof that $\sqrt{-\Delta}$ is a classical pseudodifferential operator. Let S_0 be a classical self-adjoint pseudodifferential operator of order 1/2 with principal symbol $b_0(x, \xi) = a(x, \xi)^{1/2}$, then $S_0^*S_0$ is positive-semidefinite, and $T - S_0^*S_0$ is a classical pseudodifferential operator of order zero. Let r_1 denote the principal symbol of this operator. More generally, let us suppose that $T - (S_0 + \cdots + S_n)^*(S_0 + \cdots + S_n)$ is

a classical pseudodifferential operator of order $-n$, with principal symbol r_n . If we consider a classical pseudodifferential operator S_{n+1} of order $1/2 - (n+1)$ with principal symbol $b_{n+1}(x, \xi) = r_n(x, \xi)/2b_0(x, \xi)$, then $T - (S_0 + \cdots + S_{n+1})^*(S_0 + \cdots + S_{n+1})$ is a classical pseudodifferential operator of order $-(n+1)$. Thus if we choose a self-adjoint operator S of order $1/2$ such that $S \sim \sum_{i=0}^{\infty} S_i$, then $T - S^*S$ is smoothing. \square

Lemma 13.23. *Suppose T is a classical self-adjoint elliptic pseudodifferential operator of order 1 with a non-negative principal symbol. Then for sufficiently large $\lambda_0 > 0$, we have that for $u \in H_c^1(M)$,*

$$\langle (T + \lambda_0)u, u \rangle \sim \|u\|_{H^{1/2}(M)}^2,$$

where the implicit constants are locally uniform over u supported on a common compact subset of M .

Proof. Without loss of generality, we may assume that the support of the kernel of T is compact. Choose a classical, self-adjoint, elliptic pseudodifferential operator S of order $1/2$ such that $T - S^*S$ is smoothing. Then for $u \in H_c^1(M)$, there exists $\gamma_0 > 0$ such that

$$\left| \langle Tu, u \rangle - \|Su\|_{L^2(M)}^2 \right| = \left| \int_M (T - S^*S)u(x) \bar{u}(x) dx \right| \leq \gamma_0 \|u\|_{L^2(M)}^2.$$

Thus if $\gamma > \gamma_0$,

$$\langle (T + \gamma)u, u \rangle = \langle Tu, u \rangle + \gamma \|u\|_{L^2(M)}^2 \geq \|Su\|_{L^2(M)}^2 + (\gamma - \gamma_0) \|u\|_{L^2(M)}^2.$$

Thus we conclude that $T + \gamma$ is positive definite. Since S has principal symbol $a(x, \xi)^{1/2}$, it is elliptic, and so for $u \in H^{1/2}(M)$, if $\gamma > \gamma_0$.

$$\begin{aligned} \|u\|_{H^{1/2}(M)}^2 &\lesssim \|Su\|_{L^2(M)}^2 + \|u\|_{L^2(M)}^2 \\ &\lesssim |\langle Tu, u \rangle| + \|u\|_{L^2(M)}^2 \\ &= |\langle (T + \gamma)u, u \rangle - \langle \gamma u, u \rangle| + \|u\|_{L^2(M)}^2 \\ &\leq \langle (T + \gamma)u, u \rangle + \langle \gamma u, u \rangle + \|u\|_{L^2(M)}^2 \\ &\leq \langle (T + 2\gamma + 1)u, u \rangle. \end{aligned}$$

Thus if $\lambda_0 > 2\gamma_0 + 1$, then

$$\langle (T + \lambda_0)u, u \rangle \lesssim \|Su\|_{L^2(M)}^2 + \|u\|_{L^2(M)}^2 \lesssim \|u\|_{H^{1/2}(M)}^2. \quad \square$$

Theorem 13.24. Suppose T is a classical, self-adjoint pseudodifferential operator of order one on a compact manifold M , with a non-negative principal symbol. Then for suitably large $\lambda_0 > 0$,

- The operators $(T + \lambda_0)^k$ are isomorphisms from $H^k(M)$ to $L^2(M)$ for all $k > 0$.
- There exists an increasing family of real numbers $\{\lambda_i\}$ with $\lambda_i \rightarrow \infty$, and an orthogonal basis $\{e_i\}$ for $L^2(M)$ consisting solely of elements of $C^\infty(M)$, such that $Te_i = \lambda_i e_i$ for each $i > 0$. The functions $\{e_i\}$ satisfy

$$\|e_i\|_{H^k(M)} \sim_k (\lambda_0 + |\lambda_i|)^k$$

- If $u \in H^k(M)$, then $\langle u, e_i \rangle \lesssim_k \|u\|_{H^k(M)} (\lambda_0 + \lambda_i)^{-k}$.

Proof. The last result implies that $T + \lambda_0$ is an injective operator from $H^1(M)$ to $L^2(M)$ for sufficiently large $\lambda_0 > 0$. We have already seen that, since M is compact, $T + \lambda_0$ is a closed, self-adjoint unbounded operator on $L^2(M)$. But the theory of unbounded operators thus implies that because $T + \lambda_0$ is injective, it has dense image in $L^2(M)$. Thus we can define its inverse $R = (T + \lambda_0)^{-1}$, which is a densely defined continuous operator in $L^2(M)$, and thus extends to a continuous operator on all of $L^2(M)$. Because $T + \lambda_0$ is positive-definite, i.e. $\langle (T + \lambda_0)u, u \rangle > 0$ for all non-zero $u \in H^1(M)$, density implies that $\langle Rv, v \rangle > 0$ for all $v \neq 0$. Thus R is positive definite. If $u \in H^1(M)$, and $(T + \lambda_0)u = v$, i.e. $u = Rv$, then ellipticity implies that

$$\|Rv\|_{H^1(M)} \lesssim \|v\|_{L^2(M)} + \|Rv\|_{L^2(M)} \lesssim \|v\|_{L^2(M)}.$$

Conversely, continuity of pseudodifferential operators implies that

$$\|v\|_{L^2(M)} = \|(T + \lambda_0)u\|_{L^2(M)} \lesssim \|u\|_{H^1(M)}.$$

Thus density implies that R is an *isomorphism* from $L^2(M)$ to $H^1(M)$. In particular we note that that $T + \lambda_0$ is an isomorphism from $H^1(M)$ to $L^2(M)$, i.e. it was surjective in the first place rather than merely having dense image. Moving on, The Rellich-Kondrachov theorem thus implies that R is a *compact*, positive definite operator on $L^2(M)$. Thus the spectral analysis of such operators implies that there exists an orthogonal basis $\{e_i\}$

and a family of positive, decreasing eigenvalues $\{\gamma_i\}$ such that if E_i is orthogonal projection onto the span of e_i , then $(T + \lambda_0)^{-1} = \sum_i \gamma_i^{-1} E_i$. If $\lambda_i = \gamma_i - \lambda_0$, then this means that $T = \sum_i \lambda_i E_i$.

Applying elliptic regularity, since $(T + \lambda_0)e_i = \gamma_i e_i$ lies in $L^2(M)$, we conclude that $e_i \in H^1(M)$. But iterating this argument gives that $e_i \in H^N(M)$ for any $N > 0$. Applying the Sobolev embedding theorem thus shows that $e_i \in C^\infty(M)$ for each i .

A simple generalization of the argument above (the spectral theory, ellipticity, and so on) shows that $(T + \lambda_0)^k$ is actually an isomorphism from $H^k(M)$ to $L^2(M)$ for each $k > 0$. We thus find that for any $u \in H^k(M)$,

$$\|u\|_{H^k(M)}^2 \sim_k \|(T + \lambda_0)^k u\|_{L^2(M)}^2 = \sum (\lambda_i + \lambda_0)^{2k} |\langle f, e_i \rangle|^2.$$

Thus we find that

$$|\langle u, e_i \rangle| \lesssim_k \|u\|_{H^k(M)} |\lambda_i + \lambda_0|^{-k} \lesssim_k \|u\|_{H^k(M)} |1 + \lambda_0|^{-k}. \quad \square$$

The following, known as *Hörmander's square root trick*, allows us to extend the theory above to elliptic, self-adjoint operators of arbitrary order.

Theorem 13.25. *Fix integers p and q with $q \neq 0$, and set $t = p/q$. Suppose T is a classical, positive-definite, elliptic pseudodifferential operator of order t on a compact manifold M . Then the operator $T^{q/p}$, defined using the spectral functional calculus, is a classical, positive-definite, elliptic pseudodifferential operator of order 1, with principal symbol $a(x, \xi)^{q/p}$.*

Proof. The case $t = -1$ is easily seen to be true, since the assumptions imply T is invertible, and its inverse is a pseudodifferential operator modulo smoothing. Thus, without loss of generality, we may assume that $q = 1$, and that $p > 0$. We can certainly construct an operator $S \in \Psi_{\text{cl}}^1(M)$ which is self-adjoint, with principal symbol $a(x, \xi)^{1/p}$, such that $S^p - T$ is a smoothing operator. Applying the theory above, we see that S has an eigenvalue decomposition, with eigenvectors lying in $C^\infty(M)$, and only finitely many eigenvalues being negative. Thus, modulo a smoothing operator, i.e. a finite sum of the projection operators $\{E_i\}$, we may actually replace S with a positive-definite operator. We claim that $S - T^{1/p}$ is also a smoothing operator, which would complete the proof since the principal symbol of S is equal to $a(x, \xi)^{q/p}$. Since S and T are both positive-definite, they are injective, and so by the spectral calculus of unbounded operators, T^{-1} and S^{-1}

is a well-defined bounded, positive-definite operator on $L^2(M)$. If γ is a contour in the complex plane which winds around each element of $\sigma(T^{-1})$ exactly once, then we find using the holomorphic functional calculus that

$$T^{1/p} = \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} (z - T^{-1})^{-1} dz$$

and

$$S = \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} (z - S^{-p})^{-1} dz$$

Now S^{-p} is a pseudodifferential operator of order $-p$. Since $S^p - T$ is a smoothing operator, the composition calculus tells us that so is $T^{-1}(S^p - T)S^{-p} = T^{-1} - S^{-p}$. Write $S^{-p} = T^{-1} - R$. Then

$$\begin{aligned} S - T^{1/p} &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} ((z - T^{-1} + R)^{-1} - (z - T^{-1})^{-1}) \\ &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} ((z - T^{-1} + R)^{-1} R (z - T^{-1})^{-1}). \end{aligned}$$

This is an integral over a compact curve over a continuous family of smoothing operators, which is therefore a smoothing operator. \square

Remark. As above, finding an operator S of order $t/2$ such that $T - S^*S$ is smoothing allows one to show that if T is a classical, elliptic, self-adjoint pseudodifferential operator of order t , then $T + \lambda_0$ is positive definite for suitably large $\lambda_0 > 0$, so that we may apply the theorem above to a wider situation.

Using the trick, we quickly extend the theory above to general operators of this form.

Theorem 13.26. *Suppose T is a classical, self-adjoint, elliptic pseudodifferential operator of order $t = p/q$ on a compact manifold M . Then for sufficiently large $\lambda_0 > 0$,*

- $(T + \lambda_0)^k$ is an isomorphism from $H^{tk}(M)$ to $L^2(M)$ for all $k > 0$.
- There is a basis $\{e_i\}$ for $L^2(M)$ consisting of elements of $C^\infty(M)$ and an increasing sequence of positive real numbers $\{\lambda_i\}$ with $\lambda_i \rightarrow \infty$, such that $Te_i = (\lambda_0 + \lambda_i)^t e_i$. The functions $\{e_i\}$ satisfy

$$\|e_i\|_{H^k(M)} \sim_k (\lambda_0 + |\lambda_i|)^k$$

- If $u \in H^k(M)$, then $|\langle u, e_i \rangle| \lesssim_k \|u\|_{H^k(M)} (\lambda_0 + |\lambda_i|)^{-k}$.

The special case of this result, where Δ is the Laplace-Beltrami operator on a compact Riemannian manifold, was first conjectured by Ohm and Raleigh, and proved by Hilbert in 1904.

Example. Let us explicitly check the result of this theorem in a well studied example. The operator $-\Delta$ is a classical positive semi definite elliptic pseudodifferential operator of order 2 on $\mathbf{T}^n = \mathbf{R}^n / \mathbf{Z}^n$. The theorem above, applied to $\alpha - \Delta$ for each $\alpha > 0$, shows that there is a basis $\{e_i\}$ for $L^2(\mathbf{T}^n)$, and a sequence of non-negative real numbers $\{\lambda_i\}$, such that $\Delta e_i = -\lambda_i^2 e_i$, and that if $u \in H^k(\mathbf{T}^n)$, then

$$|\langle u, e_i \rangle| \lesssim_k \|u\|_{H^k(M)} (1 + \lambda_i)^{-k}.$$

In this case, we can explicitly compute the basis $\{e_i\}$. It is precisely the functions $e_\xi(x) = e^{2\pi i \xi \cdot x}$, where $\xi \in \mathbf{Z}^n$, with corresponding eigenvalue $\lambda_\xi = 2\pi |\xi|$ since $\Delta e_\xi = -4\pi^2 |\xi|^2 e_\xi$. In this case the Fourier transform explicitly tells us the slightly stronger identity

$$|\langle u, e_\xi \rangle| = |\hat{u}(\xi)| \lesssim_k \min \left(\|u\|_{L^1(\mathbf{T}^n)}, \|D^k u\|_{L^1(\mathbf{T}^n)} |\xi|^{-k} \right).$$

13.11 Distribution of Eigenvalues and the Half Wave Operator

Consider the setup we began our study of in the last section. We have a classical, self-adjoint, elliptic pseudodifferential operator T of order one on a compact manifold M with principal symbol $p(x, \xi) \geq 0$, and we have associated with T a sequence of orthonormal smooth functions $\{e_i\}$ in $C^\infty(M)$ which form a basis for $L^2(M)$, and an increasing sequence $\{\lambda_i\}$ tending to ∞ , such that $T e_i = \lambda_i e_i$. Our goal now is to further the study of the behaviour of the eigenfunctions $\{e_i\}$, and the distributions of the eigenvalues $\{\lambda_i\}$. A key object in the study of the distribution of the eigenvalues is the function $N(\lambda)$, which gives the number of eigenvalues of T less than or equal to λ . Our results already give a very basic estimate for N , which we will considerably sharpen over the course of this section.

Theorem 13.27. *The function N is tempered in λ . More precisely, for $\lambda > 0$,*

$$N(\lambda) \lesssim_\varepsilon \lambda^{n+\varepsilon}.$$

Proof. Define the operators S_λ , which are the projections in $L^2(M)$ onto the eigenspaces corresponding to eigenvalues with value $\leq \lambda$. Then S_λ clearly has kernel

$$K_\lambda(x, y) = \sum_{\lambda_i \leq \lambda} e_i(x) \overline{e_i(y)}.$$

This function is smooth, and

$$\int K_\lambda(x, x) dx = \sum_{\lambda_i \leq \lambda} |e_i(x)|^2 = N(\lambda).$$

Thus $N(\lambda)$ is the trace of the operator S_λ . Because $\|e_i\|_{H^k(M)} \lesssim_k (1 + |\lambda|)^k$ if $\lambda_i \leq \lambda$, we have the elementary estimate

$$\|S_\lambda f\|_{H^k(M)} \lesssim_k \lambda^k \|f\|_{L^2(M)}.$$

The Sobolev embedding theorem thus implies that if $\dim(M) = d$, and if $\varepsilon > 0$, then

$$\|S_\lambda f\|_{L^\infty(M)} \lesssim_\varepsilon \|S_\lambda f\|_{H^{d/2+\varepsilon}(M)} \lesssim \lambda^{d/2+\varepsilon} \|f\|_{L^2(M)}.$$

Thus we conclude that, for any $f \in C^\infty(M)$, and any $x \in M$,

$$\left| \int K_\lambda(x, y) f(y) dy \right| \lesssim_\varepsilon \lambda^{d/2+\varepsilon} \|f\|_{L^2(M)}.$$

By density, this implies that if $g_{\lambda, x}(y) = K_\lambda(x, y)$, then

$$\|g_{\lambda, x}\|_{L^2(M)} \lesssim_\varepsilon \lambda^{d/2+\varepsilon}.$$

But $g_{\lambda, x} \in \text{span}\{e_i : \lambda_i \leq \lambda\}$, so this means that

$$\|g_{\lambda, x}\|_{L^\infty(M)} = \|S_\lambda g_{\lambda, x}\|_{L^\infty(M)} \lesssim_\varepsilon \lambda^{d/2+\varepsilon} \|g_{\lambda, x}\|_{L^2(M)} \lesssim_\varepsilon \lambda^{d+2\varepsilon}.$$

But this means we conclude that $\|K_\lambda\|_{L^\infty(M \times M)} \lesssim_\varepsilon \lambda^{d+\varepsilon}$, and thus that

$$N(\lambda) = \int K_\lambda(x, x) dx \lesssim_\varepsilon \lambda^{d+\varepsilon}. \quad \square$$

Our next goal is now to establish the *Sharp Weyl formula*. If

$$c = \int_{\{(x,\xi) \in T^*M : a(x,\xi) \leq 1\}} d\xi \, dx,$$

then the result states that

$$N(\lambda) = c\lambda^n + O(\lambda^{n-1}).$$

We will continue to use the representation $N(\lambda) = \text{Tr}(S_\lambda)$. However, a key step here is to recognize that S_λ is a *function* of T , namely $S_\lambda = \chi_\lambda(T)$, where $\chi_\lambda(\xi) = \mathbf{I}(\xi \leq \lambda)$. The next trick is to use the Fourier multiplication formula to obtain a useful representation of the structure of S_λ . For $f \in C^\infty(M)$, we can write

$$\begin{aligned} S_\lambda f &= \int_{-\infty}^{\infty} \chi_\lambda(\tau) \left(\sum_i \delta_{\lambda_i}(\tau) E_i f \right) d\tau \\ &= \int_{-\infty}^{\infty} \hat{\chi}_\lambda(t) \left(\sum_i e^{2\pi i t \lambda_i} E_i f \right) dt \\ &= \int_{-\infty}^{\infty} \hat{\chi}_\lambda(t) e^{2\pi i t T} dt \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{2\pi i t(T-\lambda)} f}{t + i0} dt. \end{aligned}$$

Thus we lead to study the *half wave operator* $\partial_t - 2\pi i T$, since one sees that $(\partial_t - 2\pi i T)\{e^{2\pi i t T} f\} = 0$ for all $f \in L^2(M)$; it is easily verified for the eigenfunctions of T , and then the result follows by the distributional continuity of all operations involved.

We will understand the operators $e^{2\pi i t T}$ by constructing a *parametrix* for the equation $\partial_t - 2\pi i T$ over small times, i.e. for $|t| \lesssim 1$. This equation is a pseudodifferential variant of a hyperbolic partial differential equation, so based on the parametrix construction in this scenario, we might hope to find a parametrix defined by an oscillatory integral of the form

$$Sf(x, t) = S(t)f(x) = \int s(t, x, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)} f(y) \, dy \, d\xi,$$

such that:

- $\Phi(t, x, y, \xi) = \phi(x, y, \xi) + tp(y, \xi)$, where ϕ is smooth away from $\xi = 0$, homogeneous of degree one, and $\phi(x, y, \xi) \approx (x - y) \cdot \xi$ in the sense that on the support of s ,

$$\partial_\xi^\beta \{ \phi(x, y, \xi) - (x - y) \cdot \xi \} \lesssim_\beta |x - y|^2 |\xi|^{1-\beta}.$$

In particular, this implies that $\phi(x, y, \xi) = 0$ when $(x - y) \cdot \xi = 0$.

- s is a symbol of order zero, supported on $|x - y| \lesssim 1$ and on $|\xi| \geq 1$, in such a way that

$$|\nabla_\xi \phi(x, y, \xi)| \gtrsim |x - y|$$

and

$$|\nabla_x \phi(x, y, \xi)| \gtrsim |\xi|$$

for $(x, y) \in \text{supp}_x(s) \times \text{supp}_y(s)$.

If $(\partial_t - 2\pi iT) \circ S = 0$ is a smoothing operator on $(-\varepsilon, \varepsilon) \times M$ and $S(0)$ differs from the identity operator by a smoothing operator, then $S - e^{2\pi itT}$ is smoothing for $|t| \leq \varepsilon$. This will therefore give us much more information about the behaviour of the propagators $e^{2\pi itT}$ over small times.

To find a choice of ϕ and s which gives us this parametrix, let us start by determining what properties these functions should satisfy. Let us fix a coordinate system (x, U) , where $x(U)$ is a precompact subset of \mathbf{R}^n . Let us assume that in these coordinates, T has symbol $a(x, \xi)$. Then the kernel of $(\partial_t - 2\pi iT) \circ S$ in this coordinate system is

$$\int (\partial_t + 2\pi iT(x, D)) \left\{ s(t, \cdot, y, \xi) e^{2\pi i \Phi(t, \cdot, y, \xi)} \right\} d\xi.$$

If we set

$$s'(t, x, y, \xi) = e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t - 2\pi iT(x, D)) \left\{ s(t, \cdot, y, \xi) e^{2\pi i \Phi(t, \cdot, y, \xi)} \right\}$$

then the kernel is

$$\int s'(t, x, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)} d\xi.$$

Provided that s' is a symbol of order $-\infty$ for $0 < |t| \leq \varepsilon$, integration by parts shows that $(\partial_t + 2\pi iT) \circ S$ is smoothing, and so we will try to choose ϕ and s so as to obtain such a result.

In our discussion of pseudodifferential operators, we have already discussed an asymptotic formula for s' , namely, if

$$r_{x,y}(z) = \nabla_x \phi(x, z, \xi) \cdot (x - z) - \{\phi(x, y, \xi) - \phi(z, y, \xi)\}.$$

then for any $N > 0$, if $a \sim \sum_{k=-\infty}^1 a_k$, where a_k is homogeneous of degree k , and if $\xi_\phi = \nabla_x \Phi(t, x, y, \xi) = \nabla_x \phi(x, y, \xi)$,

$$\begin{aligned} s'(t, x, y, \xi) &= \underbrace{(p(y, \xi) - a(x, \xi_\phi)) \cdot s(t, x, y, \xi)}_{\text{symbols of order 1}} \\ &+ \underbrace{\partial_t s(t, x, y, \xi)}_{\text{symbol of order 0}} \\ &- \sum_{1 \leq |\beta| < N} \underbrace{\frac{2\pi i}{\beta! \cdot (2\pi i)^\beta} \cdot \partial_\xi^\beta a(x, \xi_\phi) \partial_z^\beta \{e^{2\pi i r_{x,y}(z)} s(t, z, y, \xi)\}}_{\text{symbols of order } 1 - \lceil |\beta|/2 \rceil} \Big|_{z=y} \\ &+ R_N(t, x, y, \xi). \end{aligned}$$

where, because $|\nabla_x \Phi(t, x, y, \xi)| \gtrsim |\xi|$ on the support of s ,

$$\langle \xi \rangle^{t - [N/2]} R_N \in L^\infty((-\varepsilon, \varepsilon) \times U \times U \times \mathbf{R}^d).$$

It is simple to establish estimates of the form

$$|\partial_x^\alpha \partial_y^\beta \partial_\xi^\lambda s'(t, x, y, \xi)| \lesssim \langle \xi \rangle^{N_{\alpha\beta\lambda}}.$$

Thus if we can justify that $|s'(t, x, y, \xi)| \lesssim_N \langle \xi \rangle^{-N}$ for all $N > 0$, then it will follow that s' is a symbol of order $-\infty$. We now determine the properties of the symbol s and the symbol ϕ which will give us these estimates.

To begin with, let us specify the function ϕ . In order to guarantee that s' is a symbol of order zero, the expansion above shows that $(p(y, \xi) - p(x, \xi_\phi)) \cdot s(t, x, y, \xi)$ must be a symbol of order zero. This will be true if we can pick ϕ such that, on the support of s , and for $|\xi| \gtrsim 1$,

$$p(x, \nabla_x \phi(x, y, \xi)) = p(y, \xi).$$

This is an example of an *Eikonal equation*, e.g. an equation of the form

$$q(z, \nabla_z \psi(z)) = 0$$

for some function $q(z, \zeta)$. In our case, $z = (x, y, \xi)$, so $\zeta = (\zeta_x, \zeta_y, \zeta_\xi)$, and so

$$q(z, \zeta) = p(x, \zeta_x) - p(y, \xi).$$

Let us make some further remarks we desire about our choice of function ϕ :

- We want ϕ to be homogeneous and smooth away from the origin. If we solve the equation for all $|\xi| = 1$, and then extend ϕ such that for $\lambda > 0$ and $|\xi| = 1$,

$$\phi(x, y, \lambda \xi) = \lambda \phi(x, y, \xi) \psi(\lambda),$$

where ψ is smooth, equal to one for $|\lambda| \geq 3/4$, and vanishing for $|\lambda| \leq 1/2$, then ϕ will satisfy the equation for all $|\xi| \gtrsim 1$. This means that

$$p(x, \nabla_x \phi(x, y, \xi)) - p(y, \xi)$$

is smooth and supported on $|\xi| \lesssim 1$, which implies it is a symbol of order $-\infty$, which suffices for our construction. Thus it suffices to solve the equation for $|\xi| = 1$.

- Since ϕ is smooth away from the origin and homogeneous, the equation

$$|\partial_\xi^\beta \{ \phi(x, y, \xi) - (x - y) \cdot \xi \}| \lesssim_\beta |x - y|^2 \langle \xi \rangle^{1-\beta}$$

holds if, for $|\xi| = 1$, we have $\phi(x, y, \xi) = 0$ whenever $(x - y) \cdot \xi = 0$, and $\nabla_x \phi(x, y, \xi) = \xi$ whenever $x = y$. Thus we have some *initial conditions* for our Eikonal equation.

The second condition constitutes a type of initial condition for ϕ , since it specifies it's behaviour on a hypersurface, a kind of Cauchy condition, and thus we should expect these are close to the conditions that give unique solutions to the equation. And the following Lemma indeed shows that there is a unique function ϕ , defined for $|x - y| \lesssim 1$ and $|\xi| = 1$ with these properties.

Lemma 13.28. *Let Z be a smooth manifold, and let $q(z, \zeta)$ be a real-valued, smooth function defined locally around a point $(z_0, \zeta_0) \in T^*Z$. Let S be a smooth hypersurface in Z passing through z_0 with conormal vector ζ_S at z_0 , such that*

$$\frac{\partial q}{\partial \zeta_S}(z_0, \zeta_0) = \lim_{t \rightarrow 0} \frac{q(z_0, \zeta_0 + t \zeta_S) - q(z_0, \zeta_0)}{t}$$

is nonzero. Suppose that ψ is any smooth function defined on S locally about z_0 , such that $d\psi(z_0)$ agrees with the action of ζ_0 on $T_{x_0}S$. Then there exists a unique smooth function ϕ defined in a neighborhood of z_0 , which agrees with ψ on S , satisfies the Eikonal equation $q(z, \nabla_z \phi(z)) = 0$, and has $\nabla_z \phi(z_0) = \zeta_0$.

Proof. TODO: See Sogge, Theorem 4.1.1. \square

In our case,

$$Z = \{(x, y, \xi) : |\xi| = 1\}.$$

We have $z_0 = (x_0, x_0, \xi_0)$, $\zeta_0 = (\xi, \xi, 0)$, and

$$S = \{(x, y, \xi) : |\xi| = 1 \text{ and } (x - y) \cdot \xi = 0\}.$$

The conormal vector ξ_S of S at z_0 is a multiple of $(\xi_0, -\xi_0, 0)$, and so by homogeneity,

$$\frac{\partial q}{\partial \xi_S} = \lim_{t \rightarrow 0} \frac{p(x_0, (1+t)\xi_0) - p(x_0, \xi_0)}{t} = p(x_0, \xi_0),$$

which is nonvanishing because T is elliptic. If we define ψ equal to zero on S , then $d\psi = 0$, which agrees with the action of ζ_0 on S . Thus the theorem applies local uniqueness and existence to solutions to the Eikonal equation, and by compactness of Z we can patch such solutions together to find a solution defined for all $|x - y| \lesssim 1$.

We therefore conclude that there exists a unique choice of ϕ such that, if s has small enough support, $s'(t, x, y, \xi)$ is a symbol of order zero. Next, let us see what constraints are forced on us in order to ensure that $S(0)$ differs from the identity by a smoothing operator. The kernel of U is precisely

$$\int s(0, x, y, \xi) e^{2\pi i \phi(x, y, \xi)} d\xi.$$

We now show that this operator is actually a *pseudodifferential operator* of order zero, and determine its symbol up to first order.

To do this, we write $\phi_\alpha(x, y, \xi) = (1 - \alpha)\phi(x, y, \xi) + \alpha(x - y) \cdot \xi$. Let U_α be the operator with kernel

$$\int s(0, x, y, \xi) e^{2\pi i \phi_\alpha(x, y, \xi)} d\xi.$$

Assume the support of s is close enough to the diagonal such that

$$|\nabla_\xi \phi_\alpha(x, y, \xi)| \gtrsim |x - y|$$

on the support of s . Then $\partial_\alpha^n U_t$ has kernel

$$\int (2\pi i)^n (\phi_1 - \phi_0)^n s(0, x, y, \xi) e^{2\pi i \phi_\alpha(x, y, \xi)} d\xi.$$

This is an oscillatory integral defined by a symbol of order n . However, when $t = 1$, the fact that $\phi(x, y, \xi) \approx (x - y) \cdot \xi$, together with the formula for converting pseudodifferential operators with compound symbols into standard Kohn-Nirenberg type symbols shows that $\partial_\alpha^n U_1$ is actually a pseudodifferential operator of order $-n$. Integration by parts, similarly, shows that $\partial_\alpha^n U_t$ is defined by an oscillator integral against a symbol of order $-n$. But this means that if we define a pseudodifferential operator by the asymptotic formula

$$V \sim \sum \frac{(-1)^n}{n!} \partial_\alpha^n U_1,$$

then $U - V$ is smoothing. Indeed, for any n , by Taylor's formula we have

$$U = \sum_{k=0}^{n-1} \frac{(-1)^k}{k!} \partial_\alpha^k U_1 + \frac{(-1)^n}{n!} \int_0^1 \alpha^{n-1} \partial_\alpha^n U_\alpha d\alpha$$

The integral here is an oscillatory integral defined against a symbol of order $-n$, and thus taking $n \rightarrow \infty$ verifies the claim.

It is an important remark that reversing this argument shows that *any* pseudodifferential operator can be written in the form above for the particular choice of ϕ we have given. This is a special case of the *equivalence of phase functions* theorem. This in particular guarantees that we can choose a symbol $I(x, y, \xi)$ of order zero such that $U - 1$ is smoothing if and only if $s(0, x, y, \xi) - I(x, y, \xi)$ is a symbol of order $-\infty$. The symbol I can be chosen to be vanishing for $|x - y| \gtrsim 1$, since the difference will be a smoothing pseudodifferential operator.

Next, the quantity

$$\begin{aligned} & \partial_t s(t, x, y, \xi) \\ & + \sum_{k=1}^d \partial_\xi^k a(x, \xi_\phi) \partial_x^k s(t, x, y, \xi) \\ & + \left(a_0(x, \xi_\phi) + \frac{1}{2\pi} \sum_{|\beta|=2} \partial_\xi^\beta p(x, \xi_\phi) \partial_x^\beta \phi(x, y, \xi) \right) s(t, x, y, \xi). \end{aligned}$$

must be a symbol of order -1 . But because the coefficients of this equation are smooth, and all derivatives are bounded, it follows from the general theory of transport equations that there exists a unique smooth function s_0 defined for $|t| \leq \varepsilon$, which is a symbol of order zero, such that $s_0(0, x, y, \xi) = I(x, y, \xi)$, s_0 vanishes for $|x - y| \gtrsim 1$, and satisfies the transport equation

$$\begin{aligned} & \partial_t s_0(t, x, y, \xi) \\ & + \sum_{k=1}^d \partial_\xi^k a(x, \xi_\phi) \partial_x^k s_0(t, x, y, \xi) \\ & + \left(a_0(x, \xi_\phi) + \frac{1}{2\pi} \sum_{|\beta|=2} \partial_\xi^\beta p(x, \xi_\phi) \partial_x^\beta \phi(x, y, \xi) \right) s_0(t, x, y, \xi) = 0. \end{aligned}$$

We have thus justified that the quantity

$$R_0(t, x, y, \xi) = e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t - 2\pi i T)(s_0(t, \cdot, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)})$$

is a symbol of order -1 . Now we come to a quirk of this parametrix, which does not occur in the study of hyperbolic partial differential equations. Since the operator $P(x, D)$ is only *pseudolocal* rather than completely local, the remainder term R_0 is *not* necessarily supported on a neighborhood of the origin. To fix this, we now successively define the terms $\{s_k\}$ for $k < 0$, which are symbols of order $-k$, such that $s_k(0, x, y, \xi) = 0$, and

TODO : SPECIFY REQUIRED EQUATION.

Again, solutions exist for small time periods. And this implies that $e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t + 2\pi i T)((s_0 + \dots + s_{-k}) e^{2\pi i \Phi(t, x, y, \xi)})$ is a symbol of order $-k$ (TODO: Is It), and we can continue the calculation to complete the argument.

Let us now use this equation to prove the sharp Weyl formula, which immediately is given by the following, more precise estimate.

Theorem 13.29. *Let M be a compact manifold, and let T be a classical, self-adjoint, elliptic pseudodifferential operator of order one on M , with principal symbol $p(x, \xi)$. If S_λ is the projection operator onto the span of eigenfunctions with eigenvalue $\leq \lambda$, and K_λ is the kernel of the operator S_λ , then*

$$K_\lambda(x, x) = c(x)\lambda^n + O(\lambda^{n-1}),$$

where

$$c(x) = \int_{p(x, \xi) \leq 1} d\xi.$$

Proof. Recall that we have

$$S_\lambda f(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{2\pi i t(T-\lambda)} f}{t + i0} dt.$$

This integral only has a singularity when $t = 0$, so it makes sense to decompose the integral into two parts using an even function $\rho \in C_c^\infty(\mathbf{R})$ supported on $|t| \leq \varepsilon/2$ and equal to one for $|t| \leq \varepsilon/4$, i.e. writing

$$S_\lambda f = S_{\lambda, \text{low}} f + S_{\lambda, \text{high}} f,$$

where

$$S_{\lambda, \text{low}} f = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\rho(t)}{t + i0} e^{2\pi i t(T-\lambda)} f dt$$

and

$$S_{\lambda, \text{high}} f = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1 - \rho(t)}{t} e^{2\pi i t(T-\lambda)} f dt.$$

To understand $S_{\lambda, \text{low}} f$, we use the parametrix for the wave equation. If we switch to coordinates around a particular point, then we find that the kernel of $S_{\lambda, \text{low}}$ is equal to $\tilde{K}_\lambda + R_\lambda$, where \tilde{K}_λ is supported on a neighborhood of the diagonal, and locally in coordinates we can write

$$\tilde{K}_\lambda(x, y) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t + i0} s(t, x, y, \xi) e^{2\pi i (\phi(x, y, \xi) + t(p(y, \xi) - \lambda))} d\xi dt.$$

and where

$$R_\lambda(x, y) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\rho(t)}{t + i0} A(t, x, y) e^{2\pi i t \lambda} dt,$$

where A is smooth for $|t| \leq \varepsilon$ and for $x, y \in M$. But since $\rho \cdot A$ is smooth, with compact support in the t variable taking inverse Fourier transforms in the t variable shows that

$$|R_\lambda(x, y)| = \left| \int_{-\infty}^{\lambda} \mathcal{F}_t^{-1}\{\rho A\}(\lambda - \tau, x, y) d\tau \right| \lesssim 1,$$

By Taylor's formula, we can write $s(t, x, x, \xi) = s(0, x, x, \xi) + t \cdot r(t, x, \xi)$, where r is a symbol of order zero. But if N is sufficiently large, we have

$$\begin{aligned} & \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t + i0} (t \cdot r(t, x, \xi)) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \rho(t) r(t, x, \xi) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt \\ &= \frac{1}{2\pi} \int_{\mathbf{R}^n} \mathcal{F}_t^{-1}\{\rho \cdot r\}(p(x, \xi) - \lambda, x, \xi) d\xi \\ &= \int O_N(1 + |p(x, \xi) - \lambda|)^{-N} d\xi = O(\lambda^{n-1}). \end{aligned}$$

We also have $s(0, x, x, \xi) = 1 + s_{-1}(x, \xi)$, where s_{-1} is a symbol of order -1 . Now if $\tilde{\chi}_\lambda = \chi_\lambda * \mathcal{F}_t^{-1}\{\rho\}$, then

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t + i0} s_{-1}(x, \xi) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt = \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) s_{-1}(x, \xi) d\xi.$$

Then $\tilde{\chi}_\lambda(p(x, \xi)) \sim 1$ for $|\xi| \lesssim \lambda$, and for $|\xi| \gtrsim \lambda$,

$$\tilde{\chi}_\lambda(p(x, \xi)) \lesssim_N \langle |\xi| - \lambda \rangle^{-N}$$

This implies that

$$\left| \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) s_{-1}(x, \xi) d\xi \right| \lesssim \int_0^{\lambda+O(1)} r^{n-2} + \int_{\lambda+O(1)}^{\infty} r^{n-1} (r - \lambda)^{-N} \lesssim \lambda^{n-1}.$$

Thus we have

$$\begin{aligned}
\tilde{K}_\lambda(x, x) &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t + i0} e^{2\pi i t(p(x, \xi) - \lambda)} + O(\lambda^{n-1}) \\
&= \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) d\xi + O(\lambda^{n-1}) \\
&= \int_{\mathbf{R}^n} \tilde{\chi}(p(x, \xi) - \lambda) d\xi + O(\lambda^{n-1}) \\
&= c(x)\lambda^n + \int_{\mathbf{R}^n} (\tilde{\chi} - \chi)(p(x, \xi) - \lambda) d\xi + O(\lambda^{n-1}).
\end{aligned}$$

Now $\tilde{\chi} - \chi = \chi * u$, where u is the inverse Fourier transform of $\rho(t) - 1$. But then

$$(\chi * u)(\tau) = \int_{-\infty}^0 \hat{\rho}(\tau - \omega) d\omega - \mathbf{I}(\tau \geq 0),$$

from which we verify that

$$|(\chi * u)(\tau)| \lesssim_N (1 + |\tau|)^{-N}.$$

But we can use this to again show that

$$\begin{aligned}
\left| \int_{\mathbf{R}^n} (\tilde{\chi} - \chi)(p(x, \xi) - \lambda) d\xi \right| &\lesssim \int_0^{\lambda - O(1)} r^{n-1} |\xi|^{-N} dr \\
&\quad + \int_{\lambda - O(1)}^{\lambda + O(1)} (\lambda - r)^{-N} r^{n-1} dr \\
&\quad + \int_{\lambda + O(1)}^{\infty} r^{n-1} (r - \lambda)^{-N} \\
&\lesssim \lambda^{n-1}.
\end{aligned}$$

Thus we have established the result for the kernel $K_{\lambda, \text{low}}$ of the operator $S_{\lambda, \text{low}}$. The proof of this theorem would be finished if we were able to justify that

$$|K_\lambda(x, x) - K_{\lambda, \text{low}}(x, x)| \lesssim \lambda^{n-1}.$$

Define the function

$$\begin{aligned}
g(\lambda, x) &= K_\lambda(x, x) - K_{\lambda, \text{low}}(x, x) \\
&= \frac{1}{2\pi i} \int_{-\infty}^{\infty} (1 - \rho(t)) \{e^{-2\pi i t(T - \lambda)}(x, x)\} dt.
\end{aligned}$$

Then the Fourier transform in the λ variable is

$$\widehat{g}(t, x) = \frac{1}{2\pi i} (1 - \rho(t)) e^{-2\pi i t T}.$$

Thus the support of \widehat{g} is supported away from $|t| \leq \varepsilon/2$. A Lemma of Tauberian type following this proof shows that it suffices to show that for $\lambda > 0$, and $0 < \tau \leq 1$,

$$|g(\lambda + \tau, x) - g(\lambda, x)| \lesssim (1 + \lambda)^{n-1}$$

But this follows if we can show that

$$|K_{\lambda+\tau}(x, x) - K_{\lambda}(x, x)| \lesssim (1 + \lambda)^{n-1}$$

and that

$$|K_{\lambda+\tau, \text{low}}(x, x) - K_{\lambda, \text{low}}(x, x)| \lesssim (1 + \lambda)^{n-1}.$$

These inequalities are equivalent to the (L^1, L^2) *discrete restriction theorem* for eigenfunctions of the Laplacian, which we will analyze in the next section. Indeed, if we let $\chi_{\lambda, \tau}$ be the operator with kernel $K_{\lambda+\tau} - K_{\lambda}$, then the discrete restriction theorem tells us that

$$\|\chi_{\lambda, \tau} f\|_{L^2(M)} \lesssim (1 + \lambda)^{(n-1)/2} \|f\|_{L^1(M)}$$

which by Schur's test, is equivalent to us having

$$\sup_{x \in M} \int_M |K_{\lambda+\tau}(x, y) - K_{\lambda}(x, y)|^2 dy \lesssim (1 + \lambda)^{n-1}$$

But the left hand side, by orthogonality, is equal to

$$\sup_{x \in M} \sum_j \mathbf{I}(\lambda \leq \lambda_j \leq \lambda + \tau) |e_j(x)|^2 = \sup_{x \in M} |K_{\lambda+\tau}(x, x) - K_{\lambda}(x, x)|,$$

which gives the required inequality. The analysis of $K_{\lambda, \text{low}}$ follows because we can write

$$K_{\lambda+\tau, \text{low}}(x, y) - K_{\lambda, \text{low}}(x, y) = \frac{1}{2\pi i} \int \frac{\rho(t)}{t + i0} (e^{-2\pi i \tau t} - 1) e^{2\pi i (T - \lambda)t} dt.$$

The $e^{-2\pi i \tau t} - 1$ term annihilates the singularity at the origin, and one can now argue as above to show this quantity is $O(\lambda^{n-1})$. \square

Lemma 13.30. *Let $g(\lambda)$ be a piecewise continuous function on \mathbf{R} , such that for $\lambda > 0$, and $0 < \tau \leq 1$,*

$$|g(\lambda + \tau) - g(\lambda)| \lesssim (1 + \lambda)^a.$$

If $\widehat{g}(t)$ vanishes for $|t| \leq 1$, then

$$|g(\lambda)| \lesssim (1 + \lambda)^a.$$

Proof. If

$$G(\lambda) = \int_{\lambda}^{\lambda+1} g(\tau) d\tau,$$

then G is absolutely continuous, and for almost all λ , G is differentiable with

$$|G'(\lambda)| = |g(\lambda + 1) - g(\lambda)| \lesssim (1 + \lambda)^a.$$

The Fourier transform of G also vanishes for $|t| \leq 1$. But

$$|g(\lambda)| \leq |G(\lambda)| + O((1 + \lambda)^a)$$

so it suffices to prove the estimates for G . If $\eta(t) = 1$ for $|t| > 1$, and $\eta(t) = 0$ for $|t| \leq 1/2$, and if ψ has Fourier transform $\eta(t)/2\pi it$, then ψ is bounded and rapidly decreasing at infinity. But $G' * \psi = G$, which gives

$$|G(\lambda)| \lesssim (1 + \lambda)^a \int |\psi(s)|(1 + |s|)^a ds \lesssim (1 + \lambda)^a. \quad \square$$

13.12 Discrete Restriction Theory

Let T be a classical, self-adjoint, elliptic pseudodifferential operator of order one on a compact manifold M , and consider an eigenfunction decomposition $\{e_j\}$ for T , corresponding to an increasing set of eigenvalues $\{\lambda_i\}$. We are concerned with the (L^p, L^q) bounds for the *spectral band projection operators*

$$\chi_{\lambda, \varepsilon} = \sum_{\lambda \leq \lambda_j \leq \lambda + \varepsilon} E_j.$$

These operators are closely related to restriction theory. The Stein-Tomas theorem studies the spherical projection operator, defined for functions on \mathbf{R}^n by setting

$$\chi f(x) = \int_{|\xi|=1} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

We can also consider the *spherical projection operators*

$$\chi_\lambda f(x) = \int_{|\xi|=\lambda} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi = \lambda^{-1} \text{Dil}_{1/\lambda} \chi \{ \text{Dil}_\lambda f \}.$$

The Stein-Tomas theorem characterizes the mapping properties of the operator χ , showing that the only inequality the operator has are of the form

$$\|\chi f\|_{L^{p^*}(\mathbf{R}^n)} \lesssim \|f\|_{L^p(\mathbf{R}^n)}$$

for $1 \leq p \leq 2(n+1)/(n+3)$. Rescaled, one concludes that we have

$$\|\chi_\lambda f\|_{L^{p^*}(\mathbf{R}^n)} \lesssim \lambda^{n(1/p-1/p^*)-1} \|f\|_{L^p(\mathbf{R}^n)}.$$

This inequality implies that if we define the spectral band projection operators

$$\chi_{\lambda,\varepsilon} f(x) = \int_{\lambda \leq |\xi| \leq \lambda+\varepsilon} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi = \int_\lambda^{\lambda+\varepsilon} S_\tau f(x) d\tau,$$

then,

$$\begin{aligned} \|\chi_{\lambda,\varepsilon} f\|_{L^{p^*}(\mathbf{R}^n)} &\lesssim \int_\lambda^{\lambda+\varepsilon} \tau^{n(1/p-1/p^*)-1} \|f\|_{L^p(\mathbf{R}^n)} d\tau \\ &\lesssim \varepsilon (1+\lambda)^{n(1/p-1/p^*)-1} \|f\|_{L^p(\mathbf{R}^n)}. \end{aligned}$$

It is also clear that a proof of this inequality would prove that χ_λ was bounded, since $\chi_\lambda = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} \chi_{\lambda,\varepsilon}$. Thus the theory of boundedness of spherical projections is closely related to the theory of boundedness for spectral band projections.

The original context for the spherical projection operators is to apply a TT^* argument, i.e. proving the (L^p, L^2) and (L^2, L^{p^*}) boundedness of the *restriction* and *extension* operators, which map functions on \mathbf{R}^n to functions on λS^{n-1} , and vice-versa, by setting

$$R_\lambda f(\xi) = \widehat{f}(\xi) \quad \text{and} \quad E_\lambda f(x) = \int_{|\xi|=\lambda} f(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

These operators are adjoints of one another, and $S_\lambda = E_\lambda \circ R_\lambda$. Thus a T^*T argument justifies that for $1 \leq p \leq 2(n+1)/(n+3)$,

$$\|R_\lambda f\|_{L^2(\lambda S^{n-1})} \lesssim \lambda^{n(1/p-1/2)-1/2} \|f\|_{L^p(\mathbf{R}^n)}$$

and

$$\|E_\lambda f\|_{L^{p^*}(\mathbf{R}^n)} \lesssim \lambda^{n(1/p-1/2)-1/2} \|f\|_{L^2(\lambda S^{n-1})}.$$

It is difficult to find direct analogues of the operators R_λ and E_λ on a manifold, just as we cannot directly find a direct analogue of χ_λ , because our eigenfunction is discrete. But we *can* consider analogues obtained by thickening the restriction set, i.e. considering the operators $R_{\lambda,\varepsilon}$ and $E_{\lambda,\varepsilon}$, mapping functions on \mathbf{R}^n to functions on the annulus $A_{\lambda,\varepsilon} = \{\xi \in \mathbf{R}^n : \lambda \leq |\xi| \leq \lambda + \varepsilon\}$ and vice versa by setting

$$R_{\lambda,\varepsilon} f(\xi) = \hat{f}(\xi) \quad \text{and} \quad E_{\lambda,\varepsilon} f(x) = \int_{\lambda \leq |\xi| \leq \lambda + \varepsilon} f(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

We have $\chi_{\lambda,\varepsilon} = E_{\lambda,\varepsilon} \circ R_{\lambda,\varepsilon}$, and the Stein-Tomas theorem is then verified to be equivalent to bounds of the form

$$\|R_{\lambda,\varepsilon} f\|_{L^2(A_{\lambda,\varepsilon})} \lesssim \varepsilon \lambda^{n(1/p-1/2)-1/2} \|f\|_{L^p(\mathbf{R}^n)}$$

and

$$\|E_{\lambda,\varepsilon} f\|_{L^{p^*}(\mathbf{R}^n)} \lesssim \varepsilon \lambda^{n(1/p-1/2)-1/2} \|f\|_{L^2(A_{\lambda,\varepsilon})}.$$

But the orthogonality of the Fourier transform, shows that the restriction bound is equivalent to the fact that

$$\|\chi_{\lambda,\varepsilon} f\|_{L^2(\mathbf{R}^n)} \lesssim \varepsilon \lambda^{n|1/p-1/2|-1/2} \|f\|_{L^p(\mathbf{R}^n)}$$

and

$$\|\chi_{\lambda,\varepsilon} f\|_{L^{p^*}(\mathbf{R}^n)} \lesssim \varepsilon \lambda^{n|1/p-1/2|-1/2} \|f\|_{L^2(\mathbf{R}^n)}.$$

This bound is the one we will begin by focus on generalizing to arbitrary manifolds, and we will start with case $p = 1$. Since there is no ambiguity, in the sequel, we let $\chi_\lambda = \chi_{\lambda,1}$.

We note that in \mathbf{R}^n , one can prove the same result for any positive homogeneous function $p(\xi)$ of degree one such that the *cosphere* $\Sigma = \{\xi : p(\xi) = 1\}$ has non-vanishing curvature, Stein-Tomas gives results about

$$\chi_\lambda f(x) = \int_{\Sigma} \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

Returning to the analysis of the operator T with principal symbol $p(x, \xi)$. We will obtain good results about the operator χ_λ provided that the co-spheres $\Sigma(x) = \{\xi : p(x, \xi) = 1\}$ has non-vanishing curvature for all $x \in M$.

The easiest bound in the Stein-Tomas theorem are the equivalent bounds

$$\|\chi_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{(n-1)/2} \|f\|_{L^1(\mathbf{R}^n)}$$

and

$$\|\chi_\lambda f\|_{L^\infty(\mathbf{R}^n)} \lesssim \lambda^{(n-1)/2} \|f\|_{L^2(\mathbf{R}^n)},$$

which does not even utilize the curvature of the domain of projection whatsoever, solely using the fact that the sphere of radius λ has surface measure $O(\lambda^{n-1})$. Let us see if we can justify the analogous result on a compact manifold. We will find we do not need to assume curvature here either.

Lemma 13.31. *Let T be a classical, elliptic, self-adjoint pseudodifferential operator of order one on a compact manifold M of dimension n , and consider the association spectral band projection operators $\{\chi_\lambda\}$. Then*

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim \lambda^{(n-1)/2} \|f\|_{L^1(M)},$$

and thus by self-adjointness, that

$$\|\chi_\lambda f\|_{L^\infty(M)} \lesssim \lambda^{(n-1)/2} \|f\|_{L^2(M)}.$$

Proof. We want to exploit the parametrix for the wave equation. Thus we start by replacing χ_λ with an operator $\tilde{\chi}_\lambda$ given by

$$\tilde{\chi}_\lambda f(x) = \sum_j \chi(\lambda_j - \lambda) E_j,$$

where χ is a non-negative Schwartz function with $\chi(0) > 0$, and such that $\hat{\chi}(t)$ has support on $|t| \leq \varepsilon/2$. Because $\mathbf{I}(\lambda \leq \lambda_j \leq \lambda + \varepsilon) \lesssim \chi(\lambda_j - \lambda)$, orthogonality shows that

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim \|\tilde{\chi}_\lambda f\|_{L^2(M)}$$

and so it suffices to prove the result for $\tilde{\chi}_\lambda$. But if $\tilde{\chi}_\lambda$ has kernel

$$\tilde{K}_\lambda(x, y) = \sum_j \chi(\lambda_j - \lambda) e_j(x) \overline{e_j(y)}.$$

This is equivalent to showing that

$$\|\tilde{K}_\lambda\|_{L_y^\infty L_x^2} \lesssim \lambda^{(n-1)/2}.$$

But orthogonality shows that

$$\begin{aligned}\|\tilde{K}_\lambda\|_{L_y^\infty L_x^2} &= \left(\sup_{y \in M} \sum_j \chi(\lambda_j - \lambda)^2 |e_j(y)|^2 \right)^{1/2} \\ &\leq \|\chi\|_{L^\infty}^{1/2} \left(\sup_{y \in M} \sum_j \chi(\lambda_j - \lambda) |e_j(y)|^2 \right)^{1/2} \\ &\leq \|\chi\|_{L^\infty}^{1/2} \left(\sup_{y \in M} \tilde{\chi}_\lambda(y, y) \right)^{1/2}.\end{aligned}$$

Thus we need to show that for all $y \in M$,

$$\tilde{\chi}_\lambda(y, y) \lesssim \lambda^{n-1}.$$

This follows by an argument similar to the proof of the Weyl law, e.g. applying the parametrix for the wave equation. \square

Now we move onto results that require a curvature assumption.

Theorem 13.32. *Let T be a classical, elliptic, self-adjoint pseudodifferential operator of order one on a compact manifold M of dimension n and with principal symbol $p(x, \xi)$, such that the cospheres*

$$\Sigma(x) = \{\xi \in T^*M : p(x, \xi) = 1\}$$

have non-vanishing curvature for each $x \in M$. If we consider the association spectral band projection operators $\{\chi_\lambda\}$, then for $p_c = 2(n+1)/(n+3)$, we have

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim (1 + \lambda)^{(n-1)/2(n+1)} \|f\|_{L^{p_c}(M)}.$$

Remark. Combined with the (L^1, L^2) boundedness of the equation, and the fact that

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim \|f\|_{L^2(M)}$$

we conclude using interpolation that for $1 \leq p \leq 2(n+1)/(n+3)$,

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim (1 + \lambda)^{n|1/p - 1/2| - 1/2} \|f\|_{L^p(M)}$$

and for $2(n+1)/(n+3) \leq p \leq 2$,

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim (1 + \lambda)^{(\frac{n-1}{2})(1/p - 1/2)} \|f\|_{L^p(M)}.$$

All these inequalities are sharp, as we will discuss after the proof.

Proof. We start by using a strategy we began the (L^1, L^2) boundedness result with, namely, to swap χ_λ out with a more well behaved operator. The operators $\tilde{\chi}_\lambda$ used in that proof become singular near the diagonal as $\lambda \rightarrow \infty$, i.e. the kernel is $\Theta(\lambda^n)$. Thus we modify their definition slightly. We fix a small quantity $\varepsilon_0 \leq \varepsilon/2$, and define a Schwartz function χ (not necessarily positive this time), with $\chi(0) = 1$ and with Fourier support on $\varepsilon_0/2 \leq t \leq \varepsilon_0$. Nonetheless, it remains true that $\chi(x) \gtrsim 1$ for $|x| \leq \varepsilon$, so that

$$\|\chi_\lambda f\|_{L^2(M)} \lesssim \|\tilde{\chi}_\lambda f\|_{L^2(M)}.$$

It thus suffices to obtain a bound for $\tilde{\chi}_\lambda$ at the critical exponent. Applying the parametrix, and seeing that the remainder term has operator norm $O_N((1 + \lambda)^{-N})$ for all $N > 0$, it suffices to show that the operator $\tilde{\chi}_\lambda$ with kernel

$$\iint \hat{\chi}(t) s(t, x, y, \xi) e^{2\pi i [\phi(x, y, \xi) + t(p(y, \xi) - \lambda)]} dt d\xi$$

is well behaved. The kernel of this operator is an oscillatory integral, and on the support of the amplitude s , and if we set $\Phi(t, x, y, \xi) = \phi(x, y, \xi) + tp(y, \xi)$ has

$$|\nabla_\xi \Phi(t, x, y, \xi)| \gtrsim 1$$

on the support of the amplitude $\hat{\chi}s$ for $\xi \neq 0$ except if $|x - y| \sim \varepsilon_0$. Thus, applying stationary phase, if $\eta \in C^\infty(M \times M)$ is a function which equals one for $|x - y| \sim \varepsilon_0$, and vanishes everywhere else, the difference between $\tilde{\chi}_\lambda$ and the operator χ'_λ with kernel

$$\iint \hat{\chi}(t) s(t, x, y, \xi) \eta(x, y) e^{2\pi i [\Phi(t, x, y, \xi) - \lambda t]} d\xi dt$$

is an operator of the form

$$\int \hat{\chi}(t) A(t, x, y) e^{-2\pi i \lambda t},$$

where A is smooth, and it therefore follows this kernel is $O_N((1 + \lambda)^{-N})$. Let us set $a(t, x, y, \xi) = \hat{\chi}(t) s(t, x, y, \xi) \eta(x, y)$. Our proof is now completed by applying some results about non-homogeneous oscillatory integral operators, once we show that

$$\iint a(t, x, y, \xi) e^{2\pi i [\Phi(t, x, y, \xi) - \lambda t]} d\xi dt = \lambda^{(n-1)/2} a_\lambda(x, y) e^{2\pi i \lambda \psi(x, y)},$$

where ψ is non-homogeneous, and satisfies the $n \times n$ Carleson-Sjölin conditions on the support of a_λ , which is smooth, and satisfies $|\partial_x^\alpha \partial_y^\beta a_\lambda| \lesssim_{\alpha,\beta} 1$, uniformly in λ . \square

Chapter 14

Fourier Integral Operators

Pseudodifferential operators formalize the family of all operators that modulate the amplitude of wave packets. The theory of Fourier integral operators extends this theory by not only modulating the amplitude of wave packets, but also moving them around in phase space in a *symplectic manner*, i.e. in a way which, roughly speaking, obeys the uncertainty principle. Basic examples of Fourier integral operators include the translation operators

$$\text{Trans}_{x_0} f(x) = f(x - x_0),$$

modulation operators

$$\text{Mod}_{\xi_0} f(x) = e^{2\pi i \xi_0 \cdot x} f(x),$$

the change of variables operator T_A associated with an invertible linear transformation $A : \mathbf{R}^n \rightarrow \mathbf{R}^n$, i.e. such that

$$T_A f(x) = |\det(A)|^{-1/2} f(A^{-1}x),$$

and the Fourier transform

$$\mathcal{F} f(\xi) = \hat{f}(\xi).$$

These four operators are all unitary, so might be thought of as preserving the amplitude of wave packets, but they move wave packets around in phase space in different ways:

- The translation operators move wave packets in phase space according to the diffeomorphism $\Phi(x, \xi) = (x + x_0, \xi)$.

- The modulation operators move packets in phase space according to the diffeomorphism $\Phi(x, \xi) = (x, \xi + \xi_0)$.
- The change of variables move packets in space according to the diffeomorphism $\Phi(x, \xi) = (Ax, (A^T)^{-1} \xi)$.
- The Fourier transform move packets according to the diffeomorphism $\Phi(x, \xi) = (\xi, -x)$.

All four of the diffeomorphisms $\Phi : T^* \mathbf{R}^d \rightarrow T^* \mathbf{R}^d$ are *symplectomorphisms*, i.e. they preserve the symplectic form $\omega = \sum dx^i \wedge d\xi_i$, i.e. the bilinear form

$$\sigma((x_1, \xi_1), (x_2, \xi_2)) = \xi_2(x_1) - \xi_1(x_2),$$

which is the derivative of the *tautological one form* $\theta = \sum \xi_i dx^i$, a fact very important to the tractability of the study of the associated operators, because of the uncertainty principle.

Let us see why this is essential. With any diffeomorphism $\Phi : T^* \mathbf{R}^n \rightarrow T^* \mathbf{R}^n$, we can try and construct a unitary operator T_Φ from $L^2(\mathbf{R}^n)$ to itself, which roughly speaking, has the property that it maps a wave packet localized at a point (x_0, ξ_0) to a wave packet localized near $\Phi(x_0, \xi_0)$. If we consider an pseudodifferential operator $S_a = a(x, D)$ with symbol a , then S amplifies wave packets localized at (x_0, ξ_0) by a quantity $a(x_0, \xi_0)$. Thus, morally speaking, we should expect the operator $T_\Phi^{-1} \circ S_a \circ T_\Phi$ to fix the location of wave packets, and amplify wave packets localized at (x_0, ξ_0) by the quantity $a \circ \Phi$, i.e. so that

$$T_\Phi^{-1} \circ S_a \circ T_\Phi \approx (a \circ \Phi)(x, D).$$

We should at least expect this to hold up to first order. If we have another operator S_b , then we should expect that, using the formula above, up to first order we should have

$$\begin{aligned} [(a \circ \Phi)(x, D), (b \circ \Phi)(x, D)] &\approx [T_\Phi^{-1} \circ S_a \circ T_\Phi, T_\Phi^{-1} \circ S_b \circ T_\Phi] \\ &\approx T_\Phi^{-1} \circ [S_a, S_b] \circ T_\Phi \\ &\approx ((a \circ b) \circ \Phi)(x, D). \end{aligned}$$

Taking principles of either side of the equation yields to the exact equation

$$\{a \circ \Phi, b \circ \Phi\} = \{a, b\} \circ \Phi.$$

Since $\{f, g\} = \omega(\nabla f, \nabla g)$, we conclude that this is only true if $\Phi^* \omega = \omega$, i.e. Φ is a symplectomorphism. In this case, the graph of Φ , namely the set

$$\Lambda_\Phi = \{(x, y; \xi, \eta) : (x, \xi) = \Phi(y, \eta)\} \subset T^*(\mathbf{R}^n \times \mathbf{R}^n)$$

will be the *canonical relation* of Φ , i.e. a Lagrangian submanifold of $T^*\mathbf{R}_X^n \times T^*\mathbf{R}_Y^n$ with respect to the symplectic form

$$\omega_X - \omega_Y = \sum dx^i \wedge d\xi_i - dy^i \wedge d\eta_i,$$

since $T_{(x, y; \xi, \eta)} \Lambda_\Phi$ can be identified with pairs of tangent vectors

$$v \in T_{(x, \xi)}(T^*\mathbf{R}_X^n) \quad \text{and} \quad w \in T_{(y, \eta)}(T^*\mathbf{R}_Y^n)$$

such that $v = \Phi_*(w)$, and $\omega_X(\Phi_*(w_1), \Phi_*(w_2)) = \omega_Y(w_1, w_2)$.

The fact that we are reducing ourselves to the study of symplectomorphisms Φ gives us a hint as to how to define the resulting operator T_Φ , at least microlocally. Intuitively speaking, our discussion above shows that the wave front set of the kernel of Φ must be contained in Λ_Φ , because spectral singularities at a point (y, η) will be moved to singularities at a point (x, ξ) , so we should expect that $\text{WF}(T_\Phi f) = \Lambda_\Phi \circ \text{WF}(f)$. We know a family of operators of this property: in a previous section we discussed the oscillatory integral distributions

$$\int a(x, \theta) e^{2\pi i \phi(x, \theta)} d\theta.$$

The wave-front set contained in $\Lambda_\phi = \{(x, \nabla_x \phi(x, \theta)) : \nabla_\theta \phi(x, \theta) = 0\}$, and, provided ϕ is non-degenerate, Λ_ϕ is a Lagrangian manifold. Thus, given a symplectomorphism $\Phi : T^*\mathbf{R}^n \rightarrow T^*\mathbf{R}^n$, if we can find a phase $\phi : \mathbf{R}^n \times \mathbf{R}^n \rightarrow \mathbf{R}$ such that $\Lambda_\Phi = \Lambda_\phi$, then, modulo C^∞ kernels, we might expect to find a symbol a such that

$$T_\Phi f(x) = \int a(x, y, \theta) e^{2\pi i \phi(x, \theta)} f(y) d\theta dy.$$

More generally, we might not be able to find a phase ϕ that works globally for Φ , but we can localize, and such ϕ will *always* exist locally by virtue of the symplectic structure of Λ_Φ . Thus we might wish to consider the families of operators formed from finite sums of integral operators given by the right hand side of this equation. We have now reached the study of general Fourier integral operators.

14.1 Hyperbolic Equations

Fourier integral operators were initially introduced to obtain parametrices for hyperbolic equations. To see how these arise, let us begin with a constant coefficient linear differential operator on $\mathbf{R}_t \times \mathbf{R}_x^n$, given by $P(\partial_t, D_x)$, where $P(\tau, \xi)$ is a polynomial, which we will assume can be written in the form $\tau^m + \tau^{m-1}Q_1(\xi) + \cdots + Q_m(\xi)$, where $Q_i(\xi)$ is a polynomial of degree at most i . Then the hyperbolic equation is

$$L = \partial_t^m + \partial_t^{m-1}Q_1(D_x) + \cdots + \partial_t Q_{m-1}(D_x) + Q_m(D_x).$$

We note that here ∂_t is the standard derivative operator, whereas D_x^α is the derivative operator, normalized by dividing by an appropriate power of $2\pi i$ so that it is the Fourier multiplier of ξ^α . We recall the Cauchy-Kovalevskaya theorem, which gives unique analytic solutions to the Cauchy problem $Lu = f$ given initial conditions $u_0, \partial_t u_0, \dots, \partial_t^{m-1} u_0$, given that f , and the initial conditions are analytic functions on \mathbf{R}^n . But we are interested in more general existence results.

The surprising feature of this problem is that these operators need not even have solutions if we switch from studying analytical initial conditions to say, compactly supported smooth initial conditions. For instance, suppose there exists a distribution u on $\mathbf{R}_t \times \mathbf{R}_x$, tempered in the x -variable, such that $Lu = 0$, where $L = \partial_t - D_x$, and we let $u_0 \in \mathcal{S}(\mathbf{R}_x)^*$ be the initial value of the distribution. Then, taking Fourier transforms in the x variable, we conclude that $\partial_t \hat{u}(t, \xi) = \xi \hat{u}(t, \xi)$, which implies that $\hat{u}(t, \xi) = \hat{u}_0(\xi)e^{\xi t}$. But this distribution is *never* tempered in the ξ variable; if it was tempered for one positive value of t , and one negative value of t , then we could conclude that

$$|\hat{u}_0(\xi)| \lesssim e^{-\varepsilon|\xi|}$$

for some $\varepsilon > 0$. But the Paley-Wiener theorem and its variants therefore imply that u_0 is analytic, and actually extends to a holomorphic function on a small strip containing the real line. Thus the existence of solutions to the Cauchy problem $\partial_t u - Du = 0$ is very delicate; in particular, there are no solutions with initial conditions in $\mathcal{D}(\mathbf{R}_x^d)$. This hints at the fact that to make the solution to the Cauchy problem tractable, we must ensure that the polynomial $P(\tau, \xi) = 0$ have *imaginary roots*. A desire to find a more powerful existence statement for solutions to such equations, tempered in

the x -variable, will force us to choose polynomials $P(\tau, \xi)$ whose principal part has purely imaginary roots in the τ variable.

Let us discuss this situation more precisely. Let $\tilde{E}_0, \dots, \tilde{E}_{m-1} : \mathbf{R}_\xi^n \times \mathbf{R}_t \rightarrow \mathbf{C}$ be the analytic solutions to the Cauchy problem $P(\partial_t, \xi) = 0$ with initial conditions

$$\partial_t^i \tilde{E}_j(0, \xi) = \delta_{ij}$$

for $0 \leq i \leq m-1$. If we are to expect $P(\partial_t, D_x) = 0$ to be a well posed differential equation, then we should expect each solution \tilde{E}_i to be tempered in the ξ variable. We can calculate the functions $\{\tilde{E}_i\}$ explicitly. If we fix ξ_0 , and assume first that the roots of $P(\tau, \xi_0)$ in the τ variable are distinct, then the roots are distinct locally around ξ_0 . If we let $\tau_1(\xi), \dots, \tau_m(\xi)$ be the roots of the equation, then these are analytic functions in the ξ variable locally around ξ_0 . The functions $h_i(t, \xi) = e^{i\tau_i(\xi)t}$ then satisfy the Cauchy problem $P(\partial_t, \xi) = 0$ with initial conditions

$$\partial_t^i h_j(0, \xi) = \tau_j(\xi)^i$$

for $0 \leq i \leq m-1$. The uniqueness of analytic solutions guaranteed by the Cauchy-Kovalevsky theorem imply that

$$h_i(t, \xi) = \tilde{E}_0(t, \xi) + \tilde{E}_1(t, \xi)\tau_i(\xi) + \dots + \tilde{E}_{m-1}(t, \xi)\tau_i^{m-1}(\xi).$$

If $\tilde{E} = (\tilde{E}_0, \dots, \tilde{E}_{m-1})$ and $h = (h_0, \dots, h_{m-1})$, then we can summarize this in the matrix equation

$$h = \begin{pmatrix} 1 & \tau_1 & \dots & \tau_1^{m-1} \\ 1 & \tau_2 & \dots & \tau_2^{m-1} \\ \vdots & \ddots & \dots & \vdots \\ 1 & \tau_m & \dots & \tau_m^{m-1} \end{pmatrix} \tilde{E}.$$

Provided that the roots $\{\tau_i\}$ are distinct, we can solve this equation to find the functions $\{\tilde{E}_i\}$ in terms of the functions $\{h_i\}$ using Cramer's rule. Namely, if $V(\tau_1, \dots, \tau_m)$ is the Vandermonde determinant, i.e. the determinant of the matrix $\{\tau_i^j\}$, and if $V_i(\tau_1, \dots, \tau_m; t)$ is the determinant of the matrix obtained by replacing the i th column with the vector $\{e^{t\tau_j(\xi)}\}$, then

$$\tilde{E}_i(t, \xi) = \frac{V_i(\tau_1(\xi), \dots, \tau_m(\xi); t)}{V(\tau_1(\xi), \dots, \tau_m(\xi))}.$$

For instance, if $m = 2$, then

$$\tilde{E}_1(t, \xi) = \frac{e^{t\tau_1(\xi)}\tau_2(\xi) - e^{t\tau_2(\xi)}\tau_1(\xi)}{\tau_2(\xi) - \tau_1(\xi)} \quad \text{and} \quad \tilde{E}_2(t, \xi) = \frac{e^{t\tau_2(\xi)} - e^{t\tau_1(\xi)}}{\tau_2(\xi) - \tau_1(\xi)}$$

Note that the function

$$(\tau_1, \dots, \tau_m, t) \mapsto \frac{V_i(\tau_1, \dots, \tau_m; t)}{V(\tau_1, \dots, \tau_m)}$$

are analytic in t , symmetric in the variables $\{\tau_i\}$, and *entire* in the variables $(\tau_1, \dots, \tau_m) \in \mathbf{C}^n$. For instance, in the case $m = 2$, when $\tau_1(\xi_0) = \tau_2(\xi_0) = \tau$ we have

$$\tilde{E}_1(t, \xi_0) = e^{t\tau}(1 - t\tau) \quad \text{and} \quad \tilde{E}_2(t, \xi_0) = te^{t\tau}.$$

Thus the equation we have constructed continues to specify the functions $\{\tilde{E}_i\}$ even when the roots of P in the ξ variable are not distinct.

By virtue of the roots switching around, one cannot necessarily define the functions h_1, \dots, h_n globally for all $\xi \in \mathbf{R}^n$ if roots coincide. But quantities symmetric in the variables $\{h_i\}$ are well defined, for instance, the function

$$s(t, \xi) = |\operatorname{Re}(h_1(t, \xi))| + \dots + |\operatorname{Re}(h_n(t, \xi))|.$$

is globally defined. If $\tilde{E}_0, \dots, \tilde{E}_{m-1}$ are tempered in the ξ variable, then it follows that the function s is tempered in the ξ variable, and thus satisfies some equation of the form

$$|s(t, \xi)| \lesssim_t \langle \xi \rangle^{N_t}$$

for some $N_t > 0$. But this means that

$$|\operatorname{Re}(\tau_1(\xi))| + \dots + |\operatorname{Re}(\tau_m(\xi))| \lesssim 1 + \log \langle \xi \rangle.$$

But the theory of semialgebraic sets (which applies because (τ_1, \dots, τ_m) are the projections onto the τ variable of solutions to the polynomial equation $P(\tau, \xi) = 0$) implies that this can only be possible if

$$|\operatorname{Re}(\tau_1(\xi))| + \dots + |\operatorname{Re}(\tau_m(\xi))| \lesssim 1,$$

i.e. because we can only have polynomial growth on semialgebraic sets. This is actually a necessary and sufficient condition for the Cauchy problem to be solvable (a result of Garding). An operator with this property will be called *hyperbolic*.

There is an equivalent specification of being hyperbolic which is very useful to the study of such equations. If $L = P(\partial_t, D_x)$ is a hyperbolic constant-coefficient partial differential equation of order m , then we can consider the degree m polynomial $P_m(\tau, \xi)$. We claim the roots of P_m in the τ variable differ from the roots of P by at most $O(1)$ for $|\xi| \gg 1$. Note that this is the only part of the discussion where we have used the fact that the polynomials Q_i have degree at most i . This means that the condition

$$|\operatorname{Re}(\tau_1(\xi))| + \cdots + |\operatorname{Re}(\tau_m(\xi))| \lesssim 1$$

holds if and only if the roots of P_m are all purely imaginary. Thus $P(\partial_t, D_x)$ is hyperbolic if and only if the roots of P_m in the τ variable are all purely imaginary.

In addition, we say a differential operator $P(\partial_t, D_x)$ is *strongly*, or *strictly hyperbolic* if in addition to being hyperbolic, the imaginary roots of P_m are all distinct from one another for all $\xi \in \mathbf{R}^d$, which implies the roots of P are distinct for large ξ . If we label the roots of P_m as $i\lambda_1(\xi), \dots, i\lambda_m(\xi)$, then the functions $\{\lambda_i\}$ are real-valued analytic functions in $\mathbf{R}^n - \{0\}$, which are homogeneous of degree one. Thus $|\lambda_i(\xi) - \lambda_j(\xi)| \gtrsim |\xi|$ for $i \neq j$.

14.2 Local Symplectic Geometry

Before we get into the general theory of Fourier integral operators, let's recall some results in the theory of symplectic geometry. We recall that a *symplectic vector space* V is a finite dimensional vector space equipped with a non-degenerate, skew-symmetric bilinear form ω . A fundamental example is that for any vector space W , $V = W^* \oplus W$ is naturally a symplectic vector space with the symplectic form $\omega((x_1^*, x_1), (x_2^*, x_2)) = x_2^*(x_1) - x_1^*(x_2)$. Spectral theory can be used to show that for any symplectic vector space V , V is even dimensional, and that we can find a pair of independent sets $\{e_i\}$ and $\{f^i\}$, forming a basis of V , such that $\omega(e_i, e_j) = \omega(f^i, f^j) = 0$, and $\omega(e_i, f^i) = 1$. Such a basis is called a *Darboux basis*.

A *Lagrangian subspace* of a symplectic vector space V is a subspace W of V such that $W^\perp = W$. This is equivalent to $\dim(W) = \dim(V)/2$, and for any $w_1, w_2 \in W$, $\omega(w_1, w_2) = 0$. As an example, if we write $V = V_e \oplus V_f$, where V_e and V_f are the spans of the separate parts of a Darboux basis, then V_e and V_f are Lagrangian subspaces. If $A : W \rightarrow W^*$ is a linear map, then the graph $\Gamma_A = \{(x, Ax) : x \in W\}$ is a Lagrangian submanifold of $W \oplus$

W^* if and only if A is self-adjoint, where we identify W^{**} with W , i.e. if $\langle Ax_1, x_2 \rangle = \langle x_1, Ax_2 \rangle$ for all $x_1, x_2 \in W$.

Lemma 14.1. *Let V be a symplectic vector space, and let*

$$\{e_1, \dots, e_{n_1}\} \cup \{f^1, \dots, f^{n_2}\}$$

be linearly independent vectors such $\omega(e_i, e_j) = \omega(f^i, f^j) = 0$, and $\omega(e_i, f^j) = \delta(i, j)$. Then we can extend these sets to a full Darboux basis for V .

Proof. Suppose first that $n_1 < n_2$. Since ω is non-degenerate, we can find a vector e such that $\omega(e, e_i) = 0$ for $1 \leq i \leq n_1$, and $\omega(e, f^i) = \delta(i, n_1 + 1)$. It follows that e is linearly independent from the previously selected elements of the basis and we can set $e_{n_1+1} = e$. Thus we can increase n_1 by one. A similar argument works for choosing f_{n_2+1} if $n_2 < n_1$. If $n_1 = n_2$, and we don't yet have a basis, then we choose $e = e_{n_1+1}$ linearly independent to the previous set such that $\omega(e, e_i) = \omega(e, f^i) = 0$. Iterating this selection procedure yields the required Darboux basis. \square

Lemma 14.2. *If V_0 and V_1 are Lagrangian subspaces of a symplectic vector space V , then we can find a third Lagrangian subspace V_2 which is transverse to both V_0 and V_1 .*

Proof. Let V have dimension $2n$. It suffices to find a Darboux basis in which

$$V_0 = \text{span}(e_1, \dots, e_n) \quad \text{and} \quad V_1 = \text{span}(e_1, \dots, e_k, f^{k+1}, \dots, f^n),$$

since we can then set V_2 to be the space spanned by $\{e_i + f^i : k+1 \leq i \leq n\} \cup \{f^1, \dots, f^k\}$. To do this, we pick $\{e_1, \dots, e_n\}$ such that $\{e_1, \dots, e_k\}$ spans $V_0 \cap V_1$, and $\{e_1, \dots, e_n\}$ spans V_1 . Next, we note that the map $T : V_0 \rightarrow \mathbf{R}^{n-k}$ given by

$$Tv = (\omega(e_{k+1}, v), \dots, \omega(e_n, v))$$

has kernel equal to $V_0 \cap V_1$. This is because $V_0^\perp = V_0$, so if $v \in V_1$, then we automatically obtain that $\omega(e_1, v) = \dots = \omega(e_k, v) = 0$, so that if $Tv = 0$, then $v \in V_0^\perp = V_0$. But this means that T is surjective, so we can find $\{f^{k+1}, \dots, f^n\} \subset V_1$ such that $\omega(e_i, f_j) = \delta(i, j)$. It follows that $\{e_1, \dots, e_k\} \cup \{f^{k+1}, \dots, f^n\}$ are linearly independent, and thus form a basis for V_1 . And we can now use the previous Lemma to extend these vectors to a Darboux basis. \square

If V_X and V_Y are symplectic vector spaces, then $V_X \oplus V_Y$ can be made into a symplectic vector space, if we equip it either with the symplectic form $\omega_X - \omega_Y$, or with $\omega_Y - \omega_X$. In either case, we call a Lagrangian submanifold C of $V_X \oplus V_Y$ a *linear canonical relation*. We recall that a *symplectic linear map* $A : V_1 \rightarrow V_2$ being symplectic vector spaces is a map preserving the symplectic form.

Lemma 14.3. *Let $C \subset V_X \oplus V_Y$ be a linear canonical relation. Then we can find orthogonal decompositions $V_X = V_{X,1} \oplus V_{X,2}$ and $V_Y = V_{Y,1} \oplus V_{Y,2}$ such that*

$$C = C_X \oplus \Gamma \oplus C_Y,$$

where C_X is a Lagrangian submanifold of $V_{X,1}$, C_Y is a Lagrangian submanifold of $V_{Y,1}$, and Γ is the graph of a symplectic isomorphism $A : V_{X,2} \rightarrow V_{Y,2}$.

Proof. Let

$$C_X = \{x \in V_X : (x, 0) \in C\} \quad \text{and} \quad C_Y = \{y \in V_Y : (0, y) \in C\}.$$

Because C is Lagrangian, C is contained in $C_X^{\omega_X} \oplus C_Y^{\omega_Y}$, i.e. if $(x_1, 0), (0, y_1)$, and (x_2, y_2) are in C , then $\omega(x_1, x_2) = \omega(y_1, y_2) = 0$. In particular, $C_X \subset C_X^{\omega_X}$ and $C_Y \subset C_Y^{\omega_Y}$. Find orthogonal $V_{X,2}$ and $V_{Y,2}$ such that

$$C_X^{\omega_X} = C_X \oplus V_{X,2} \quad \text{and} \quad C_Y^{\omega_Y} = C_Y \oplus V_{Y,2}.$$

Then $C \subset C_X \oplus C_Y \oplus V_{X,2} \oplus V_{Y,2}$. We can thus find $\Gamma \subset V_{X,2} \oplus V_{Y,2}$ such that $C = C_X \oplus C_Y \oplus \Gamma$. We claim Γ projects bijectively onto $V_{X,2}$ and $V_{Y,2}$. For instance, suppose $x \in V_{X,2}$ and $y_1, y_2 \in V_{Y,2}$ are such that (x, y_1) and (x, y_2) lie in Γ . Then $(0, y_1 - y_2)$ lies on Γ and on C_Y , so by orthogonality, $y_1 = y_2$. Thus Γ is the graph of an isomorphism $A : V_{X,2} \rightarrow V_{Y,2}$. It is easy to see that C_X and C_Y are symplectic manifolds. And Γ is also Lagrangian since C is Lagrangian, which implies A is symplectic, completing the proof. \square

A *symplectic manifold* M is a manifold equipped with a symplectic two form ω , i.e. a two form which gives each of the tangent spaces of M a symplectic structure. The basic example here is $M = T^*X$, where X is any smooth manifold - the natural two form here is the two form $\omega = dx \wedge d\xi = d\theta$, where $\theta = \sum \xi_i dx^i$ is the *tautological one form* on M .

Like with a Riemannian manifold, the symplectic structure gives us a natural bundle isomorphism $J : T^*M \rightarrow TM$. In particular, given a function $f : M \rightarrow \mathbf{R}$, we can define the *symplectic gradient* $\nabla_{\Xi} f$ as the vector

field which is identified under the bundle isomorphism with the covector field df . As an example, if $M = T^*X$, then in local coordinates (x, ξ) on T^*X , the identification is given by

$$J(d\xi^i) = \frac{\partial}{\partial x^i} \quad \text{and} \quad J(dx^i) = -\frac{\partial}{\partial \xi^i}.$$

Thus for a function $f : T^*X \rightarrow \mathbf{R}$, we have

$$\nabla_{\Xi} f = \sum \frac{\partial f}{\partial \xi^i} \frac{\partial}{\partial x^i} - \frac{\partial f}{\partial x^i} \frac{\partial}{\partial \xi^i}.$$

This gradient is closely related to the theory of Hamiltonian vector fields, i.e. since, given a Hamiltonian H on some phase space representing a physical system, $\nabla_{\Xi} H$ gives the flow of that physical system. We can define the *Poisson bracket* of two functions $f, g : M \rightarrow \mathbf{R}$ by setting $\{f, g\} = \nabla_{\Xi} f(g) = dg(\nabla_{\Xi} f)$. For $M = T^*X$, we can write this in coordinates as

$$\{f, g\} = \sum \frac{\partial f}{\partial \xi_i} \frac{\partial g}{\partial x_i} - \frac{\partial f}{\partial x_i} \frac{\partial g}{\partial \xi_i},$$

which agrees with the classical Poisson bracket.

A (immersed) *Lagrangian submanifold* of a symplectic manifold M is a submanifold N such that its tangent space is a Lagrangian subspace of the tangent space of M at each point. To check that an immersion $i : X \rightarrow M$ gives a Lagrangian submanifold, it suffices to show that $i^*\omega = 0$, and that $\dim(X) = \dim(M)/2$. We have already seen these kinds of subspaces in our discussion of oscillatory integral distributions, since the wavefront sets of these distributions form Lagrangian submanifolds of the cotangent space of the space the distributions live on. Another example includes particular kinds of sections $X \rightarrow T^*X$.

Lemma 14.4. *The image of a section $s : X \rightarrow T^*X$ is an immersed Lagrangian submanifold if and only if locally we can write $s = df$ for some function $f : X \rightarrow \mathbf{R}$.*

Proof. If $\theta = \sum \xi^i dx^i$, then $d\theta$ is the symplectic form ω , and $s^*\theta = s$. Thus s gives a Lagrangian manifold if and only if $s^*\omega = s^*(d\theta) = d(s^*\theta) = ds = 0$. The result above now follows by Poincaré's Lemma. \square

A similar result holds if Λ is a conic Lagrangian submanifold of T^*X .

Lemma 14.5. *Suppose that $\Lambda \subset T^*X - \{0_X\}$ is a conic Lagrangian manifold containing some covector $v \in T_{x_0}^*X$. Then local coordinates (x, U) can be chosen on X , centered at x_0 , inducing coordinates (x, ξ) on T^*U , such that the map $(x, \xi) \mapsto \xi$ is a diffeomorphism with domain $\Lambda \cap T^*U$, and there exists a smooth, homogeneous function H such that*

$$\Lambda \cap T^*U = \{(\nabla H(\xi), \xi) : \xi \in \mathbf{R}^d\}.$$

Proof. We first choose coordinates (x, U) such that $(x, \xi) \mapsto \xi$ is a diffeomorphism. Begin by choosing coordinates (y, V) centered at x_0 such that $v = dy_1$. The tangent plane V_0 to Λ at v must be Lagrangian. If V_0 is transverse to the Lagrangian plane given in (y, η) coordinates by

$$V_1 = \{(0, a) : a \in \mathbf{R}^d\},$$

then we can set x to be y . Otherwise, we find V_2 Lagrangian and transverse to both V_0 and V_1 . Since V_2 is transverse to V_1 , it can be identified with a linear section, and the last result thus implies that we can find a quadratic form $Q : \mathbf{R}^d \rightarrow \mathbf{R}^d$ such that

$$V_2 = \{(a, dQ(a)) : a \in \mathbf{R}^d\}.$$

If we set $x_1 = y_1 + Q(y)$, and $x_i = y_i$ for $2 \leq i \leq n$, then these are the required coordinates.

We claim now that if (x, U) gives a diffeomorphism, then we can find H . Shrinking U if necessary, there exists a radial $\phi : \mathbf{R}^d \rightarrow U$ such that

$$\Lambda \cap U = \{(\phi(\xi), \xi) : \xi \in \mathbf{R}^d\}.$$

Since Λ is Lagrangian, if $\psi(\xi) = (\phi(\xi), \xi)$, then

$$\psi^*\theta = \sum \xi_i d\phi_i = 0.$$

If we set $H(\xi) = \sum \xi_i \phi_i(\xi)$, then $\nabla H = \phi$, giving the required result. \square

The structure of linear canonical relations can give us results about ‘nonlinear’ canonical relations, i.e. a conic Lagrangian submanifold of $T^*X \times T^*Y$ for two manifolds X and Y .

Lemma 14.6. *Let X and Y be smooth manifolds, and let \mathcal{C} be a conic Lagrangian submanifold of $T^*X \times T^*Y$. Fix $(x_0, y_0; \xi_0, \eta_0) \in \mathcal{C}$, and assume that the vector*

$$\xi_0 \frac{\partial}{\partial \xi} + \eta_0 \frac{\partial}{\partial \eta}$$

is not tangent to \mathcal{C} . Then X and Y have coordinate systems $x = (x', x'')$ and $y = (y', y'')$ centered at x_0 and y_0 , such that $\xi_0 = (1, \dots, 0)$, $\eta_0 = (1, \dots, 0)$, and the tangent plane to \mathcal{C} is given by

$$dx' = dy' \quad \text{and} \quad d\xi' = d\eta' \quad \text{and} \quad d\xi'' = 0 \quad \text{and} \quad d\eta'' = 0.$$

Then (x', x'', η', y'') can be used as local coordinates for \mathcal{C} , and we can find a phase function $\phi(x', x'', y'', \eta')$ such that \mathcal{C} is parameterized by ϕ

14.3 Local Theory

Let us recall the theory of oscillatory integral distributions. Let $U \subset \mathbf{R}^n$ be open. Consider a real-valued phase function $\phi \in C^\infty(U_x \times \mathbf{R}_\theta^p)$, homogeneous in the θ variable, and such that $\nabla_{x,\theta} \phi$ is non-vanishing on the support of a symbol $a \in S^m(U \times \mathbf{R}^p)$. Then we can define a distribution u , formally speaking, by the equation

$$u(x) = \int_{\mathbf{R}^p} a(x, \theta) e^{2\pi i \phi(x, \theta)} d\theta.$$

If we consider the conic set

$$\Sigma_\theta = \{(x, \theta) \in U \times \mathbf{R}^p : \nabla_\theta \phi(x, \theta) = 0\},$$

then $\text{WF}(u) \subset \Sigma_\theta$. Let us now assume the phase is *nondegenerate*, in the sense that whenever $\nabla_\theta \phi = 0$, $D_{x,\theta}(\nabla_\theta \phi)$ is an invertible matrix. Then Σ_θ will be an n dimensional manifold in $U \times \mathbf{R}^p$, and the map $\Sigma_\theta \rightarrow T^*U$ given by $(x, \theta) \mapsto \nabla_x \phi(x, \theta)$ will be an immersion, which we will denote by Λ_θ . This immersed manifold will be a *Lagrangian submanifold* of T^*U , in the sense that the Tangent spaces W at each point of Λ_θ will satisfy $W^\perp = W$ with respect to the symplectic form $d\xi \wedge dx$.

A *Fourier integral operator* is a continuous operator $T : C_c^\infty(Y) \rightarrow \mathcal{D}'(X)$ whose Schwartz kernel K is given, modulo a smoothing kernel, by a sum

of oscillatory integral distributions of the form

$$K(x, y) = \int_{\mathbf{R}^p} a(x, y, \theta) e^{2\pi i \phi(x, y, \theta)} d\theta,$$

for some symbol $a \in S^\mu(X \times Y \times \mathbf{R}^N)$. It's *Canonical relation* is

$$C_\phi = \{(x, y; \nabla_x \phi(x, y, \theta), -\nabla_y \phi(x, y, \theta)) : \nabla_\theta \phi(x, y, \theta) = 0\},$$

which roughly speaking, tells us how T moves wave packets around in phase space. The *order* of the operator is $\mu - n/4 + p/2$ (which in particular, agrees with the previously understood case where $\phi(x, y, \xi) = (x - y) \cdot \xi$, i.e. with the theory of pseudodifferential operators.

Example. The solution to the half-wave equation $\partial_t f = \sqrt{-\Delta} f$ is given by the equation

$$e^{it\sqrt{-\Delta}} f = \int e^{2\pi i(\xi \cdot x + t|\xi|)} \hat{f}(\xi) = \int e^{2\pi i(\xi \cdot (x-y) + t|\xi|)} f(y) d\xi dy,$$

which is a Fourier integral operator which phase $\phi_t(x, y, \xi) = \xi \cdot (x - y) + t|\xi|$ and symbol $a_t(x, y, \xi) = 1$. Given two input functions f_0 and f_1 , if we write

$$f_+ = \frac{f_0 - i\sqrt{-\Delta}^{-1} f_1}{2} \quad \text{and} \quad f_- = \frac{f_0 + i\sqrt{-\Delta}^{-1} f_1}{2},$$

then the function

$$u(x, t) = e^{it\sqrt{-\Delta}} f_+ + e^{-it\sqrt{-\Delta}} f_-$$

solves the wave equation $\partial_t^2 u = \Delta u$, with initial conditions

$$u(x, 0) = f_+ + f_- = f_0$$

and

$$\partial_t u(x, 0) = i\sqrt{-\Delta} f_+ - i\sqrt{-\Delta} f_- = f_1.$$

Expanding out the definition of u , we obtain a sum of operators applied to f_0 and f_1 , of the form

$$Tf(x, t) \mapsto \int e^{2\pi i(\xi \cdot x \pm t|\xi|)} |\xi|^{-j} \hat{f}(\xi).$$

Each of these operators is a Fourier integral operator with canonical relation

$$C_\phi = \{y \mp t\xi/|\xi|, y; \xi, \xi\}.$$

Similarly, one may define wave propagators on a compact Riemannian manifold M . Locally in coordinates, these propagators can be expressed as Fourier integral operators, with a very similar canonical relation, i.e.

$$C_\phi = \{\exp_y(t\xi/|\xi|), y; \xi, \xi\},$$

so that singularities travel along geodesics.

Example. Let A_t be the spherical averaging operator, i.e. $A_t f(x)$ is the average of f on a sphere of radius t centered at x . We can write $A_t = f * \sigma_t$, where σ_t is the surface measure of the sphere of radius t . Recall that stationary phase tells us that

$$\hat{\sigma}(\xi) = \sum_{\pm} e^{\pm 2\pi i |\xi|} a_{\pm}(\xi)$$

for symbols a_{\pm} of order $-(n-1)/2$. Thus we can write

$$A_t f(x) = \sum_{\pm} \int e^{2\pi i (\xi \cdot x \pm t|\xi|)} a_{\pm}(t\xi) \hat{f}(\xi),$$

which relates the study of averaging operators to the theory of Fourier integral operators. Alternatively, if $\Phi(x, y, t) = |x - y|^2/t^2 - 1$, then

$$\begin{aligned} A_t f(x) &= \int \delta(\Phi(x, y, t)) f(y) dy \\ &= \int e^{2\pi i \lambda \Phi(x, y, t)} f(y) d\lambda dy. \end{aligned}$$

Thus we can express A_t in many different ways as Fourier integral operators.

The last example indicates an important problem with the theory of Fourier integral operators; there is a lack of uniqueness in the choice of phase defining the operator. Nonetheless, we have seen that the *wavefront set* is essentially uniquely verified, because it is connected to the relation between $\text{WF}(Tu)$ and $\text{WF}(u)$, which do not need a phase representation to be defined. The *equivalence of phase theorem* says that the canonical relation is essentially the only invariant depending on the phase; if two

phases have the same canonical relations, then Fourier integral operators defined in terms of one phase can be converted into an operator defined in terms of the other operator, and one has an asymptotic formula relating the two symbols of the associated

TODO: Merge these notes with later notes.

Let us now get to the new theory. Let M^n be a manifold, and let $\Lambda \subset T^*M - \{0_M\}$ be a conic Lagrangian submanifold. A *Lagrangian distribution* of order m on M associated with Λ is a distribution u such that for any $N > 0$, and any properly supported pseudodifferential operators P_1, \dots, P_N of order one with principal symbols vanishing on Λ , we have

$$(P_1 \circ \dots \circ P_N)u \in H_{-m-n/4, \text{loc}}^\infty(M).$$

The set of all Lagrangian distributions of order m is denoted $I^m(M, \Lambda)$. The condition above means, roughly, that one can differentiate u in directions away from Λ while maintaining the regularity of u . In particular, Sobolev embedding results imply that if $u \in I^m(M, \Lambda)$, then $\text{WF}(u) \subset \Lambda$. It will be easy to see what kinds of distributions are Lagrangian once we prove the equivalence of phase theorem, which allows one to reduce the study of Lagrangian distributions to oscillatory integrals like the ones above.

The next lemma is useful for the proof of the equivalence of phase theorem, and is, in fact, a special case.

Lemma 14.7. *Suppose that u is a compactly supported Lagrangian distribution in $I^m(\mathbf{R}^n, \Lambda)$, where*

$$\Lambda = \{(\nabla H(\xi), \xi) : \xi \in \mathbf{R}^n\}$$

for some smooth, homogeneous function H . Then for $|\xi| \geq 1$, there exists a symbol $v \in S^{m-n/4}(\mathbf{R}^n)$ such that

$$\hat{u}(\xi) = e^{-2\pi i H(\xi)} v(\xi),$$

i.e. so that formally speaking, u is given by the oscillatory integral distribution.

$$u(x) = \int v(\xi) e^{2\pi i (\xi \cdot x - H(\xi))} d\xi$$

Proof. TODO: 6.1.1 of Sogge. □

Theorem 14.8. Let ϕ be a non-degenerate phase function defined on an open conic neighborhood of a point $(x_0, \xi_0) \in T^*\mathbf{R}^n$, and consider the conic Lagrangian manifold Λ_ϕ . If $a \in S^\mu(\mathbf{R}^n \times \mathbf{R}^p)$ is supported on a sufficiently small conic neighborhood of (x_0, ξ_0) , then the oscillatory integral distribution

$$u(x) = \int a(x, \theta) e^{2\pi i \phi(x, \theta)} d\theta$$

lies in $I^m(\mathbf{R}^n, \Lambda)$, where $m = \mu - n/4 + p/2$. Moreover, there exists $v \in S^{m-n/4-1}$ such that

$$\hat{u}(\xi) = (2\pi)^{n/2-p/2} e^{-2\pi i H(\xi)} \left(a(x, \theta) | \text{Det}(H\phi) |^{-1/2} e^{(i\pi/4) \text{sgn}(H\phi)} + v(\xi) \right),$$

where (x, θ) is the unique solution to the differential equation $\nabla_\theta \phi(x, \theta) = 0$ and $\nabla_x \phi(x, \theta) = \xi$. Conversely, every Lagrangian distribution $u \in I^m(\mathbf{R}^n, \Lambda)$ with $\text{WF}(u)$ contained in a small enough neighborhood of (x_0, ξ_0) can be written in this form.

Example. Let $\phi(x, y, \xi)$ be smooth away from $\xi = 0$, homogeneous of degree one, and satisfy

$$\partial_\xi^\beta \{ \phi(x, y, \xi) - (x - y) \cdot \xi \} \lesssim_\beta |x - y|^2 |\xi|^{1-\beta}.$$

For any symbol $a \in S^m(\mathbf{R}^n \times \mathbf{R}^n)$ supported on $|x - y| \lesssim 1$ and $|\xi| \gtrsim 1$ in such a way that $|\nabla_\xi \phi| \gtrsim |x - y|$ and $|\nabla_x \phi| \gtrsim |\xi|$ on the support of a , define

$$K(x, y) = \int a(x, y, \xi) e^{[2\pi i \phi(x, y, \xi)]} d\theta$$

We have seen one proof that K is a pseudodifferential operator. Let's give another proof using the equivalence of phase theorem. The distribution K is defined by an oscillatory integral distribution with Σ_ϕ contained in $\{(x, x; \xi) : x \in \mathbf{R}^n, \xi \in \mathbf{R}^n\}$. The fact that $\phi(x, y, \xi) \approx (x - y) \cdot \xi$ means that the resulting Lagrangian manifold is then

$$\Lambda_\phi = \{(x, x; \xi, -\xi)\},$$

which is the same Lagrangian manifold associated pseudodifferential operators. To determine the Fourier transform of K using equivalence of phase, fix (x_0, ξ_0) . Then at $p = (x_0, x_0; \xi_0, -\xi_0)$, the tangent space to Λ_ϕ is the span of

$$\{\partial_{x_i} + \partial_{y_i}\} \cup \{\partial_{\xi_i} - \partial_{\eta_i}\}.$$

Let V be the Lagrangian subspace of $T_p(T^*M)$ spanned by

$$\{(a, 0, c, a + c)\}$$

$$\{\partial_{\eta_i}\}\cup\{\partial_{x_i}+\partial_{\xi_i}-\partial_{\eta_i}\}$$

$$e_1,\ldots,e_n=\partial_{x_i}+\partial_{y_i}$$

$$e'_1,\ldots,e'_n=\partial_{x_i}$$

$$f'_i=\partial_{\xi_i}-\partial_{\eta_i}$$

$$f_i=\partial_{\eta_i}$$

$$z^1=\xi_0dx-\xi_0dy$$

$$\text{Choose coordinates } \xi_0dx-\xi_0dy=dz^1$$

Now $u\in I^m(\mathbf{R}^n,\Lambda_\phi)$. The result above says that

$$\widehat{K}(\xi,\eta)=(2\pi)^{n/2}e^{-2\pi iH(\xi)}$$

Part III

Calderon-Zygmund Theory

Here, we try and describe the more modern approaches to real-variable harmonic analysis, as developed by the *Calderon-Zygmund school* in the 1960s and 1970s. Almost all of the problems we consider can be phrased as showing some operator is bounded as a map between functions spaces. Given some function f lying in a space V , we have an associated function Tf lying in some space W . The main goal of the techniques in this part of the book attempt to understand how quantitative control on certain properties of f imply quantitative control on properties of Tf . In particular, given some quantity $A(f)$ associated with each $f \in V$, and a quantity $B(g)$ defined for all $g \in W$, our goal is to understand whether a general bound $B(Tf) \lesssim A(f)$ is possible for all functions $f \in V$, i.e. whether there exists a universal constant $C > 0$ such that $B(Tf) \leq C \cdot A(f)$ for all $f \in V$.

A core technique we employ here is the method of *decomposition*. We write $f = \sum_k f_k$, where the functions f_k have particular properties, perhaps being concentrated in a particular region of space, or having a Fourier transform concentrated in a particular region. These concentration properties often simplify the analysis of the operator T , enabling us to obtain bounds $B(Tf_k) \lesssim A(f_k)$ for each n . Provided that the operator T , and the quantities A and B are ‘stable under addition’, we can then obtain the bound $B(Tf) \leq A(f)$ by ‘summing’ up the related quantities. The stability of A and B is often obtained by assuming these quantities are *norms* on their respective function spaces, i.e. that there exists norms $\|\cdot\|_V$ and $\|\cdot\|_W$ such that $A(f) = \|f\|_V$ for each $f \in V$ and $B(g) = \|g\|_W$ for each $g \in W$. The stability of T under addition is obtained by assuming linearity, or at least sub-linearity, in the sense that for each $f_1, f_2 \in V$,

$$\|T(f_1 + f_2)\|_W \leq \|Tf_1\|_W + \|Tf_2\|_W.$$

We can then use the triangle inequality to conclude that

$$\|Tf\|_W \leq \sum_k \|Tf_k\|_W \lesssim \sum_k \|f_k\|_V.$$

Thus if $\sum_k \|f_k\|_V \lesssim \|f\|_V$, our argument is complete. This will be true, for instance, if there exists $\varepsilon > 0$ such that $\|f_k\|_V \lesssim 2^{-\varepsilon k} \|f\|_V$. This can often be obtained if we employ one of a family of *dyadic decomposition techniques*. For such decompositions, it is also possible to generalize our techniques not only to norms, but also to *quasinorms*, i.e. maps $\|\cdot\|$ which are homogeneous and satisfy a *quasi-triangle inequality* $\|v + w\| \lesssim \|v\| + \|w\|$.

Lemma 14.9. Suppose $\|\cdot\|_V$ is a quasi-norm on a vector space V , and under the topology induced by $\|\cdot\|_V$, we can write $f = \sum_{k=1}^{\infty} f_k$, where there is $\varepsilon > 0$ and $C > 0$ such that for each n , $\|f_k\|_V \leq C \cdot 2^{-\varepsilon k}$. Then $\|f\|_V \lesssim_{\varepsilon} C$.

Remark. Thus if T is sublinear and we have $\|Tf_k\|_W \lesssim \|f_k\|_V$ and $\|f_k\|_V \lesssim 2^{-\varepsilon k} \|f\|_V$, we conclude $\|Tf_k\|_W \lesssim 2^{-\varepsilon k} \|f\|_V$, and then by sublinearity and the lemma applied to $\|\cdot\|_W$, we conclude

$$\|Tf\|_W \leq \left\| \sum_k Tf_k \right\|_W \lesssim_{\varepsilon} \|f\|_V.$$

A slight modification of the proof below even gives this claim provided T is *quasi sublinear*, in the sense that for all $f_1, f_2 \in V$, $\|T(f_1 + f_2)\|_W \lesssim \|Tf_1\|_V + \|Tf_2\|_V$ for all $f_1, f_2 \in V$. However, such operators occur so rarely in practice that it isn't worth concentrating on them.

Proof. Pick $A > 0$ such that $\|f_1 + f_2\|_V \leq A \cdot (\|f_1\|_V + \|f_2\|_V)$ for all f_1 and f_2 . If $A < 2^{\varepsilon}$, we can write apply the quasitriangle inequality iteratively to conclude

$$\|f\| \leq C \cdot \sum_{k=1}^{\infty} A^k \|f_k\|_V \leq C \cdot \left(\sum_{k=1}^{\infty} (A2^{-\varepsilon})^k \right) \leq C \cdot \left(\frac{1}{1 - A2^{-\varepsilon}} \right) \lesssim_{\varepsilon} C.$$

In general, fix N , and write $f = f^1 + \cdots + f^N$, where $f^m = \sum_{k=0}^{\infty} f_{m+Nk}$. Then $\|f_{m+Nk}\|_V \leq C \cdot 2^{-N\varepsilon k}$, and if N is chosen large enough that $A < 2^{N\varepsilon}$, we can apply the previous case to conclude that $\|f^m\|_V \lesssim_{\varepsilon} C$. Then we can apply the quasi-triangle inequality to conclude that $\|f\| \lesssim_{\varepsilon} C$. \square

We can even apply the method of decomposition in the presence of suitably large polynomial decay.

Lemma 14.10. Suppose $\|\cdot\|_V$ is a quasinorm on a function space V . Then there exists t such that for all $s > t$, if $f = \sum_{k=1}^{\infty} f_k$, and if $\|f_k\|_V \leq C \cdot k^{-s}$, for $s > t$, then $\|f\|_V \lesssim_s C$.

Proof. As in the previous lemma, pick $A > 0$ such that $\|f_1 + f_2\|_V \leq A(\|f_1\|_V + \|f_2\|_V)$ for all $f_1, f_2 \in V$. We perform a decomposition of dyadic type, writing $f = \sum_{m=0}^{\infty} f^m$, where

$$f^m = \sum_{k=2^m}^{2^{m+1}-1} f_k.$$

By applying the divide and conquer approach, splitting up the indices $2^m \leq k \leq 2^{m+1} - 1$ via a binary tree with depth at most $O(m)$, we can ensure that

$$\|f^m\|_V \lesssim A^{m+1} \sum_{k=2^m}^{2^{m+1}-1} \|f_k\|_V \leq C \cdot A^{m+1} \sum_{k=2^m}^{2^{m+1}-1} k^{-s} \lesssim C(A2^{1-s})^m.$$

If $s > 1 + \lg(A)$, the previous lemma implies that $\|f\|_V \lesssim C$. □

In this part of the notes, we define the various classes of quasi-norms we will study, describe the general methods which make up the Calderon-Zygmund theory, and find applications to geometric measure theory, complex analysis, partial differential equations, and analytic number theory.

Chapter 15

Monotone Rearrangement Invariant Norms

In this chapter, we discuss common families of *monotone, rearrangement invariant quasinorms* that occur in harmonic analysis. The general framework is as follows. For each function f , we associate its *distribution function* $F : [0, \infty) \rightarrow [0, \infty]$ given by $F(t) = |\{x : |f(x)| > t\}|$. A *rearrangement invariant space* is a subspace V of the collection of measurable complex-valued functions on some measure space X , equipped with a quasi-norm $\|\cdot\|$, satisfying the following two properties:

- *Monotonicity*: If $|f(x)| \leq |g(x)|$ for all $x \in X$, then $\|f\| \leq \|g\|$.
- *Rearrangement-Invariance*: If f and g have the same distribution function, then $\|f\| = \|g\|$.

A monotone rearrangement-invariant norm essentially provides a way of quantifying the height and width of functions on X . It has no interest in the ‘shape’ of the objects studied, because of the property of rearrangement invariance. In a particular problem, one picks the norm best emphasizing a particular family of features useful in the problem. Height and width are of course, vague properties; but roughly speaking, the canonical example of a function with height H and width W is a function $f = H \cdot \mathbf{I}_E$, where $|E| = W$. More general functions can be thought of as being composed from a combination of functions with various heights and widths, and since a unique way to quantify the behaviour of these combinations is not obvious, we will obtain various different ways to quantify how various heights and widths come together.

There are two very useful classes of functions useful for testing the behaviour of translation invariant norms:

- The *indicator functions* $\mathbf{I}_E(x) = \mathbf{I}(x \in E)$, for a measurable set E .
- The *simple functions* $f = \sum_{i=1}^n a_i \mathbf{I}_{E_i}$, for disjoint sets E_i .

The class of all simple functions forms a vector space, and for almost all the monotone rearrangement invariant norm we consider in this section, this vector space will form a dense subspace of the class of all functions. This means that when we want to study how an operator transforms the height and width of functions, the behaviour of the operator on simple functions often reflects the behaviour of an arbitrary function.

15.1 The L^p norms

We begin by introducing the most fundamental monotone, rearrangement invariant norms. For $p \in (0, \infty)$, we define the L^p norm of a measurable function f on a measure space X by

$$\|f\|_{L^p(X)} = \left(\int_X |f(x)|^p dx \right)^{1/p}.$$

For $p = \infty$, we define

$$\|f\|_{L^\infty(X)} = \min \{t \geq 0 : |f(x)| \leq t \text{ almost surely}\},$$

a quantity often called the *essential* supremum. If the measure space X is implicit, these quantities are also denoted by $\|f\|_p$, as we do often in this chapter. The space of functions f with $\|f\|_p < \infty$ is denoted by $L^p(X)$. The most important spaces to consider here are the space $L^1(X)$, consisting of absolutely square integrable functions, $L^\infty(X)$, consisting of almost-everywhere bounded functions, and $L^2(X)$, consisting of square integrable functions. The main motivation for the introduction of the other L^p spaces is that much of the quantitative theory in harmonic analysis for $p = 1$ and $p = \infty$ is rather trivial, in the sense that for most operators that occur in practice, it is simple to determine whether these operators are bounded on these spaces; obtaining L^p bounds of an operator for $1 < p < \infty$ reflects a deeper understanding of the quantitative properties of an operator.

As p increases, the L^p norm of a particular function f gives more control over the height of the function f , and weaker control on values where f is particularly small. At one extreme, $L^\infty(X)$ only has control over the height of a function, and no control over its width. Conversely, as $p \rightarrow 0$, $L^p(X)$ has more control over the width of functions. It is therefore natural to introduce the space $L^0(X)$ as the space of measurable functions with finite measure support. But there is no natural norm on $L^0(X)$ which can classify the support of functions. After all, such a quantity couldn't be homogeneous, since the width of f and αf are the same for each $\alpha \neq 0$.

Example. If $f(x) = |x|^{-s}$ for $x \in \mathbf{R}^d$ and $s > 0$, then integration by radial coordinates shows that

$$\int_{\varepsilon \leq |x| \leq M} \frac{1}{|x|^{sp}} dx \approx \int_{\varepsilon}^M r^{d-1-ps} dr = \frac{M^{d-ps} - \varepsilon^{d-ps}}{d-ps}.$$

This quantity remains finite as $\varepsilon \rightarrow 0$ if and only if $d > ps$, and finite as we let $M \rightarrow \infty$ if and only if $d < ps$. Thus if $p < d/s$, f is locally in L^p , in the sense that $f \in L^p(B)$ for every bounded $B \in \mathbf{R}^d$. The class of functions for which this condition holds is denoted $L_{loc}^p(X)$. Conversely, if $p > d/s$, then for every closed set B not containing the origin, $f \in L^p(B)$. For $p = d/s$, the function f fails to be $L^p(\mathbf{R}^d)$, but only 'by a logarithm', which manifests in the fact that

$$\int_{\varepsilon \leq |x| \leq M} \frac{1}{|x|^{sp}} dx \approx \int_{\varepsilon}^M \frac{dr}{r} = \log(M/\varepsilon).$$

We will later introduce Lorentz norms $L^{p,q}$, which one can think of as differing from the standard L^p norms 'by a logarithmic factor', and the $L^{p,q}$ norm of f will be finite if q is chosen appropriately.

The last example shows that, roughly speaking, control on the L^p norm of a function for large values of p prevents the formation of sharp singularities, and control of an L^p norm for small values of p ensures that functions have large decay at infinity.

Example. If $s = A\chi_E$, and we set $H = |A|$ and $W = |E|$, then $\|s\|_p = W^{1/p}H$. As $p \rightarrow \infty$, the value of $\|s\|_p$ depends more and more on H , and less on W , and in fact $\lim_{p \rightarrow \infty} \|s\|_p = H$. If $s = \sum A_n \chi_{E_n}$, and $|A_m|$ is the largest constant from all other values A_n , then as p becomes large, $|A_m|^p$ overwhelms all other terms. We calculate that as $p \rightarrow \infty$,

$$\|s\|_p = \left(\sum |E_n| |A_n|^p \right)^{1/p} = |A_m|^p (|E_m| + o(1))^{1/p} = |A_m| (1 + o(1)).$$

This implies $\|s\|_p \rightarrow |A_m|$ as $p \rightarrow \infty$. But as $p \rightarrow 0$, $\lim_{p \rightarrow 0} \|f\|_p$ does not in general exist, even for step functions with finite measure support. Nonetheless, we can conclude that $\lim_{p \rightarrow 0} \|s\|_p^p = \sum |E_n|$.

As $p \rightarrow \infty$, the last example shows the width of a function is disregarded completely by the L^p norm, from which it follows that $\|s\|_{L^p(X)} \rightarrow \|s\|_{L^\infty(X)}$ as $p \rightarrow \infty$. The same is true for more general functions, which we can prove using a density argument.

Theorem 15.1. *Let $p \in (0, \infty)$. If $f \in L^p(X) \cap L^\infty(X)$, then*

$$\lim_{t \rightarrow \infty} \|f\|_t = \|f\|_\infty.$$

Proof. Without loss of generality, assume $p \geq 1$. Consider the norm $\|\cdot\|$ on $L^p(X) \cap L^\infty(X)$ given by

$$\|f\| = \|f\|_p + \|f\|_\infty.$$

Then $L^p(X) \cap L^\infty(X)$ is complete with respect to this metric. For each $t \in [p, \infty)$, define $T_t(f) = \|f\|_t$. Then the functions $\{T_t\}$ are uniformly bounded in the norm $\|\cdot\|$, since if $p = \theta t$, then

$$|T_t(f)| = \|f\|_t \leq \|f\|_p^\theta \|f\|_\infty^{1-\theta} \leq \|f\|^\theta \|f\|^{1-\theta} = \|f\|.$$

For any $\varepsilon > 0$, we can find a step function s with $\|s - f\|_p, \|s - f\|_\infty \leq \varepsilon$. This means that for all $t \in (p, \infty)$, $\|s - f\|_t \leq \varepsilon$. And so

$$|T_t(f) - \|f\|_\infty| \leq |T_t(f) - T_t(s)| + |T_t(s) - \|s\|_\infty| + \|\|s\|_\infty - \|f\|_\infty\| \leq 2\varepsilon + o(1).$$

Taking $\varepsilon \rightarrow 0$ gives the result. \square

Abusing notation, we define $\|f\|_0^0 = |\text{supp } f| = |\{x : f(x) \neq 0\}|$, and let $L^0(X)$ be the space of functions with finite support. We know that for any simple function s , $\|s\|_p^p \rightarrow \|s\|_0^0$ as $p \rightarrow 0$. If $f \in L^0(X) \cap L^p(X)$ for some $p \in (0, \infty)$, then the monotone and dominated convergence theorems implies that

$$\|f\|_0^0 = \int \mathbf{I}(f(x) \neq 0) = \int \left(\lim_{t \rightarrow 0} |f(x)|^t \right) dx = \lim_{t \rightarrow 0} \int |f(x)|^t dx = \lim_{t \rightarrow 0} \|f\|_t^t.$$

Thus the space $L^0(X)$ lies at the opposite end of the spectrum to L^∞ .

The fact that $\|f\|_0^0$ is a norm taken to the ‘power of zero’ implies that many nice norm properties of the L^p spaces fail to hold for $L^0(X)$. For instance, homogeneity no longer holds; in fact, for each $\alpha \neq 0$,

$$\|\alpha f\|_0^0 = \|f\|_0^0.$$

It does, however, satisfy the triangle inequality $\|f + g\|_0^0 \leq \|f\|_0^0 + \|g\|_0^0$, which follows from a union bound on the supports of the functions.

Example. Let $p < q$, and suppose $f \in L^p(X) \cap L^q(X)$. For any $r \in (p, q)$, the L^r norm emphasizes the height of f less than the L^q norm, and emphasizes the width of f less than the L^p norm. In particular, we find that for any $\lambda \geq 0$,

$$\begin{aligned} \|f\|_r^r &= \int_{\mathbf{R}} |f(x)|^r dx = \int_{|f(x)| \leq 1} |f(x)|^r dx + \int_{|f(x)| > 1} |f(x)|^r dx \\ &\leq \int_{|f(x)| \leq 1} |f(x)|^p dx + \int_{|f(x)| > 1} |f(x)|^q dx \\ &\leq \|f\|_p^p + \|f\|_q^q < \infty. \end{aligned}$$

In particular, this shows $f \in L^r(X)$.

Remark. The bound obtained in the last example can be improved by using scaling symmetries. For any $A > 0$,

$$\|f\|_r^r = \frac{\|Af\|_r^r}{A^r} \leq \frac{\|Af\|_p^p + \|Af\|_q^q}{A^r} \leq \frac{A^p \|f\|_p^p + A^q \|f\|_q^q}{A^r}.$$

If $1/r = \theta/p + (1-\theta)/q$, and we set $A = \|f\|_q^{q/(p-q)} / \|f\|_p^{p/(p-q)}$, then the above inequality implies $\|f\|_r \leq 2 \|f\|_p^\theta \|f\|_q^{1-\theta}$, which is a homogenous equality. The constant 2 can be removed in the equation using the *tensor power trick*. If we consider the function on X^n defined by $f^{\otimes n}(x_1, \dots, x_n) = f(x_1) \dots f(x_n)$, then $\|f^{\otimes n}\|_r = \|f\|_r^n$, and so

$$\|f\|_r = \|f^{\otimes n}\|_r^{1/n} \leq \left(2 \|f^{\otimes n}\|_p^\theta \|f^{\otimes n}\|_q^{1-\theta} \right)^{1/n} = 2^{1/n} \|f\|_p^\theta \|f\|_q^{1-\theta}.$$

We can then take $n \rightarrow \infty$ to conclude that $\|f\|_r \leq \|f\|_p^\theta \|f\|_q^{1-\theta}$, a special case of Hölder’s Inequality.

The argument in the last remark is an instance of *real interpolation*; In order to conclude some fact about a function which lies ‘between’ two other functions we know how to deal with, we split the function up into two parts lying in the other spaces, deal with them separately, and then put them back together to get some equality. This often introduces some extraneous (though not too inefficient) constants. But if these constants are unnecessary, one can often apply various symmetry considerations (homogeneity and the tensor power trick being two examples) to eliminate extraneous constants. We now also show how to prove this inequality using convexity, which illustrates another core technique. In the next theorem, $1/\infty = 0$.

Theorem 15.2 (Hölder). *If $0 < p, q \leq \infty$ and $1/p + 1/q = 1/r$,*

$$\|fg\|_r \leq \|f\|_p \|g\|_q.$$

Proof. The case where p or q is ∞ is left as an exercise to the reader. In the other case, by moving around exponents, we may simplify to the case where $r = 1$. The theorem depends on the log convexity inequality, such that for $A, B \geq 0$ and $0 \leq \theta \leq 1$, $A^\theta B^{1-\theta} \leq \theta A + (1 - \theta)B$. But since the logarithm is concave, we calculate

$$\log(A^\theta B^{1-\theta}) = \theta \log A + (1 - \theta) \log B \leq \log(\theta A + (1 - \theta)B),$$

and we can then exponentiate. To prove Hölder’s inequality, by scaling f and g , which is fine by homogeneity, we may assume that $\|f\|_p = \|g\|_q = 1$. Then we calculate

$$\begin{aligned} \|fg\|_1 &= \int |f(x)| |g(x)| = \int |f(x)|^{p/p} |g(x)|^{q/q} \\ &\leq \int \frac{|f(x)|^p}{p} + \frac{|g(x)|^q}{q} = \frac{1}{p} + \frac{1}{q} = 1 = \|f\|_p \|g\|_q. \end{aligned}$$

If $p = \infty$, $q = 1$, then the inequality is trivial, since we have the pointwise inequality $|f(x)g(x)| \leq \|f\|_\infty |g(x)|$ almost everywhere, which we can then integrate. \square

Remark. Note that $A^\theta B^{1-\theta} \leq \theta A + (1 - \theta)B$ is an *equality* if and only if $A = B$, or $\theta \in \{0, 1\}$. In particular, following through the proof above shows that if $\|f\|_p = \|g\|_q = 1$, we must have $|f(x)|^{1/p} = |g(x)|^{1/q}$ almost everywhere. In general, this means Hölder’s inequality is sharp if and only if $|f(x)|^{1/p}$ is a constant multiple of $|g(x)|^{1/q}$.

The next inequality is known as the *triangle inequality*.

Corollary 15.3. *Given f, g , and $p \geq 1$, $\|f + g\|_p \leq \|f\|_p + \|g\|_p$.*

Proof. The inequality when $p = 1$ is obtained by integrating the inequality $|f(x) + g(x)| \leq |f(x)| + |g(x)|$, and the case $p = \infty$ is equally trivial. When $1 < p < \infty$, by scaling we can assume that $\|f\|_p + \|g\|_p = 1$. Then we can apply Hölder's inequality combined with the $p = 1$ case to conclude

$$\begin{aligned} \int |f(x) + g(x)|^p &\leq \int |f(x)| |f(x) + g(x)|^{p-1} + \int |g(x)| |f(x) + g(x)|^{p-1} \\ &\leq \|f\|_p \|f + g\|_q^{p-1} + \|g\|_p \|f + g\|_q^{p-1} = \|f + g\|_p^{p-1} \end{aligned}$$

Thus $\|f + g\|_p^p \leq \|f + g\|_p^{p-1}$, and simplifying gives $\|f + g\|_p \leq 1$. \square

Remark. Suppose $\|f + g\|_p = \|f\|_p + \|g\|_p$. Following through the proof given above shows that both applications of Hölder's inequality must be sharp. And this is true if and only if $|f(x)|^p$ and $|g(x)|^p$ are scalar multiples of $|f(x) + g(x)|^p$ almost everywhere. But this means $|f(x)|$ and $|g(x)|$ are scalar multiples of $|f(x) + g(x)|$. If $|f(x)| = A|f(x) + g(x)|$ and $|g(x)| = B|f(x) + g(x)|$. If $g \neq 0$, this implies there is C such that $|f(x)| = C|g(x)|$ for some $C > 0$. Thus we can write $f(x) = Ce^{i\theta(x)}g(x)$, and we must have

$$\|f + g\|_p^p = \int |1 + Ce^{i\theta(x)}|^p |g(x)|^p = (1 + C)^p \int |g(x)|^p$$

so $|1 + Ce^{i\theta(x)}| = |1 + C|$ almost everywhere but this can only be true if $e^{i\theta(x)} = 1$ almost everywhere, so $f = Cg$. Thus the triangle inequality is only sharp if f and g are positive scalar multiples of one another.

This discussion leads to a useful heuristic: Unless f and g are 'aligned' in a certain way, the triangle inequality is rarely sharp. For instance, if f and g have disjoint support, we calculate that

$$\|f + g\|_p = \left(\|f\|_p^p + \|g\|_p^p \right)^{1/p}$$

For $p > 1$, this is always sharper than the triangle inequality.

If $p < 1$, then the proof of Corollary 15.3 no longer works, and in fact, is no longer true. In fact, if f and g are non-negative functions, then we actually have the *anti* triangle inequality

$$\|f + g\|_p \geq \|f\|_p + \|g\|_p,$$

as proved in the next theorem.

Theorem 15.4. If $p \geq 1$, then for any functions $f_1, \dots, f_N \geq 0$,

$$(\|f_1\|_p^p + \dots + \|f_N\|_p^p)^{1/p} \leq \|f_1 + \dots + f_N\|_p \leq \|f_1\|_p + \dots + \|f_N\|_p. \quad (15.1)$$

If $p \leq 1$, then the inequality reverses, i.e. for any positive functions f_1, \dots, f_N ,

$$\|f_1\|_p + \dots + \|f_N\|_p \leq \|f_1 + \dots + f_N\|_p \leq (\|f_1\|_p^p + \dots + \|f_N\|_p^p)^{1/p} \quad (15.2)$$

Proof. The upper bound in (15.1) is just obtained by applying the triangle inequality iteratively. To obtain the lower bound, we note that for $A_1, \dots, A_N \geq 0$,

$$(A_1 + \dots + A_N)^p \geq A_1^p + \dots + A_N^p,$$

One can prove this from induction from the inequality $(A_1 + A_2)^p \geq A_1^p + A_2^p$, which holds when $A_2 = 0$, and the derivative of the left hand side is greater than the right hand side for all $A_2 \geq 0$. But then setting $A_k = f_k$ and then integrating gives

$$\|f_1 + \dots + f_N\|_p^p \geq \|f_1\|_p^p + \dots + \|f_N\|_p^p.$$

Now assume $0 < p < 1$. We begin by proving the lower bound in 15.2. We can assume $N = 2$, and $\|f_1\|_p + \|f_2\|_p = 1$, and then it suffices to show $\|f_1 + f_2\|_p \geq 1$. For any $\theta \in (0, 1)$, and $A, B \geq 0$, concavity implies

$$(A + B)^p = (\theta(A/\theta) + (1 - \theta)(B/(1 - \theta)))^p \geq \theta^{1-p} A^p + (1 - \theta)^{1-p} B^p.$$

Thus setting $A = f_1(x)$, $B = f_2(x)$, and $\theta = \|f_1\|_p$, so that $1 - \theta = \|f_2\|_p$, and then integrating, we find

$$\|f_1 + f_2\|_p^p \geq \theta + (1 - \theta) = 1.$$

On the other hand, the inequality $(A_1 + \dots + A_N)^p \leq A_1^p + \dots + A_N^p$, which holds for $A_1, \dots, A_N \geq 0$, can be applied with $f_k = A_k$ and integrated to yield

$$\|f_1 + \dots + f_N\|_p^p \leq \|f_1\|_p^p + \dots + \|f_N\|_p^p. \quad \square$$

Thus the triangle inequality is not satisfied for the L^p norms when $p < 1$. However, for $p < 1$, we do have a *quasi* triangle inequality.

Theorem 15.5. For $f_1, \dots, f_N \in L^p(X)$, with $0 < p < 1$,

$$\|f_1 + \dots + f_N\|_p \leq N^{1/p-1} (\|f_1\|_p + \dots + \|f_N\|_p).$$

Proof. By Hölder's inequality applied to sums,

$$\|f_1 + \cdots + f_N\|_p \leq (\|f\|_p^p + \cdots + \|f_N\|_p^p)^{1/p} \leq N^{1/p-1}(\|f_1\|_p + \cdots + \|f_N\|_p). \quad \square$$

This result is sharp, i.e. if we take a disjoint family of sets $\{E_1, E_2, \dots\}$ with $|E_i| = 1$ for each i , and then set $f_i = \mathbf{I}_{E_i}$, then the inequality is sharp for each N .

Remark. When $p < 1$, the space $L^p(X)$ is *not* normable. To see why, we look at the topological features of $L^p(X)$. Fix $\varepsilon > 0$, and let C be a convex set containing all functions f with $\|f\|_p < \varepsilon$. Thus, in particular, C contains all step functions $H\mathbf{I}_E$ where $H|E|^{1/p} < \varepsilon$. But if we now find a countable sequence of disjoint sets $\{E_k\}$, each with positive measure, and for each k , define $H_k = (\varepsilon/2)|E_k|^{-1/p}$, then for any N , the function

$$f_N = (H_1/N)\mathbf{I}_{E_1} + \cdots + (H_N/N)\mathbf{I}_{E_N}$$

lies in C , and

$$\|f_N\|_p = (1/N)(H_1^p|E_1| + \cdots + H_N^p|E_N|)^{1/p} = (\varepsilon/2)N^{1/p-1}$$

as $N \rightarrow \infty$, the L^p norm of f_N becomes unbounded. In particular, this means that we have proven that every bounded convex subset of $L^p(X)$ has empty interior, and a norm space certainly does not have this property.

As we have mentioned, as $p \rightarrow \infty$, the L^p norm excludes functions with large peaks, or large height, and as $p \rightarrow 0$, the L^p norm excludes functions with large tails, or large width. They form a continuously changing family of functions as p ranges over the positive numbers. In general, there is no inclusion of $L^p(X)$ in $L^q(X)$ for any p, q , except in two circumstances which occur often enough to be mentioned.

Example. If X is a finite measure space, and $0 < p \leq q \leq \infty$, $L^q(X) \subset L^p(X)$, because Hölder's inequality implies

$$\|f\|_p = \|f\chi_X\|_p \leq \|f\|_q |X|^{1/p-1/q}.$$

Taking $q \rightarrow \infty$, we conclude $\|f\|_p \leq |X|^{1/p}\|f\|_\infty$. One can best remember the constants here by the formula

$$\left(\int |f(x)|^p\right)^{1/p} \leq \left(\int |f(x)|^q\right)^{1/q}.$$

In particular, when X is a probability space, the L^p norms are increasing.

Example. On the other hand, suppose the measure space is granular, in the sense that there is $\varepsilon > 0$ such that either $|E| = 0$ or $|E| \geq \varepsilon$ for any measurable set E . Then $L^q(X) \subset L^p(X)$ for $0 < p \leq q \leq \infty$. First we check the $q = \infty$ case, which follows by the trivial estimate

$$\int |f(x)|^p \geq \varepsilon \|f\|_\infty^p,$$

so $\|f\|_\infty \leq \|f\|_p \varepsilon^{-1/p}$. But then applying log convexity, if $p \leq q < \infty$, we can write $1/q = \theta/p$ for $0 < \theta \leq 1$, and then log convexity shows

$$\|f\|_q = \|f\|_p^\theta \|f\|_\infty^{1-\theta} \leq \varepsilon^{-(1-\theta)/p} \|f\|_p = \varepsilon^{-1/p-1/q} \|f\|_p.$$

If $\varepsilon = 1$, which occurs if $X = \mathbf{Z}$, then the L^p norms are decreasing in p . This gives the best way to remember the constants involved, since the measure $\mu(E) = |E|/\varepsilon$ is one granular, and so

$$\left(\frac{1}{\varepsilon} \int |f(x)|^q dx \right)^{1/q} \leq \left(\frac{1}{\varepsilon} \int |f(x)|^p dx \right)^{1/p}.$$

Remark. We can often use such results in spaces which are not granular by coarsening the sigma algebra. For instance, the Lebesgue measure is ε^d granular over the sigma algebra generated by the length ε cubes whose corner's lie on the lattice $(\mathbf{Z}/\varepsilon)^d$, and if a function is measurable with respect to such a σ algebra we call the function ε -granular.

One can also often obtain analogous results when dealing with functions which are roughly constant at a scale ε , rather than literally constant at this scale. Basic examples of this include Bernstein's inequality; the Sobolev embeddings are also of this flavor. But this is a topic for another section.

Remark. If we let $X = \{1, \dots, N\}$, then X is both finite and granular, so all L^p norms are comparable. In particular, if $p \leq q$,

$$\|f\|_q \leq \|f\|_p \leq N^{1/p-1/q} \|f\|_q.$$

The left hand side of this inequality becomes sharp when f is concentrated at a single point, i.e. $f(n) = \mathbf{I}(n=1)$. On the other hand, the right hand side becomes sharp when f is constant, i.e. $f(n) = 1$ for all n .

Example. We can obtain similar L^p bounds by controlling the functions f involved, rather than the measure space. For instance, if $|f(x)| \leq M$, and $p \leq q$, then $\|f\|_q \leq \|f\|_p^{p/q} M^{1-p/q}$, which follows by log convexity. On the other hand, if $|f(x)| \geq M$ on the support of f , then $\|f\|_p \leq \|f\|_q^{q/p} M^{1-q/p}$.

Theorem 15.6. If p_θ lies between p_0 and p_1 , then

$$L^{p_0}(X) \cap L^{p_1}(X) \subset L^{p_\theta}(X) \subset L^{p_0}(X) + L^{p_1}(X)$$

Proof. If $\|f\|_{p_0}, \|f\|_{p_1} < \infty$, then for any p_θ between p_0 and p_1 ,

$$\begin{aligned} \|f \chi_{|f| \leq 1}\|_{p_\theta}^{p_\theta} &= \int_{|f| \leq 1} |f|^{p_\theta} \leq \int_{|f| \leq 1} |f|^{p_0} < \infty \\ \|f \chi_{|f| > 1}\|_{p_\theta}^{p_\theta} &= \int_{|f| > 1} |f|^{p_\theta} \leq \int_{|f| > 1} |f|^{p_1} < \infty \end{aligned}$$

Applying the triangle inequality, we conclude that $\|f\|_{p_\theta} < \infty$. In the case where $p_1 = \infty$, then $f \chi_{|f| > 1}$ is bounded, and must have finite support if $p_0 < \infty$, which shows this integral is bounded. Note the inequalities above show that we can split any function with finite L^{p_θ} norm into the sum of a function with finite L^{p_0} norm and another with finite L^{p_1} norm. \square

Remark. This theorem is important in the study of interpolation theory, because if we have two linear operators T_{p_0} defined on $L^{p_0}(X)$ and T_{p_1} on $L^{p_1}(X)$, and they agree on $L^{p_0}(X) \cap L^{p_1}(X)$, then there is a unique linear operator T_{p_θ} on $L^{p_\theta}(X)$ which agrees with these two functions, and we can consider the boundedness of such a function with respect to the L^{p_θ} norms.

The last property of the L^p norms we want to focus on is the principle of *duality*. Given any values of p and q with $1/p + 1/q = 1$, Hölder's inequality implies that if $f \in L^p(X)$ and $g \in L^q(X)$, then $fg \in L^1(X)$. In particular, for each function $g \in L^q(X)$, the map

$$\lambda : f \mapsto \int f(x)g(x) dx$$

is a linear functional on $L^p(X)$. Hölder's inequality implies that $\|\lambda\| \leq \|g\|_q$. But this is actually an *equality*. In particular, if $1 < p < \infty$, one can show these are *all* linear functionals. For $p \in \{1, \infty\}$, the dual space of $L^p(X)$ is more subtle. But, since in harmonic analysis we concentrate on quantitative bounds, the following theorem often suffices as a replacement.

Theorem 15.7. *If $1 \leq p < \infty$, and $f \in L^p(X)$, then*

$$\|f\|_p = \sup \left\{ \int f(x)g(x) : \|g\|_q = 1 \right\}.$$

If the underlying measure space is σ finite, then this claim also holds for $p = \infty$.

Proof. Suppose that $1 \leq p < \infty$. Given f , we define

$$g(x) = \frac{1}{\|f\|_p^{p-1}} \operatorname{sgn}(f(x)) |f(x)|^{p-1}.$$

If $\|f\|_p < \infty$, then

$$\|g\|_q^q = \frac{1}{\|f\|_p^{pq-q}} \int |f(x)|^{pq-q} = \frac{1}{\|f\|_p^p} \|f\|_p^p = 1,$$

and

$$\int f(x)g(x) = \frac{1}{\|f\|_p^{p-1}} \int |f(x)|^p = \|f\|_p.$$

On the other hand, suppose $\|f\|_p = \infty$. Then there exists a sequence of step functions $s_1 \leq s_2 \leq \dots \rightarrow |f|$. Each s_k lies in $L^p(X)$, but the monotone convergence theorem implies that $\|s_k\|_p \rightarrow \infty$. For each k , find a function $g_k \geq 0$ with $\|g_k\|_q = 1$, and $\int g_k(x)s_k(x) \geq \|s_k\|_p/2$. Then

$$\int g_k(x) \cdot \operatorname{sgn}(f(x))f(x) = \int g_k(x)|f(x)| \geq \int g_k(x)s_k(x) \geq \|s_k\|_p/2 \rightarrow \infty,$$

this completes the proof in this case.

Now we take the case $p = \infty$. Given any f , fix $\varepsilon > 0$. Then we can find a set E with $0 < |E| < \infty$ such that $|f(x)| \geq \|f\|_\infty - \varepsilon$ for $x \in E$. If $g(x) = \operatorname{sgn}(f(x))\mathbf{I}_E/|E|$, then $\|g\|_1 = 1$, and

$$\int f(x)g(x) = \frac{1}{|E|} \int_E |f(x)| \geq \|f\|_\infty - \varepsilon.$$

Taking $\varepsilon \rightarrow 0$ completes the claim. □

15.2 Decreasing Rearrangements

The measure-theoretic properties of a function's distribution are best reflected quite simply in the *distribution function* of the function f , i.e. the function $F : [0, \infty) \rightarrow [0, \infty)$ given by $F(t) = |\{x : |f(x)| > t\}|$, and any rearrangement invariant norm on f should be a function of F . The function F is right-continuous and decreasing, but has a jump discontinuity whenever $\{x : |f(x)| = t\}$ is a set of positive measure. We denote distributions of functions g and h by G and H .

Lemma 15.8. *Given a function f and g , $\alpha \in \mathbb{C}$, and $t, s > 0$, then*

- *If $|g| \leq |f|$, then $G \leq F$.*
- *If $g = \alpha f$, then $G(t) = F(t/|\alpha|)$.*
- *If $h = f + g$, then $H(t + s) \leq F(t) + G(s)$.*
- *If $h = fg$, then $H(ts) \leq F(t) + G(s)$.*

Proof. The first point follows because $\{x : |g(x)| > t\} \subset \{x : |f(x)| > t\}$, and the second because $\{x : |\alpha f(x)| > t\} = \{x : |f(x)| > t/|\alpha|\}$. The third point follows because if $|f(x) + g(x)| \geq t + s$, then either $|f(x)| \geq t$ or $|g(x)| \geq s$. Finally, if $|f(x)g(x)| \geq ts$, then $|f(x)| \geq t$ or $|g(x)| \geq s$. \square

We can simplify the study of the distribution of f even more by defining the *decreasing rearrangement* of f , a decreasing function $f^* : [0, \infty) \rightarrow [0, \infty)$ such that $f^*(s)$ is the *smallest* number t such that $F(t) \leq s$. Effectively, $f^*(s)$ is the inverse of F :

- If there is a unique t with $F(t) = s$, then $f^*(s) = t$.
- If there are multiple values t with $F(t) = s$, let $f^*(s)$ be the *smallest* such value.
- If there are no values t with $F(t) = s$, then we pick the first value t with $F(t) < s$.

We find

$$\{s : f^*(s) > t\} = \{s : s < F(t)\} = [0, F(t)),$$

which has measure $F(t)$. This is the most important property of f^* ; it is a decreasing function on the line which has the same distribution as the

function $|f|$. It is also the unique such function which is right continuous. Thus our intuition when analyzing monotone, rearrangement invariant norms is not harmed if we focus on right continuous decreasing functions.

Theorem 15.9. *The function f^* is right continuous.*

Proof. We note that $F(t) > s$ if and only if $t < f^*(s)$. Since f^* is decreasing, for any $s \geq 0$, we automatically have $f^*(s^+) \leq f^*(s)$. If $f^*(s^+) < f^*(s)$, then

$$s < F(f^*(s^+)) \leq F(f^*(s)) \leq s,$$

which gives a contradiction, so $f^*(s) = f^*(s^+)$. \square

Remark. We have a jump discontinuity at a point s wherever F is flat, and f^* is flat wherever F has a jump discontinuity.

In particular, when understanding intuition about monotone rearrangement invariant norms, one is allowed to focus on non-increasing, right continuous functions on $(0, \infty)$. For instance, this means that these norms do not care about the number of singularities that a function has, since all these singularities ‘pile up’ in the decreasing rearrangement. The ‘mass’ of these singularities, of course, is important.

15.3 Weak Norms

The weak L^p norms are obtained as a slight ‘refinement’ of the L^p norms.

Theorem 15.10. *If ϕ is an increasing, differentiable function on the real line with $\phi(0) = 0$, then*

$$\int_X \phi(|f(x)|) = \int_0^\infty \phi'(t) F(t) dt$$

Proof. An application of Fubini’s theorem is all that is needed to show

$$\begin{aligned} \int_X \phi(|f(x)|) dx &= \int_X \int_0^{|f(x)|} \phi'(t) dt dx \\ &= \int_0^\infty \phi'(t) \int_{|f(x)| > t} dx du \\ &= \int_0^\infty \phi'(t) F(t) dt. \end{aligned} \quad \square$$

As a special case we find

$$\|f\|_p = \left(p \int_0^\infty F(t) t^p \frac{dt}{t} \right)^{1/p}.$$

For this to be true, $F(t)$ must tend to zero ‘logarithmically faster’ than $1/t^p$. Indeed, we find

$$F(t) = |\{|f|^p > t^p\}| \leq \frac{1}{t^p} \int |f|^p = \frac{\|f\|_p^p}{t^p},$$

a fact known as *Chebyshev’s inequality*. But a bound $F(t) \lesssim 1/t^p$ might be true even if $f \notin L^p(\mathbf{R}^d)$. This leads to the *weak L^p norm*, denoted by $\|f\|_{p,\infty}$, which is defined to be the smallest value A such that $F(t) \leq (A/t)^p$ for all t . We let $L^{p,\infty}(X)$ denote the space of all functions f for which $\|f\|_{p,\infty} < \infty$. By Chebyshev’s inequality, $\|f\|_{p,\infty} \leq \|f\|_p$ for any function f . The reason that the value A occurs within the brackets is so that the norm is homogenous; if $g = \alpha f$, and $\|f\|_{p,\infty} = A$, then

$$G(t) = F(t/|\alpha|) \leq \left(\frac{A|\alpha|}{t} \right)^p,$$

so $\|\alpha f\|_{p,\infty} = |\alpha| \|f\|_{p,\infty}$. The weak norms do not satisfy a triangle inequality, but they do satisfy a quasitriangle inequality. This can be proven quite simply from the property that if $f = f_1 + \dots + f_N$, and $\alpha_1, \dots, \alpha_N \in [0, 1]$ satisfy $\alpha_1 + \dots + \alpha_N = 1$, then

$$F(t) = F_1(\alpha_1 t) + \dots + F_N(\alpha_N t).$$

Thus if $f = g + h$, then

$$F(t) \leq G(t/2) + H(t/2) \leq \frac{\|g\|_{p,\infty}^p + \|h\|_{p,\infty}^p}{t^p} \lesssim_p \left(\frac{\|g\|_{p,\infty} + \|h\|_{p,\infty}}{t} \right)^p.$$

Thus $\|f + g\|_{p,\infty} \lesssim \|f\|_{p,\infty} + \|g\|_{p,\infty}$. We can measure the degree to which the weak L^p norm fails to be a norm by determining how much the triangle inequality fails for the sum of N functions, instead of just one function.

Theorem 15.11 (Stein-Weiss Inequality). *Let f_1, \dots, f_N be functions. If $p > 1$, then*

$$\|f_1 + \dots + f_N\|_{p,\infty} \lesssim_p \|f_1\|_{p,\infty} + \dots + \|f_N\|_{p,\infty}.$$

If $p = 1$, then

$$\|f_1 + \cdots + f_N\|_{1,\infty} \lesssim \log N [\|f_1\|_{1,\infty} + \cdots + \|f_N\|_{1,\infty}].$$

If $0 < p < 1$, then

$$\|f_1 + \cdots + f_N\|_{p,\infty} \lesssim_p \left(\|f_1\|_{p,\infty}^p + \cdots + \|f_N\|_{p,\infty}^{1/p} \right)^{1/p}$$

Proof. Begin with the case $p \geq 1$. Without loss of generality, assume $\|f_1\|_{p,\infty} + \cdots + \|f_N\|_{p,\infty} = 1$. Fix $t > 0$. For each $k \in [1, N]$, define

$$g_k(x) = \begin{cases} f_k(x) & : |f_k(x)| \geq t/2, \\ 0 & : \text{otherwise,} \end{cases}$$

and

$$h_k(x) = \begin{cases} f_k(x) & : |f_k(x)| \leq \|f_k\|_{p,\infty} \cdot (t/2), \\ 0 & : \text{otherwise.} \end{cases}$$

Also define $j_k = f_k - g_k - h_k$. Then write $f = f_1 + \cdots + f_N$, $g = g_1 + \cdots + g_N$, $h = h_1 + \cdots + h_N$, and $j = j_1 + \cdots + j_N$. Note that $\|h\|_\infty \leq t/2$, so

$$\{x : |f(x)| \geq t\} \subset \{x : |g(x)| \geq t/4\} \cup \{x : |j(x)| \geq t/4\}.$$

Each g_k is supported on a set of measure at most $\|f_k\|_{p,\infty}^p \cdot (2/t)^p$. We conclude that g is supported on a set of measure at most

$$(2/t)^p \sum_{k=1}^N \|f_k\|_{p,\infty}^p \leq (2/t)^p.$$

If $p > 1$, then the measure of $\{x : |j(x)| \geq t/4\}$ is bounded by

$$\begin{aligned}
\frac{4}{t} \int |j(x)| dx &\leq \frac{4}{t} \sum_{k=1}^N \int |j_k(x)| \\
&= \frac{4}{t} \sum_{k=1}^N \int_{\|f_k\|_{p,\infty}(t/2)}^{t/2} \frac{\|j_k\|_{p,\infty}^p}{s^p} ds \\
&= \frac{2^{p+1}}{p-1} \frac{1}{t^p} \sum_{k=1}^N \|j_k\|_{p,\infty}^p \left(\frac{1}{\|f_k\|_{p,\infty}^{p-1}} - 1 \right) \\
&\leq \frac{2^{p+1}}{p-1} \frac{1}{t^p} \sum_{k=1}^N \|f_k\|_{p,\infty}^p \left(\frac{1}{\|f_k\|_{p,\infty}^{p-1}} - 1 \right) \\
&\leq \frac{2^{p+1}}{p-1} \frac{1}{t^p}.
\end{aligned}$$

Thus in total, we conclude the measure of $\{x : |f(x)| \geq t\}$ is at most

$$\frac{2^p}{t^p} + \frac{2^{p+1}}{p-1} \frac{1}{t^p} \lesssim_p \frac{1}{t^p}.$$

If $p = 1$, then the measure of $\{x : |j(x)| \geq t/4\}$ is bounded

$$\begin{aligned}
(4/t) \int |j(x)| dx &\leq (4/t) \sum_{k=1}^N \int |j_k(x)| \\
&= (4/t) \sum_{k=1}^N \int_{\|f_k\|_{1,\infty}(t/2)}^{t/2} \frac{\|j_k\|_{1,\infty}}{s} ds \\
&= (4/t) \sum_{k=1}^N \|f_k\|_{1,\infty} \log(1/\|f_k\|_{1,\infty}).
\end{aligned}$$

Now the maximum of $x_1 \log(1/x_1) + \dots + x_N \log(1/x_N)$, subject to the constraint that $x_1 + \dots + x_N = 1$, is maximized by taking $x_k = 1/N$ for all N , which gives a maximal bound of $\log(N)$. In particular, we find that

$$(2/t) \sum_{k=1}^N \|f_k\|_{1,\infty} \log(1/\|f_k\|_{1,\infty}) \leq (2 \log N)/t.$$

Thus in total, we conclude the measure of $\{x : |f(x)| \geq t\}$ is at most

$$2(1 + \log N)/t \lesssim \log N/t.$$

If $p < 1$, we may assume without loss of generality that

$$\|f_1\|_{p,\infty}^p + \cdots + \|f_N\|_{p,\infty}^p = 1.$$

Then, we perform the same decomposition as before, with functions $\{g_k\}$, $\{h_k\}$, and $\{j_k\}$, defined the same as before, except that

$$h_k(x) = \begin{cases} f_k(x) & : |f_k(x)| \leq \|f_k\|_{p,\infty}^p \cdot (t/2), \\ 0 & : \text{otherwise.} \end{cases}$$

The function g_k has support at most $\|f_k\|_{p,\infty}^p \cdot (2/t)^p$, and thus g has total support

$$\sum \|f_k\|_{p,\infty}^p (2/t)^p = (2/t)^p.$$

The measure of $\{x : |j(x)| \geq t/4\}$ is bounded by

$$\begin{aligned} \frac{4}{t} \int |j(x)| \, dx &\leq \frac{4}{t} \sum_{k=1}^N \int_{\|f_k\|_{p,\infty}^p (t/2)}^{t/2} \frac{\|f_k\|_{p,\infty}^p}{s^p} \, ds \\ &\leq \frac{2^{p+1}}{t^p} \frac{1}{1-p} \sum_{k=1}^N \|f_k\|_{p,\infty}^{p+p(1-p)} \\ &= \frac{2^{p+1}}{t^p} \frac{1}{1-p} \max \|f_k\|_{p,\infty}^{p(1-p)} \lesssim_p \frac{1}{t^p}, \end{aligned}$$

Combining the two bounds gives that $\|f_1 + \cdots + f_N\|_{p,\infty} \lesssim_p 1$. \square

Remark. For $p = 1$, compare this *logarithmic* failure to be a norm with the *polynomial* failure to be a norm found in the norms $\|\cdot\|_p$, when $p < 1$, in Theorem 15.5.

For $p = 1$, the Stein-Weiss inequality is asymptotically tight in N .

Example. Let $X = \mathbf{R}$. For each k , let

$$f_k(x) = \frac{1}{|x - k|}.$$

Then $\|f_k\|_{1,\infty} \lesssim 1$ is bounded independantly of k . If $|x| \leq N$, there are integers $k_1, \dots, k_N > 0$ such that $|x - k_i| \leq 2i$, so

$$f(x) \geq \sum_{i=1}^N \frac{1}{|x - k_i|} \geq \sum_{i=1}^N \frac{1}{2i} \gtrsim \log(N).$$

Thus $\|f\|_{1,\infty} \gtrsim N \log N \gtrsim \log N \sum \|f_k\|_{1,\infty}$.

The weak L^p norms provide another monotone translation invariant norm, and it oftens comes up when finer tuning is needed in certain interpolation arguments, especially when dealing with maximal functions.

Example. If $f = H\mathbf{I}_E$, with $|E| = W$, then

$$F(t) = W \cdot \mathbf{I}_{[0,H]}.$$

Thus

$$\|f\|_{p,\infty} = \left(\sup_{0 \leq t < H} W t^p \right)^{1/p} = W^{1/p} H^p = \|f\|_p.$$

If $f = H_1\mathbf{I}_{E_1} + H_2\mathbf{I}_{E_2}$, with $|E_1| = W_1$ and $|E_2| = W_2$, with $H_1 \leq H_2$, then

$$F(t) = \begin{cases} W_1 + W_2 & : t < H_1, \\ W_2 & : t < H_2, \\ 0 & : \text{otherwise.} \end{cases}$$

Thus

$$\|f\|_{p,\infty} = \left(\max((W_1 + W_2)H_1^p, W_2H_2^p) \right)^{1/p} = \max((W_1 + W_2)^{1/p}H_1, W_2^{1/p}H_2).$$

Example. The function $f(x) = 1/|x|^s$ does not lie in any $L^p(\mathbf{R}^d)$, but lies in $L^{p,\infty}$ precisely when $p = d/s$, since

$$|\{1/|x|^{ps} > t\}| = \left| \left\{ |x| \leq \frac{1}{t^{1/ps}} \right\} \right| \propto_d \frac{1}{t^{d/ps}}.$$

Before we move on, we consider a form of duality for the weak L^p norm, at least when $p > 1$.

Theorem 15.12. *If $p > 1$, and X is σ -finite, then*

$$\|f\|_{p,\infty} \sim_p \sup_{|E|<\infty} \frac{1}{|E|^{1-1/p}} \int_E |f(x)| \, dx$$

Proof. Suppose $\|f\|_{p,\infty} < \infty$. If we write $f = \sum f_k$, where $f_k = \mathbf{I}_{F_k} f$, and $F_k = \{x : 2^{k-1} < |f(x)| \leq 2^k\}$, then $|F_k| \leq \|f\|_{p,\infty}^p 2^{-kp}$. Thus

$$\left| \int_E |f_k(x)| \, dx \right| \leq 2^k \|f\|_{p,\infty}^p 2^{-kp} = \|f\|_{p,\infty}^p 2^{k(1-p)}.$$

Fix some integer n . Then

$$\begin{aligned} \int_E |f(x)| \, dx &\leq \sum_{k=-\infty}^{n-1} \int_E |f_k(x)| \, dx + \sum_{k=n}^{\infty} \int_E |f_k(x)| \, dx \\ &\leq |E| 2^{n-1} + \|f\|_{p,\infty}^p \sum_{k=n}^{\infty} 2^{k(1-p)} \\ &\lesssim_p |E| 2^n + \|f\|_{p,\infty}^p 2^{-n(1-p)}. \end{aligned}$$

If we let $2^n \sim \|f\|_{p,\infty} |E|^{1/p}$, then we conclude

$$\int_E |f(x)| \, dx \lesssim_p |E|^{1-1/p} \|f\|_{p,\infty}.$$

Conversely, write

$$A = \sup_{|E|<\infty} \frac{1}{|E|^{1-1/p}} \int_E |f(x)| \, dx$$

If $G_t = \{x : |f(x)| \geq t\}$, then

$$|G_t| \leq \frac{1}{t} \int_{G_t} |f(x)| \, dx \leq \frac{A |G_t|^{1-1/p}}{t},$$

so

$$|G_t| \leq \frac{A^p}{t},$$

which gives $\|f\|_{p,\infty} \leq A$. □

For $p \leq 1$, the spaces $L^{p,\infty}(X)$ are not normable, as seen by the tightness of the Stein-Weiss inequality. Nonetheless, we still have a certain ‘duality’ property, that is often useful in the analysis of operators on these spaces. Most useful is its application when $p = 1$.

Theorem 15.13. *Let $0 < p < \infty$, and let $f \in L^{p,\infty}(X)$, and let $\alpha \in (0, 1)$. Then the following are equivalent:*

- $\|f\|_{p,\infty} \lesssim_{\alpha,p} A$.
- For any set $E \subset X$ with finite measure, there is $E' \subset E$ with $|E'| \geq \alpha|E|$ such that

$$\int_{E'} |f(x)| dx \lesssim_{\alpha,p} A |E'|^{1-1/p}.$$

Proof. By homogeneity, assume $\|f\|_{p,\infty} \leq 1$, so that if F is the distribution of f , $F(t) \leq 1/t^p$. If $|E| = (1 - \alpha)^{-1}/t_0^p$, and we set

$$E' = \{x : |f(x)| \leq t_0\},$$

then

$$|E'| \geq |E| - F(t_0) = \frac{(1 - \alpha)^{-1} - 1}{t_0^p} = \alpha|E|,$$

and

$$\int_{E'} |f(x)| \leq t_0 |E'| \lesssim_{\alpha} |E'|^{1-1/p}.$$

Conversely, suppose Property (2) holds. For each k , set

$$E_k = \{x : 2^k \leq |f(x)| < 2^{k+1}\}.$$

Then there exists E'_k with $|E'_k| \geq \alpha|E_k|$ and

$$\int_{E'_k} |f(x)| dx \leq |E'_k|^{1-1/p}$$

On the other hand,

$$\int_{E'_k} |f(x)| dx \geq 2^k |E'_k|.$$

Rearranging this equation gives $|E'_k| \leq 2^{-pk}$, and so $|E_k| \lesssim_{\alpha} 2^{-pk}$. But this means

$$F(2^N) = \sum_{k=N}^{\infty} |E_k| \lesssim_{\alpha,p} 2^{-Np},$$

and this implies $\|f\|_{p,\infty} \lesssim_{\alpha,p} 1$. □

15.4 Lorentz Spaces

Recall that we can write

$$\|f\|_p = \left(p \int_0^\infty F(t) t^p \frac{dt}{t} \right)^{1/p}.$$

Thus $F(t)t^p$ is integrable with respect to the Haar measure on \mathbf{R}^+ . But if we change the integrability condition to the condition that $F(t)t^p \in L^q(\mathbf{R}^+)$ for some $0 < q \leq \infty$, we obtain a different integrability condition, giving rise to a monotone, translation-invariant norm. This leads us to the definition of the *Lorentz norms*. For each $0 < p, q < \infty$, we define the Lorentz norm

$$\|f\|_{p,q} = p^{1/q} \|tF^{1/p}\|_{L^q(\mathbf{R}^+)}$$

The *Lorentz space* $L^{p,q}(X)$ as the space of functions f with $\|f\|_{p,q} < \infty$. We can define the norm in terms of f^* as well.

Lemma 15.14. *For any measurable $f : X \rightarrow \mathbf{R}$, $\|f(t)\|_{p,q} = \|s^{1/p} f^*(s)\|_{L^q(\mathbf{R}^+)}$.*

Proof. First, assume f^* has non-vanishing derivative on $(0, \infty)$, and that f is bounded, with finite support. An integration by parts gives

$$\|f\|_{p,q} = p^{1/q} \left(\int_0^\infty t^{q-1} F(t)^{q/p} dt \right)^{1/q} = \left(\int_0^\infty t^q F(t)^{q/p-1} (-F'(t)) dt \right)^{1/q}.$$

If we set $s = F(t)$, then $f^*(s) = t$, and $ds = F'(t)dt$, and so

$$\left(\int_0^\infty t^q F(t)^{q/p-1} F'(t) dt \right)^{1/q} = \left(\int_0^\infty f^*(s)^q s^{q/p-1} ds \right)^{1/q} = \|s^{1/p} f^*\|_{L^q(\mathbf{R}^+)}.$$

This gives the result in this case. The general result can then be obtained by applying the monotone convergence theorem to an arbitrary f^* with respect to a family of smooth functions. \square

The definition of the Lorentz space may seem confusing, but we really only require various special cases in most applications. Aside from the weak L^p norms $\|\cdot\|_{p,\infty}$ and the L^p norms $\|\cdot\|_p = \|\cdot\|_{p,p}$, the $L^{p,1}$ norms and $L^{p,2}$ norms also occur, the first, because of the connection with integrability, and the second because we may apply orthogonality techniques. As $q \rightarrow 0$, the norms $\|\cdot\|_{p,q}$ give stronger control over the function f .

Theorem 15.15. For $q < r$, $\|f\|_{p,r} \lesssim_{p,q,r} \|f\|_{p,q}$.

Proof. First we treat the case $r = \infty$. We have

$$\begin{aligned} s_0^{1/p} f^*(s_0) &= \left((p/q) \int_0^{s_0} [s^{1/p} f^*(s)]^q \frac{ds}{s} \right)^{1/q} \\ &\leq \left((p/q) \int_0^{s_0} [s^{1/p} f^*(s)]^q \frac{ds}{s} \right) \\ &\leq (p/q)^{1/q} \|f\|_{p,q}. \end{aligned}$$

When $r < \infty$, we can interpolate, calculating

$$\begin{aligned} \|f\|_{p,r} &= \left(\int_0^\infty [s^{1/p} f^*(s)]^r \frac{ds}{s} \right)^{1/r} \\ &\leq \|f\|_{p,\infty}^{1-q/r} \|f\|_{p,q}^{q/r} \leq (p/q)^{p(1/q-1/r)} \|f\|_{p,q}. \end{aligned} \quad \square$$

The fact that multiplying a function by a constant dilates the distribution implies that the Lorentz norm is homogeneous. We do not have a triangle inequality for the Lorentz norms, but we have a quasi triangle inequality.

Theorem 15.16. For each $p, q > 0$, $\|f_1 + f_2\|_{p,q} \lesssim_{p,q} \|f_1\|_p + \|f_2\|_q$.

Proof. We calculate that if $g = f_1 + f_2$,

$$\begin{aligned} \|g\|_{p,q} &= \left(q \int_0^\infty \left[t G(t)^{1/p} \right]^q \frac{dt}{t} \right)^{1/q} \\ &\leq \left(q \int_0^\infty \left[t (F_1(t/2) + F_2(t/2))^{1/p} \right]^q \frac{dt}{t} \right)^{1/q} \\ &\lesssim \left(q \int_0^\infty \left[t (F_1(t) + F_2(t))^{1/p} \right]^q \frac{dt}{t} \right)^{1/q} \\ &\lesssim_p \left(q \int_0^\infty t^q \left(F_1(t)^{q/p} + F_2(t)^{q/p} \right) \frac{dt}{t} \right)^{1/q} \\ &\lesssim_q \left(q \int_0^\infty t^q F_1(t)^{q/p} \frac{dt}{t} \right)^{1/q} + \left(q \int_0^\infty t^q F_2(t)^{q/p} \frac{dt}{t} \right)^{1/q} \\ &= \|f_1\|_{p,q} + \|f_2\|_{p,q}. \end{aligned} \quad \square$$

An important trick to utilizing Lorentz norms is by utilizing a dyadic layer cake decomposition. The dyadic layer cake decompositions enable us to understand a function by breaking it up into parts upon which we can control the height or width of a function. We say f is a *sub step function* with height H and width W if f is supported on a set E with $|E| \leq W$, and $|f(x)| \leq H$. A *quasi step function* with height H and width W if f is supported on a set E with $|E| \sim W$ and on E , $|f(x)| \sim H$.

Remark. It might seem that sub step functions of height H and width W can take on a great many different behaviours, rather than that of a step function with height H and width W . However, from the point of view of monotone, translation invariant norms, this isn't so. This is because using the binary expansion of real numbers, for every sub-step function f of height H and width W , we can find sets $\{E_k\}$ such that

$$f(x) = H \sum_{k=1}^{\infty} 2^{-k} \mathbf{I}_{E_k},$$

where $|E_k| = 1$. Thus bounds on step functions that are stable under addition tend to automatically imply bounds on substep functions.

We start by discussing the *vertical dyadic layer cake decomposition*. We define, for each $k \in \mathbf{Z}$,

$$f_k(x) = f(x) \mathbf{I}(2^{k-1} < |f(x)| \leq 2^k)$$

Then we set $f = \sum f_k$. Each f_k is a quasi step function with height 2^k and width $F(2^{k-1}) - F(2^k)$. We can also perform a *horizontal layer cake decomposition*. If we define $H_k = f^*(2^k)$, and set

$$f_k(x) = f(x) \mathbf{I}(H_{k-1} < |f(x)| \leq H_k),$$

then f_k is a substep function with height H_k and width 2^k . These decompositions are best visualized with respect to the representation f^* of f , in which case the decomposition occurs over particular intervals.

Theorem 15.17. *The following values A_1, \dots, A_4 are all comparable up to absolute constant depending only on p and q :*

1. $\|f\|_{p,q} \leq A_1.$

2. We can write $f = \sum_{k \in \mathbf{Z}} f_k$, where f_k is a quasi-step function with height 2^k and width W_k , and

$$\left(\sum_{k \in \mathbf{Z}} [2^k W_k^{1/p}]^q \right)^{1/q} \leq A_2.$$

3. We can write $f = \sum_{k \in \mathbf{Z}} f_k$, where f_k is a sub-step function with height 2^k and width W_k , and

$$\left(\sum_{k \in \mathbf{Z}} [2^k W_k^{1/p}]^q \right)^{1/q} \leq A_3.$$

4. We can write $f(x) = \sum_{k \in \mathbf{Z}} f_k$, where f_k is a sub-step function with width 2^k and height H_k , where $\{H_k\}$ is decreasing in k , and

$$\left(\sum_{k \in \mathbf{Z}} [H_k 2^{k/p}]^q \right)^{1/q} \leq A_4.$$

Proof. It is obvious that we can always select $A_3 \leq A_2$. Next, we bound A_2 in terms of A_1 by performing a vertical layer cake decomposition on f . If we write $f = \sum_{k \in \mathbf{Z}} f_k$, then f_k is supported on a set with measure $W_k = F(2^{k-1}) - F(2^k) \leq F(2^{k-1})$, and so

$$\begin{aligned} \sum_{k \in \mathbf{Z}} [2^k W_k^{1/p}]^q &\leq \sum_{k \in \mathbf{Z}} [2^k F(2^{k-1})^{1/p}]^q \\ &\lesssim_q \sum_{k \in \mathbf{Z}} [2^{k-1} F(2^k)^{1/p}]^q \\ &\lesssim \sum_{k \in \mathbf{Z}} \int_{2^{k-1}}^{2^k} [t F(t)^{1/p}]^q \frac{dt}{t} \lesssim_q \|f\|_{p,q}^q \leq A_1^q. \end{aligned}$$

Thus $A_2 \lesssim_q A_1$. Next, we bound A_4 in terms of A_1 . Perform a horizontal layer cake decomposition, writing $f = \sum f_k$, where f_k is supported on a set with measure $W_k \leq 2^k$, and $H_{k+1} \leq |f_k(x)| \leq H_k$. Then a telescoping sum

shows

$$\begin{aligned}
H_k 2^{k/p} &= \left(\sum_{m=0}^{\infty} (H_{k+m}^q - H_{k+m+1}^q) 2^{kq/p} \right)^{1/q} \\
&\lesssim_q \left(\sum_{m=0}^{\infty} \int_{H_{k+m+1}}^{H_{k+m}} [t 2^{k/p}]^q \frac{dt}{t} \right)^{1/q} \\
&\leq \left(\sum_{m=0}^{\infty} \int_{H_{k+m+1}}^{H_{k+m}} [t F(t)^{1/p}]^q \frac{dt}{t} \right)^{1/q}
\end{aligned}$$

Thus

$$\left(\sum_{k \in \mathbf{Z}} [H_k 2^{k/p}]^q \right)^{1/q} \leq \left(\int_0^\infty [t F(t)^{1/p}]^q \frac{dt}{t} \right)^{1/q} \lesssim_q A_1.$$

Thus $A_4 \lesssim_q A_1$. It remains to bound A_1 by A_4 and A_3 . Given A_3 , we can write $|f(x)| \leq \sum 2^k \mathbf{I}_{E_k}$, where $|E_k| \leq W_k$. We then find

$$F(2^k) \leq \sum_{m=1}^{\infty} W_{k+m}.$$

Thus

$$\int_{2^{k-1}}^{2^k} [t F(t)^{1/p}]^q \frac{dt}{t} \lesssim \left[2^k \left(\sum_{m=0}^{\infty} W_k \right)^{1/p} \right]^q.$$

Thus if $q \leq p$,

$$\begin{aligned}
\|f\|_{p,q} &\lesssim_q \left(\sum_{k \in \mathbf{Z}} \left[2^k \left(\sum_{m=0}^{\infty} W_{k+m} \right)^{1/p} \right]^q \right)^{1/q} \\
&\leq \left(\sum_{k \in \mathbf{Z}} \sum_{m=0}^{\infty} [2^k W_{k+m}^{1/p}]^q \right)^{1/q} \\
&\leq \left(\sum_{m=0}^{\infty} 2^{-qm} \sum_{k \in \mathbf{Z}} [2^{k+m} W_{k+m}^{1/p}]^q \right)^{1/q} \\
&\leq \left(A_3^q \sum_{m=0}^{\infty} 2^{-mq} \right)^{1/q} \lesssim_q A_3.
\end{aligned}$$

If $q \geq p$, we can employ the triangle inequality for $l^{q/p}$ to write

$$\begin{aligned} \|f\|_{p,q} &\lesssim_q \left(\sum_{k \in \mathbf{Z}} \left[2^k \left(\sum_{m=0}^{\infty} W_{k+m} \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq \left(\sum_{m=0}^{\infty} \left(\sum_{k \in \mathbf{Z}} 2^{kq} W_{k+m}^{q/p} \right)^{p/q} \right)^{1/p} \\ &\leq \left(A_3^p \sum_{m=0}^{\infty} 2^{-mq} \right)^{1/p} \lesssim_{p,q} A_3. \end{aligned}$$

The bound of A_1 in terms of A_4 involves the same ‘shifting’ technique, and is left to the reader. \square

Remark. Heuristically, the theorem above says that if $f = \sum_{k \in \mathbf{Z}} f_k$, where f_k is a quasi-step function with width H_k and width W_k , and if either $\{H_k\}$ and $\{W_k\}$ grow faster than powers of two, then

$$\|f\|_{p,q} \sim_{p,q} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p}]^q \right)^{1/q}.$$

Thus the $L^{p,q}$ norm has little interaction between elements of the sum when the sum occurs over dyadically different heights or width. This is one reason why we view the q parameter as a ‘logarithmic’ correction of the L^p norm. In particular, if we can write $f = f_1 + \cdots + f_N$, and $q_1 < q_2$, then the last equation, combined with a l^{q_1} to l^{q_2} norm bound, gives

$$\left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p}]^{q_1} \right)^{1/q_1} \leq N^{1/q_1 - 1/q_2} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p}]^{q_2} \right)^{1/q_2}$$

This implies

$$\|f\|_{p,q_2} \lesssim_{p,q_1,q_2} \|f\|_{p,q_1} \lesssim_{p,q_1,q_2} N^{1/q_1 - 1/q_2} \|f\|_{p,q_2}.$$

In particular, this occurs if there exists a constant C such that $C \leq |f(x)| \leq C \cdot 2^N$ for all x . On the other hand, if we vary the p parameter, we find that

for $p_1 < p_2$,

$$\begin{aligned} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_1}]^q \right)^{1/q} &\leq \max(W_k)^{1/p_1 - 1/p_2} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^q \right)^{1/q}, \\ \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^q \right)^{1/q} &\leq \left(\frac{1}{\min(W_k)} \right)^{1/p_1 - 1/p_2} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^q \right)^{1/q}. \end{aligned}$$

which gives

$$\min(W_k)^{1/p_1 - 1/p_2} \|f\|_{p_2, q} \lesssim_{p_1, p_2, q} \|f\|_{p_1, q} \lesssim_{p_1, p_2, q} \max(W_k)^{1/p_1 - 1/p_2} \|f\|_{p_2, q}.$$

Both of these inequalities can be tight. Because of the dyadic decomposition of f , we find $\max(W_k) \geq 2^N \min(W_k)$, so these two norms can differ by at least $2^{N(1/p_1 - 1/p_2)}$, and at *most* if the f_k occur over consecutive dyadic values, which is *exponential* in N . Conversely, if the heights change dyadically, we find that

$$\begin{aligned} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^q \right)^{1/q} &\leq \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^{qp_2/p_1} \right)^{(p_1/p_2)/q} \\ &\leq \max(H_k)^{1 - p_1/p_2} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_1}]^q \right)^{(p_1/p_2)/q} \\ \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_1}]^q \right)^{1/q} &\lesssim \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_1}]^{qp_1/p_2} \right)^{(p_2/p_1)/q} \\ &\leq \left(\frac{1}{\min(H_k)} \right)^{p_2/p_1 - 1} \left(\sum_{k \in \mathbf{Z}} [H_k W_k^{1/p_2}]^q \right)^{(p_2/p_1)/q} \end{aligned}$$

where \lesssim denotes a factor ignoring polynomial powers of N occurring from the estimate. Thus

$$\min(H_k)^{p_2 - p_1} \|f\|_{p_1, q}^{p_1} \lesssim_{p_1, p_2, q} \|f\|_{p_2, q}^{p_2} \lesssim_{p_1, p_2, q} \max(H_k)^{p_2 - p_1} \|f\|_{p_1, q}^{p_1}$$

again, these inequalities can be both tight, and $\max(H_k) \geq 2^N \min(H_k)$, with equality if the quasi step functions from which f is composed occur consecutively dyadically.

Example. Consider the function $f(x) = |x|^{-s}$. For each k , let

$$E_k = \{x : 2^{-(k+1)/s} \leq |x| < 2^{-k/s}\}$$

and define $f_k = f \mathbf{I}_{E_k}$. Then f_k is a quasi-step function with height 2^k , and width $1/2^{dk/s}$. We conclude that if $p = d/s$, and $q < \infty$,

$$\|f\|_{p,q} \sim_{p,q,d} \left(\sum_{k=-\infty}^{\infty} 2^{qk(1-d/ps)} \right)^{1/q} = \infty.$$

Thus the function f lies exclusively in $L^{p,\infty}(\mathbf{R}^d)$.

A simple consequence of the layer cake decomposition is Hölder's inequality for Lorentz spaces.

Theorem 15.18. If $0 < p_1, p_2, p < \infty$ and $0 < q_1, q_2, q < \infty$ with

$$1/p = 1/p_1 + 1/p_2 \quad \text{and} \quad 1/q \geq 1/q_1 + 1/q_2,$$

then

$$\|fg\|_{p,q} \lesssim_{p_1,p_2,q_1,q_2} \|f\|_{p_1,q_1} \|g\|_{p_2,q_2}.$$

Proof. Without loss of generality, assume $\|f\|_{p_1,q_1} = \|g\|_{p_2,q_2} = 1$ and that $1/q = 1/q_1 + 1/q_2$. Perform horizontal layer cake decompositions of f and g , writing $|f| \leq \sum_{k \in \mathbf{Z}} H_k \mathbf{I}_{E_k}$ and $|g| \leq \sum_{k \in \mathbf{Z}} H'_k \mathbf{I}_{F_k}$, where $|E_k|, |F_k| \leq 2^k$. Then

$$|fg| \leq \sum_{k,k' \in \mathbf{Z}} H_k H'_k \mathbf{I}_{E_k \cap F_{k'}}$$

For each fixed k , $|E_{k+m} \cap F_m| \leq 2^m$, and so

$$\begin{aligned} \left\| \sum_{m \in \mathbf{Z}} H_{k+m} H'_m \mathbf{I}_{E_{k+m} \cap F_m} \right\|_{p,q} &\lesssim_{p,q} \left(\sum_{m \in \mathbf{Z}} [H_{k+m} H'_m 2^{m/p}]^q \right)^{1/q} \\ &= \left(\sum_{m \in \mathbf{Z}} \left[(H_{k+m} 2^{m/p_1}) (H'_m 2^{m/p_2}) \right]^q \right)^{1/q} \\ &\leq \left(\sum_{m \in \mathbf{Z}} [H_{k+m} 2^{m/p_1}]^{q_1} \right)^{1/q_1} \left(\sum_{m \in \mathbf{Z}} [H'_m 2^{m/p_2}]^{q_2} \right)^{1/q_2} \\ &\lesssim_{p,q,p_1,q_1,p_2,q_2} 2^{-k/p_1} \end{aligned}$$

Summing over $k > 0$ gives that

$$\left\| \sum_{k \geq 0} \sum_{m \in \mathbf{Z}} H_{k+m} H'_m \mathbf{I}_{E_{k+m} \cap F_m} \right\| \lesssim_{p,q,p_1,q_1,p_2,q_2} 1$$

By the quasitriangle inequality, it now suffices to obtain a bound

$$\left\| \sum_{k < 0} \sum_{m \in \mathbf{Z}} H_{k+m} H'_m \mathbf{I}_{E_{k+m} \cap F_m} \right\|_{p,q}.$$

This is done similarly, but using the bound $|E_{k+m} \cap F_m| \leq 2^{k+m}$ instead of the other bound. \square

Corollary 15.19. *If $p > 1$ and $q > 0$, $L^{p,q}(X) \subset L^1_{loc}(X)$.*

Proof. Let E have finite measure and let $f \in L^{p,q}(X)$. Then the Hölder's inequality for Lorentz spaces shows

$$\|f\|_{L^1(E)} = \|\mathbf{I}_E f\|_{L^1(X)} \lesssim_{p,q} |E|^{1-1/p} \|f\|_{p,q} < \infty. \quad \square$$

A consequence of Hölder's inequality is a duality of the $L^{p,q}$ norms. If $1 < p < \infty$, and $1 < q < \infty$, then $L^{p,q}(X)^* = L^{p',q'}(X)$. When $q = 1$ or $q = \infty$, things are more complex, but the following theorem often suffices. When $p = 1$, things get more tricky, so we leave this case out.

Theorem 15.20. *Let $1 < p < \infty$ and $1 \leq q < \infty$. Then if $f \in L^{p,q}(X)$,*

$$\|f\|_{p,q} \sim \sup \left\{ \int f g : \|g\|_{p',q'} \leq 1 \right\}.$$

Proof. Without loss of generality, we may assume $\|f\|_{p,q} = 1$. We may perform a vertical layer cake decomposition, writing $f = \sum_{k \in \mathbf{Z}} f_k$, where $2^{k-1} \leq |f_k(x)| \leq 2^k$, is supported on a set with width W_k , and

$$\left((2^k W_k^{1/p})^q \right) \sim_{p,q} 1.$$

Define $a_k = 2^k W_k^{1/p}$, and set $g = \sum_{k \in \mathbf{Z}} g_k$, where $g_k(x) = a_k^{q-p} \operatorname{sgn}(f_k(x)) |f_k(x)|^{p-1}$. Then

$$\begin{aligned} \int f(x) g(x) &= \sum_{k \in \mathbf{Z}} \int f_k(x) g_k(x) = \sum_{k \in \mathbf{Z}} a_k^{q-p} \int |f_k(x)|^p \\ &\gtrsim_p \sum_{k \in \mathbf{Z}} a_k^{q-p} W_k 2^{kp} = \sum_{k \in \mathbf{Z}} a_k^q \gtrsim_{p,q} 1. \end{aligned}$$

We therefore need to show that $\|g\|_{p',q'} \lesssim 1$. We note $|g_k(x)| \lesssim a_k^{q-p} 2^{kp}$, and has width W_k . This gives a decomposition of g , but neither the height nor the widths necessarily in powers of two. Still, we can fix this since the heights increase exponentially; define

$$H_k = \sup_{l \geq 0} a_{k-l}^{q-p} 2^{kp} 2^{-lp/2}.$$

Then $|g_k(x)| \lesssim_{p,q} H_k$, and $H_{k+1} \geq 2^{p/2} H_k$. In particular, if we pick m such that $2^{mp/2} \geq 1$, then for any $l \leq m$, the sequence H_{km+l} , as k ranges over values, increases dyadically, and so by the quasitriangle inequality for the $L^{p',q'}$ norm, and then the triangle inequality in l^q , we find

$$\begin{aligned} \|g\|_{p',q'} &\lesssim_{m,p,q} \left(\sum [H_k W_k^{1/p'}]^{q'} \right)^{1/q'} \\ &\lesssim \left(\sum_{k \in \mathbf{Z}} \left[\left(\sup_{l \geq 0} a_{k-l}^{q-p} 2^{kp} 2^{-lp/2} \right) (a_k 2^{-k})^{p-1} \right]^{q'} \right)^{1/q'} \\ &\lesssim_p \left(\sum_{k \in \mathbf{Z}} \left[a_k^{p-1} \sum_{l=0}^{\infty} a_{k-l}^{q-p} 2^{-lp/2} \right]^{q'} \right)^{1/q'} \\ &\lesssim \sum_{l=0}^{\infty} 2^{-lp/2} \left(\sum_{k \in \mathbf{Z}} [a_k^{p-1} a_{k-l}^{q-p}]^{q'} \right)^{1/q'}. \end{aligned}$$

Applying Hölder's inequality shows

$$\begin{aligned} \left(\sum_{k \in \mathbf{Z}} [a_k^{p-1} a_{k-l}^{q-p}]^{q'} \right)^{1/q'} &\leq \left(\sum_{k \in \mathbf{Z}} a_k^q \right)^{(p-1)/q} \left(\sum_{k \in \mathbf{Z}} a_{k-l}^q \right)^{(q-p)/q} \\ &\lesssim_{p,q} \|f\|_{p,q}^{q-1} \lesssim_{p,q} 1. \end{aligned} \quad \square$$

Remark. This technique shows that if $f = \sum f_k$, where f_k is a quasi-step function with measure W_k and height 2^{ck} , then we can find m such that $cm > 1$, and then consider the m functions f^1, \dots, f^m , where $f_i = \sum f_{km+i}$. Then the functions f_{km+i} have heights which are separated by powers of

two, and so the quasi-triangle inequality implies

$$\begin{aligned}\|f\|_{p,q} &\lesssim_m \sum_{i=1}^m \|f^i\|_{p,q} \\ &\lesssim_{p,q} \sum_{i=1}^m \left(\sum \left[H_{km+i} W_{km+i}^{1/p} \right]^q \right)^{1/q} \\ &\lesssim_m \left(\sum \left[H_k W_k^{1/p} \right]^q \right)^{1/q}\end{aligned}$$

On the other hand,

$$\begin{aligned}\|f\|_{p,q} &\gtrsim \max_{1 \leq i \leq m} \|f^i\|_{p,q} \\ &\sim \max_{1 \leq i \leq m} \left(\sum \left[H_{km+i} W_{km+i}^{1/p} \right]^q \right)^{1/q} \\ &\gtrsim_m \left(\sum \left[H_k W_k^{1/p} \right]^q \right)^{1/q}.\end{aligned}$$

Thus the dyadic layer cake decomposition still works in this setting.

We remark that if $1 < p < \infty$ and $1 \leq q \leq \infty$, then for each $f \in L^{p,q}$, the value

$$\sup \left\{ \int f g : \|g\|_{p',q'} \leq 1 \right\}$$

gives a norm on $L^{p,q}(X)$ which is comparable with the $L^{p,q}$ norm. In particular, this implies that for $p > 1$ and $q \geq 1$,

$$\|f_1 + \cdots + f_N\|_{p,q} \lesssim_{p,q} \|f_1\|_{p,q} + \cdots + \|f_N\|_{p,q},$$

so that the triangle inequality has constants independent of N . We can also use a layer cake decomposition to get a version of the Stein-Weiss inequality for Lorentz norms.

Theorem 15.21. *For each $1 < q < \infty$, there is $\alpha(q) > 0$ such that for any functions f_1, \dots, f_N ,*

$$\|f_1 + \cdots + f_N\|_{1,q} \lesssim (\log N)^{\alpha(q)} (\|f_1\|_{1,q} + \cdots + \|f_N\|_{1,q}).$$

Proof. For values A and B in this argument, we write $A \lesssim B$ if there exists α such that $A \lesssim (\log N)^\alpha B$. Given f_1, \dots, f_N , write $f_i = \sum_{j=-\infty}^{\infty} f_{ij}$, where f_{ij}

has width W_{ij} and height 2^j . If we assume, without loss of generality, that $\|f_1\|_{1,q} + \dots + \|f_N\|_{1,q} = 1$, then

$$\sum_{i=1}^N \left(\sum_{j=-\infty}^{\infty} (2^j W_{ij})^q \right)^{1/q} \lesssim_q 1$$

Thus we want to show $\|f_1 + \dots + f_N\|_{1,q} \lesssim_q 1$. Our first goal is to upper bound the measure of the set

$$E = \{x : 2^{k-1} < |f_1(x) + \dots + f_N(x)| \leq 2^k\}$$

The measure of the set E is upper bounded by the measure of the set

$$E' = \left\{ x : 2^{k-2} < \left| \sum_{j=k-\lg(N)}^k f_{1j}(x) + \dots + f_{Nj}(x) \right| \leq 2^{k+1} \right\}$$

Applying the usual Stein-Weiss inequality, we have

$$\left\| \sum_{i=1}^N \sum_{j=k-\lg N}^k f_{ij} \right\|_{1,\infty} \lesssim \sum_{i=1}^N \sum_{j=k-\lg N}^k \|f_{ij}\|_{1,\infty} \lesssim \sum_{i=1}^N \sum_{j=k-\lg N}^k \|f_{ij}\|_{1,\infty} \lesssim_q \sum_{i=1}^N \sum_{j=k-\lg N}^k W_{ij} 2^j$$

Thus we conclude

$$|E'| \lesssim_q 2^{-k} \sum_{i=1}^N \sum_{j=k-\lg N}^k W_{ij} 2^j$$

This implies that

$$\|f_1 + \dots + f_N\|_{1,q} \lesssim_q \left(\sum_{k=-\infty}^{\infty} \left(\sum_{i=1}^N \sum_{j=k-\lg N}^k W_{ij} 2^j \right)^q \right)^{1/q}.$$

Applying Minkowski's inequality, we conclude

$$\begin{aligned}
\left(\sum_{k=-\infty}^{\infty} \left(\sum_{i=1}^N \sum_{j=k-\lg N}^k W_{ij} 2^j \right)^q \right)^{1/q} &\lesssim \sum_{i=1}^N \left(\sum_{k=-\infty}^{\infty} \left(\sum_{j=k-\lg N}^k W_{ij} 2^j \right)^q \right)^{1/q} \\
&\lesssim \sum_{i=1}^N \left(\sum_{k=-\infty}^{\infty} \sum_{j=k-\lg N}^k W_{ij}^q 2^{qj} \right)^{1/q} \\
&\lesssim \sum_{i=1}^N \left(\sum_{j=-\infty}^{\infty} W_{ij}^q 2^{qj} \right)^{1/q} \lesssim 1. \quad \square
\end{aligned}$$

Here is an interesting weighted inequality whose proof utilizes the layer cake decomposition. I first encountered this inequality in Heo, Nazarov, and Seeger's paper *Radial Fourier Multipliers in High Dimensions*.

Theorem 15.22. *Suppose that $\{f_n\}$ is a family of functions, fix $p_0 < p_\theta < p_1$. Then*

$$\left\| \sum_n f_n \right\|_{L^p(X)} \lesssim_{p_0, p, p_1} \left(\sum_n \max_i \left\{ 2^{n(p-p_i)} \|f_n\|_{p_i, \infty}^{p_i} \right\} \right)^{1/p}$$

In particular, if $\|f_n\|_{p_i, \infty} \leq C 2^n W_n^{1/p_i}$ for each n , then

$$\left\| \sum_n f_n \right\|_{L^p(X)} \lesssim_{p_0, p, p_1} C \left(\sum_n 2^{pn} W_n \right)^{1/p}.$$

This might hold, for instance, if f_n is a sub step function with height 2^n and width W_n for each n .

Proof. Define $f_{nm} = f_n \cdot \mathbf{I}(2^{n+m} \leq |f_n| < 2^{n+m+1})$. Then $f_n = \sum_m f_{nm}$. For each fixed m , define $\tilde{f}_m = \sum_n f_{nm}$. Since $f_{nm} \approx 2^{n+m}$, \tilde{f}_m is defined by a sum over different dyadic scales, and so we have a pointwise bound

$$\left| \sum_n f_{nm} \right| \sim \max\{2^m : f_{nm} \neq 0\} \sim_p \left(\sum_n |f_{nm}|^p \right)^{1/p}.$$

Thus we find

$$\|\tilde{f}_m\|_{L^p(X)} \lesssim_p \left\| \left(\sum_n |f_{nm}|^p \right)^{1/p} \right\|_{L^p(X)} = \left(\sum_n \|f_{nm}\|_{L^p(X)}^p \right)^{1/p}.$$

Chebyshev's inequality implies that

$$\begin{aligned} \|f_{nm}\|_{L^p(X)}^p &\leq 2^{(n+m)p} \cdot \min_i \{ 2^{-(n+m)p_i} \|f_n\|_{L^{p_i,\infty}(X)}^{p_i} \} \\ &\lesssim \min_i \{ 2^{(n+m)(p-p_i)} \|f_n\|_{L^{p_i,\infty}(X)}^{p_i} \}. \end{aligned}$$

But this means that if $m \geq 0$,

$$\|\tilde{f}_m\|_{L^p(X)} \lesssim_p 2^{-|m|(p_1/p-1)} \left(\sum_n 2^{-n(p_1-p)} \|f_n\|_{L^{p_1,\infty}(X)}^{p_1} \right)^{1/p}$$

and for $m \leq 0$,

$$\|\tilde{f}_m\|_{L^p(X)} \lesssim_p 2^{-|m|(1-p_0/p)} \left(\sum_n 2^{n(p-p_0)} \|f_n\|_{L^{p_0,\infty}(X)}^{p_0} \right)^{1/p}$$

Applying the triangle inequality to $\|\sum f_n\|_{L^p(X)} = \|\sum \tilde{f}_m\|_{L^p(X)}$ and summing over m completes the proof. \square

15.5 Mixed Norm Spaces

Given two measure spaces X and Y , we can form the product measure space $X \times Y$. If we have a norm space V of functions on X , with norm $\|\cdot\|_V$ and a norm space W of functions on Y , with norm $\|\cdot\|_W$, we can consider a 'product norm'; for each function f on $X \times Y$, we can consider the function $y \mapsto \|f(\cdot, y)\|_V$, and take the norm of this function over Y , i.e. $\|\|f(\cdot, y)\|_V\|_W$. The most important case of this process is where we fix $0 < p, q \leq \infty$, and consider

$$\|f\|_{L^p(X)L^q(Y)} = \left(\int \left(\int |f(x, y)|^q dy \right)^{p/q} dx \right)^{1/p}.$$

Similarly, we can define $\|f\|_{L^q(Y)L^p(X)}$. The notation is justified by the fact that the first case is the L^p norm of the function f , viewed as a map from

X to $L^q(Y)$, and the latter the L^q norm of f , viewed as a map from Y to $L^p(X)$. We have a duality theory here; for each $1 \leq p, q \leq \infty$ and any f with $\|f\|_{L^p(X)L^q(Y)} < \infty$, the standard L^p and L^q duality gives

$$\|f\|_{L^p(X)L^q(Y)} = \sup \left\{ \int_{X \times Y} f(x, y) h(x, y) \, dx \, dy : \|h\|_{L^{p^*}(X)L^{q^*}(Y)} \leq 1 \right\}.$$

It is often important to interchange norms, and we find the biggest quantity obtained by interchanging norms is always obtained with the largest exponents on the inside, i.e. in the notation we use, the exponents are monotonically increasing from left to right.

Theorem 15.23. *If $p \geq q$, $\|f\|_{L^p(X)L^q(Y)} \leq \|f\|_{L^q(Y)L^p(X)}$.*

Proof. If $p = q$, then the Fubini-Tonelli theorem implies that

$$\|f\|_{L^p(X)L^q(Y)} = \|f\|_{L^q(Y)L^p(X)}.$$

If $q = 1$, then this result is precisely the Minkowski inequality. But not complex interpolation justifies the result in general. More precisely, a variation of the proof of Riesz-Thorin using the duality established above gives the result. \square

Let us consider two special cases. Firstly, bounds for the pointwise maxima of functions dominate the maximum L^p norms

$$\sup_n \|f_n\|_{L^p(X)} \leq \left\| \sup_n f_n \right\|_{L^p(X)}.$$

Another special case is the triangle inequality

$$\left\| \sum_n f_n \right\|_{L^p(X)} \leq \sum_n \|f_n\|_{L^p(X)}$$

for $p \geq 1$.

The next result shows that if $p < q$ and $\|f\|_{L^p(X)L^q(Y)} = \|f\|_{L^q(Y)L^p(X)}$, then $|f|$ is a tensor product. Thus switching mixed norms is likely only efficient if we think the functions we are working with are close to tensor products.

Theorem 15.24. Suppose $p > q$, f is a function on $X \times Y$, and

$$\|f\|_{L^p(X)L^q(Y)} = \|f\|_{L^q(Y)L^p(X)} < \infty.$$

Then there exists $f_1(x)$ and $f_2(y)$ such that for any $x \in X$ and $y \in Y$, $|f(x, y)| = |f_1(x)||f_2(y)|$.

Proof. Expanding this equation out, we conclude

$$\left(\int_Y \left(\int_X |f(x, y)|^p dx \right)^{q/p} dy \right)^{1/q} = \left(\int_X \left(\int_Y |f(x, y)|^q dy \right)^{p/q} dx \right)^{1/p}.$$

Setting $g(x, y) = |f(x, y)|^p$, we see that Minkowski's integral inequality is tight for g , i.e.

$$\left(\int_Y \left(\int_X |g(x, y)| dx \right)^{q/p} dy \right)^{p/q} = \left(\int_X \left(\int_Y |g(x, y)|^{q/p} dy \right)^{p/q} dx \right).$$

Thus it suffices to show that to show the theorem for $p = 1$ and $q > 1$. Recall the standard proof of Minkowski's inequality, i.e. that by Hölder's inequality

$$\begin{aligned} & \int_Y \left(\int_X |f(x, y)| dx \right)^p dy \\ &= \int_X \left[\int_Y |f(x_1, y)| \left(\int_X |f(x_2, y)| dx_2 \right)^{p-1} dy \right] dx_1 \\ &\leq \int_X \left[\left(\int_Y |f(x_1, y)|^p dy \right)^{1/p} \left(\int_Y \left(\int_X |f(x_2, y)| dx_2 \right)^{(p-1)p^*} dy \right)^{1/p^*} \right] dx_1 \\ &= \left[\int_X \left(\int_Y |f(x_1, y)|^p dy \right)^{1/p} dx_1 \right] \left[\int_Y \left(\int_X |f(x_2, y)| dx_2 \right)^p \right]^{1/p^*}. \end{aligned}$$

and rearranging gives Minkowski's inequality. If this inequality is tight, then our application of Hölder's inequality is tight for almost every $x_1 \in X$. Since $\int |f(x_2, y)| dx_2 \neq 0$ for all y unless $f = 0$, it follows that there exists $\lambda(x_1)$ for almost every $x_1 \in X$ such that for almost every $y \in Y$,

$$|f(x_1, y)|^p = |\lambda(x_1)| \left(\int_X |f(x_2, y)| dx_2 \right)^{p^*(p-1)} = |\lambda(x_1)| \left(\int_X |f(x_2, y)| dx_2 \right)^p.$$

Setting $f_1(x) = |\lambda(x)|^{1/p}$ and $f_2(y) = \int_X |f(x,y)| dx$ thus completes the proof. \square

15.6 Orlicz Spaces

To develop the class of Orlicz spaces, we note that if $\|f\|_p \leq 1$, and we set $\Phi(t) = t^p$, then

$$\int \Phi(|f(x)|) dx = 1.$$

More generally, given any function $\Phi : [0, \infty) \rightarrow [0, \infty)$, we might ask if we can define a norm $\|\cdot\|_\Phi$ such that if $\|f\|_\Phi \leq 1$, then

$$\int \Phi(|f(x)|) dx = 1.$$

Since a norm would be homogenous, this would imply that if $\|f\|_\Phi \leq A$, then

$$\int \Phi\left(\frac{|f(x)|}{A}\right) dx \leq 1.$$

If we want these norms to be monotone, we might ask that if $A < B$, then

$$\int \Phi\left(\frac{|f(x)|}{B}\right) dx \leq \int \Phi\left(\frac{|f(x)|}{A}\right) dx,$$

and the standard way to ensure this is to ask the Φ is an increasing function. To deal with the property that $\|0\| = 0$, we set $\Phi(0) = 0$. In order for $\|\cdot\|_\Phi$ to be a norm, the set of functions $\{f : \|f\|_\Phi \leq 1\}$ needs to be convex, and the standard way to obtain this is to assume that Φ is convex.

In short, we consider an increasing, convex function Φ with $\Phi(0) = 0$. We then define

$$\|f\|_\Phi = \inf \left\{ A > 0 : \int \Phi\left(\frac{|f(x)|}{A}\right) dx \leq 1 \right\}.$$

This function is a norm on the space of all f with $\|f\|_\Phi < \infty$. It is easy to verify that $\|f\|_\Phi = 0$ if and only if $f = 0$ almost everywhere, and that $\|\alpha f\|_\Phi = |\alpha| \|f\|_\Phi$. To justify the triangle inequality, we note that if

$$\int \Phi\left(\frac{|f(x)|}{A}\right) dx \leq 1 \quad \text{and} \quad \int \Phi\left(\frac{|f(x)|}{B}\right) dx \leq 1,$$

then applying convexity gives

$$\begin{aligned} \int \Phi \left(\frac{|f(x) + g(x)|}{A + B} \right) &\leq \int \Phi \left(\frac{|f(x)| + |g(x)|}{A + B} \right) \\ &\leq \int \left(\frac{A}{A + B} \right) \Phi \left(\frac{|f(x)|}{A} \right) + \left(\frac{B}{A + B} \right) \Phi \left(\frac{|g(x)|}{B} \right) \leq 1. \end{aligned}$$

Thus we obtain the triangle inequality.

The spaces $L^p(X)$ for $p \in [1, \infty)$ are Orlicz spaces with $\Phi(t) = t^p$. The space $L^\infty(X)$ is not really an Orlicz space, but it can be considered as the Orlicz function with respect to the ‘convex’ function

$$\Phi(t) = \begin{cases} \infty & t > 1, \\ t & t \leq 1. \end{cases}$$

More interesting examples of Orlicz spaces include

- $L \log L$, given by the Orlicz norm induced by $\Phi(t) = t \log(2 + t)$.
- e^L , defined with respect to $\Phi(t) = e^t - 1$.
- e^{L^2} , defined with respect to $\Phi(t) = e^{t^2} - 1$.

One should not think too hard about the constants in the functions defined above, which are included to make $\Phi(0) = 0$. When we are dealing with a finite measure space (often the case, since these norms often occur in probability theory), they are irrelevant.

Lemma 15.25. *If $\Phi(x) \lesssim \Psi(x)$ for all x , then $\|f\|_{\Phi(L)} \lesssim \|f\|_{\Psi(L)}$. If X is finite, and $\Phi(x) \lesssim \Psi(x)$ for sufficiently large x , then $\|f\|_{\Phi(L)} \lesssim \|f\|_{\Psi(L)}$.*

Proof. The first proposition is easy, and we now deal with the finite case. We note that the condition implies that for each $\varepsilon > 0$, there exists C_ε such that $\Phi(x) \leq C_\varepsilon \Psi(x)$ if $|x| \geq \varepsilon$. Assume that $\|f\|_{\Psi(L)} \leq 1$, so that

$$\int \Psi(|f(x)|) dx \leq 1.$$

Then convexity implies that for each $A > 0$,

$$\int \Psi \left(\frac{|f(x)|}{A} \right) \leq \frac{1}{A}.$$

Thus

$$\begin{aligned} \int \Phi\left(\frac{|f(x)|}{A}\right) dx &\leq \Phi(\varepsilon)|X| + C_\varepsilon \int \Psi\left(\frac{|f(x)|}{A}\right) \\ &\lesssim \Phi(\varepsilon)|X| + \frac{C_\varepsilon}{A}. \end{aligned}$$

If $\Phi(\varepsilon) \leq 2/|X|$, and $A \geq 2C_\varepsilon$, then we conclude that

$$\int \Phi\left(\frac{|f(x)|}{A}\right) dx \leq 1.$$

Thus $\|f\|_{\Phi(L)} \lesssim 1$. □

The Orlicz spaces satisfy an interesting duality relation. Given a function Φ , which we assume is *superlinear*, in the sense that $\Phi(x)/x \rightarrow \infty$ as $x \rightarrow \infty$, define its *Young dual*, for each $y \in [0, \infty)$, by

$$\Psi(y) = \sup\{xy - \Phi(x) : x \in [0, \infty)\}.$$

Then Ψ is the smallest function such that $\Phi(x) + \Psi(y) \geq xy$ for each x, y . This quantity is finite for each y because Φ is superlinear; for each $y \geq 0$, there exists $x(y)$ such that $\Phi(x(y)) \geq xy$, and thus the maximum of $xy - \Phi(x)$ is attained for $x \leq x(y)$. In particular, since Φ is continuous, the supremum is actually attained. Conversely, for each $x_0 \in [0, \infty)$, convexity implies there exists a largest y such that the line $y(x - x_0) + f(x_0) \leq f(x)$ for all $x \in [0, \infty)$. This means that $\Psi(y) = x_0 y - x_0$.

We note also that $\Psi(0) = 0$, and Ψ is increasing. Most importantly, the function is convex. Given any $y, z \in [0, \infty)$, and any $x \in [0, \infty)$,

$$\begin{aligned} x(\alpha y + (1 - \alpha)z) - \Phi(x) &\leq \alpha(xy - \Phi(x)) + (1 - \alpha)(xz - \Phi(x)) \\ &\leq \alpha\Psi(y) + (1 - \alpha)\Psi(z). \end{aligned}$$

Taking infimum over all x gives convexity. The function Ψ is also superlinear, since for any $x \in [0, \infty)$,

$$\lim_{y \rightarrow \infty} \frac{\Psi(y)}{y} \geq \lim_{y \rightarrow \infty} \frac{xy - \Phi(x)}{y} = x.$$

In particular, we can consider the Young dual of Ψ .

Lemma 15.26. *If Ψ is the Young dual of Φ , then Φ is the Young dual of Ψ .*

Proof. Π is the smallest function such that $\Pi(x) + \Psi(y) \geq xy$. Since $\Phi(x) + \Psi(y) \geq xy$ for each x and y , we conclude that $\Pi(x) \leq \Phi(x)$ for each x . For each x , there exists y such that $\Psi(y) = yx - \Phi(x)$. But this means that $\Phi(x) = yx - \Psi(y) \leq \Pi(x)$. \square

Given the Orlicz space $\Phi(L)$ for superlinear Φ , we can consider the Orlicz space $\Psi(L)$, where Ψ is the Young dual of Φ . The inequality $xy \leq \Phi(x) + \Psi(y)$, then

$$|f(x)g(x)| \leq \Phi(|f(x)|) + \Psi(|g(x)|),$$

so if $\|f\|_{\Phi(L)}, \|g\|_{\Psi(L)} \leq 1$, then

$$\left| \int f(x)g(x) \right| \leq \int |f(x)||g(x)| \leq \int \Phi(|f(x)|) + \int \Psi(|g(x)|) \leq 2.$$

Thus in general, we have

$$\left| \int f(x)g(x) \right| \leq 2\|f\|_{\Phi(L)}\|g\|_{\Psi(L)},$$

a form of Hölder's inequality. The duality between convex functions extends to a duality between the Orlicz spaces.

Theorem 15.27. *For any superlinear Φ with Young dual Ψ ,*

$$\|f\|_{\Phi(L)} \sim \sup \left\{ \int fg : \|g\|_{\Psi(L)} \leq 1 \right\}.$$

Proof. Without loss of generality, assume $\|f\|_{\Phi(L)} = 1$. The version of Hölder's inequality proved above shows that

$$\|f\|_{\Phi(L)} \lesssim 1.$$

Conversely, for each x , we can find $g(x)$ such that $f(x)g(x) = \Phi(|f(x)|) + \Psi(|g(x)|)$. Provided $\|g\|_{\Psi(L)} < \infty$, we have

$$\int fg = \int \Phi(|f(x)|) + \int \Psi(|g(x)|) \geq 1 + \|g\|_{\Psi(L)}.$$

Assuming $f \in L^\infty(X)$, we may choose $g \in L^\infty(X)$. For such a choice of function, $\|g\|_{\Psi(L)} < \infty$, which implies the result. Taking an approximation argument then gives the result in general. \square

Let us now consider some examples of duality.

Example. If $\Phi(x) = x^p$, for $p \geq 1$, and $1 = 1/p + 1/q$, then its Young dual Ψ satisfies

$$\Psi(y) = \sup_{x \geq 0} xy - x^p = y^{1+q/p}/p^{q/p} - y^q/p^q = y^q[p^{-q/p} - p^{-q}].$$

Thus the Young dual corresponds, up to a constant, to the conjugate dual in the L^p spaces.

Example. Suppose X has finite measure. If $\Phi(t) = e^t - 1$, then its dual satisfies, for large y ,

$$\begin{aligned} \Psi(y) &= \sup_{x \geq 0} xy - (e^x - 1) \\ &= y \log y - (y - 1) \sim y \log y. \end{aligned}$$

This is comparable to $y \log(y + 2)$ for large y . Thus $L \log L$ is dual to e^L .

Example. Suppose X has finite measure. If $\Phi(x) = e^{x^2} - 1$, then for $y \geq 2$,

$$\Psi(y) = \sup_{x \geq 0} xy - (e^{x^2} - 1) \sim y \log(y/2)^{1/2}.$$

Thus the dual of e^{L^2} is the space $L(\log L)^{1/2}$.

There is a generalization of both the Lorentz spaces and the Orlicz spaces, known as the Lorentz-Orlicz spaces, but these come up so rarely in analysis that we do not dwell on these norms.

Chapter 16

Sobolev Spaces

TODO: Treves, Chapter 31 for cool discussion of Sobolev spaces of negative index.

Previously, we discussed the rearrangement invariant spaces, which quantify properties of a function $f : X \rightarrow Y$ in terms of its ‘width’ and ‘height’. Here we discuss the ‘regularity’ of functions, i.e. how gradually they change, and the related topic (from a Fourier analytic perspective) of how fast the function ‘oscillates’ (inversely proportional to the ‘wavelength’ of a function). Such properties are certainly not quantified in terms of rearrangement invariant properties, since change and oscillation necessarily imply some topological features of the spaces involved. Here are some canonical, intuitive examples. To introduce them, we fix some function $\phi \in C_c^\infty(\mathbf{R})$, which we intuitively think of as having some height H and width W :

- If $\phi \in C_c^\infty(\mathbf{R})$, then the function $f(t) = \phi(t)e^{iNt}$ is the canonical example of a function oscillating at a frequency N . It has height H and width W . However, its derivative $f'(t) = \phi'(t)e^{iNt} + iN\phi(t)e^{iNt}$ for large N has height roughly proportional to $N \cdot H$ and width roughly proportional to W , and as we successively differentiate, the height grows even faster in N . This means that the function fails to be ‘regular’ for large N .
- Let $f(t) = N^{-s}\phi(t)e^{iNt}$, or let $f(t) = N^{-s}\phi(Nt)$. Both of these functions has height $N^{-s}H$ and width W , and oscillates at a frequency N . But unlike the last examples, the height (e.g. the L^∞ norm) of the

function $f^{(k)}(t)$ is uniformly bounded in t provided that $k \leq s$. We therefore think of f as having ‘ s degrees of regularity’.

- Let us consider the more complex function $f(x) = \phi(x)|x|^s \mathbf{I}_{x>0}$. One can break this function up into a sum $\sum_{n=0}^{\infty} \phi_n(x)|x|^s$, where $\phi_n(x) = \phi(2^n x) - \phi(2^{n+1} x)$. The function $f_n(x) = \phi_n(x)|x|^s$ behaves very similarly to the function $\phi_n(x)2^{-ns}$, an example considered in the last bullet with $N = 2^n$. In particular, the function oscillates at a frequency 2^n , and has s degrees of regularity. Thus we would image f itself has s degrees of regularity, but is composed of many different frequency scales.

There are various norms that quantify height, width, frequency scale, and regularity. The most common is the Sobolev norm, though there are various refinements, like the Hölder norms, Besov norms, Triebel-Lizorkin norms.

16.1 Hölder Spaces

Hölder Spaces are a simplified version of Sobolev spaces which account for the height, regularity, and oscillation of a function, but not the width. For $f : \mathbf{R}^d \rightarrow \mathbf{C}$, and an integer $k \geq 0$, we define

$$\|f\|_{C^k(\mathbf{R}^d)} = \max_{1 \leq i \leq k} \|D^i f\|_{L^\infty(\mathbf{R}^d)}.$$

Note that a C^k function might not necessarily lie have finite $C^k(\mathbf{R}^d)$ norm if it's derivatives are unbounded. We let C_{loc}^k denote the space of all C^k functions. For instance, $e^x \in C_{\text{loc}}^\infty(\mathbf{R})$, but $e^x \notin C^\infty(\mathbf{R})$. Each of the space $C^k(\mathbf{R}^d)$, where k is finite, forms a Banach space. The space $C^\infty(\mathbf{R}^d)$ is a Fréchet space.

It is a useful heuristic that only the smallest and largest derivatives really matter when measuring regularity in a space, a fact also tied up to interpolation. For instance, we have the following result in Hölder spaces.

Theorem 16.1. *For any function $f : \mathbf{R}^d \rightarrow \mathbf{C}$,*

$$\|f\|_{C^k(\mathbf{R}^d)} \sim_{k,d} \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^k f\|_{L^\infty(\mathbf{R}^d)}.$$

Proof. Certainly the right hand side is upper bounded by the left hand side. On the other hand, suppose both f and all its k th order derivatives are bounded by A . We will prove by induction that for each $1 \leq i < k$, $\|D^i f\|_{L^\infty(\mathbf{R}^d)} \lesssim_{i,k,d} \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)}$. To do this, we perform a Taylor expansion, writing

$$\left| f(x+y) - f(x) - \sum_{j=1}^i D^j f(x)(y, \dots, y) \right| \lesssim \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)} \cdot |y|^{i+1}.$$

We have $|f(x+y) - f(x)| \leq 2\|f\|_{L^\infty(\mathbf{R}^d)}$. For $i = 1$, this means that

$$|Df(x)(y)| \lesssim \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^2 f\|_{L^\infty(\mathbf{R}^d)} |y|^2,$$

and since y was arbitrary, letting y range over all $|y| = 1$, we have

$$\|Df\|_{L^\infty(\mathbf{R}^d)} \lesssim \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^2 f\|_{L^\infty(\mathbf{R}^d)}.$$

For $i \geq 2$, and for each $1 \leq j \leq i-1$, we have by induction,

$$\|D^j f(x)(y, \dots, y)\| \lesssim_d \left(\|f\|_{L^\infty(\mathbf{R}^d)} + \|D^{j+1} f\|_{L^\infty(\mathbf{R}^d)} \right) |y|^j \lesssim_d \left(\|f\|_{L^\infty(\mathbf{R}^d)} + \|D^i f\|_{L^\infty(\mathbf{R}^d)} \right) |y|^j.$$

Carrying these sums up the Taylor series, we conclude that Plugging these values in gives that

$$|D^i f(x)(y, \dots, y)| \lesssim_d \|f\|_{L^\infty(\mathbf{R}^d)} \left(1 + |y|^{i-1} \right) + \|D^i f\|_{L^\infty(\mathbf{R}^d)} \left(|y| + |y|^{i-1} \right) + \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)} \cdot |y|^{i+1}.$$

Pick a constant $C \geq 1$ such that the left hand side is upper bounded by C times the right hand side. Then

$$|D^i f(x)(y, \dots, y)| - C \|D^i f\|_{L^\infty(\mathbf{R}^d)} \left(|y| + |y|^{i-1} \right) \leq \|f\|_{L^\infty(\mathbf{R}^d)} \left(1 + |y|^{i-1} \right) + C \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)} \cdot |y|^{i+1}$$

Choosing $y = 10C \cdot z$ for $|z| = 1$ gives

$$(10C)^i \cdot D^i f(x)(z, \dots, z) - (20C)^{i-1} \|D^i f\|_{L^\infty(\mathbf{R}^d)} \leq (10C)^{i+2} \left(\|f\|_{L^\infty(\mathbf{R}^d)} + \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)} \right).$$

Our proof is completed when we notice that

$$\|D^i f(x)\|_{L^\infty(\mathbf{R}^d)} \sim_n \sup_{|z|=1} |D^i f(x)(z, \dots, z)|,$$

so if we pick C large enough depending solely on d , we conclude that

$$\|D^i f\|_{L^\infty(\mathbf{R}^d)} \lesssim_d \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^{i+1} f\|_{L^\infty(\mathbf{R}^d)}.$$

This completes the induction. □

More generally, for any $0 \leq \alpha \leq 1$, we set $C^{k,\alpha}(\mathbf{R}^d)$ to be the set of all C^k functions such that $D^k f$ satisfies a Hölder condition of order α , i.e.

$$\|f\|_{C^{k,\alpha}(\mathbf{R}^d)} = \|f\|_{L^\infty(\mathbf{R}^d)} + \|D^k f\|_{L^\infty(\mathbf{R}^d)} + \sup_{x \neq y} \frac{|D^k f(x) - D^k f(y)|}{|x - y|^\alpha}.$$

For $\alpha > 1$, this norm is finite if and only if $D^k f$ is constant, which is why only the case $\alpha \leq 1$ is interesting. It is simple to see that $C^k(\mathbf{R}^d)$ is contained in $C^{k,\alpha}(\mathbf{R}^d)$ for each such α . We also have $C^{k,1}(\mathbf{R}^d) \subset C^{k+1}(\mathbf{R}^d)$, because of Taylor's formula.

Example. To see the relation between the Hölder norm and the height, width, frequency scale, etc, consider a bump function $\phi \in C_c^\infty(\mathbf{R}^d)$, and let $f(t) = A\phi(t/R)e^{Nt}$. Provided that $R \geq 1/N$, we think of this function as having height A , width R , and frequency N (if $R < 1/N$, the frequency term does not have enough 'space' to oscillate). Then $\|f\|_{C^{k,\alpha}(\mathbf{R}^d)} \lesssim_{\phi,d,k} AN^{k+\alpha}$. Thus R is not measured at all.

Lemma 16.2. An element of $C^{k,1}(\mathbf{R}^d)$ is precisely an element $f \in C^k(\mathbf{R}^d)$ such that the distributional derivative $D^{k+1}f$ is an L^∞ tensor.

Proof. It suffices to prove the case $k = 0$. If $f \in C^{0,1}(\mathbf{R}^d)$ then f is precisely a bounded, uniformly Lipschitz function. Such functions are absolutely continuous with uniformly bounded derivative defined almost everywhere, and thus the distributional derivative Df lies in L^∞ . Conversely, if $f \in L^\infty(\mathbf{R}^d)$ and $f' \in L^\infty(\mathbf{R}^d)$, then we have a representation formula

$$f(x) = \int_0^x f'(y) dy,$$

which implies f is Lipschitz. □

Chapter 17

Sobolev Spaces

Let Ω be an open subset of \mathbf{R}^d . A natural problem when studying smooth functions $\phi \in C_c^\infty(\Omega)$ is to obtain estimates on the partial derivatives of ϕ . For instance, one can consider the norms

$$\|\phi\|_{C^n(\Omega)} = \max_{|\alpha| \leq n} \|D^\alpha \phi\|_{L^\infty(\Omega)}.$$

The space $C_c^\infty(\Omega)$ is not complete with respect to this norm, but its completion is the space $C_b^n(\Omega)$ of n times bounded continuously differentiable functions on Ω , which still consists of regular functions. Unfortunately, such estimates are only encountered in the most trivial situations. As in the non-smooth case, one can often get much better estimates using the L^p norms of the derivatives, i.e. considering the norms

$$\|\phi\|_{W^{n,p}(\Omega)} = \left(\sum_{|\alpha| \leq n} \|D^\alpha \phi\|_{L^p(\Omega)}^p \right)^{1/p}.$$

As might be expected, $C_c^\infty(\Omega)$ is not complete with respect to the $W^{n,p}(\Omega)$ norm. However, its completion cannot be identified with a family of n times differentiable functions. Instead, to obtain a satisfactory picture of the completion under this norm, a Banach space we will denote by $W^{n,p}(\Omega)$, we must take a distribution approach.

For each multi-index α , if f and f_α are locally integrable functions on Ω , we say f_α is a weak derivative for f if for any $\phi \in C_c^\infty(\Omega)$,

$$\int_{\Omega} f_\alpha(x) \phi(x) \, dx = (-1)^{|\alpha|} \int_{\Omega} f(x) \phi_\alpha(x) \, dx.$$

In other words, this is the same as the derivative of f viewed as a distribution on Ω . We define $W^{n,p}$ to be the space of all functions $f \in L^p(\Omega)$ such that for each $|\alpha| \leq n$, a weak derivative f_α exists and is an element of $L^p(\Omega)$. We then define

$$\|f\|_{W^{n,p}(\Omega)} = \left(\sum_{|\alpha| \leq n} \|f_\alpha\|_{L^p(\Omega)} \right)^{1/p}.$$

Where this sum is treated as a maximum in the case $p = \infty$. Later on we will be able to show this space is a complete Banach space.

Example. Let B be the open unit ball in \mathbf{R}^d , and let $u(x) = |x|^{-s}$, where $s < n-1$. For which p is $u \in W^{1,p}(B)$? We calculate by an integration by parts that if $\phi \in C_c^\infty(B)$, we fix $\varepsilon > 0$ and write

$$\int_B \phi_i(x) u(x) dx = \int_{|x| \leq \varepsilon} \phi_i(x) u(x) + \int_{\varepsilon < |x| \leq 1} \phi_i(x) u(x).$$

The integral on the ε ball is negligible since $s < n$. Since u is smooth away from the origin, its distributional derivative agrees with its standard derivative, which is

$$u_i(x) = \frac{-\alpha x_i}{|x|^{s+2}}.$$

Thus $|u_i| \lesssim 1/|x|^{s+1}$. An integration by parts gives

$$\int_{\varepsilon < |x| \leq 1} \phi_i(x) u(x) = \int_{|x|=\varepsilon} \phi(x) u(x) v_i dS + \int_{\varepsilon < |x| \leq 1} \frac{s \phi(x) x_i}{|x|^{s+2}} dx,$$

where v_i is the normal vector to the sphere pointing inward. Since $s < n-1$, the surface integral tends to zero as $\varepsilon \rightarrow 0$. Thus the weak derivative of u is equal to the standard derivative. Consequently, $u \in W^{1,p}(B)$ if $s < n/p - 1$.

Example. If $\{r_k\}$ is a countable, dense subset of B , then we can define

$$u(x) = \sum_{k=1}^{\infty} \frac{|x - r_k|^{-s}}{2^k}$$

Then $u \in W^{1,p}(B)$ if $0 < \alpha < n/p - 1$, yet u has a dense family of singularities, and thus does not behave like any differentiable function we would think of.

Theorem 17.1. *For each $k \in \mathbf{N}$ and $1 \leq p \leq \infty$, $W^{k,p}(\Omega)$ is a Banach space.*

Proof. It is easy to verify that $\|\cdot\|_{W^{k,p}}$ is a norm on $W^{k,p}(\Omega)$. Let $\{u_n\}$ be a Cauchy sequence in $W^{k,p}(\Omega)$. In particular, this means that $\{D^\alpha u_n\}$ is a Cauchy sequence in $L^p(\Omega)$ for each multi-index α with $|\alpha| \leq k$. In particular, these are functions v_α such that $D^\alpha u_n$ converges to v_α in the L^p norm for each α . Thus it suffices to prove that if $v = \lim u_n$, then $D^\alpha v = v_\alpha$ for each α . But this follows because the Hölder inequality implies that for each fixed $\phi \in C_c^\infty(\Omega)$,

$$\begin{aligned} (-1)^{|\alpha|} \int \phi_\alpha(x) v(x) dx &= \lim_{n \rightarrow \infty} (-1)^{|\alpha|} \int \phi_\alpha(x) u_n(x) dx \\ &= \lim_{n \rightarrow \infty} \int \phi(x) (D^\alpha u_n)(x) dx \\ &= \int \phi(x) v_\alpha(x) dx. \end{aligned}$$

Thus $W^{k,p}(\Omega)$ is complete. □

17.1 Smoothing

It is often useful to be able to approximate elements of $W^{k,p}(\Omega)$ by elements of $C^\infty(\Omega)$. This is mostly possible. If $u \in W^{k,p}(\Omega)$, and $\{\eta_\varepsilon\}$ is a family of smooth mollifiers, then, viewing u as a function on \mathbf{R}^n supported on Ω , we can consider the convolution $u^\varepsilon = u * \eta_\varepsilon$, i.e. the function defined by setting

$$u^\varepsilon(x) = \int_{\Omega} u(x-y) \eta_\varepsilon(y) dy.$$

This is just normal convolution, where we identify the function u with the function $u\mathbf{I}_\Omega$ on \mathbf{R}^d . Then u^ε is a smooth function on \mathbf{R}^d supported on a ε thickening of Ω . However, u^ε does not necessarily converge to u in $W^{k,p}(\Omega)$ as $\varepsilon \rightarrow 0$, since the behaviour of the convolution can cause issues at the boundary of Ω , where the distributional derivative $D^\alpha(u\mathbf{I}_\Omega)$ does not behave like a locally integrable function. This is the only problem, however.

Theorem 17.2. *If $U \Subset \Omega$, then $\lim_{\varepsilon \rightarrow 0} \|u^\varepsilon - u\|_{L^p(U)} = 0$.*

Proof. For each $\varepsilon > 0$, let $U^\varepsilon = \{x \in \Omega : d(x, \partial\Omega) > \varepsilon\}$. If $x \in \Omega^\varepsilon$, then

$$((D^\alpha u) * \eta_\varepsilon)(x) = (u_\alpha \mathbf{I}_\Omega * \eta_\varepsilon)(x),$$

since the convolution only depends on the behaviour of $D^\alpha u$ on a ε ball around x , which is contained in the interior of Ω . We can apply standard results about mollifiers to conclude that $u_\alpha \mathbf{I}_\Omega * \eta_\varepsilon$ converges to $u_\alpha \mathbf{I}_\Omega$ in $L^p(\mathbf{R}^d)$ as $\varepsilon \rightarrow 0$. Since $U \subseteq \Omega$, we have $U \subset U^\varepsilon$ for small enough ε , and so $(D^\alpha u) * \eta_\varepsilon$ converges to u_α in $L^p(U)$ as $\varepsilon \rightarrow 0$. Since this is true for each α with $|\alpha| \leq k$, we obtain the result. \square

If we are a little more careful, then we can fully approximate elements of $W^{k,p}(\Omega)$ by smooth functions on U .

Theorem 17.3. $C_c^\infty(\Omega) \cap W^{k,p}(\Omega)$ is dense in $W^{k,p}(\Omega)$.

Proof. Consider a family of open sets $\{V_n\}$ such that $V_n \subseteq \Omega$ for each n , and $U = \bigcup V_n$. Then we can consider a smooth partition of unity $\{\xi_n\}$ subordinate to the cover $\{V_n\}$. For each $u \in W^{k,p}(\Omega)$, we can write $u = \sum_n u \xi_n$. In particular, this means that for each $\varepsilon > 0$, there is N such that $\|\sum_{n=N+1}^\infty u \xi_n\|_{W^{k,p}(\Omega)} \leq \varepsilon$. For each $n \in \{1, \dots, N\}$, we can find δ_n small enough that the δ_n thickening of V_n is compactly contained in Ω . If ε_n is small enough, we find $(u \xi_n)^{\varepsilon_n}$ is supported on the δ_n thickening of V_n , and $\|(u \xi_n)^{\varepsilon_n} - u \xi_n\|_{W^{k,p}(V_n)} \leq \varepsilon/N$. But we then find

$$\|u - \sum_{n=1}^N (u \xi_n)^{\varepsilon_n}\|_{W^{k,p}(\Omega)} \leq \varepsilon + \sum_{n=1}^N \|u \xi_n - (u \xi_n)^{\varepsilon_n}\|_{W^{k,p}(\Omega)} \leq 2\varepsilon.$$

Thus $C_c^\infty(\Omega)$ is dense in $W^{k,p}(\Omega)$. \square

Approximation by elements of $C^\infty(\overline{\Omega})$ requires some more care, and additional assumptions on the behaviour of $\partial\Omega$.

Chapter 18

Interpolation Theory

One of the most fundamental tools in the ‘hard style’ of mathematical analysis, involving explicit quantitative estimates on quantities that arises in basic methods of mathematics, is the theory of interpolation. The main goal of interpolation is to take two estimates, and blend them together to form a family of intermediate estimates. Often each estimate will focus on one component of the problem at hand (an estimate in terms of the decay of the function at ∞ , an estimate involving the growth of the derivative, or the low frequency the function is, etc). By interpolating, we can optimize and obtain an estimate which simultaneously takes into account multiple features of the function. As should be expected, our main focus will be on the *interpolation of operators*. There are both complex methods of interpolation, and real methods of interpolation. These methods are interchangeable, *most of the time*, except when dealing with the boundedness of operators at endpoints. Sometimes complex interpolation is more useful, and sometimes real interpolation works better. So it is useful to know both types of interpolation.

18.1 Interpolation of Functions

The most basic way to interpolate is using the notion of convexity. Given two inequalities $A_0 \leq B_0$ and $A_1 \leq B_1$, for any parameter $0 \leq \theta \leq 1$, if we define the additive weighted averages $A_\theta = (1 - \theta)A_0 + \theta A_1$ and $B_\theta = (1 - \theta)B_0 + \theta B_1$, then we conclude $A_\theta \leq B_\theta$ for all θ . Similarly, we can consider the weighted multiplicative averages $A_\theta = A_0^{1-\theta} A_1^\theta$ and

$B_\theta = B_0^{1-\theta} B_1^\theta$, in which case we still have $A_\theta \leq B_\theta$. Note that the additive averages are obtained by taking the unique linear function between two values, and the multiplicative averages are obtained by taking the unique log-linear function between two values. In particular, if A_θ is defined to be any convex function, then $A_\theta \leq (1-\theta)A_0 + \theta A_1$, and if B_θ is logarithmically convex, so that $\log B_\theta$ is convex, then $B_\theta \leq B_0^{1-\theta} B_1^\theta$. Thus convexity provides us with a more general way of interpolating estimates, which is what makes this property so useful in analysis, enabling us to simplify estimates.

Example. For a fixed, measurable function f , the map $p \mapsto \|f\|_p$ is a log convex function. This statement is precisely Hölder's inequality, since the inequality

$$\|f\|_{\theta p + (1-\theta)q} \leq \|f\|_p^\theta \|f\|_q^{1-\theta}$$

says

$$\| |f|^{\theta p} |f|^{(1-\theta)q} \|_1^{1/(\theta p + (1-\theta)q)} \leq \|f^{\theta p}\|_{1/\theta}^\theta \|f^{(1-\theta)q}\|_{1/(1-\theta)}^{1-\theta}$$

which is precisely Hölder's inequality. Note this implies that if $p_0 < p_\theta < p_1$, then $L^{p_0}(X) \cap L^{p_1}(X) \subset L^{p_\theta}(X)$.

Example. The weak L^p norm is log convex, because if $F(t) \leq A_0^{p_0}/t^{p_0}$, and $F(t) \leq A_1^{p_1}/t^{p_1}$, then we can apply scalar interpolation to conclude that if $p_\theta = (1-\alpha)p_0 + \alpha p_1$,

$$F(t) \leq \frac{A_0^{(1-\alpha)p_0} A_1^{\alpha p_1}}{t^{(1-\alpha)p_0 + \alpha p_1}} = \frac{A_\theta^{p_\theta}}{t^{p_\theta}}$$

where p_θ is the harmonic weighted average between p_0 and p_1 , and A_θ the geometric weighted average. Using this argument, interpolating slightly to the left and right of p_θ , we can conclude that if $p_0 < p_\theta < p_1$, then $L^{p_0, \infty}(X) \cap L^{p_1, \infty}(X) \subset L^{p_\theta}(X)$.

18.2 Complex Interpolation

Another major technique to perform an interpolation is to utilize the theory of complex analytic functions to obtain estimates. The core idea of this technique is to exploit the maximum principle, which says that bounding an analytic function at its boundary enables one to obtain bounds everywhere in the domain of the function. The next result, known as Lindelöf's

theorem, is one of the fundamental examples of the application of complex analysis.

Theorem 18.1 (The Three Lines Lemma). *If f is a holomorphic function on the strip $S = \{z : \operatorname{Re}(z) \in [a, b]\}$ and there exists constants $A, B, \delta > 0$ such that for all $z \in S$,*

$$|f(z)| \leq Ae^{Be^{(\pi-\delta)|z|}}.$$

Then the function $M : [a, b] \rightarrow [0, \infty]$ given by

$$M(s) = \sup_{t \in \mathbf{R}} |f(s + it)|$$

is log convex on $[a, b]$.

Proof. By a change of variables, we can assume that $a = 0$, and $b = 1$, and we need only show that if there are $A_0, A_1 > 0$ such that

$$|f(it)| \leq A_0 \quad \text{and} \quad |f(1 + it)| \leq A_1 \quad \text{for all } t \in \mathbf{R},$$

then for any $s \in [a, b]$ and $t \in \mathbf{R}$,

$$|f(s + it)| \leq A_0^{1-s} A_1^s.$$

By replacing $f(z)$ with the function $A_0^{1-z} A_1^z f(z)$, we may assume without loss of generality that $A_0 = A_1 = 1$, and we must show that $\|f\|_{L^\infty(S)} \leq 1$. If $|f(s + it)| \rightarrow 0$ as $|t| \rightarrow \infty$, then for large N , we can conclude that $|f(s + it)| \leq 1$ for $s \in [a, b]$ and $|t| \geq N$. But then the maximum principle entails that $|f(s + it)| \leq 1$ for $s \in [a, b]$ and $|t| \leq N$, which completes the proof in this case. In the general case, for each $\varepsilon > 0$, define

$$u_\varepsilon(z) = \exp(-2\varepsilon \sin((\pi - \varepsilon)z + \varepsilon/2)).$$

Then if $z = s + it$,

$$|u_\varepsilon(z)| = \exp(-\varepsilon[e^{(\pi-\varepsilon)t} + e^{-(\pi-\varepsilon)t}]\sin((\pi - \varepsilon)s + \varepsilon/2)),$$

So, in particular, $|u_\varepsilon(z)| \leq 1$, and there exists a constant C such that if $z \in S$,

$$|u_\varepsilon(z)| \leq e^{-C\varepsilon^2 e^{(\pi-\varepsilon)|z|}}$$

Note that if $\varepsilon < \delta$, then as $|\operatorname{Im}(z)| \rightarrow \infty$,

$$|f(z)u_\varepsilon(z)| \leq Ae^{Be^{(\pi-\delta)|z|} - C\varepsilon^2 e^{(\pi-\varepsilon)|z|}} \rightarrow 0.$$

Applying the previous case to the function $|f(z)u_\varepsilon(z)|$, we conclude that for any $\varepsilon > 0$,

$$|f(z)| \leq \frac{1}{|u_\varepsilon(z)|}.$$

Thus

$$|f(z)| \leq \lim_{\varepsilon \rightarrow 0} \frac{1}{|u_\varepsilon(z)|} = 1,$$

which completes the proof. \square

Remark. The function $e^{-ie^{\pi i}s}$ shows that the assumption of the three lines lemma is essentially tight. In particular, this means there is no family of holomorphic functions g_ε which decays faster than double exponentially, and pointwise approximates the identity as $\varepsilon \rightarrow 0$.

Remark. Similar variants can be used to show that if f is a holomorphic function on an annulus, then the supremum over circles centered around the origin is log convex in the radius of the circle (a result often referred to as the three circles lemma).

Example. Here we show how we can use the three lines lemma to prove that the L^p norms are log convex. If $f = \sum a_n \chi_{E_n}$ is a simple function, then the function

$$g(s) = \int |f|^s = \sum |a_n|^s |E_n|$$

is analytic in s , and satisfies the growth condition of the three lines lemma because each term of the sum is exponential in growth. Since $|g(s)| \leq |g(\sigma)|$, the three lines lemma implies that g is log convex on the real line. By normalizing the function f and the underlying measure, given p_0, p_1 , we may assume $\|f\|_{p_0} = \|f\|_{p_1} = 1$, and it suffices to prove that $\|f\|_{p_\theta} \leq 1$ for all $p_\theta \in [p_0, p_1]$. But the log convexity of g guarantees this is true, since $|g(p)| = \|f\|_p^p$. A standard limiting argument then gives the inequality for all functions f .

Example. Let f be a holomorphic function on a strip $S = \{z : \operatorname{Re}(z) \in [a, b]\}$, such that if $z = a + it$, or $z = b + it$, for some $t \in \mathbf{R}$,

$$|f(z)| \leq C_1(1 + |z|)^\alpha.$$

Then there exists a constant C' such that for any $z \in S$,

$$|f(z)| \leq C_2(1 + |z|)^\alpha.$$

Proof. The function

$$g(z) = \frac{f(z)}{(1 + z)^\alpha}$$

is holomorphic on S , and if $z = a + it$ or $z = b + it$,

$$|g(z)| \leq \frac{C_1(1 + |z|)^\alpha}{|1 + z|^\alpha} \lesssim 1.$$

Thus the three lines lemma implies that $|g(z)| \lesssim 1$ for all $z \in S$, so

$$|f(z)| \lesssim |1 + z|^\alpha \lesssim (1 + |z|)^\alpha. \quad \square$$

TODO: Is Lemma 20.1 of Treves Distribution Theory an example of interpolation?

18.3 Interpolation of Operators

A major part of modern harmonic analysis is the study of operators, i.e. maps from function spaces to other function spaces. We are primarily interested in studying *linear operators*, i.e. operators T such that $T(f + g) = T(f) + T(g)$, and $T(\alpha f) = \alpha T(f)$, and also *sublinear operators*, such that $|T(\alpha f)| = |\alpha| |T(f)|$ and $|T(f + g)| \leq |Tf| + |Tg|$. Even if we focus on linear operators, it is still of interest to study sublinear operators because one can study the *uniform boundedness* of a family of operators $\{T_k\}$ by means of the function $T^*(f)(x) = \max(T_k f)(x)$. This is the method of *maximal functions*. Another important example are the l^p sums

$$(S^p f)(x) = \left(\sum |T_k(x)|^p \right)^{1/p}.$$

These two examples are specific examples where we have a family of operators $\{T_y\}$, indexed by a measure space Y , and we define an operator S by taking Sf to be the norm of $\{T_y f\}$ in the variable y .

Here we address the most basic case of operator interpolation. As we vary p , the L^p norms provide different ways of measuring the height and

width of functions. Let us consider a simple example. Suppose that for an operator T , we have a bound

$$\|Tf\|_{L^1(Y)} \leq \|f\|_{L^1(X)} \quad \text{and} \quad \|Tf\|_{L^\infty(Y)} \leq \|f\|_{L^\infty(X)}.$$

The first inequality shows that the width of Tf is controlled by the width of f , and the second inequality says the height of Tf is controlled by the height of f . If we take a function $f \in L^p(X)$, for some $p \in (1, \infty)$, then we have some control over the height of f , and some control of the width. In particular, this means we might expect some control over the width and height of Tf , i.e. for each p , a bound

$$\|Tf\|_{L^p(Y)} \leq \|f\|_{L^p(X)}.$$

This is the idea of interpolation on the $L^p(X)$ spaces.

18.4 Complex Interpolation of Operators

The first theorem we give is the Riesz-Thorin theorem, which utilizes complex interpolation to give such a result. In the next theorem, we work with a linear operator T which maps simple functions f on a measure space X to functions on a measure space Y . For the purposes of applying duality, we make the mild assumption that for each simple function g ,

$$\int |(Tf)(y)| |g(y)| dy < \infty.$$

Our goal is to obtain L^p bounds on the function T . The Hahn-Banach theorem then guarantees that T has a unique extension to a map defined on all L^p functions.

Theorem 18.2 (Riesz-Thorin). *Let $p_0, p_1 \in (0, \infty]$ and $q_0, q_1 \in [1, \infty]$. Suppose that*

$$\|Tf\|_{L^{q_0}(Y)} \leq A_0 \|f\|_{L^{p_0}(X)} \quad \text{and} \quad \|Tf\|_{L^{q_1}(Y)} \leq A_1 \|f\|_{L^{p_1}(X)}.$$

Then for any $\theta \in (0, 1)$, if

$$1/p_\theta = (1 - \theta)/p_0 + \theta/p_1 \quad \text{and} \quad 1/q_\theta = (1 - \theta)/q_0 + \theta/q_1,$$

then

$$\|Tf\|_{L^{q_\theta}(Y)} \leq A_\theta \|f\|_{L^{p_\theta}(X)},$$

where $A_\theta = A_0^{1-\theta} A_1^\theta$.

Proof. If $p_0 = p_1$, the proof follows by the log convexity of the L^p norms of a function. Thus we may assume $p_0 \neq p_1$, so p_θ is finite in any case of interest. By normalizing the measures on both spaces, we may assume $A_0 = A_1 = 1$. By duality and homogeneity, it suffices to show that for any two simple functions f and g such that $\|f\|_{q_\theta} = \|g\|_{q_\theta^*} = 1$,

$$\left| \int_Y (Tf)g \, dy \right| \leq 1.$$

Our challenge is to make this inequality complex analytic so we can apply the three lines lemma. We write $f = F_0^{1-\theta} F_1^\theta a$, where F_0 and F_1 are non-negative simple functions with $\|F_0\|_{L^{p_0}(X)} = \|F_1\|_{L^{p_1}(X)} = 1$, and a is a simple function with $|a(x)| = 1$. Similarly, we can write $g = G_0^{1-\theta} G_1^\theta b$. We now write

$$H(s) = \int_Y T(F_0^{1-s} F_1^s a) G_0^{1-s} G_1^s b \, dy.$$

Since all functions involved here are simple, $H(s)$ is a linear combination of positive numbers taken to the power of $1-s$ or s , and is therefore obviously an entire function in s . Now for all $t \in \mathbf{R}$, we have

$$\|F_0^{1-it} F_1^{it} a\|_{L^{p_0}(X)} = \|F_0\|_{L^{p_0}(X)} = 1,$$

$$\|G_0^{1-it} G_1^{it} b\|_{L^{q_0}(Y)} = \|G_0\|_{L^{q_0}(Y)} = 1.$$

Therefore

$$|H(it)| = \left| \int_Y T(F_0^{1-it} F_1^{it} a) G_0^{1-it} G_1^{it} b \, dy \right| \leq 1.$$

Similarly, $|H(1+it)| \leq 1$ for all $t \in \mathbf{R}$. An application of Lindelöf's theorem implies $|H(s)| \leq 1$ for all s . Setting $s = \theta$ completes the argument. \square

If, for each p, q , we let $F(1/p, 1/q)$ to be the operator norm of a linear operator T viewed as a map from $L^p(X)$ to $L^q(Y)$, then the Riesz-Thorin theorem says that F is a log-convex function. In particular, the set of $(1/p, 1/q)$ such that T is bounded as a map from $L^p(X)$ to $L^q(Y)$ forms a convex set. If this is true, we often say T is of *strong type* (p, q) .

Example. For any two integrable functions $f, g \in L^1(\mathbf{R}^d)$, we can define an integrable function $f * g \in L^1(\mathbf{R}^d)$ almost everywhere by the integral formula

$$(f * g)(x) = \int f(y)g(x - y) dy.$$

If $f \in L^1(\mathbf{R}^d)$ and $g \in L^p(\mathbf{R}^d) \cap L^1(\mathbf{R}^d)$, for some $p \geq 1$, then Minkowski's integral inequality implies

$$\begin{aligned} \|f * g\|_p &= \left(\int |f * g(x)|^p dx \right)^{1/p} \leq \int \left(\int |f(y)g(x - y)|^p dx \right)^{1/p} dy \\ &= \int |f(y)| \|g\|_{L^p(\mathbf{R}^d)} dy = \|f\|_{L^1(\mathbf{R}^d)} \|g\|_{L^p(\mathbf{R}^d)}. \end{aligned}$$

Hölder's inequality implies that if $f \in L^p(\mathbf{R}^d)$ and $g \in L^q(\mathbf{R}^d)$, where p and q are conjugates of one another, then

$$\left| \int f(y)g(x - y) dy \right| \leq \int |f(y - x)| |g(x)| \leq \|f\|_{L^p(\mathbf{R}^d)} \|g\|_{L^q(\mathbf{R}^d)}.$$

Thus we have the bound

$$\|f * g\|_{L^\infty(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)} \|g\|_{L^q(\mathbf{R}^d)}.$$

Now that these mostly trivial results have been proved, we can apply convolution. For each $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$, we have a convolution operator $T : L^1(\mathbf{R}^d) \rightarrow L^1(\mathbf{R}^d)$ defined by $Tg = f * g$. We know that T is of strong type $(1, p)$, and of type (q, ∞) , where q is the harmonic conjugate of p , and T has operator norm 1 with respect to each of these types. But the Riesz Thorin theorem then implies that if $1/r = \theta + (1 - \theta)/q$, then T is bounded as a map from $L^r(\mathbf{R}^d)$ to $L^{p/\theta}(\mathbf{R}^d)$ with operator norm one. Reparameterizing gives Young's convolution inequality. Note that we never really used anything about \mathbf{R}^d here other than its translational structure, and as such Young's inequality continues to apply in the theory of any modular locally compact group. In particular, the Haar measure μ on such a group is only defined up to a scalar multiple, and if we swap μ with $\alpha\mu$, for some $\alpha > 0$, then Young's inequality for this measure implies

$$\lambda^{1+1/r} \|f * g\|_r = \lambda^{1/p+1/q} \|f\|_p \|g\|_p$$

which is a good way of remembering that we must have $1 + 1/r = 1/p + 1/q$.

Example. Let X be a measure space with σ algebra Σ_0 , and let $\Sigma \subset \Sigma_0$ be a σ finite sub σ algebra. Then $L^2(X, \Sigma)$ is a closed subspace of $L^2(X, \Sigma_0)$, and so there is an orthogonal projection operator $\mathbf{E}(\cdot|\Sigma) : L^2(X, \Sigma_0) \rightarrow L^2(X, \Sigma)$, which we call the conditional expectation operator. The properties of the projection operator imply that for any $f, g \in L^2(X, \Sigma_0)$,

$$\int \mathbf{E}(f|\Sigma) \bar{g} = \int f \bar{g} = \int \mathbf{E}(f|\Sigma) \overline{\mathbf{E}(g|\Sigma)}.$$

If $g \in L^2(X, \Sigma)$, then

$$\int \mathbf{E}(f|\Sigma) \bar{g} = \int f \bar{g}.$$

This gives a full description of $\mathbf{E}(f|\Sigma)$. In particular, if $u \in L^\infty(X, \Sigma_0)$, then for each $g \in L^2(X, \Sigma)$

$$\int \mathbf{E}(uf|\Sigma) \bar{g} = \int f [u \bar{g}] = \int u \mathbf{E}(f|\Sigma) \bar{g}.$$

Since this is true for all $g \in L^2(X, \Sigma)$, we find $\mathbf{E}(uf|\Sigma) = u \mathbf{E}(f|\Sigma)$. Moreover, if $0 \leq f \leq g$, then $\mathbf{E}(f|\Sigma) \leq \mathbf{E}(g|\Sigma)$. This is easy to see because if $f \geq 0$, and $F = \{x : \mathbf{E}(f|\Sigma) < 0\}$, then if $|F| \neq 0$,

$$0 > \int \mathbf{E}(f|\Sigma) \mathbf{I}_F = \int f \mathbf{I}_F \geq 0.$$

Thus $|F| = 0$, and so $\mathbf{E}(f|\Sigma) \geq 0$ almost everywhere.

Like all other orthogonal projection operators, conditional expectation is a contraction in the L^2 norm, i.e. $\|\mathbf{E}(f|\Sigma)\|_{L^2(X)} \leq \|f\|_{L^2(X)}$. We now use interpolation to show that conditional expectation is strong (p, p) , for all $1 \leq p \leq \infty$. It suffices to prove the operator is strong $(1, 1)$ and strong (∞, ∞) . So suppose $f \in L^2(X, \Sigma_0) \cap L^\infty(X, \Sigma_0)$. If $|E| < \infty$, then $\mathbf{I}_E \in L^2(X)$, so

$$|\mathbf{E}(f|\Sigma)| \mathbf{I}_E = |\mathbf{E}(\mathbf{I}_E f|\Sigma)| \leq \mathbf{E}(\mathbf{I}_E |f| |\Sigma) \leq \|f\|_\infty \mathbf{E}(\mathbf{I}_E |\Sigma) = \|f\|_\infty \mathbf{I}_E.$$

Since Σ is a sigma finite sigma algebra, we can take $E \rightarrow \infty$ to conclude $\|\mathbf{E}(f|\Sigma)\|_\infty \leq \|f\|_\infty$. The case $(1, 1)$ can be obtained by duality, since conditional expectation is self adjoint, or directly, since if $f \in L^1(X, \Sigma_0) \cap L^2(X, \Sigma_0)$, then for any set $E \in \Sigma$ with $|E| < \infty$,

$$\int |\mathbf{E}(f|\Sigma)| \mathbf{I}_E \leq \int \mathbf{E}(|f| |\Sigma) \mathbf{I}_E = \int_E |f| \mathbf{I}_E \leq \|f\|_1.$$

Since Σ is σ finite, we can take $E \rightarrow \infty$ to conclude $\|\mathbf{E}(f|\Sigma)\|_1 \leq \|f\|_1$. Thus the Riesz interpolation theorem implies that for each $1 \leq p \leq \infty$, $\|\mathbf{E}(f|\Sigma)\|_p \leq \|f\|_p$.

Since $L^2(X, \Sigma_0)$ is dense in $L^p(X, \Sigma_0)$ for all $1 \leq p < \infty$, there is a unique extension of the conditional expectation operator from $L^p(X, \Sigma_0)$ to $L^p(X, \Sigma_0)$. For $p = \infty$, there are infinitely many extensions of the conditional expectation operator from $L^\infty(X, \Sigma_0)$ to $L^\infty(X, \Sigma_0)$. However, there is a unique extension such that for each $f \in L^2(\Sigma_0)$ and $g \in L^\infty(\Sigma)$, $\mathbf{E}(fg|\Sigma) = g\mathbf{E}(f|\Sigma)$. This is because for any $E \in \Sigma$ with $|E| < \infty$, $\mathbf{E}(f\mathbf{I}_E|\Sigma) = \mathbf{I}_E\mathbf{E}(f|\Sigma)$ is uniquely defined since $f\mathbf{I}_E \in L^2(\Sigma_0)$, and taking $E \rightarrow \infty$ by σ finiteness.

A simple consequence of the uniform boundedness of these operators on the various L^p spaces is that if $\Sigma_1, \Sigma_2, \dots$ are a family of σ algebras, and Σ_∞ is the smallest σ algebra containing all sets in $\bigcup_{i=1}^\infty \Sigma_i$, then for each $1 \leq p < \infty$, and for each $f \in L^p(\Sigma_0)$, $\lim_{i \rightarrow \infty} \mathbf{E}(f|\Sigma_i) = \mathbf{E}(f|\Sigma_\infty)$. This is because the operators $\{\mathbf{E}(\cdot|\Sigma_i)\}$ are uniformly bounded. The limit equation holds for any simple function f composed of sets in $\bigcup_{i=1}^\infty \Sigma_i$, and a σ algebra argument can then be used to show this family of simple functions is dense in $L^p(\Sigma_0)$.

It was an important observation of Elias-Stein that complex interpolation can be used not only with a single operator T , but with an ‘analytic family’ of operators $\{T_s\}$, one for each s , such that for each pair of simple functions f and g , the function

$$\int (T_s f)(y) g(y)$$

is analytic in s . Thus bounds on T_{0+it} and T_{1+it} imply intermediary bounds on all other operators, provided that we still have at most doubly exponential growth. The next theorem gives an example application.

Theorem 18.3 (Stein-Weiss Interpolation Theorem). *Let T be a linear operator, and let $w_0, w_1 : X \rightarrow [0, \infty)$ and $v_0, v_1 : Y \rightarrow [0, \infty)$ be weights which are integrable on every finite-measure set. Suppose that*

$$\|Tf\|_{L^{q_0}(X, v_0)} \leq A_0 \|f\|_{L^{p_0}(X, w_0)} \quad \text{and} \quad \|Tf\|_{L^{q_1}(X, v_1)} \leq A_1 \|f\|_{L^{p_1}(X, w_0)}.$$

Then for any $\theta \in (0, 1)$,

$$\|Tf\|_{L^{q_\theta}(X, v_\theta)} \leq A_\theta \|f\|_{L^{p_\theta}(X, w_\theta)},$$

where $w_\theta = w_0^{1-\theta} w_1^\theta$ and $v_\theta = v_0^{1-\theta} v_1^\theta$.

Proof. Fix a simple function f with $\|f\|_{L^{p_\theta}(X, w_\theta)}$. We begin with some simplifying assumptions. A monotone convergence argument, replacing $w_i(t)$ with

$$w'_i(y) = \begin{cases} w_i(y) & : \varepsilon \leq w_i(t) \leq 1/\varepsilon, \\ 0 & : \text{otherwise,} \end{cases}$$

and then taking $\varepsilon \rightarrow 0$, enables us to assume without loss of generality that w_0 and w_1 are both bounded from below and bounded from above. Truncating the support of Tf enables us to assume that Y has finite measure. Since f has finite support, we may also assume without loss of generality that X has finite support, and by applying the dominated convergence theorem we may replace the weights v_i with

$$v'_i(x) = \begin{cases} v_i(x) & : \varepsilon \leq v_i(x) \leq 1/\varepsilon, \\ 0 & : \text{otherwise,} \end{cases}$$

and then take $\varepsilon \rightarrow 0$. Thus we can assume that the v_i are bounded from above and below. Restricting to the support of X , we can also assume X has finite measure.

For each s , consider the operator T_s defined by

$$T_s f = w_0^{\frac{1-s}{q_0}} w_1^{\frac{s}{q_1}} T \left(f v_0^{-\frac{1-s}{p_0}} v_1^{-\frac{s}{p_1}} \right).$$

The fact that all functions involved are simple means that the family of operators $\{T_s\}$ is analytic. Now for all $t \in \mathbf{R}$

$$\|T_{it} f\|_{L^{q_0}(Y)} = \|Tf\|_{L^{q_0}(Y, w_0)} \leq A_0 \|f v_0^{-1/p_0}\|_{L^{p_0}(X, v_0)} = A_0 \|f\|_{L^{p_0}(X)}.$$

For similar reasons, $\|T_{1+it} f\|_{L^{q_1}(Y)} \leq A_1 \|f\|_{L^{p_0}(X, v_0)}$. Thus the Stein variant of the Riesz-Thorin theorem implies that

$$\|T_\theta f\|_{L^{q_\theta}(Y)} \leq A_\theta \|f\|_{L^{p_\theta}(X)}.$$

But this, of course, is equivalent to the bound we set out to prove. \square

18.5 Real Interpolation of Operators

Now we consider the case of real interpolation. One advantage of real interpolation is that it can be applied to sublinear as well as linear operators,

and requires weaker endpoint estimates than the complex case. A disadvantage is that, usually, the operator under study cannot vary, and we lose out on obtaining explicit bounds.

A strong advantage to using real interpolation is that the criteria for showing boundedness at the endpoints can be reduced considerably. Let us give names for the boundedness we will want to understand for a particular operator T .

- We say T is *strong type* (p, q) if $\|Tf\|_{L^q(Y)} \lesssim \|f\|_{L^p(X)}$.
- We say T is *weak type* (p, q) if $\|Tf\|_{L^{q,\infty}(Y)} \lesssim \|f\|_{L^p(X)}$.
- We say T is *restricted strong type* (p, q) if we have a bound

$$\|Tf\|_{L^q(Y)} \lesssim HW^{1/p}$$

for any sub-step functions with height H and width W . Equivalently, for any set E ,

$$\|T(\mathbf{I}_E)\|_{L^q(Y)} \lesssim |E|^{1/p}.$$

The equivalence is proven by breaking any sub-step function f with height H and width W into a dyadic sum $\sum_{k=1}^{\infty} H \mathbf{I}_{E_k} 2^{-k}$, where $|E_k| \leq W$.

- We say T is *restricted weak type* (p, q) if we have a bound

$$\|Tf\|_{L^{q,\infty}(Y)} \lesssim HW^{1/p}$$

for all sub-step functions with height H and width W . Equivalently, for any set E ,

$$\|T(\mathbf{I}_E)\|_{L^{q,\infty}(Y)} \lesssim |E|^{1/p}.$$

An important tool for us will be to utilize duality to make our interpolation argument ‘bilinear’. Let us summarize this tool in a lemma. Proving the lemma is a simple application of Theorem 15.13.

Lemma 18.4. *Let $0 < p < \infty$ and $0 < q < \infty$. Then an operator T is restricted weak-type (p, q) if and only if for any finite measure sets $E \subset X$ and $F \subset Y$, there is $F' \subset Y$ with $|F'| \geq \alpha |F|$ such that*

$$\int_{F'} |T(\mathbf{I}_E)| \lesssim_{\alpha} |E|^{1/p} |F|^{1-1/q}.$$

Scalar interpolation leads to a simple version of real interpolation, which we employ as a subroutine to obtain a much more powerful real interpolation principle.

Lemma 18.5. *Let $0 < p_0, p_1 < \infty$, $0 < q_0, q_1 < \infty$. If T is restricted weak type (p_0, q_0) and (p_1, q_1) , then T is restricted weak type (p_θ, q_θ) for all $\theta \in (0, 1)$.*

Proof. By assumption, if $E \subset X$ and $F \subset Y$, then there is $F_0, F_1 \subset Y$ with $|F_i| \geq (3/4)|F|$ such that

$$\int_{F_i} |T(\mathbf{I}_E)| \lesssim |E|^{1/p_i} |F_i|^{1-1/q_i}.$$

If we let $F_\theta = F_0 \cap F_1$, then $|F_\theta| \geq |F|/2$, and for each i ,

$$\int_{F_\theta} |T(\mathbf{I}_E)| \lesssim |E|^{1/p_i} |F_\theta|^{1-1/q_i}.$$

Scalar interpolation implies

$$\int_{F_\theta} |T(\mathbf{I}_E)| \lesssim |E|^{1/p_\theta} |F_\theta|^{1-1/q_\theta},$$

and thus we have shown

$$\|T(\mathbf{I}_E)\|_{q_\theta, \infty} \lesssim |E|^{1/p_\theta}.$$

This is sufficient to show T is restricted weak type (p_θ, q_θ) . \square

Theorem 18.6 (Marcinkiewicz Interpolation Theorem). *Let $0 < p_0, p_1 < \infty$, $0 < q_0, q_1 < \infty$, and suppose T is restricted weak type (p_i, q_i) , with constant A_i , for each i . Then, for any $\theta \in (0, 1)$, if $q_\theta > 1$, then for any $0 < r < \infty$, then*

$$\|Tf\|_{L^{q_\theta, r}(Y)} \lesssim A_\theta \|f\|_{L^{p_\theta, r}(X)},$$

with implicit constants depending on p_0, p_1, q_0 , and q_1 .

Proof. By scaling T , and the measures on X and Y , we may assume that $\|f\|_{L^{p_\theta, r}(X)} \leq 1$, and that T is restricted type (p_i, q_i) with constant 1, so that for any step function f with height H and width W ,

$$\|Tf\|_{L^{q_i, \infty}(Y)} \leq \|f\|_{L^{p_i}(X)}.$$

By duality, using the fact that $q_\theta > 1$, it suffices to show that for any simple function g with $\|g\|_{L^{q'_\theta, r'}(Y)} = 1$,

$$\int |Tf||g| \leq 1.$$

Using the previous lemma, we can ‘adjust’ the values q_0, q_1 so that we can assume $q_0, q_1 > 1$. We can perform a horizontal layer decomposition, writing

$$f = \sum_{i=-\infty}^{\infty} f_i, \quad \text{and} \quad g = \sum_{i=-\infty}^{\infty} g_i,$$

where f_i and g_i are sub-step functions with width 2^i and heights H_i and H'_i respectively, and if we write $A_i = H_i 2^{i/p_\theta}$, and $B_i = H'_i 2^{i/q_\theta}$, then

$$\|A\|_{l^r(\mathbf{Z})}, \|B\|_{l^{r'}(\mathbf{Z})} \lesssim 1.$$

Applying the restricted weak type inequalities, we know for each i and j ,

$$\int |Tf_i||g_j| \lesssim H_i H_j \min_{k \in \{0,1\}} \left[2^{i/p_k + j(1-1/q_k)} \right].$$

Applying sublinearity (noting that really, the decomposition of f and g is finite, since both functions are simple). Thus

$$\begin{aligned} \int |Tf||g| &\leq \sum_{i,j} \int |Tf_i||g_j| \\ &\lesssim \sum_{i,j} H_i H'_j \min_{k \in \{0,1\}} \left[2^{i/p_k + j(1-1/q_k)} \right] \\ &\lesssim \sum_{i,j} A_i B_j \min_{k \in \{0,1\}} \left[2^{i(1/p_k - 1/p_\theta) + j(1/q_\theta - 1/q_k)} \right]. \end{aligned}$$

If $i(1/p_k - 1/p_\theta) + j(1/q_\theta - 1/q_k) = \varepsilon(i + \lambda j)$, where $\varepsilon = (1/p_k - 1/p_\theta)$. We then have

$$\sum_{i,j} A_i B_j \min_{k \in \{0,1\}} \left[2^{i(1/p_k - 1/p_\theta) + j(1/q_\theta - 1/q_k)} \right] \sim \sum_{k=-\infty}^{\infty} \min(2^{\varepsilon k}, 2^{-\varepsilon k}) \sum_i A_i B_{k - \lfloor i/\lambda \rfloor}.$$

Applying Hölder's inequality,

$$\begin{aligned} \sum_i A_i B_{k-\lfloor i/\lambda \rfloor} &\leq \|A\|_{l^r(\mathbf{Z})} \left(\sum_i |B_{k-\lfloor i/\lambda \rfloor}|^{r'} \right)^{1/r'} \\ &\lesssim \lambda^{1/r'} \|A\|_{l^r(\mathbf{Z})} \|B\|_{l^{r'}(\mathbf{Z})} \lesssim 1. \end{aligned}$$

Thus we conclude that

$$\sum_{k=-\infty}^{\infty} \min(2^{\varepsilon k}, 2^{-\varepsilon k}) \sum_i A_i B_{k-\lfloor i/\lambda \rfloor} \lesssim \sum_{k=-\infty}^{\infty} \min(2^{\varepsilon k}, 2^{-\varepsilon k}) \lesssim_{\varepsilon} 1. \quad \square$$

There are many variants of the real interpolation method, but the general technique almost always remains the same: incorporate duality, decompose inputs, often dyadically, bound these decompositions, and then sum up.

Chapter 19

Basic Integral Operator Estimates

We now consider a very general class of operators, which can be seen as the infinite dimensional analogue of matrix multiplication, replacing summation over coordinates by integration. We fix two measure spaces X and Y , and consider a function (or, if X and Y are smooth manifolds, distributions) $K : X \times Y \rightarrow \mathbf{C}$, which we call a *kernel*. From this kernel, if K is suitably regular, we obtain an induced operator T_K taking functions on X to functions on Y , given, heuristically at least, by the integral formula

$$(T_K f)(y) = \int_X K(x, y) f(x) dx.$$

Our goal is to relate properties of the kernel K to the behaviour of the operator T_K .

Example. Let $X = \{1, \dots, N\}$ and $Y = \{1, \dots, M\}$, each equipped with the counting measure. Then each kernel K corresponds to an $M \times N$ matrix A , with $A_{ij} = K(j, i)$. For any $f : X \rightarrow \mathbf{C}$ we can define a vector $v \in \mathbf{R}^N$ by setting $v_i = f(i)$, and then

$$(T_K f)(m) = \sum_{n=1}^N f(n) K(n, m) = \sum_{n=1}^N A_{mn} v_n = (Av)_m.$$

Thus with respect to the standard basis, T_K is just given by matrix multiplication by A .

Example. Let $X = Y = \mathbf{R}^d$, and let $K(x, y) = e^{2\pi i x \cdot y}$, then using this function as a kernel we can obtain an integral operator

$$(Tf)(y) = \int f(x) e^{2\pi i x \cdot y} dx.$$

This is just the Fourier transform in disguise. One can define Tf directly by this integral (viewed as a Lebesgue integral) for any $f \in L^1(\mathbf{R}^d)$, in which the integrand will be absolutely integrable. We also know that for any $f \in L^1(\mathbf{R}^d)$,

$$\|Tf\|_{L^\infty(\mathbf{R}^d)} \leq \|f\|_{L^1(\mathbf{R}^d)}.$$

We also know from the classical Hausdorff-Young inequality that if $1 \leq p \leq 2$, then for any $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$,

$$\|Tf\|_{L^{p^*}(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)}.$$

In particular, this means that there exists a unique extension of T to a bounded operator from $L^p(\mathbf{R}^d)$ to $L^{p^*}(\mathbf{R}^d)$; note, however, that for a general element $f \in L^p(\mathbf{R}^d)$, the integral formula

$$\int f(x) e^{2\pi i \xi \cdot x} dx$$

is not well-defined in the Lebesgue sense. Thus we can only heuristically view the integral formula as defining the integral operator. There are two approaches to defining an integral operator T on more general classes of functions. The method above, is to work with a dense family of more regular functions for which the integral formula makes sense, and then to prove uniform estimates in this family of inputs. Another method is to replace the kernel in the integral formula T with a family of kernels converging to our original kernel, and then to prove uniform estimates in this family. For instance, one might instead study the family of truncated Fourier integrals

$$T_R f(y) = \int_{|x| \leq R} e^{2\pi i x \cdot y} f(x) dx$$

which are well defined for any $f \in L^1_{loc}(\mathbf{R}^d)$, and then prove that for $1 \leq p \leq 2$,

$$\|T_R f\|_{L^{p^*}(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)}$$

uniformly in R , which is sufficient justification to define Tf as the $L^p(\mathbf{R}^d)$ limit of $T_R f$ as $R \rightarrow \infty$, for $f \in L^p(\mathbf{R}^d)$, since this limit can be easily justified to exist for $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$.

19.1 Schur's Lemma

It is a useful heuristic that determining the boundedness of the operator T mapping *from* $L^1(X)$, or *into* $L^\infty(Y)$ is almost always trivial. This is one motivation for introducing the intermediate L^p norms, since these norms enable us to extract more features out of the kernel operator K . Before we discuss this, we must first reflect on the fact that without even qualitative knowledge of the kernel K besides its measurability, it is difficult to know how one might interpret the integral formula defining the operator. A natural trick to begin with is to introduce the sublinear analogue of the kernel operator, i.e. the operator S_K defined by setting

$$(S_K f)(y) = \int_X |K(x, y)| |f(x)| dx$$

The flexibility of the theory of non-negative Lebesgue integrals means this operator is well defined for *any* measurable f . Moreover, if we are to interpret the integral formula for $(T_K f)(y)$ in the Lebesgue sense, it is necessary and sufficient that $(S_K f)(y) < \infty$.

Theorem 19.1. *Fix $q \geq 1$. Then*

$$\|S_K f\|_{L^q(Y)} \leq \|K\|_{L^\infty(X)L^q(Y)} \|f\|_{L^1(X)}.$$

Thus $T_K f(y)$ are well defined by a Lebesgue integral for almost every $y \in Y$, and

$$\|T_K f\|_{L^q(Y)} \leq \|K\|_{L^\infty(X)L^q(Y)} \|f\|_{L^1(X)}.$$

Proof. The proof is a simple consequence of Minkowski's inequality, i.e.

$$\|S_K f\|_{L^q(Y)} = \|Kf\|_{L^q(Y)L^1(X)} \leq \|Kf\|_{L^1(X)L^q(Y)} \leq \|K\|_{L^\infty(X)L^q(Y)} \|f\|_{L^1(X)}. \square$$

Remark. In many situations, this result is tight. For instance, suppose

$$K = \sum_{i=1}^N \sum_{j=1}^M a_{ij} \mathbf{I}_{E_i \times F_j}$$

where E_1, \dots, E_N and F_1, \dots, F_N are disjoint finite measure sets. Then there exists $i \in \{1, \dots, N\}$ such that for each $x \in E_i$,

$$\left(\int |K(x, y)|^q dy \right)^{1/q} = \left(\sum_{j=1}^M |a_{ij}|^q |F_j| \right)^{1/q} = \|K\|_{L^\infty(X)L^q(Y)}.$$

If $f = \mathbf{I}_{E_i}$, then $\|f\|_{L^1(X)} = |E_i|$, and $T_K f = \sum_{j=1}^M a_{ij} \mathbf{I}_{F_j}$, so

$$\|T_K f\|_{L^q(Y)} = \left(\sum_{j=1}^M |a_{ij}|^q |F_j| \right)^{1/q} = \|K\|_{L^\infty(X)L^q(Y)} \|f\|_{L^1(X)}.$$

Thus we conclude that for a certain ‘dense’ family of kernels K , the inequality above is tight. This gives a strong heuristic that the inequality above is tight for a great many operators K , which trivializes the analysis of $L^1(X) \rightarrow L^q(Y)$ estimates. It is difficult to come up with a general statement of this form, because we require some regularity in the kernel to even formulate the operators under study. But by taking monotone limits we can at least justify that the norm of S_K from $L^1(X)$ to $L^q(Y)$ is given by $\|K\|_{L^\infty(X)L^q(Y)}$.

A dual statement trivializes the analysis of bounds from $L^p(X)$ to $L^\infty(Y)$.

Theorem 19.2. *Suppose $1 \leq p \leq \infty$. Then*

$$\|S_K f\|_{L^\infty(Y)} \leq \|K\|_{L^\infty(Y)L^{p^*}(X)} \|f\|_{L^p(X)}.$$

Thus if $\|K\|_{L^{p^}(X)L^\infty(Y)} < \infty$, then $T_K f(y)$ is well defined for almost every $y \in Y$, and*

$$\|T_K f\|_{L^\infty(Y)} \leq \|K\|_{L^\infty(Y)L^{p^*}(X)} \|f\|_{L^p(X)}.$$

Proof. One option to proving this bound is to take the adjoint of the kernel operator T_K and rely on previous estimates, but we can work more directly. Applying Hölder’s inequality, we conclude that

$$\|S_K f\|_{L^\infty(Y)} = \|K f\|_{L^\infty(Y)L^1(X)} \leq \|K\|_{L^\infty(Y)L^{p^*}(X)} \|f\|_{L^p(X)}. \quad \square$$

Though trivial, the two kernel bounds can often be applied together with an interpolation argument to give more sophisticated bounds.

Theorem 19.3 (Schur’s Test). *Fix $1 \leq r \leq \infty$, and suppose that*

$$\|K\|_{L^\infty(X)L^r(Y)} \leq A$$

and

$$\|K\|_{L^\infty(Y)L^r(X)} \leq B.$$

Then if $1 \leq p \leq q \leq \infty$ satisfy $1/p + 1/r = 1/q + 1$, then for $f \in L^p(X)$, $(T_K f)(y)$ is well defined by an absolutely convergent integral for almost every y , and

$$\|T_K f\|_{L^q(Y)} \leq A^{r/q} B^{1-r/q} \|f\|_{L^p(X)}.$$

Proof. The previous two results imply that S_K is bounded from $L^1(X)$ to $L^r(Y)$ and from $L^{r^*}(X)$ to $L^\infty(Y)$. Real interpolation (we cannot use complex interpolation since S_K is sublinear) shows that S_K is bounded from $L^p(X)$ to $L^q(Y)$ for the p and q satisfying the conditions above. This means $S_K f(x)$ is well defined for almost every $x \in X$, so the operator T_K is well defined by Lebesgue integrals for $f \in L^p(X)$. Applying the Riesz-Thorin interpolation theorem to T_K , which satisfies the bounds

$$\|T_K f\|_{L^r(Y)} \leq A \|f\|_{L^1(X)}$$

and

$$\|T_K f\|_{L^\infty(Y)} \leq B \|f\|_{L^{r^*}(X)},$$

we obtain the required result. \square

Example. Here is a version of Schur's Lemma that is more hands on and illustrative of the result. Let M be an $m \times n$ matrix, such that the sums of the absolute values of the entries of M on each row are upper bounded by A , and the sums of the absolute values of the entries of M on each column are upper bounded by B . Then for $x \in \mathbf{R}^n$, we have

$$|Mx| \leq \sqrt{AB} \cdot |x|.$$

Example. Young's Inequality can be justified by Schur's lemma, over any locally compact group G equipped with a Haar measure. Indeed, for a fixed g , the kernel of the operator $Cf = f * g$ is given by

$$K(x, y) = g(x^{-1}y)$$

and then

$$\|K\|_{L_x^\infty(G)L_y^r(G)} = \|g\|_{L^r(G)}$$

and

$$\|K\|_{L_y^\infty(G)L_x^r(G)} = \|g\|_{L^r(G)}$$

so Schur's Lemma shows that for $1/p + 1/r = 1/q + 1$,

$$\|f * g\|_{L^q(G)} = \|Cf\|_{L^q(G)} \leq \|f\|_{L^p(G)} \|g\|_{L^r(G)}.$$

Example. We cannot use Schur's Lemma to bound the fractional integration operators

$$T_1 f(y) = \int \frac{f(x)}{|x - y|^{d-s}} dx$$

since the kernel $K_1(x, y) = 1/|x - y|^{d-s}$ does not lie in any of the spaces $L^p(X)$. On the other hand, we can use Schur's Lemma to bound the modified fractional integration operators

$$T_2 f(y) = \int \frac{f(x)}{\langle x - y \rangle^{d-s}} dx,$$

where $\langle t \rangle = (1 + |t|^2)^{1/2}$ is the Japanese bracket. The kernel

$$K_2(x, y) = 1/\langle x - y \rangle^{d-s}$$

lies in $L_x^\infty(\mathbf{R}^d)L_y^r(\mathbf{R}^d)$ and $L_y^\infty(\mathbf{R}^d)L_x^r(\mathbf{R}^d)$ provided that $r > d/(d - s)$. Thus Schur's Lemma implies that for $1/p - 1/q > s/d$,

$$\|T_2 f\|_{L^q(\mathbf{R}^d)} \lesssim_{d,s,p,q} \|f\|_{L^p(X)}.$$

We will obtain a weak-type version of Schur's test which will enable us to bound T_1 later in this section.

For $1 < p < \infty$, we do not expect Schur's test to be sharp in general. But a good heuristic is that a variant of the inequality can be made to be sharp provided that two properties hold:

- For all $y \in Y$,

$$\int |K(x, y)| dx \approx A$$

and for all $x \in X$,

$$\int |K(x, y)| dy \approx B.$$

This kind of homogeneity will be present if $K(x, y) = k(x - y)$ is a convolution kernel.

- There is little oscillation in the kernel K .

Assuming for simplicity that X and Y have finite measure, from the first property we conclude that

$$A|Y| \approx B|X|.$$

Thus if we set $f = \mathbf{I}_X$, then $Tf(y) \approx A$ for all $y \in Y$, hence

$$\|Tf\|_{L^p(Y)} \approx A|Y|^{1/p} \approx A^{1-1/p} B^{1/p} |X|^{1/p}.$$

Thus we have tightness. If the second property remains true, but the first property fails, Schur's lemma still may remain sharp if we consider a weighted inequality, or alternatively, if we decompose the operator into components on which the marginal is approximately constant.

In some senses, if we are allowed to work with arbitrary weights, and if $K \geq 0$, Schur's test is always sharp. Suppose that

$$\|T_K f\|_{L^p(Y)} \leq A \|f\|_{L^p(X)}$$

for all $f \in L^p(X)$, and this inequality is sharp for some particular function f_0 . We may assume without loss of generality that $\|f_0\|_{L^p(X)} = 1$ and, since K is non-negative, that $f \geq 0$. Thus

$$\int_Y (T_K f_0(y))^p dy = A^p.$$

An application of Lagrangian multipliers and basic calculus of variations then shows that there exists a scalar λ such that

$$T_K^*((T_K f)^{p-1})(x) = \lambda f(x)^{p-1}.$$

But this means that

$$\begin{aligned} \lambda &= \int \lambda f(x)^p dx \\ &= \int f(x) T_K^*((T_K f)^{p-1})(x) dx \\ &= \int (T_K f)^p(x) = A^p. \end{aligned}$$

Thus if we set $w(x) = f(x)$ and $v(y) = (T_K f(y))^{p-1}$, then

$$\int_X K(x, y) w(x) dx = v(y)^{1/(p-1)}$$

and

$$\int_Y K(x, y) v(y) dy = A^p w(x)^{p-1}.$$

Using these estimates, a weighted variant of Schur's lemma, carried out in particular for the next example, gives the bound

$$\|T_K f\|_{L^p(X)} \leq A \|f\|_{L^p(X)},$$

which shows that the two weighted identities above contain as much information as the original bound.

Example. Let us consider an example where a weighted estimate can give better results about a particular operator. Consider the operator

$$Tf(y) = \frac{1}{\pi} \int_0^\infty \frac{f(x)}{x+y} dx.$$

The kernel of this operator is

$$K(x, y) = \frac{1}{\pi} \frac{1}{x+y}.$$

Since K is monotonically decreasing in y , maximizers for this inequality must be monotonically decreasing, with a singularity at the origin. The worst case singularity we can have near the origin while remaining close to being in L^2 is a singularity of the form $x^{-1/2}$. Thus we plug in $f(x) = x^{-1/2}$ into the operator T . Then we calculate that

$$\begin{aligned} Tf(y) &= \frac{1}{\pi} \int_0^\infty \frac{1}{x^{1/2}(x+y)} dx \\ &\sim \int_0^y \frac{1}{x^{1/2}y} + \int_y^\infty \frac{1}{x^{3/2}} dx \sim y^{-1/2}. \end{aligned}$$

Thus if we let $w(x) = x^{-1/2}$, and $v(y) = y^{-1/2}$, then we have already calculated that

$$\int K(x, y)[w(x)/v(y)] dx \lesssim 1.$$

Similarly,

$$\int K(x, y)[v(y)/w(x)] dy \lesssim 1.$$

Define

$$K_t(x, y) = K(x, y)w(x)^{2t-1}v(y)^{1-2t}.$$

Then $\{K_t\}$ is an analytic family of kernels giving us an analytic family of operators $\{T_t\}$, to which we can apply complex interpolation. The oscillation does not cause us a problem. For $t = 0$, we obtain that

$$\|T_0f\|_{L^1(\mathbf{R}^+)} \lesssim \|f\|_{L^1(\mathbf{R}^+)}.$$

For $t = 1$, we obtain that

$$\|T_1f\|_{L^\infty(\mathbf{R}^+)} \lesssim \|f\|_{L^\infty(\mathbf{R}^+)}.$$

For $t = 1/2$, we therefore obtain by interpolation that

$$\|T_{1/2}f\|_{L^2(\mathbf{R}^+)} \lesssim \|f\|_{L^\infty(\mathbf{R}^+)}.$$

But $T_{1/2} = T$, which means we have proved the required boundedness statement.

We can also get L^p bounds for all $1 < p < \infty$ (the case $p = 1$ or $p = \infty$ is not possible). To do this, we obtain a weak type bound from $L^1(\mathbf{R}^+)$ to $L^{1,\infty}(\mathbf{R}^+)$ by proving that for $f \in L^1(\mathbf{R}^+)$,

$$Tf(y) \lesssim \|f\|_{L^1(\mathbf{R}^+)}/y,$$

from which it follows that $\|Tf\|_{L^{1,\infty}(\mathbf{R}^+)} \lesssim \|f\|_{L^1(\mathbf{R}^+)}$. To prove this result, we note that for $f \geq 0$,

$$Tf(y) = \int \frac{f(x)}{x+y} \lesssim \int_0^y \frac{f(x)}{y} dx + \int_y^\infty \frac{f(x)}{x} dx \lesssim \|f\|_{L^1(\mathbf{R}^+)}/y + \|f\|_{L^1(\mathbf{R}^+)}/y.$$

Real interpolation gives bounds from $L^p(\mathbf{R}^+)$ to $L^p(\mathbf{R}^+)$ for $1 < p \leq 2$, and then duality (and the fact T is self-adjoint, apply boundedness for $2 \leq p < \infty$).

As another example of the trivial nature of the boundedness of kernel operators from $L^1(X)$, let us consider a version of Schur's Lemma that characterizes the boundedness from the $L^1(X)$ norm to a suitable family of Lorentz spaces, and from those Lorentz spaces to $L^\infty(Y)$.

Theorem 19.4. Fix $1 < q \leq \infty$ and $1 \leq s \leq \infty$. Then

$$\|S_K f\|_{L^{q,s}(Y)} \lesssim_{q,s} \|K\|_{L^\infty(Y)L^{q,s}(X)} \|f\|_{L^1(X)},$$

which means $T_K f$ is well-defined as a Lebesgue integral for $f \in L^1(X)$, and satisfies equivalent bounds. If $0 < p \leq \infty$, and $1 \leq s \leq \infty$, then

$$\|S_K f\|_{L^\infty(Y)} \lesssim_{p,s} \|K\|_{L^\infty(X)L^{p^*,s^*}(Y)} \|f\|_{L^{p,s}(X)}.$$

Thus $T_K f$ is well defined as a Lebesgue integral for $f \in L^{p,s}(X)$, and satisfies equivalent bounds. In particular, if $1 < r < \infty$, and

$$\|K\|_{L^\infty(Y)L^{r,\infty}(X)} \leq A,$$

and

$$\|K\|_{L^\infty(X)L^{r,\infty}(Y)} \leq B,$$

for some $1 < r < \infty$, then for $1 < p < q < \infty$ with $1/p + 1/r = 1/q + 1$, and any $0 < s \leq \infty$,

$$\|S_K f\|_{L^{q,s}(X)} \lesssim_{p,q,r,s} A^{r/q} B^{1-r/q} \|f\|_{L^{p,s}(Y)}.$$

Proof. The space $L^{q,s}(Y)$ is a norm space for $q > 1$, so we can employ Minkowski's inequality to conclude that

$$\|S_K f\|_{L^{q,s}(Y)} = \|S_K f\|_{L^{q,s}(Y)L^1(X)} \lesssim \|S_K f\|_{L^1(X)L^{q,s}(Y)}.$$

And then we conclude

$$\|S_K f\|_{L^1(X)L^{q,s}(Y)} \leq \|K\|_{L^\infty(X)L^{q,s}(Y)} \|f\|_{L^1(X)}$$

which proves the first claim. To obtain the second claim, we apply Hölder's inequality in the $L^{p,q}$ norms, which shows that

$$\|S_K f\|_{L^\infty(Y)} = \|K f\|_{L^\infty(Y)L^1(X)} \lesssim_{p,s} \|K\|_{L^\infty(Y)L^{p^*,s^*}(X)} \|f\|_{L^{p,s}(X)}. \quad \square$$

Remark. Establishing bounds from $L^1(X)$ to $L^{1,\infty}(Y)$ can be much more subtle than indicated by the other results here, e.g. for the Hardy-Littlewood Maximal function or for the Hilbert transform.

Example. For $s > 0$, and $f \in L^1(\mathbf{R}^d) \cap L^\infty(\mathbf{R}^d)$, we can define

$$Tf(x) = \int \frac{f(y)}{|y-x|^{d-s}} dy,$$

because $z \mapsto 1/|z|^{d-s}$ is integrable for $|z| \leq 1$, and bounded for $|z| \geq 1$. Applying the result above, since $1/|z|^{d-s}$ lies in $L^r(\mathbf{R}^d)$ for $r = d/(d-s)$, we conclude that for $1 < p < q < \infty$ with $s = d(1/p - 1/q)$,

$$\|Tf\|_{L^q(\mathbf{R}^d)} \lesssim_{d,s,p,q} \|f\|_{L^p(\mathbf{R}^d)}.$$

This is the Hardy-Littlewood-Sobolev inequality.

If T has kernel $K : X \times Y \rightarrow \mathbf{C}$ which is square integrable, then we find via Hölder's inequality that

$$\begin{aligned} \|Tf\|_{L^2(Y)}^2 &= \int_Y \left| \int_X K(x,y) f(x) dx \right|^2 dy \\ &\leq \int_Y \|K\|_{L_x^2}^2 \|f\|_{L_x^2}^2 dy \\ &= \|K\|_{L_x^2 L_y^2}^2 \|f\|_{L_x^2}^2. \end{aligned}$$

Thus $\|T\|_{L^2 \rightarrow L^2} \leq \|K\|_{L^2(X \times Y)}$. The quantity $\|K\|_{L^2(X \times Y)}$ is the *Hilbert-Schmidt*, or *Frobenius* norm of the operator T . If this inequality is tight at all steps for some input f , then tightness for Hölder's inequality implies that there exists $a(y)$ for almost every y such that $K(x, y) = a(y)\overline{f(x)}$, so that

$$Tg(y) = a(y)\langle g, f \rangle$$

is a rank one operator. Heuristically, this means that employing the Hilbert-Schmidt / Frobenius norm will only yield good results when T is a low rank operator. One of the main importances of the family of Hilbert-Schmidt integral operators is that they are *compact*, since

19.2 Change of Variables

TODO

On the other hand, it is not necessarily possible to change all variables in $\mathbf{R}^n \times \mathbf{R}^m$ without affecting the mapping properties of the operator. For instance, if T is the circular means operator from \mathbf{R}^d to \mathbf{R}^d , then T has kernel $K(x, y) = \delta(1 - |x - y|)$. If we change variables simultaneously, for instance, letting $x = z + w$ and $y = w$, then we obtain the kernel $K'(z, w) = K(z + w, w) = \delta(1 - |z|)$ is independent of w , which has completely different mapping properties, in particular, the output of the kernel does not actually depend on the input.

19.3 Localization In Space

Localization is a fundamental technique in analysis, since it enables us to isolate certain parts of a function or operator. If we understand these localized parts, one can then often recover results about the original result using a partition of unity. By doing this, we can isolate different features of a function to be controlled.

Many of the techniques of harmonic analysis are only possible once localized. In other words, rather than proving estimates of the form

$$\|Tf\|_{L^q(Y)} \lesssim \|f\|_{L^p(X)},$$

we only prove estimates of the form

$$\|T_{\phi_1, \phi_2} f\|_{L^q(Y)} \lesssim \|f\|_{L^p(X)}$$

where ϕ_1 and ϕ_2 are supported on finite measure subsets of X and Y , and $T_{\phi_1, \phi_2} f = \phi_1 T(\phi_2 f)$. Because of the localized nature of the problem, the difficulty of the bound increases as q becomes larger, and as p becomes smaller. If T has a kernel $K \in L^\infty(\text{supp}(\phi_1) \times \text{supp}(\phi_2))$, then T_{ϕ_1, ϕ_2} maps $L^1(\mathbf{R}^d)$ to $L^\infty(\mathbf{R}^d)$, which makes the study of the qualitative behaviour of T_{ϕ_1, ϕ_2} with respect to the convex norms trivial. Working quantitatively, we should therefore expect studying localized estimates to depend on quantitative measures of the singular nature of K , e.g. most crudely, the L^∞ norm. Because of the sensitivity to the singular nature of K , choosing ϕ_1 and ϕ_2 to be non-singular often makes the problem more amenable to analysis, e.g. if X and Y are smooth manifolds, it is often useful to assume $\phi_1 \in C_c^\infty(Y)$ and $\phi_2 \in C_c^\infty(X)$.

Intuitively, the smaller we make the kernel, the more well behaved the resulting operator will be. If K is a non-negative kernel, then it is certainly true that the L^p to L^q boundedness of an operator T with kernel K implies the boundedness of an operator T_Ω with kernel $K\mathbf{I}_\Omega$ for any $\Omega \subset X \times Y$, since we will then have a pointwise bound $|T_\Omega(f)| \leq T(|f|)$. But if K is not necessarily non-negative, then this is not necessarily true anymore, since T may only be bounded because of more subtle cancellation properties.

Example. Set $K(x, y) = e^{-2\pi i x \cdot y}$, and let $\Omega \subset \mathbf{R}^d \times \mathbf{R}^d$ be the set of all pairs (x, y) such that $\text{Re}(e^{-2\pi i x \cdot y}) \geq 0$. Then the operator T with kernel K is unitary, and thus bounded from $L^2(\mathbf{R}^d)$ to $L^2(\mathbf{R}^d)$. The operator T_Ω with kernel $K\mathbf{I}_\Omega$, on the other hand, is not bounded from $L^2(\mathbf{R}^d)$ to $L^2(\mathbf{R}^d)$, since, if $f = \mathbf{I}(|x| \leq R)$, then we will have $|T_\Omega f(y)| \gtrsim_d R^d$ for all $y \in \mathbf{R}^d$, so that $T_\Omega f \notin L^2(\mathbf{R}^d)$.

There are several cases where we can perform an arbitrary truncation. If $p = 1$, or $q = \infty$, then the operator norm of T with kernel K depends solely on mixed L^p norms of the kernel K , and thus behave well under truncation. Another case is if $\Omega = E \times F$, for then

$$\|T_\Omega f\|_{L^q(Y)} = \|\mathbf{I}_E T(\mathbf{I}_F f)\|_{L^q(Y)} \leq \|T(\mathbf{I}_F f)\|_{L^q(Y)} \leq \|\mathbf{I}_F f\|_{L^p(X)} \leq \|f\|_{L^p(X)}.$$

Thus $\|T_\Omega\|_{L^p \rightarrow L^q} \leq \|T\|_{L^p \rightarrow L^q}$. Similarly, if $X = \mathbf{R}^n$, $Y = \mathbf{R}^m$, $\Omega_1 \subset \mathbf{R}^n$ and $\Omega_2 \subset \mathbf{R}^m$ are precompact open sets, $L_1 : \mathbf{R}^n \rightarrow \mathbf{R}^n$ and $L_2 : \mathbf{R}^m \rightarrow \mathbf{R}^m$ are invertible linear maps, and $\phi : \mathbf{R}^n \times \mathbf{R}^m \rightarrow \mathbf{C}$ is a bump function adapted to $L_1(\Omega_1) \times L_2(\Omega_2)$, then

$$\|T_K \phi\|_{L^p \rightarrow L^q} \lesssim_{\Omega_1, \Omega_2} \|T_K\|_{L^p \rightarrow L^q}.$$

The trick here is to replace ϕ with a tensor product using a density argument / Fourier series which reduces our study to block diagonal type truncations.

More generally, if we have a family of disjoint sets $\{E_n\}$ and $\{F_n\}$, then we can truncate to the *block diagonal* region $\Omega = \bigcup (E_n \times F_n) = \bigcup \Omega_n$. The norm of such an operator can be calculated exactly from the behaviour on each subregion, as the next lemma shows.

Lemma 19.5. *Consider a family of Banach spaces $\{X_n\}$ and $\{Y_n\}$, and a family of bounded operators $T_n: X_n \rightarrow Y_n$. Then for $p \leq q$,*

$$\sup_{x_n \in X_n} \frac{\|T_n(x_n)\|_{l_n^q}}{\|x_n\|_{l_n^p}} = \sup_n \|T_n\|.$$

and if $p > q$, then

$$\sup_{x_n \in X_n} \frac{\|T_n(x_n)\|_{l_n^q}}{\|x_n\|_{l_n^p}} = \|T_n\|_{l_n^r},$$

where $1/p + 1/r = 1/q$.

Proof. We calculate that for $p \leq q$,

$$\|T_n(x_n)\|_{l_n^q} \leq \|T_n(x_n)\|_{l_n^p} \leq \|T_n\|_{l_n^\infty} \|x_n\|_{l_n^p}.$$

The converse bound for the supremum is simple. For $p > q$, Hölder's inequality implies that

$$\|T_n(x_n)\|_{l_n^q} \leq \| \|T_n\| \|x_n\| \|_{l_n^q} \leq \|T_n\|_{l_n^r} \|x_n\|_{l_n^p}.$$

Conversely, if we pick x_n with $\|x_n\| = 1$ for each n such that $\|T_n(x_n)\| \geq (1 - \varepsilon) \|T_n\|$, and if we pick $a_n = \|T_n\|^{r/q-1} = \|T_n\|^{r/p}$, then

$$\|T_n(a_n x_n)\|_{l_n^q} \geq (1 - \varepsilon) \| \|T_n\|^{r/q} \|_{l_n^q} \geq (1 - \varepsilon) \|T_n\|_{l_n^r}^{r/q},$$

whereas

$$\|a_n x_n\|_{l_n^p} = \| \|T_n\|^{r/p} \|_{l_n^p} = \|T_n\|_{l_n^r}^{r/p}.$$

Thus

$$\frac{\|T_n(a_n x_n)\|_{l_n^q}}{\|a_n x_n\|_{l_n^p}} \geq (1 - \varepsilon) \|T_n\|_{l_n^r}^{r/q-r/p} = (1 - \varepsilon) \|T_n\|_{l_n^r}.$$

Taking $\varepsilon \rightarrow 0$ gives the lower bound. □

A similar result holds for restriction to *almost* block diagonal operators, i.e. operators of the form

$$T = \sum_{(n,m) \in G} T_{nm}$$

where $T_{nm} : X_n \rightarrow Y_m$, and G is a graph on the index set such that each node has degree at most $O(1)$. One can then break the graph into $O(1)$ many block diagonal operators T_1, \dots, T_K , from which we obtain that e.g. for $p \leq q$,

$$\|T_{nm}x_n\|_{l_n^q} \lesssim \sup \|T_{nm}\|_{l_{nm}^\infty} \|x_n\|_{l_n^p}.$$

and for $p < q$,

$$\|T_{nm}x_n\|_{l_n^q} \lesssim \|T_{nm}\|_{l_{nm}^r} \|x_n\|_{l_n^p}.$$

On the other hand, we certainly have

$$\frac{\sup_x \|T_{nm}x_n\|_{l_n^q}}{\|x_n\|_{l_n^p}} \geq \|T_{nm}\|_{l_{nm}^\infty}.$$

TODO: Do we have a l^r lower bound as well, or not?

19.4 The Christ-Kiselev Lemma

We can also truncate to ‘upper triangular’ diagonals rather than just block diagonals, provided that $q > p$. For $q = p$, we ‘lose a logarithm’.

Theorem 19.6 (Christ-Kiselev). *Consider an operator T mapping functions on X to functions on Y , consider subsets $\{E_n\}$ of X and subsets $\{F_n\}$ of Y , and let*

$$\Omega = \bigcup_{n \leq m} E_n \times F_m.$$

Then if $1 \leq p < q \leq \infty$,

$$\|T_\Omega\|_{L^p \rightarrow L^q} \lesssim_{p,q} \|T\|_{L^p \rightarrow L^q}.$$

If $1 \leq p \leq \infty$, and n ranges over $\{1, \dots, N\}$, then

$$\|T_\Omega\|_{L^p \rightarrow L^p} \lesssim \log(N) \cdot \|T\|_{L^p \rightarrow L^p}.$$

Proof. Let us first address the case where $q > p$. We assume the index set is finite, i.e. $1 \leq n \leq N$ for some large $N > 0$, and prove the result by induction, though our constant will be independant of N and thus give us results for infinite index sets by taking limits. Fix a large constant $A_{p,q}$. Then the base case $N = 1$ is automatically satisfied if $A_{p,q}$ is large enough. Without loss of generality, assume $\|T\|_{L^p \rightarrow L^q} = 1$, and consider $f \in L^p(X)$ with $\|f\|_{L^p(X)} = 1$. Our proof is complete if we can show $\|T_\Omega f\|_{L^q(Y)} \leq A_{p,q}$. The idea here is to divide and conquer in an intelligent way. We can find an index $0 \leq n_0 \leq N$ such that

$$\|f \mathbf{I}_{E_{\leq n_0}}\|_{L^p(X)}^p \leq 1/2 < \|f \mathbf{I}_{E_{\leq n_0+1}}\|_{L^p(X)}^p.$$

Write $f_0 = f \mathbf{I}_{E_{\leq n_0}}$, $f_1 = f \mathbf{I}_{E_{n_0+1}}$, and $f_2 = f \mathbf{I}_{E_{\geq n_0+2}}$. Then $\|f_0\|_{L^p(X)}, \|f_2\|_{L^p(X)} \leq 1/2^{1/p}$. Applying induction, we have

$$\|\mathbf{I}_{F_{\leq n_0}} T_\Omega f_0\|_{L^q(Y)} \leq A_{p,q} 2^{-1/p}$$

and

$$\|T_\Omega f_2\|_{L^q(Y)} \leq A_{p,q} 2^{-1/p}.$$

On the other hand, the other restrictions are block diagonals, which gives

$$\|\mathbf{I}_{F_{\geq n_0+1}} T_\Omega f_0\|_{L^q(Y)} \leq 2^{-1/p},$$

and

$$\|T_\Omega f_1\|_{L^q(Y)} \leq \|f_1\|_{L^p(X)} \leq \|f\|_{L^p(X)} \leq 1.$$

The triangle inequality implies for $q > p$ that if $A_{p,q}$ is chosen large enough, depending on p and q ,

$$\|\mathbf{I}_{F_{\leq n_0}} T_\Omega f\|_{L^q(Y)} \leq (1 + A_{p,q}) 2^{-1/p} \leq 2^{(1/2)(1/p-1/q)} A_{p,q} 2^{-1/p},$$

$$\|\mathbf{I}_{F_{n_0+1}} T_\Omega f\|_{L^q(Y)} \leq 1 + 2^{1-1/p} \leq 3,$$

and

$$\|\mathbf{I}_{F_{\geq n_0+2}} T_\Omega f\|_{L^q(Y)} \leq 1 + (1 + A_{p,q}) 2^{-1/p} \leq 2^{(1/2)(1/p-1/q)} A_{p,q} 2^{-1/p}.$$

Putting these three estimates together, using the disjoint support, gives, again if $A_{p,q}$ is suitably large, depending on p and q , that

$$\begin{aligned} \|T_\Omega f\|_{L^q(Y)} &\leq (2(2^{(1/2)(1/p-1/q)})^q A_{p,q}^q 2^{-q/p} + 3^q)^{1/q} \\ &\leq (2^{-(1/2)(q/p-1)} A_{p,q}^q + 3^q)^{1/q} \leq A_{p,q}. \end{aligned}$$

This completes the proof in the case $q > p$.

Now consider the case $q = p$, and continue an induction on N , assuming without loss of generality that $\|T\|_{L^p \rightarrow L^q} = 1$. Write $A_p(N)$ for the optimal constant such that

$$\|T_\Omega\|_{L^p \rightarrow L^p} \leq A_p(N).$$

over all upper diagonal block operators of size $\leq N$. Consider f with $\|f\|_{L^p(X)} = 1$, and break $f = f_0 + f_1$, where $f_0 = f\mathbf{I}_{E_{\leq n_0}}$ and $f_1 = f\mathbf{I}_{E_{> n_0}}$, where $n_0, N - n_0 \leq N/2$. Let $\|f_0\|_{L^p(X)} = r$ and $\|f_1\|_{L^p(X)} = s$, so $r^p + s^p = 1$. Applying induction, we have

$$\|\mathbf{I}_{F_{\leq n_0}} T_\Omega f_0\|_{L^p(Y)} \leq A_p(N/2)r$$

and

$$\|T_\Omega f_1\|_{L^p(Y)} \leq A_p(N/2)s.$$

We also have

$$\|\mathbf{I}_{F_{\geq n_0}} T_\Omega f_0\|_{L^p(Y)} \leq 1.$$

Putting this together yields

$$\begin{aligned} \|T_\Omega f\|_{L^p(Y)} &\leq (A_p(N/2)^p r^p + (1 + A_p(N/2)s^p)^{1/p})^{1/p} \\ &\leq (A_p(N/2)^p (r^p + s^p) + O_p(A_p(N/2)^{p-1} s^{p-1}))^{1/p} \\ &= (A_p(N/2)^p + O_p(A_p(N/2)^{p-1} s^{p-1}))^{1/p} \\ &\leq (A_p(N/2)^p + O_p(A_p(N/2)^{p-1}))^{1/p} \\ &\leq A_p(N/2) + O_p(1). \end{aligned}$$

Thus we find $A_p(N) \leq A_p(N/2) + O_p(1)$, and this implies that $A_p(N) \lesssim_p \log(N)$, completing the proof. \square

The decomposition is arbitrary, which implies (via linearization) a variant by maximal functions.

Theorem 19.7 (Christ-Kiselev). *Consider $E_1 \subset E_2 \subset \dots \subset X$, let $T : L^p(X) \rightarrow L^q(Y)$, and consider the maximal operator*

$$Mf = \sup_n |T(\mathbf{I}_{E_n} f)|.$$

For $1 \leq p < q \leq \infty$,

$$\|M\|_{L^p \rightarrow L^q} \lesssim_{p,q} \|T\|_{L^p \rightarrow L^q}$$

and for $1 \leq p \leq \infty$, and $E_1 \subset E_2 \subset \cdots \subset E_N \subset X$ are given,

$$\|M\|_{L^p \rightarrow L^p} \lesssim \log(N) \|T\|_{L^p \rightarrow L^p}.$$

Proof. By monotone convergence, we may assume without loss of generality that the index set is finite. For $f \in L^p(X)$, and $y \in Y$, let $n(y)$ be an index such that $Mf(y) = T(\mathbf{I}_{E_{n(y)}}f)(y)$. Let $X_n = E_n - E_{n-1}$, and let $Y_n = \{y \in Y : n(y) = n\}$. Then for each $y \in Y$, if $\Omega = \bigcup_{n \leq m} X_n \times Y_m$,

$$Mf(y) = T(\mathbf{I}_{E_{n(y)}}f)(y) = T_\Omega f(y).$$

We can then apply the Christ-Kiselev Lemma above, since Ω is upper triangular. \square

We can apply this result to study the *maximal Fourier transform*

$$\mathcal{F}^*f(\xi) = \sup_I \int_I f(x) e^{-2\pi i \xi \cdot x} dx = \sup_I \widehat{f \mathbf{I}_I},$$

where I ranges over all compact intervals.

Theorem 19.8 (Menshov-Paley-Zygmund). *For $1 \leq p < 2$,*

$$\|\mathcal{F}^*f\|_{L^{p^*}(\mathbf{R})} \lesssim_p \|f\|_{L^p(\mathbf{R})}.$$

Proof. Applying the triangle inequality, we may assume one endpoint of each interval in the supremum lies at the origin. By symmetry, we may assume that this is the left hand endpoint. A convergence argument allows us to assume that the other endpoint is a positive rational number, and we may restrict ourselves by monotone convergence to a finite subset $\{q_1 < q_2 < \cdots < q_N\}$. But then the result follows by the Christ-Kiselev Maximal Function Lemma above, and the Hausdorff-Young inequality. \square

A qualitative consequence is that if $f \in L^p(\mathbf{R})$ for some $1 \leq p < 2$, then

$$\widehat{f}(\xi) = \lim_{a,b \rightarrow \infty} \int_{-a}^b f(x) e^{-2\pi i \xi \cdot x} dx,$$

for almost every $\xi \in \mathbf{R}$.

Remark. The result is also true for $p = 2$, but is a *much harder theorem*, due to Carleson.

The Christ-Kiselev Lemma and its ‘vector valued estimates’ becomes very useful in the study of nonlinear dispersive partial differential equations.

Chapter 20

Maximal Averages

This chapter is an introduction to the behaviour of basic averaging operators. A classical example, given a function $f \in L^1_{\text{loc}}(\mathbf{R})$, are the averaging operators

$$A_\delta f(x) = \frac{1}{2\delta} \int_{x-\delta}^{x+\delta} f(y) dy.$$

If $f \in C(\mathbf{R})$, then for each $x \in \mathbf{R}$, $\lim_{\delta \rightarrow 0} A_\delta f(x) = f(x)$. This fact is fundamentally connected to differentiation under the integral sign; if we define the function

$$F(x) = \int_0^x f(y) dy$$

then for each $x \in \mathbf{R}$,

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(y) dy = f(x).$$

Our main goal will be study whether pointwise convergence of the averages $A_\delta f$ hold in higher dimensions, for a more general family of functions that are not necessarily continuous, and for a more general family of averaging operators, thus testing the extent to which the ‘fundamental theorem of calculus’ holds.

The classical family of averaging operators on \mathbf{R}^d are defined for $\delta > 0$, $f \in L^1_{\text{loc}}(\mathbf{R}^d)$, and $x \in \mathbf{R}^d$ by setting

$$A_\delta f(x) = \oint_{B(x,\delta)} f(y) dy,$$

where $B(x, \delta)$ is the ball of radius δ centred at x . A simple application of Schur's lemma shows that $\|A_\delta f\|_{L^p(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)}$ for all $1 \leq p \leq \infty$, uniformly in p . This uniform bound in δ is strong enough, together with the density of compactly supported continuous functions is enough to conclude that for any $f \in L^p(\mathbf{R}^d)$, for $1 \leq p < \infty$, $A_\delta f$ converges to f in L^p norm. This implies that for any $f \in L^p(\mathbf{R}^d)$, there exists a sequence δ_i converging to zero such that $A_{\delta_i} f$ converges to f pointwise almost everywhere. In this chapter, we would like to show $A_\delta f$ converges to f pointwise almost everywhere *without taking a subsequence of values δ_i* . To do this, we introduce the fundamental tool of *maximal functions*.

20.1 The Method of Maximal Functions

Hardy and Littlewood introduced a powerful technique to study such pointwise convergence problems, known as the *method of maximal functions*. For each $f \in L^1_{\text{loc}}(X)$, we define a sublinear operator M by setting

$$Mf = \sup_{\delta > 0} A_\delta |f|.$$

The next theorem indicates why obtaining bounds on the operator M gives pointwise convergence results.

Theorem 20.1. *Let V be a quasinorm space, let $0 < q < \infty$, and consider a family of bounded operators $\{T_t : V \rightarrow L^{q,\infty}(X)\}$. Then we can define the pointwise maximal operator*

$$T_* f(x) = \sup_t |T_t f(x)|.$$

Suppose that for every $f \in L^p(X)$,

$$\|T_* f\|_{L^{q,\infty}(X)} \lesssim \|f\|_V.$$

Then for any bounded operator $S : V \rightarrow L^{q,\infty}(X)$, the set

$$\{f \in V : \lim_{t \rightarrow \infty} T_t f(y) = Sf(y) \text{ for a.e } y\}$$

is closed in V .

Proof. Fix $\{u_n\}$ in V converging to $u \in V$, and suppose for each n ,

$$\lim_{t \rightarrow \infty} (T_t u_n)(x) = S u_n(x)$$

holds for almost every $x \in X$. For each $\lambda > 0$, we find

$$\begin{aligned} & |\{x \in X : \limsup_{t \rightarrow \infty} |T_t u(x) - S u(x)| > \lambda\}| \\ & \leq |\{x \in X : \limsup_t |T_t(u - u_n)(x) - S(u - u_n)(x)| > \lambda\}| \\ & \leq |\{x \in X : |T_*(u - u_n)(x)| > \lambda/2\}| + |\{x : |S(u - u_n)(x)| > \lambda/2\}| \\ & \lesssim_{p,q} \frac{\|u - u_n\|_V^q}{\lambda^q} + \frac{\|u - u_n\|_V^p}{\lambda^p}. \end{aligned}$$

as $n \rightarrow \infty$, this quantity tends to zero. Thus for all $\lambda > 0$,

$$|\{x : \limsup_{t \rightarrow \infty} |T_t u(x) - S u(x)| > \lambda\}| = 0$$

Taking $\lambda \rightarrow 0$ gives that $\limsup_t |T_t u(x) - S u(x)| = 0$ for almost every $x \in X$. But this means precisely that $T_t u(x) \rightarrow S u(x)$ for almost every $x \in X$. \square

Since we know $A_\delta f$ converges to f pointwise for any $f \in C(\mathbf{R}^d)$, we see from the result above that taking $T_t = A_\delta$ and S the identity map gives almost everywhere convergence if we can obtain bounds for the maximal operator M of the form

$$\left\| \sup_{\delta > 0} A_\delta f \right\|_{L^{q,\infty}(\mathbf{R}^d)} \lesssim \|f\|_V$$

for an appropriate norm $\|\cdot\|_V$ and $0 < q < \infty$ in which $C(\mathbf{R}^d)$ is dense. We have already obtained a bound

$$\sup_{\delta > 0} \|A_\delta f\|_{L^{q,\infty}(\mathbf{R}^d)} \leq \sup_{\delta > 0} \|A_\delta f\|_{L^q(\mathbf{R}^d)} \leq \|f\|_{L^q(\mathbf{R}^d)}$$

but moving the supremum inside the L^q norm is nontrivial. One way to think about the difference between the two bounds is that the latter uniformly controls the height and width of the functions $A_\delta f$, whereas the former inequality not only controls the height and width of functions, but also shows that the main contributions to the height and widths of the functions $A_\delta f$ are supported on common regions of space.

20.2 Covering Methods

The bound $\|Mf\|_{L^\infty(\mathbf{R}^d)} \leq \|f\|_{L^\infty(\mathbf{R}^d)}$ follows from a direct calculation. This it is trivial to control the height of the function Mf in terms of the height of the function f . The difficult part is obtaining control of the width of Mf in terms of the width of f . Control on width can only be obtained up to a certain degree, because for any locally integrable $f \neq 0$, $\text{supp}(Mf) = \mathbf{R}^d$, so the width of f ‘explodes’. A slightly more technical calculation shows that we cannot even have a bound of the form

$$\|Mf\|_{L^1(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}.$$

In fact, Mf is not integrable for any nonzero $f \in L^1(\mathbf{R}^d)$.

Lemma 20.2. *If $f \in L^1(\mathbf{R}^d)$ is nonzero, then Mf is not integrable. Moreover, there exists $f \in L^1(\mathbf{R}^d)$ such that $Mf \notin L^1_{\text{loc}}(\mathbf{R}^d)$.*

Proof. Fix a nonzero $f \in L^1(\mathbf{R}^d)$. By rescaling, we may assume without loss of generality that $\|f\|_{L^1(\mathbf{R}^d)} = 2$. Then, for suitably large $R \geq 1$,

$$\int_{B_R(0)} |f(x)| \, dx \geq 1.$$

For each $x \in \mathbf{R}^d$, $B_R(0) \subset B_{|x|+R}(x)$ and so

$$Mf(x) \geq \int_{B_{|x|+R}(x)} |f(y)| \, dy \geq \frac{1}{(|x|+R)^d} \gtrsim \frac{1}{|x|^d}$$

But this means that

$$\int_{\mathbf{R}^d} |Mf(x)| \, dx \gtrsim \int_{\mathbf{R}^d} \frac{1}{|x|^d} = \infty.$$

Thus $Mf \notin L^1(\mathbf{R}^d)$.

To find f such that $Mf \notin L^1_{\text{loc}}(\mathbf{R}^d)$ take

$$f(x) = \frac{1}{|x|(\log|x|)^2}.$$

Then for $x \geq 0$,

$$\begin{aligned} \frac{1}{2h} \int_{x-h}^{x+h} \frac{dy}{|y| \log |y|^2} &= \frac{1}{2h} \left(\frac{1}{\log(x-h)} - \frac{1}{\log(x+h)} \right) \\ &= \frac{1}{2x \log x} + O\left(\frac{h}{\log x}\right) \end{aligned}$$

implies that

$$Mf(x) \geq \frac{1}{2x \log x}.$$

Thus Mf isn't integrable near the origin. \square

The last lemma shows that if $f \in L^1(\mathbf{R}^d)$, then we have a pointwise bound $|Mf(x)| \gtrsim_f \langle x \rangle^{-d}$. Note, however, that $|x|^{-d}$ is only *barely* nonintegrable. We will also show that Mf is barely nonintegrable by obtaining a bound

$$\|Mf\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim_d \|f\|_{L^1(\mathbf{R}^d)}.$$

Interpolation thus shows that $\|Mf\|_{L^p(\mathbf{R}^d)} \lesssim_{d,p} \|f\|_{L^p(\mathbf{R}^d)}$ for all $1 < p \leq \infty$.

The standard real-variable technique of obtaining the $L^1(\mathbf{R}^d) \rightarrow L^{1,\infty}(\mathbf{R}^d)$ bound of the maximal function is geometric, applying a covering argument. To obtain the weak-type bound, we must show that the set

$$E_\lambda = \{x \in \mathbf{R}^d : |Mf(x)| > \lambda\}$$

is small. If $|Mf(x)| > \lambda$, there is a ball B around x such that

$$\int_B |f(y)| dy > \lambda |B|.$$

Clearly $B \subset E_\lambda$. If we could find a large family of *disjoint balls* B_1, \dots, B_N such that this inequality held, such that $\sum |B_i| \gtrsim_d |E_\lambda|$, then we would conclude that

$$\|f\|_{L^1(\mathbf{R}^d)} \geq \sum_{i=1}^N \int_{B_i} |f(y)| dy > \lambda \sum_{i=1}^N |B_i| \gtrsim_d \lambda |E_\lambda|$$

which would show $|E_\lambda| \lesssim_d \|f\|_{L^1(\mathbf{R}^d)}/\lambda$, i.e. that $\|Mf\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim_d \|f\|_{L^1(\mathbf{R}^d)}$. This intuition is true, and the process through which we obtain the family of disjoint balls B_1, \dots, B_N is through the *Vitali covering lemma*.

This particular technique has been shown to generalize to a wide variety of situations including the maximal ball average. All that is really required for the basic theory is a basic ‘covering type argument’ that holds in a great many situations. We consider a locally compact space X equipped with a nonzero Radon measure. For each $x \in X$ and $\delta > 0$, we fix an open, precompact set $B(x, \delta)$, which we assume to be monotonically increasing in δ . The fundamental property we require of these sets is that there is $c > 0$ such that for any $x \in X$ and $\delta > 0$, if we set

$$B^*(x, \delta) = \bigcup \{B(x', \delta) : B(x, \delta) \cap B(x', \delta) \neq \emptyset\},$$

then $|B^*(x, \delta)| \leq c|B(x, \delta)|$.

In the case of balls in \mathbf{R}^d , $B^*(x, \delta) \subset B^*(x, 3\delta)$, and so $|B^*(x, \delta)| \leq 3^d|B(x, \delta)|$, so we can set $c = 3^d$. More generally, if we are working in any metric space X , where $B(x, \delta)$ are the balls of radius δ in this metric space, and our measure satisfies a *doubling condition*

$$|B(x, 3\delta)| \leq c|B(x, \delta)|$$

for all $x \in X$ and $\delta > 0$, then our assumption holds. The following are not strictly necessary for the argument to work, but are technical assumptions which make certain parts of the argument cleaner, and hold in most cases of application:

- For any $x \in X$,

$$\bigcap_{\delta>0} \overline{B}(x, \delta) = \{x\} \quad \text{and} \quad \bigcup_{\delta>0} B(x, \delta) = X$$

- For any open set $U \subset X$ and $\delta > 0$, the function

$$x \mapsto |B(x, \delta) \cap U|$$

is a continuous function of x .

These are fairly easily verifiable in most particular instance, but in some situations they can be worked around to obtain similar results that we will obtain here. It follows from these technical assumptions that $|B(x, \delta)| > 0$ for each $x \in X$ and $\delta > 0$, and moreover, for each $\delta > 0$, and $f \in L^1_{\text{loc}}(X)$, the averaged function $A_\delta f$ given by setting

$$A_\delta f(x) = \frac{1}{|B(x, \delta)|} \int_{B(x, \delta)} f(y) \, dy,$$

is measurable.

Lemma 20.3. *If $f \in L_1^{loc}(X)$, then $A_\delta f$ is a measurable function.*

Proof. If $f = a_1 \mathbf{I}_{U_1} + \cdots + a_N \mathbf{I}_{U_N}$ is a simple function, where U_1, \dots, U_N are open sets, then

$$A_\delta f(x) = a_1 \frac{|B(x, \delta) \cap U_1|}{|B(x, \delta)|} + \cdots + a_N \frac{|B(x, \delta) \cap U_N|}{|B(x, \delta)|}$$

is a continuous function by our technical assumptions. Next, if $f \geq 0$ is a step function, then there exists a monotonically decreasing family of simple functions $\{f_n\}$ such that $f_n \rightarrow f$ pointwise, then the monotone convergence theorem implies that $A_\delta f_n \rightarrow A_\delta f$ pointwise, so $A_\delta f$ is measurable. Finally, decomposing any measurable function into the difference of non-negative measurable functions and then considering pointwise limits of step functions completes the proof. \square

It also follows from our technical assumptions that for any open set U containing x , there exists δ_0 such that for $\delta \leq \delta_0$, $\overline{B(x, \delta)} \subset U$. It follows that for any $f \in C(X)$ and $x \in X$,

$$\lim_{\delta \rightarrow 0} A_\delta f(x) = f(x).$$

If $Mf = \sup_{\delta > 0} A_\delta f$, then we will show

$$\|Mf\|_{L^{1,\infty}(X)} \lesssim_c \|f\|_{L^1(X)}.$$

In particular, this shows that for any $f \in L^1(X)$,

$$\lim_{\delta \rightarrow 0} A_\delta f(x) = f(x)$$

for almost every $x \in X$. Since this result is a *local result*, it is easy to verify that the result also holds for any $f \in L_{loc}^1(X)$, i.e. it also holds for any $f \in L^p(X)$ for $1 \leq p \leq \infty$.

Lemma 20.4 (Vitali Covering Lemma). *If B_1, \dots, B_n is a finite collection of balls in X , then there is a disjoint subcollection B_{i_1}, \dots, B_{i_M} such that*

$$\left| \bigcup_{i=1}^N B_i \right| \leq c \sum_{j=1}^M |B_{i_j}|.$$

Proof. Consider the following greedy selection procedure. Let B_{i_1} be the ball in our collection of maximal radius. Given that we have selected B_{i_1}, \dots, B_{i_k} , let $B_{i_{k+1}}$ be the ball of largest radius not intersecting previous balls selected if possible. Continue doing this until we cannot select any further balls. If B_j is any ball not chosen by this procedure, it must intersect a ball with radius at least as big as B_j itself. But this means that

$$\bigcup_{i=1}^N B_i \subset \bigcup_{j=1}^M B_{i_j}^*.$$

Thus

$$\left| \bigcup_{i=1}^N B_i \right| \leq \sum_{j=1}^M |B_{i_j}^*| \leq c \sum_{j=1}^M |B_{i_j}|. \quad \square$$

We have already indicated our proof strategy for proving a weak type bound for the maximal operator, but let us now do things more rigorously.

Theorem 20.5. *For any $f \in L^1(X)$,*

$$\|Mf\|_{L^{1,\infty}(X)} \leq c \|f\|_{L^1(X)}.$$

Proof. Set

$$E_\lambda = \{x \in \mathbf{R}^d : Mf(x) > \lambda\}.$$

Fix a compact subset K of E_λ of finite measure. Then K is covered by finitely many balls B_1, \dots, B_N such that on each ball B_i ,

$$\int_{B_i} |f(y)| \, dy > \lambda |B_i|.$$

Using the Vitali lemma, extract a disjoint subfamily B_{i_1}, \dots, B_{i_M} with

$$\left| \sum_{j=1}^M B_{i_j} \right| \leq c \sum_{j=1}^M |B_{i_j}|.$$

Then

$$\|f\|_{L^1(X)} > \lambda \sum_{j=1}^M |B_{i_j}| \geq \frac{\lambda}{c} \left| \bigcup_{j=1}^M B_{i_j} \right| \geq \frac{\lambda |K|}{c}.$$

Rearranging gives

$$|K| \leq \frac{c\|f\|_{L^1(X)}}{\lambda}.$$

Since K was arbitrary, inner regularity gives

$$|E_\lambda| \leq \frac{c\|f\|_{L^1(X)}}{\lambda}.$$

Since λ was arbitrary, the proof is complete. \square

Remark. The same covering-type argument also gives the boundedness of the *uncentered* Hardy-Littlewood maximal function

$$M'f(x) = \sup_{x \in B} \frac{1}{|B|} \int_B |f(y)| \, dy$$

where B ranges over all balls.

The maximal operators, in the general assumptions above, can behave quite differently. We have the pointwise inequality $Mf \leq M'f$, but in general $M'f$ can be significantly bigger than Mf . However, we have $M'f \lesssim Mf$ under the following slightly stronger assumption on the family of balls we are working with: that there exists two constants $c_1, c_2 > 0$ such that the following two properties hold:

- (The Engulfing Condition) If $B(x, \delta) \cap B(x', \delta) \neq \emptyset$, then $B(x', \delta) \subset B(x, c_1 \delta)$.
- (The Doubling Condition) $|B(x, c_1 \delta)| \leq c_2 |B(x, \delta)|$.

If $B = B(x', \delta)$ contains x , then B is contained in $B(x, c_1 \delta)$, and conversely, $B(x, \delta)$ is contained in $B(x', c_1 \delta)$. Thus

$$|B| = |B(x', \delta)| \geq c_2^{-1} |B(x', c_1 \delta)| \geq c_2^{-1} |B(x, \delta)| \geq c_2^{-2} |B(x, c_1 \delta)|.$$

Thus

$$\frac{1}{|B|} \int_B |f(y)| \, dy \leq \frac{c_2^2}{|B(x, c_1 \delta)|} \int_{B(x, c_1 \delta)} |f(y)| \, dy \leq c_2^2 Mf(x).$$

Taking suprema gives $M'f(x) \leq c_2^2 Mf(x)$. Thus under these slightly stronger conditions, M and M' are roughly comparable operators. These assumptions will reappear when we analyze Calderon-Zygmund decomposition type arguments that come up in the real-variable analysis of singular integrals.

Remark. We can exploit the ordering of the real line to show that for any family of intervals $\{I_\alpha\}$ covering a compact set K , there is a sub-cover I_1, \dots, I_N such that any point in \mathbf{R} is contained in at most two of the intervals. A modification of the argument above shows this gives the slightly better bound $\|Mf\|_{L^{1,\infty}(\mathbf{R})} \leq 2\|f\|_{L^1(\mathbf{R})}$, rather than the bound $\|Mf\|_{L^{1,\infty}(\mathbf{R})} \leq 3\|f\|_{L^1(\mathbf{R})}$.

With the result proved, we obtain the Lebesgue differentiation theorem: If $f \in L^{1,\text{loc}}(\mathbf{R}^d)$, then for almost every $x_0 \in \mathbf{R}^d$,

$$f(x_0) = \lim_{\delta \rightarrow 0} \int_{|x-x_0| \leq \delta} f(x) dx.$$

For $d = 1$, we can directly connect this to the fundamental theorem of calculus; if $f \in L^{1,\text{loc}}(\mathbf{R}^d)$, then the function

$$g(x) = \int_0^x f(y) dy$$

is locally Lipschitz, and conversely we can find such f for any g by the Radon-Nikodym theorem. The Lebesgue differentiation theorem then says precisely that if g is locally Lipschitz, then g is differentiable almost everywhere, and that

$$g(x) = g(0) + \int_0^x g'(x) dx.$$

This result is called the *Rademacher differentiation theorem*. We develop similar ideas, and in higher dimensions, in the next chapter.

20.3 Dyadic Methods

There are many different techniques for showing the boundedness of the maximal operator. Let us consider some *dyadic methods* for proving the inequality. Recall that the set of dyadic cubes is

$$\{Q_{n,k} : n \in \mathbf{Z}, k \in 2^n \mathbf{Z}^d\}$$

where $Q_{n,k}$ is the cube $[k_1, k_1 + 2^n] \times \dots \times [k_d, k_d + 2^n]$. We note that dyadic cubes nest within one another much more easily than balls do (cubes are either nested or disjoint). In particular, if Q_1, \dots, Q_N is any collection of

dyadic cubes, there exists an almost disjoint subcollection Q_{i_1}, \dots, Q_{i_k} with $Q_{i_1} \cup \dots \cup Q_{i_k} = Q_1 \cup \dots \cup Q_N$. In particular, this operates as a Vitali-type covering lemma with a constant independent of d , so if we define the *dyadic* Hardy-Littlewood maximal operator

$$M_{\Delta}f(x) = \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f(y)| dy$$

then we easily obtain the bound $\|M_{\Delta}f\|_{L^{1,\infty}(\mathbf{R}^d)} \leq \|f\|_{L^1(\mathbf{R}^d)}$, with no implicit constant depending on d . The bound $\|M_{\Delta}f\|_{L^{\infty}(\mathbf{R}^d)} \leq \|f\|_{L^{\infty}(\mathbf{R}^d)}$ is easy, so interpolation gives $\|M_{\Delta}f\|_{L^p(\mathbf{R}^d)} \lesssim_p \|f\|_{L^p(\mathbf{R}^d)}$ for all $1 < p \leq \infty$, with a constant now *independent of dimension*.

Two families of sets $\{B(x, \delta) : x \in \mathbf{R}^d, \delta > 0\}$ and $\{B'(x, \delta) : x \in \mathbf{R}^d, \delta > 0\}$ are *equivalent* if there exists c_1, c_2 such that

$$B'(x, c_1\delta) \subset B(x, \delta) \subset B(x, c_2\delta).$$

It follows that the resultant maximal averages from the two sets pointwise differ from one another by a universal constant. This allows us to obtain bounds for maximal averages over cubes centred at a point, and ellipses with bounded eccentricity, etc. If for $2^k \leq \delta \leq 2^{k+1}$, we set $B(x, \delta)$ to be a dyadic cube with sidelength $1/2^k$, then the family $\{B(x, \delta)\}$ is *not* equivalent to the usual family of cubes, but below we show that bounds on the two maximal operators are still equivalent.

If Q is a dyadic cube, then it is contained in a ball B with $|Q| \lesssim_d |B|$. It follows that for any function f and $x \in \mathbf{R}^d$,

$$M_{\Delta}f(x) \lesssim_d Mf(x).$$

Thus bounds on M automatically give bounds on M_{Δ} . The opposite pointwise inequality is unfortunately, *not true*. For instance, if f is the indicator function on $[0, 1]$. Then $M_{\Delta}f$ is supported on $[0, 1]$, but Mf is positive on the entirety of \mathbf{R} . To reduce the study of M to the study of M_{Δ} , we must instead rely on the *1/3 translation trick* of Michael Christ.

Lemma 20.6. *Let $I \subset [0, 1]$ be an interval. Then there exists an interval J , which is either a dyadic interval, or a dyadic interval shifted by $1/3$, such that $I \subset J$ and $|J| \lesssim |I|$.*

Proof. Let $I = [a, b]$. Perform a binary expansion of a and b , writing

$$a = 0.a_1a_2\dots \quad \text{and} \quad b = b_1b_2\dots$$

Let n be the first value where $a_n \neq b_n$. Then $a_n = 0$ and $b_n = 1$. Then $[a, b]$ is contained in the dyadic interval

$$Q_1 = [0.a_1\dots a_{n-1}, 0.a_1\dots a_{n-1} + 1/2^{n-1}]$$

which has length $1/2^{n-1}$. Find $0 \leq i < \infty$ such that

$$a = 0.a_1\dots a_{n-1}01^i0\dots$$

and $0 \leq j < \infty$ such that

$$b = 0.a_1\dots a_{n-1}10^j1.$$

If no such j exists, then $b = 0.a_1\dots a_{n-1}1$, and so $[a, b]$ is contained in the rational interval

$$Q_2 = [0.a_1\dots a_{n-1}01^i, 0.a_1\dots a_{n-1}01^i + 1/2^{n+i}]$$

and $b - a \geq 1/2^{n+i+1}$, so $|Q_2| \leq 2(b - a)$. Now if $i \leq 5$ or $j \leq 5$, then $b - a \geq 1/2^{n+5}$, so $|Q_1| \leq 2^5(b - a)$. On the other hand, if $i \geq 5$ and $j \geq 5$, we find $b - a \geq 1/2^{n+\min(i,j)}$. Then we can find a dyadic interval Q_3 and $2 \leq r \leq 5$ such that

$$1/3 + Q_3 = [0.a_1\dots a_{n-1}01^{\min(i,j)-r}1010\dots, 0.a_1\dots a_{n-1}01^{i-r}1010\dots + 1/2^{n+\min(i,j)-r}]$$

and so $1/3 + Q_3$ contains $[a, b]$ and $|Q_3| = 1/2^{n+\min(i,j)-r} \leq 2^5(b - a)$. \square

It follows that for each $x \in \mathbf{R}^d$, and any function f ,

$$Mf(x) \lesssim_d (M_\Delta f)(x) + (M_\Delta \text{Trans}_{1/3} f)(x).$$

Since the L^p norms are translation invariant, this implies that the dyadic maximal operator and the maximal operator satisfy equivalent bounds, with operator norms differing by a constant depending on n . Since we independently obtained bounds on M_Δ , this section provides an alternate proof to the boundedness of M .

There is an alternate way to view the operator M_Δ . For each integer n , we let $\mathcal{B}(n)$ denote the family of all sidelength $1/2^n$ dyadic cubes. Thus $\mathcal{B}(n)$ gives a decomposition of \mathbf{R}^d into an almost disjoint union of cubes. If we define the conditional expectation operators

$$E_n f(x) = \sum_{Q \in \mathcal{B}(n)} \left(\frac{1}{|Q|} \int_Q f \right) \cdot \mathbf{I}_Q$$

then $M_\Delta f = \sup_{n \in \mathbf{Z}} E_n f$. In particular, it is easy to see from the bounds on M_Δ that for any $f \in L^1_{\text{loc}}(\mathbf{R}^d)$, $\lim_{n \rightarrow \infty} E_n f(x) = f(x)$ holds for almost every $x \in \mathbf{R}^d$. It is simple to conclude from this result a very useful technique, known as the *Calderón-Zygmund decomposition*.

Theorem 20.7. *Given $f \in L^1(\mathbf{R}^d)$ and $\lambda > 0$, we can write $f = g + b$, where $\|g\|_{L^\infty(\mathbf{R}^d)} \lesssim_d \lambda$, and there is an almost disjoint family of dyadic cubes $\{Q_i\}$ such that g is supported on $\bigcup_i Q_i$,*

$$\sum_i |Q_i| \leq \frac{\|f\|_{L^1(\mathbf{R}^d)}}{\lambda},$$

and for each i ,

$$\int_{Q_i} f(y) dy = 0.$$

We also have $\|g\|_{L^1(\mathbf{R}^d)}, \|b\|_{L^1(\mathbf{R}^d)} \lesssim_d \|f\|_{L^1(\mathbf{R}^d)}$.

Proof. Write $E = \{x : M_\Delta f(x) > \lambda\}$. By the dyadic Hardy-Littlewood maximal inequality,

$$|E| \leq \frac{\|f\|_{L^1(\mathbf{R}^d)}}{\lambda}.$$

Because f is integrable, $E \neq \mathbf{R}^d$. Thus we can write E as the almost disjoint union of dyadic cubes $\{Q_i\}$, such that for each i ,

$$\int_{Q_i} |f(x)| dx > \lambda |Q_i|,$$

and also, if R_i is the parent cube of Q_i ,

$$\int_{R_i} |f(x)| dx \leq \lambda |R_i|.$$

This can be done by a greedy strategy, taking the union of dyadic cubes of largest sidelength contained in E . This means

$$\int_{Q_i} |f(x)| dx \leq \int_{R_i} |f(x)| dx \leq \lambda |R_i| \leq 2^d \lambda |Q_i|.$$

Define

$$g(x) = \begin{cases} f(x) & : x \notin E, \\ \frac{1}{|Q_i|} \int_{Q_i} f(x) dx & : x \in Q_i \text{ for some } i. \end{cases}$$

For almost every $x \in E^c$, $|f(x)| \leq \lambda$, since $E_n f(x) \leq \lambda$ for each n , and $E_n f(x) \rightarrow f(x)$ as $n \rightarrow \infty$ for almost every x . Conversely, if $x \in Q_i$ for some i , then

$$\left| \frac{1}{|Q_i|} \int_{Q_i} f(x) dx \right| \leq \frac{1}{|Q_i|} \int_{Q_i} |f(x)| dx \leq 2^d \lambda.$$

Thus $\|g\|_{L^\infty(\mathbf{R}^d)} \lesssim_d \lambda$. If we define $b = f - g$, then b is supported on $\bigcup Q_i = E$, and for each i ,

$$\int_{Q_i} b(x) dx = \int_{Q_i} \left(f(x) - \frac{1}{|Q_i|} \int_{Q_i} f(y) dy \right) dx = 0. \quad \square$$

The Calderon-Zygmund theorem will be very useful to us in the sequel, especially when we analyze the theory of singular integrals.

20.4 Linearization of Maximal Operators

It is often a very useful technique to study a maximal operator by reducing its analysis to studying a family of linear operators. The technique is very general, and applies to virtually all the maximal functions, so we give the general discussion here. Fix a family of linear operators $\{T_t\}$, and consider the associated maximal operator $Mf = \sup_t |T_t f|$. By monotone convergence, we can obtain $L^p \rightarrow L^q$ operator norm bounds on the maximal operator M if we can prove uniform $L^p \rightarrow L^q$ operator norm bounds on the maximal operators $Mf = \sup_{t \in T} |T_t f|$, where $T = \{t_1, \dots, t_N\}$ is a finite subset of indices. For such an operator, and each $y \in Y$, there exists

$t(y) \in T$ such that $Mf(y) = T_{t(y)}f(y)$. Thus we can divide Y into N disjoint sets $Y_1 \cup \dots \cup Y_N$, such that

$$Mf = \sum_{i=1}^N \mathbf{I}_{Y_i} T_i f.$$

To bound the *sublinear* operator M , it therefore suffices to obtain uniform bounds on the behaviours of the family of *linear operators* of the form

$$\sum_{i=1}^N \mathbf{I}_{Y_i} T_i.$$

This is the method of linearization. TODO: Are there any measurability issues here?

20.5 TT^* Arguments

The method of TT^* arguments enables us to obtain bounds on a linear operator T by exploiting cancellation when the operator is composed with its adjoint. The main calculation from which the method follows is that if H is a Hilbert space, and X is a Banach space, and $T : H \rightarrow X$ is a bounded operator, then

$$\|TT^*\|_{X^* \rightarrow X} = \|T\|_{H \rightarrow X}^2,$$

because certainly $\|T^*\|_{X^* \rightarrow H} = \|T\|_{H \rightarrow X}$. For any $f \in X^*$,

$$\|T^*f\|_H^2 = \langle T^*f, T^*f \rangle = \langle TT^*f, f \rangle \leq \|TT^*\|_{X^* \rightarrow X} \|f\|_{X^*}^2,$$

so $\|T\|^2 \leq \|TT^*\|$, and conversely,

$$\|TT^*\| \leq \|T\| \|T^*\| = \|T\|^2.$$

Thus we study the boundedness of an operator T via the boundedness of the associated operator TT^* .

By linearity, to obtain bounds on the Hardy-Littlewood-Maximal operator it suffices to obtain uniform bounds on the behaviour of the linear operators of the form

$$T_r f(y) = A_{r(y)} f(y),$$

where $r(y) \in \{r_1, \dots, r_N\} \subset (0, \infty)$ are radii. Now

$$T_r^* g(x) = \int_{|x-y| \leq r(y)} \frac{g(y)}{|B(y, r(y))|} dy$$

Thus

$$\begin{aligned} T_r T_r^* g(y) &= A_{r(y)} T^* g(y) \\ &= \frac{1}{V_d^2} \int_{|x-y| \leq r(y)} \int_{|x-z| \leq r(z)} r(y)^{-d} r(z)^{-d} g(z) dx dz. \end{aligned}$$

On the integrand, we must have $|z - y| \leq r(y) + r(z)$, and x is constrained to lie in a ball of radius $\min(r(y), r(z))$. Thus

$$|T_r T_r^* g(y)| \leq \frac{1}{V_d} \int_{|y-z| \leq r(y)+r(z)} \frac{g(z)}{\max(r(y), r(z))^d} dz.$$

We can split this integral into two parts, depending on whether $r(y) \geq r(z)$ or $r(y) \leq r(z)$, which yields

$$\begin{aligned} |T_r T_r^* g(y)| &\leq \frac{1}{V_d} \int_{|y-z| \leq 2r(y)} \frac{g(z)}{r(y)^d} dy + \int_{|y-z| \leq 2r(z)} \frac{g(z)}{r(z)^d} dy \\ &\leq 2^d (T_{2r} g(y) + T_{2r}^* g(y)). \end{aligned}$$

Thus if $C_N = \sup_r \|T_r\|_{L^2 \rightarrow L^2}$, where the supremum is over functions taking over at most N values, then we conclude that

$$\|T_r T_r^* g(y)\|_{L^2} \leq 2^d (\|T_{2r} g\|_{L^2} + \|T_{2r}^* g\|_{L^2}) \leq 2^{d+1} C_N,$$

so that for any r , we find that $\|T_r\|_{L^2 \rightarrow L^2}^2 = \|T_r T_r^*\|_{L^2 \rightarrow L^2}^{1/2} \leq 2^{d+1} C_N$. Thus $C_N^2 \leq 2^{d+1} C_N$, and since $C_N < \infty$ (we have a trivial bound $C_N \lesssim N$ because the operator norm of the averages is uniformly bounded), this gives $C_N \leq 2^{d+1}$, which shows $\|Mf\|_{L^2(\mathbf{R}^d)} \leq 2^{d+1} \|f\|_{L^2(\mathbf{R}^d)}$.

20.6 Lebesgue Density Theorem

If E is a measurable subset of \mathbf{R}^d , and $x \in \mathbf{R}^d$, we say x is a point of *Lebesgue density* of E , or has *full metric density* if

$$\lim_{\delta \rightarrow 0} \frac{|B(x, \delta) \cap E|}{|B(x, \delta)|} = 1$$

This means that for any $\varepsilon > 0$, the inequality $|B(x, \delta) \cap E| \geq (1 - \varepsilon)|B(x, \delta)|$ holds for suitably small δ , so E asymptotically contains as large a fraction of the local points around x as is possible. Since $\chi_E \in L^1_{\text{loc}}(\mathbf{R}^d)$, we can apply the Lebesgue differentiation theorem to immediately obtain an interesting result.

Theorem 20.8 (Lebesgue Density Theorem). *If E is a measurable subset, then almost every point in E is a point of Lebesgue density, and almost every point not in E is not a point of Lebesgue density.*

The fact that a point is a point of Lebesgue density implies the existence of large sets of rigid patterns in E . A simple corollary is that any set of positive Lebesgue measure contains arbitrarily long arithmetic progressions.

Theorem 20.9. *Let $E \subset \mathbf{R}^d$ be a set of nonzero Lebesgue measure. Then for any non-zero $a_1, \dots, a_N \in \mathbf{R}$ there exists $x \in E$ and $c \in \mathbf{R}$ such that*

$$a_1x + c, \dots, a_Nx + c \in E.$$

Proof. Without loss of generality, by translation we may assume 0 is a point of Lebesgue density of E . We then claim that we can set $c = 0$. It is simple to see that if t_0 is a point of Lebesgue density for a set E and a set F , then it is also a point of Lebesgue density for $E \cap F$. In particular 0 is a point of Lebesgue density for $E \cap a_1^{-1}E \cap \dots \cap a_N^{-1}E$, which means the set is nonempty. If $y \in E \cap a_1^{-1}E \cap \dots \cap a_N^{-1}E$, then $y, a_1y, \dots, a_Ny \in E$. \square

If f is locally integrable, the *Lebesgue set* of f consists of all points $x \in \mathbf{R}^d$ such that $f(x)$ is finite and

$$\lim_{\delta \rightarrow 0} \frac{1}{|B_\delta|} \int_{B(x, \delta)} |f(y) - f(x)| \, dy = 0.$$

If f is continuous at x , it is obvious that x is in the Lebesgue set of f , and if x is in the Lebesgue set of f , then $A_\delta f(x) \rightarrow f(x)$ as $\delta \rightarrow 0$.

Theorem 20.10. *If $f \in L^1_{\text{loc}}(\mathbf{R}^d)$, almost every point is in the Lebesgue set of f .*

Proof. For each rational number p , the function $|f - p|$ is measurable, so that there is a set E_p of measure zero such that for $x \in E_p^c$,

$$\lim_{\delta \rightarrow 0} \int_{B(x, \delta)} |f(y) - p| \, dy \rightarrow |f(x) - p|.$$

Taking unions, we conclude that $E = \bigcup E_p$ is a set of measure zero. Suppose $x \in E^c$, and $f(x)$ is finite. For any $\varepsilon > 0$, there is a rational p such that $|f(x) - p| < \varepsilon$, and we know the equation above holds, so

$$\lim_{\delta \rightarrow 0} \int_{B(x, \delta)} |f(y) - f(x)| \, dy \leq \limsup_{\delta \rightarrow 0} \int_{B(x, \delta)} (|f(y) - p| + |p - f(x)|) \, dy \leq 2\varepsilon.$$

We then let $\varepsilon \rightarrow 0$. Since $f(x)$ is finite for almost all x , this completes the proof. \square

It is interesting to note that for any $f \in L^1_{\text{loc}}(\mathbf{R}^d)$, there is $g \in L^1_{\text{loc}}(\mathbf{R}^d)$ such that $f = g$ almost everywhere, and the Lebesgue set of g is maximal. One choice is to define

$$g(x) = \limsup_{\delta \rightarrow 0} \int_{B(x, \delta)} f(y) \, dy.$$

Often the Lebesgue set of f is defined to be the Lebesgue set of g , when one wants to think of the Lebesgue set as a distributional invariant of f rather than depending solely on the pointwise behaviour.

20.7 Ergodic Averages

Ergodic theory studies the behaviour of measure preserving dynamical systems. By this, we mean, for a probability space X , a bimeasurable bijection $T : X \rightarrow X$ such that $|T^{-1}(E)| = |E|$ for all $E \subset X$. A classic example, for $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, is the shift map $T : \mathbf{T} \rightarrow \mathbf{T}$ given by $T(x) = x + \alpha$. The map T induces a linear operator T , such that $Tf(x) = f(Tx)$. That T is measure preserving implies that T is an isometry of $L^p(X)$ for any p . We begin with a result of Von Neumann.

Theorem 20.11 (Von-Neumann Ergodic Theorem). *Let $T : H \rightarrow H$ be a unitary operator. Then for any $f \in H$, the averages*

$$A_N f = \frac{f + \dots + T^{N-1} f}{N}$$

converge as $N \rightarrow \infty$ in H .

Proof. If $Tf = f$, then it is clear that $A_N f \rightarrow f$ as $N \rightarrow \infty$. Conversely, if $f = Tg - g$ for some $g \in H$, then the averages telescope, and we obtain that $A_N f = N^{-1}(T^N g - g)$, which converges to zero since T is unitary. But these two cases turn out to be orthogonal complements of one another, i.e. we can write H as the orthogonal sum of the closures of the two subspaces

$$H_1 = \{f \in H : Tf = f\} \quad \text{and} \quad H_2 = \{f \in H : f = Tg - g \text{ for some } g \in H\}.$$

These spaces are clearly orthogonal; if $Tf_1 = f_1$ and $f_2 = Tg - g$, then

$$\langle f_1, g \rangle = \langle Tf_1, Tg \rangle = \langle f_1, Tg \rangle$$

and thus

$$\langle f_1, f_2 \rangle = \langle f_1, Tg \rangle - \langle f_1, g \rangle = 0.$$

To show these spaces are orthogonal complements, let f be orthogonal to the closure of H_2 . Then $\langle f, Tf - f \rangle = 0$. But this means that

$$\langle f, Tf \rangle = \langle f, f \rangle = \langle Tf, Tf \rangle,$$

and so $\langle Tf, Tf - f \rangle = 0$. But this means that $\langle Tf - f, Tf - f \rangle = 0$, so $Tf = f$. \square

Now we move onto the mean ergodic theorem.

Theorem 20.12. *Suppose $T : X \rightarrow X$ is measure preserving and $1 \leq p < \infty$. Then if $f \in L^p(X)$, then $\{A_N f\}$ converges in $L^p(X)$.*

Proof. The operators $\{A_N\}$ are uniformly bounded on $L^p(X)$. Thus it suffices to verify the theorem for a dense subclass of functions f . So suppose $f \in L^\infty(X)$. The functions $\{A_N f\}$ converge in $L^2(X)$ and thus converge in $L^p(X)$ for any $1 \leq p < \infty$ by Hölder's inequality. The functions are also *bounded* in $L^\infty(X)$, so interpolation gives convergence in $L^p(X)$ for any $1 \leq p < \infty$. \square

The functions $A_N f$ do not necessarily converge in $L^\infty(X)$ if $f \in L^\infty(X)$. But they do converge *almost everywhere*, as we will prove using our theory of maximal functions we have developed. For functions f on X , define

$$Mf(x) = \sup_{N \geq 1} \left| \frac{1}{N} \sum_{n=0}^{N-1} T^n f(x) \right|.$$

We trivially have $\|Mf\|_{L^\infty(X)} \leq \|f\|_{L^\infty(X)}$, and we will now justify that $\|Mf\|_{L^{1,\infty}(X)} \lesssim \|f\|_{L^1(X)}$. The space X has no metric properties, so it seems surprising that we can use our previous theory of maximal functions here. But the idea is to *lift* our transform to a space where we *do* have such properties, thus obtaining a *transference principle* reducing the study of general measure preserving systems to particular ones.

Theorem 20.13. *Let $T : X \rightarrow X$ be a measure preserving system. Then for $f \in L^1(X)$,*

$$\|Mf\|_{L^{1,\infty}(X)} \lesssim \|f\|_{L^1(X)}.$$

Proof. We have $M = \lim_{K \rightarrow \infty} M_K$, where

$$M_K f(x) = \sup_{1 \leq N < K} \frac{1}{N} \sum_{n=1}^N T^n f(x).$$

For any $K > 0$, define $F : X \times [2K] \rightarrow \mathbb{C}$ by setting

$$F(x, n) = T^n f(x).$$

The space $[K]$ is a space to which we can apply the Hardy-Littlewood theory of maximal functions we have developed, *uniformly in K* , if we equip it with the uniform measure, which is doubling with respect to the standard metric on $[K]$. Thus we obtain that for each $x \in X$,

$$\left\| \sup_{1 < N \leq K} N^{-1} \sum_{n=1}^N F(x, n+k) \right\|_{L_k^{1,\infty}[K]} \lesssim \|F(x, k)\|_{L_k^1[2K]}.$$

Apply the $L_x^{1,\infty}$ norm to the left hand side of the equation, and the L^1 norm to the right hand side. Since T is measure preserving, $\|F(x, k)\|_{L_x^1(X) L_k^1[2K]} = \|f\|_{L^1(X)}$. Thus

$$\left\| \sup_{1 < N \leq K} \frac{1}{N} \sum_{n=1}^N F(x, n+k) \right\|_{L^{1,\infty}(X \times [K])} \lesssim \|f\|_{L^1(X)}.$$

To work with the left hand side, notice the identity

$$\sup_{1 \leq N < K} \frac{1}{N} \sum_{n=1}^N T^{n+k} f(x) = \sup_{1 \leq N < K} \frac{1}{N} \sum_{n=1}^N T^n (T^k f)(x).$$

Thus the left hand side is equal to $\|M_K f\|_{L^{1,\infty}(X)} \lesssim \|f\|_{L^1(X)}$. Taking $K \rightarrow \infty$ yields the required inequality. \square

Corollary 20.14 (The Pointwise Ergodic Theorem). *Let $T : X \rightarrow X$ be a measure preserving system. If $f \in L^1(X)$, then $A_N f(x)$ converges for almost every $x \in X$.*

Proof. Our maximal function bounds thus imply that the result holds for a dense subclass of $L^1(X)$. The result is obvious for a function $f \in L^2(X)$ with $Tf = f$. The theorem is also true for $f = g - Tg$, where $g \in L^\infty(X)$. But the sums of such functions are dense in $L^2(X)$ by the orthogonal decomposition we just proved, and thus dense in $L^1(X)$. \square

20.8 Approximations to the Identity

We now switch to the study of how we can approximate functions by convolutions of concentrated functions around the origin. In this section we define the various classes of such functions which give convergence results, to various degrees of strength. We say a family of integrable functions $\{K_\alpha : \alpha > 0\}$ in \mathbf{R}^d is a *good kernel* if it is bounded in the L^1 norm, for every $\alpha > 0$,

$$\int K_\alpha(x) dx = 1$$

and if for every $\delta > 0$,

$$\lim_{\alpha \rightarrow 0} \int_{|x| \geq \delta} |K_\alpha(x)| dx \rightarrow 0.$$

It requires only basic analysis to verify good kernel convergence.

Theorem 20.15. *If $\{K_\alpha\}$ is a good kernel, then for any absolutely integrable function f , $\lim_{\alpha \rightarrow 0} f * K_\alpha = f$ in $L^1(\mathbf{R}^d)$, and for any continuity point x of f , $\lim_{\alpha \rightarrow 0} (f * K_\alpha)(x) = f(x)$.*

Proof. Note that

$$\begin{aligned} \|(f * K_\alpha) - f\|_1 &= \int |(f * K_\alpha)(x) - f(x)| dx \\ &= \int \left| \int K_\alpha(y) [f(x-y) - f(x)] dy \right| dx \\ &\leq \int |K_\alpha(y)| \|T_y f - f\|_1 dy \end{aligned}$$

where $(T_y f)(x) = f(x - y)$. We know that $\|T_y f - f\|_1 \rightarrow 0$ as $y \rightarrow 0$. Thus, for each ε , we can pick δ such that if $|y| < \delta$, $\|T_y f - f\|_1 \leq \varepsilon$, and if we pick α large enough that $\int_{|y| \geq \delta} |K_\alpha(y)| dy \leq \varepsilon$, and then

$$\|(f * K_\alpha) - f\|_1 \leq \varepsilon \int_{|y| < \delta} |K_\alpha(y)| dy + 2\|f\|_1 \int_{|y| \geq \delta} |K_\alpha(y)| dy \leq \varepsilon[\|K_\alpha\|_1 + 2\|f\|_1]$$

Since $\|K_\alpha\|_1$ is universally bounded over α , we can let $\varepsilon \rightarrow 0$ to obtain convergence. If x is a fixed point of continuity, and for a given $\varepsilon > 0$, we pick $\delta > 0$ with $|f(y) - f(x)| \leq \varepsilon$ for $|y - x| < \delta$, then

$$\begin{aligned} |(f * K_\alpha)(x) - f(x)| &= \left| \int_{-\infty}^{\infty} f(y) K_\alpha(x - y) dy - f(x) \right| \\ &= \left| \int_{-\infty}^{\infty} [f(y) - f(x)] K_\alpha(x - y) dy \right| \\ &= \left| \int_{-\delta}^{\delta} [f(y) - f(x)] K_\alpha(x - y) dy \right| \\ &\quad + \left| \int_{|y| \geq \delta} [f(y) - f(x)] K_\alpha(x - y) dy \right| \\ &\leq \varepsilon \|K_\alpha\|_1 + [\|f\|_1 + f(x)] \int_{|y| \geq \delta} |K_\alpha(y)| dy \end{aligned}$$

If $\|K_\alpha\|_1 \leq M$ for all α , and we choose α large enough that $\int_{|y| \geq \delta} |K_\alpha(y)| \leq \varepsilon$, then we conclude the value above is bounded by $\varepsilon[M + \|f\|_1 + f(x)]$, and we can then let $\varepsilon \rightarrow 0$. \square

To obtain almost sure pointwise convergence of $f * K_\alpha$ to f , we must place stronger conditions on our family. We say a family $K_\delta \in L^1(\mathbf{R}^d)$, is an *approximation to the identity* if $\int K_\delta = 1$, and

$$|K_\delta(x)| \lesssim \frac{\delta}{|x|^{d+1}} \quad |K_\delta(x)| \lesssim \frac{1}{\delta^d}$$

where the constant bound is independent of x and δ . These assumptions are stronger than being a good kernel, because if K_δ is an approximation to the identity, then

$$\int_{|x| \geq \varepsilon} |K_\delta(x)| \leq \int_{\varepsilon}^{\infty} \int_{S^{d-1}} \frac{C\delta}{r} d\sigma dr = C\delta |S^{d-1}| \int_{\varepsilon}^{\infty} \frac{dr}{r} \leq \frac{C\delta |S^{d-1}|}{\varepsilon}$$

which converges to zero as $\delta \rightarrow 0$. Combined with

$$\int_{|x|<\varepsilon} |K_\delta(x)| \leq C \int_0^\varepsilon \int_{S^{d-1}} \frac{r^{d-1}}{\delta^d} d\sigma dr = \frac{C\varepsilon^d |S^{n-1}|}{d\delta^d}$$

This calculation also implies

$$\|K_\delta\|_1 \leq C |S^{n-1}| \left[\frac{\delta}{\varepsilon} + \frac{\varepsilon^d}{\delta^d} \right]$$

Setting $\varepsilon = \delta$ optimizes this value, and gives a bound

$$\|K_\delta\|_1 \leq 2C |S^{n-1}|$$

So an approximation to the identity is a stronger version of a good kernel.

Example. If φ is a bounded function in \mathbf{R}^d supported on the closed ball of radius one with $\int \varphi(x) dx = 1$, then $K_\delta(x) = \delta^{-d} \varphi(\delta^{-1}x)$ is an approximation to the identity, because by a change of variables, we calculate

$$\int_{\mathbf{R}^d} \frac{\varphi(\delta^{-1}x)}{\delta^d} = \int_{\mathbf{R}^d} \varphi(x) = 1$$

Because φ is bounded, we find

$$|K_\delta(x)| \leq \frac{\|\varphi\|_\infty}{\delta^d}$$

Now K_δ is supported on a disk of radius δ , this bound also shows

$$|K_\delta(x)| \leq \frac{\delta \|\varphi\|_\infty}{|x|^{d+1}}$$

and so K_δ is an approximation to the identity. If φ is an arbitrary integrable function, then K_δ will only be a good kernel.

Example. The Poisson kernel in the upper half plane is given by

$$P_y(x) = \frac{1}{\pi} \frac{y}{x^2 + y^2}$$

where $x \in \mathbf{R}$, and $y > 0$. It is easy to see that

$$P_y(x) = y^{-1} P_1(xy^{-1})$$

And

$$\int \frac{1}{1+x^2} = \arctan(\infty) - \arctan(-\infty) = \pi$$

We easily obtain the bounds

$$|P_y(x)| \leq \frac{\|P_1\|_\infty}{y} \quad |P_y(x)| \leq \frac{y}{\pi|x|^2}$$

so the Poisson kernel is an approximation to the identity.

Example. The heat kernel in \mathbf{R}^d is defined by

$$H_t(x) = \frac{e^{-|x|^2/4t}}{(4\pi t)^{d/2}}$$

where $\delta = t^{1/2} > 0$. Then $H_t(x) = \delta^{-d} H_1(x\delta^{-1})$, and

$$\int e^{-|x|^2/4} = \frac{1}{2^d} \int e^{-|x|^2} = \frac{|S^{n-1}|}{2^d} \int_0^\infty r^{d-1} e^{-r^2} dr$$

Example. The Poisson kernel for the disk is

$$\frac{P_r(x)}{2\pi} = \begin{cases} \frac{1}{2\pi} \frac{1-r^2}{1-2r\cos x+r^2} & : |x| \leq \pi \\ 0 & : |x| > \pi \end{cases}$$

where $0 < r < 1$, and $\delta = 1 - r$.

Example. The Féjer kernel is

$$\frac{F_N(x)}{2\pi} = \begin{cases} \frac{1}{2\pi N} \frac{\sin^2(Nx/2)}{\sin^2(x/2)} \end{cases}$$

where $\delta = 1/N$.

As $\delta \rightarrow 0$, we may think of the K_δ as ‘tending to the unit mass’ Dirac delta function δ at the origin. δ may be given a precise meaning, either in the theory of Lebesgue-Stieltjes measures or as a ‘generalized function’, but we don’t need it to discuss the actual convergence results of the functions K_δ .

Theorem 20.16. *If $\{K_\delta\}$ is an approximation to the identity, and f is integrable on $L^1(\mathbf{R}^d)$, then $(f * K_\delta)(x) \rightarrow f(x)$ for every x in the Lebesgue set of f , and $f * K_\delta$ converges to f in the L^1 norm.*

Proof. We rely on the fact that if x is in the Lebesgue set, then the function

$$A(r) = \frac{1}{r^d} \int_{|y| \leq r} |f(x-y) - f(x)| \, dy$$

is a bounded continuous function of $r > 0$, converging to 0 as $r \rightarrow 0$. This means that if $\Delta(y) = |f(x-y) - f(x)| |K_\delta(y)|$, then

$$\int \Delta(y) \, dy = \int_{|y| \leq \delta} \Delta(y) + \sum_{k=0}^{\infty} \int_{2^k \delta \leq |y| \leq 2^{k+1} \delta} \Delta(y)$$

The first term is easily upper bounded by $CA(\delta)$, and the k 'th term of the sum by $C'2^{-k}A(2^{k+1}\delta) \leq C''2^{-k}$ for constants C', C'' that do not depend on δ . Letting $\delta \rightarrow 0$ gives us the convergence result. \square

20.9 H^p Spaces

We briefly describe how the Hardy-Littlewood maximal principle can be used to develop the theory of Hardy spaces. Recall that if $f : \mathcal{D} \rightarrow \mathbf{C}$ is a harmonic function on the interior of the unit disk, and we define $f_r : \mathbf{T} \rightarrow \mathbf{C}$ by setting $f_r(t) = f(re^{2\pi i t})$, then for the *Poisson kernel*

$$P_t(x) = \frac{1-t^2}{1+t^2-2t\cos(2\pi x)} = \operatorname{Re} \left(\frac{e^{2\pi i x} + t}{e^{2\pi i x} - t} \right),$$

we have for $0 \leq s < r < 1$, $f_s = f_r * P_{s/r}$. Applying Young's inequality with the fact $\{P_t\}$ is positive and with unit mass, we thus conclude that $s \mapsto \|f_s\|_{L^p(\mathbf{T})}$ is a monotone increasing function of s . We define $H^p(\mathcal{D})$ to be the space of all *holomorphic* (and thus harmonic) functions for which the norm

$$\|f\|_{H^p(\mathcal{D})} = \lim_{t \rightarrow 1} \|f_t\|_{L^p(\mathbf{T})}.$$

We have seen that the Poisson kernel is an approximation to the identity, so that we have a pointwise estimate $|f * P_t| \lesssim Mf$.

Theorem 20.17. *If $f \in H^p(\mathcal{D})$, then for $1 < p \leq \infty$, the function*

$$f_1 = \lim_{t \rightarrow 1} f_t$$

exists pointwise almost everywhere, and lies in $L^p(\mathbf{T})$. If $1 < p < \infty$, then $\{f_t\}$ also converges to f_1 in L^p norm.

Proof. Fix $0 < t < 1$. For $g \in L^{p^*}(\mathbf{T})$, we verify that

$$\langle f_r, g * P_t \rangle = \langle f_r * P_t, g \rangle = \langle f_{rt}, g \rangle = \langle f_t * K_r, g \rangle.$$

As $r \rightarrow 1$, $f_t * K_r$ converges to f_t in the L^p norm, so we conclude that there exists a quantity A_t such that as $r \rightarrow 1$,

$$\langle f_r, g * P_t \rangle \rightarrow A_t.$$

We have $|A_t| \lesssim \|f\|_{H^p(\mathcal{D})} \|g\|_{L^{p^*}(\mathbf{T})}$, uniformly in t . As $t \rightarrow 1$, $g * P_t$ converges to g in the $L^{p^*}(\mathbf{T})$ norm, and thus because $\{f_r\}$ is bounded in $L^p(\mathbf{T})$, uniformly in r , we conclude that $\langle f_r, g * P_t \rangle$ converges to $\langle f_r, g \rangle$ uniformly in r . But this is enough to justify the convergence of $\langle f_r, g \rangle$. Thus we have justified that there exists $f_1 \in L^p(\mathbf{T})$ such that $f_r \rightarrow f_1$ weakly in $L^p(\mathbf{T})$. But this means that

$$\langle f_t, g \rangle = \lim_{r \rightarrow 1} \langle f_t * P_r, g \rangle = \lim_{r \rightarrow 1} \langle f_r, g * P_t \rangle = \langle f_1, g * P_t \rangle = \langle f_1 * P_t, g \rangle.$$

Thus $f_t = f_1 * P_t$. From our results on approximations to the identity proved in this chapter, the remaining results are now immediate. \square

20.10 The Strong Maximal Function

TODO

20.11 The Tangential Poisson Maximal Function

TODO

Chapter 21

Differentiability of Measurable Functions

A simple consequence of the results above maximal functions is a kind of fundamental theorem of calculus. If $f \in L^1(\mathbf{R})$, then we can define $F \in C(\mathbf{R})$ by setting

$$F(t) = \int_{-\infty}^t f(s) \, ds.$$

It follows from the maximal functions bounds we've established that F is differentiable almost everywhere, and $F'(t) = f(t)$ for almost every t . Thus the fundamental theorem of calculus holds in this setting, i.e.

$$F(t) = \int_{-\infty}^t F'(s) \, ds.$$

Let us now consider *what* conditions we can assume on a measurable function f such that f is differentiable almost everywhere, such that $f' \in L^1(\mathbf{R})$, and such that

$$f(t) = \int_{-\infty}^t f'(s) \, ds$$

holds for almost every $t \in \mathbf{R}$. Clearly this is equivalent to finding which functions are expressed as the indefinite integral of an integrable function.

We shall find that if f has *bounded variation*, then most of these problems are answered. If f is a complex valued function on $[a, b]$, and P is a partition, we can consider its variation on a partition $P = a \leq t_0 < \cdots <$

$t_n \leq b$ to be

$$V(f, P) = \sum_{k=1}^n |f(t_k) - f(t_{k-1})|$$

we say f has *bounded variation* if there is a constant M such that for any partition P , $V(f, P) \leq M$. This implies that, since the net $P \mapsto V(f, P)$ is increasing, the net converges to a value $V(f) = V(f, a, b)$, the *total variation* of f on $[a, b]$.

The problem of the variation of a function is very connected to the problem of the *rectifiability of curves*. If $x : [a, b] \rightarrow \mathbf{R}^d$ parameterizes a continuous curve in the plane, then, for a given partition $P = a \leq t_0 \leq \dots \leq t_n$, we can consider an approximate length

$$L_P(x) = \sum_{k=1}^n |x(t_k) - x(t_{k-1})|$$

If x has a reasonable notion of length, then the straight lines between $x(t_{i-1})$ and $x(t_i)$ should be shorter than the length of x between t_{i-1} and t_i . It therefore makes sense to define the *length* of x as

$$L(x) = \sup L_P(x)$$

The triangle inequality implies that the map $P \mapsto L_P(x)$ is an increasing net, so L is also the limit of the meshes as they become finer and finer. If $L(x) < \infty$, we say x is a *rectifiable curve*. One problem is to determine what analytic conditions one must place on x in order to guarantee regularity, and what further conditions guarantee that, if x_i is differentiable almost everywhere,

$$L(x) = \int_a^b \sqrt{x_1'(t)^2 + \dots + x_n'(t)^2} dt$$

Considering rectifiable curves leads directly to the notion of a function with bounded variation.

Theorem 21.1. *A curve x is rectifiable iff each x_i has bounded variation.*

Proof. We can find a universal constants $A, B > 0$ such that for any $x, y \in \mathbf{R}^d$,

$$A \sum |x_i - y_i| \leq |x - y| \leq B \sum |x_i - y_i|$$

This means that if P is a partition of $[a, b]$, then

$$A \sum_{ij} |x_j(t_i) - x_j(t_{i-1})| \leq \sum |x(t_i) - x(t_{i-1})| \leq B \sum_{ij} |x_j(t_i) - x_j(t_{i-1})|$$

So $A \sum V(x_i, P) \leq L_P(x) \leq B \sum V(x_i, P)$ gives the required result. \square

Example. If f is a real-valued, monotonic, increasing function on $[a, b]$, then f has bounded variation, and one can verify that $V(f) = f(b) - f(a)$.

Example. If f is differentiable at every point, and f' is bounded, then f has bounded variation. The mean value theorem implies that if $|f'| \leq M$, then for all $x, y \in [a, b]$,

$$|f(x) - f(y)| \leq M|x - y|$$

This implies that $V(f, P) \leq M(b - a)$ for all partitions P .

Example. Consider the functions f defined on $[0, 1]$ with

$$f(x) = \begin{cases} x^a \sin(x^{-b}) & : 0 < x \leq 1 \\ 0 & : x = 0 \end{cases}$$

Then f has bounded variation on $[0, 1]$ if and only if $a > b$. The function oscillates from increasing to decreasing on numbers of the form $x = (n\pi)^{-1/b}$, so the total variation is described as

$$V(f) = 1 + \sum_{n=1}^{\infty} (n\pi)^{-a/b} + ((n+1)\pi)^{-a/b}$$

This sum is finite precisely when $a/b > 1$. Thus functions of bounded variation cannot oscillate too widely, too often.

The next result is a decomposition theorem for bounded variation functions into bounded increasing and decreasing functions. We define the *positive variation* of a real valued function f on $[a, b]$ to be

$$P(f, a, b) = \sup_P \sum_{f(t_i) \geq f(t_{i-1})} f(t_i) - f(t_{i-1})$$

The *negative variation* is

$$N(f, a, b) = \sup_P \sum_{f(t_i) \leq f(t_{i-1})} -[f(t_i) - f(t_{i-1})]$$

Note that for each partition P , the sums of the two values above add up to the variation with respect to the partition.

Lemma 21.2. *If f is real-valued and has bounded variation on $[a, b]$, then for all $a \leq x \leq b$,*

$$\begin{aligned} f(x) - f(a) &= P(f, a, x) - N(f, a, x) \\ V(f) &= P(f, a, b) + N(f, a, b) \end{aligned}$$

Proof. Given ε , there exists a partition $a = t_0 < \cdots < t_n = x$ such that

$$\begin{aligned} \left| P(f, a, x) - \sum_{f(t_i) \geq f(t_{i-1})} f(t_i) - f(t_{i-1}) \right| &< \varepsilon \\ \left| N(f, a, x) + \sum_{f(t_i) \leq f(t_{i-1})} f(t_i) - f(t_{i-1}) \right| &< \varepsilon \end{aligned}$$

It follows that

$$|f(x) - f(a) - [P(f, a, x) - N(f, a, x)]| < 2\varepsilon$$

and we can then take $\varepsilon \rightarrow 0$. The second identity follows the same way. \square

A real function f on $[a, b]$ has bounded variation if and only if f is the difference of two increasing bounded functions, because if f has bounded variation, then

$$f(x) = [f(a) + P(f, a, x)] - N(f, a, x)$$

is the difference of two bounded increasing functions. On the other hand, the difference of two bounded increasing functions is clearly of bounded variation. A complex function has bounded variation if and only if it is the linear combination of four increasing functions in each direction.

Theorem 21.3. *If f is a continuous function of bounded variation, then*

$$x \mapsto V(f, a, x) \quad x \mapsto V(x, b)$$

are continuous functions.

Proof. $V(f, a, x)$ is an increasing function of x , so for continuity on the left it suffices to prove that for each x and ε , there is $x_1 < x$ such that $V(f, a, x_1) \geq V(f, a, x) - \varepsilon$. If we consider a partition

$$P = \{a = t_0 < \cdots < t_n = x\}$$

where $|V(f, P) - V(f, a, x)| \leq \varepsilon$, then by continuity of f at x , there is $t_{n-1} < x_1 < x$ with $|f(x) - f(x_1)| < \varepsilon$, and then if we modify P to obtain Q by swapping t_n with x_1 , we find

$$\begin{aligned} V(f, a, x_1) &\geq V(f, Q) = V(f, P) - |f(x) - f(t_{n-1})| + |f(x_1) - f(t_{n-1})| \\ &\geq V(f, P) - \varepsilon \geq V(f, a, x) - \varepsilon \end{aligned}$$

A similar argument gives continuity on the right, and the continuity as the left bound of the interval changes. \square

To obtain the differentiation theorem for functions of bounded variation, we require a lemma of F. Riesz.

Lemma 21.4 (Rising Sun lemma). *If f is real-valued and continuous on \mathbf{R} , and E is the set of points x where there exists $h > 0$ such that $f(x + h) > f(x)$, then, provided E is non-empty, it must be open, and can be written as a union of disjoint intervals (a_n, b_n) , where $f(b_n) = f(a_n)$. If f is continuous on $[a, b]$, then E is still an open subset of $[a, b]$, and can be written as the disjoint union of countably many intervals, with $f(b_n) = f(a_n)$ except if $a_n = a$, where we can only conclude $f(a_n) \leq f(b_n)$.*

Proof. The openness is clear, and the fact that E can be broken into disjoint intervals follows because of the characterization of open sets in \mathbf{R} . If

$$E = \bigcup (a_n, b_n)$$

Then $f(a_n + h) \leq f(a_n)$ and $f(b_n + h) \leq f(b_n)$, implying in particular that $f(b_n) \leq f(a_n)$. If $f(b_n) < f(a_n)$, then choose $f(b_n) < c < f(a_n)$. The intermediate value theorem implies there is x with $f(x) = c$, and we may choose the largest $x \in [a_n, b_n]$ for which this is true. Then since $x \in (a_n, b_n)$, there is $y \in (x, b_n)$ with $f(x) < f(y)$, and by the intermediate value theorem, since $f(b_n) < f(x) < f(y)$, there must be $x' \in (y, b_n)$ with $f(x') = c$, contradicting that x was chosen maximally. The proof for closed intervals operates on the same principles. \square

Theorem 21.5. *If f is increasing and continuous on $[a, b]$, then f is differentiable almost everywhere. That is,*

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists for almost every $x \in [a, b]$, f' is measurable, and

$$\int_a^b f'(x) \leq f(b) - f(a)$$

In particular, if f is bounded on \mathbf{R} , then f' is integrable on \mathbf{R} .

Proof. the theorem in the It suffices to assume that f is increasing, and we shall start by proving case assuming f is continuous. We define

$$\Delta_h f(x) = \frac{f(x+h) - f(x)}{h}$$

and the four *Dini derivatives*

$$D_+ f(x) = \liminf_{h \downarrow 0} \Delta_h f(x) \quad D^+ f(x) = \limsup_{h \downarrow 0} \Delta_h f(x)$$

$$D_- f(x) = \liminf_{h \uparrow 0} \Delta_h f(x) \quad D^- f(x) = \limsup_{h \uparrow 0} \Delta_h f(x)$$

Clearly, $D_+ f \leq D^+ f$ and $D_- f \leq D^- f$, It suffices to show $D^+ f(x) < \infty$ for almost every x , and $D^+ f(x) \leq D_- f(x)$ for almost every x , because if we consider the function $g(x) = -f(-x)$, then we obtain $D^- f(x) \leq D_+ f(x)$ for almost every x , so

$$D^+ f(x) \leq D_- f(x) \leq D^- f(x) \leq D_+ f(x) \leq D^+ f(x) < \infty$$

for almost every x , implying all values are equal, and that the derivative exists at x .

For a fixed $\gamma > 0$, consider $E_\gamma = \{x : D^+ f(x) > \gamma\}$. Since each $\Delta_h f$ is continuous, the supremum of the $\Delta_h f$ over any index set is lower semi-continuous, and since

$$D^+ f(x) = \lim_{h \rightarrow 0} \sup_{0 \leq s \leq h} \Delta_h f(x+s)$$

can be expressed as the countable limit of these lower semicontinuous functions, $D^+ f$ is measurable, hence E_γ is measurable. Now consider the shifted function $g(x) = f(x) - \gamma x$. If $\bigcup (a_i, b_i)$ is the set obtainable from g from the rising sun lemma, then $E_\gamma \subset \bigcup (a_i, b_i)$, for if $D^+ f(x) > \gamma$, then there is $h > 0$ arbitrarily small with $\Delta_h f(x) > \gamma$, hence $f(x+h) - f(x) > \gamma h$,

hence $g(x+h) > g(x)$. We know that $g(a_k) \leq g(b_k)$, so $f(b_k) - f(a_k) \geq \gamma(b_k - a_k)$, so

$$|E_\gamma| \leq \sum (b_k - a_k) \leq \frac{1}{\gamma} \sum f(b_k) - f(a_k) \leq \frac{f(b) - f(a)}{\gamma}$$

Thus $|E_\gamma| \rightarrow 0$ as $\gamma \downarrow 0$, implying $D^+f(x) = \infty$ only on a set of measure zero.

Now for two real numbers $r < R$, we will now show

$$E = \{a \leq x \leq b : D^+f(x) > R \quad D_-f(x) < r\}$$

is a set of measure zero. Letting r and R range over all rational numbers establishes that $D^+f(x) \leq D_-f(x)$ almost surely. We assume $|E| > 0$ and derive a contradiction. By regularity, we may consider an open subset U in $[a, b]$ containing E such that $|U| < |E|(R/r)$. We can write U as the union of disjoint intervals I_n . For a fixed I_N , apply the rising sun lemma to the function $rx - f(-x)$ on the interval $-I_N$, yielding a union of intervals (a_n, b_n) . If we now apply the rising sun lemma to the function $f(x) - Rx$ on (a_n, b_n) , we get intervals (a_{nm}, b_{nm}) , whose union we denote U_N . Then

$$R(b_{nm} - a_{nm}) \leq f(b_{nm}) - f(a_{nm}) \quad f(b_n) - f(a_n) \leq r(b_n - a_n)$$

then, because f is increasing,

$$\begin{aligned} |U_N| &= \sum_{nm} (b_{nm} - a_{nm}) \leq \frac{1}{R} \sum_{nm} (f(b_{nm}) - f(a_{nm})) \\ &\leq \frac{1}{R} \sum f(b_n) - f(a_n) \leq \frac{r}{R} \sum_n (b_n - a_n) \leq \frac{r}{R} |I_N| \end{aligned}$$

Now $E \cap I_N$ is contained in U_N , because if $x \in E \cap I_N$, then $D^+f(x) > R$ and $D_-f(x) < r$, so we can sum in N to conclude that

$$|E| \leq \sum \frac{r}{R} |I_N| = \frac{r}{R} |U_N| < |E|$$

a contradiction proving the claim. \square

Corollary 21.6. *If f is increasing and continuous, then f' is measurable, non-negative, and*

$$\int_a^b f'(x) dx \leq f(b) - f(a)$$

Proof. The fact the f' is measurable and non-negative results from the fact that the functions $g_n(x) = \Delta_{1/n}f(x)$ are non-negative and continuous, and $g_n \rightarrow f'$ almost surely. We know

$$\begin{aligned} \int_a^b f'(x) &\leq \liminf_{n \rightarrow \infty} \int_a^b g_n(x) = \liminf_{n \rightarrow \infty} n \int_a^b [f(x + 1/n) - f(x)] \\ &= \liminf_{n \rightarrow \infty} n \left[\int_b^{b+1/n} f(x) - \int_a^{a+1/n} f(x) \right] = f(b) - f(a) \end{aligned}$$

where the last equality follows because f is continuous. \square

Even for increasing continuous functions, the inequality in the theorem above need not be an equality, as the next example shows, so we need something stronger to obtain our differentiation theorem.

Example. The Cantor-Lebesgue function is a continuous increasing function f from $[0, 1]$ to itself, with $f(0) = 0$, and $f(1) = 1$, but with $f'(x) = 0$ almost everywhere. This means

$$\int_0^1 f'(x) = 0 < 1 = f(1) - f(0)$$

so we cannot obtain equality in general. To construct f , consider the Cantor set $C = \bigcap C_k$, where C_k is the disjoint union of 2^k closed intervals. Set $f_0 = 0$, and $f_1(0) = 0$, $f_1(x) = 1/2$ on $[1/3, 2/3]$, $f_1(1) = 1$, and f linear between $[0, 1/3]$ and $[2/3, 1]$. Similarly, set $f_2(0) = 0$, $f_2(x) = 1/4$ on $[1/9, 2/9]$, $f_2(x) = 1/2$ on $[1/3, 2/3]$, $f_2(x) = 3/4$ on $[7/9, 8/9]$, and $f_2(1) = 1$. The functions f_i are increasing and cauchy in the uniform norm, so they converge to a continuous function f called the Cantor function. f is constant on each interval in the complement of the cantor set, so $f'(x) = 0$ almost everywhere.

To obtain equality in the integral formula, we require additional conditions on our increasing functions, provided by absolute continuity.

21.1 Absolute Continuity

A function $f : [a, b] \rightarrow \mathbf{R}$ is *absolutely continuous* if for every $\varepsilon > 0$, there is $\delta > 0$ such that whenever $(a_1, b_1), \dots, (a_n, b_n)$ are disjoint intervals with

$\sum(b_i - a_i) < \delta$, $\sum|f(b_i) - f(a_i)| < \varepsilon$. Thus the function should be ‘essentially constant’ over every set of zero measure. It is easy to see from this that absolutely continuous functions must be uniformly continuous, and have bounded variation. Thus f has a decomposition into the difference of two continuous increasing functions, and one can see quite easily that these functions are also absolutely continuous. Most promising to us, if f is a function defined by $f(x) = \int_a^x g(t) dt$, where $g \in L^1[a, b]$, then f is absolutely continuous. This shows that absolute continuity is necessary in order to hope for the integral formula

$$\int_a^b f'(x) dx = f(b) - f(a)$$

The Cantor function is *not* absolutely continuous, since it is constant except on the Cantor set, and we can cover the Cantor set by intervals with total length $(2/3)^n$ for each n . Thus it is impossible for the Cantor function to satisfy the fundamental theorem of calculus.

Theorem 21.7. *If $g \in L^1(\mathbf{R})$, and*

$$f(x) = \int_a^x g(t) dt$$

then f is absolutely continuous.

Proof. Fix $\varepsilon > 0$. We claim that there is δ such that if $|E| < \delta$, then $\int_E |g| < \varepsilon$. Otherwise there are sets E_n with $|E_{n+1}| \leq |E_n|/3$ and with $\int_{E_n} |g| \geq \varepsilon$. Thus if we define the sets $E'_m = E_m - \bigcup_{n>m} E_n$ then the E'_m and we have $|E_m| \sim |E'_m|$. Since g is integrable, we must have $\sum \int_{E'_n} |g| < \infty$, so we conclude that as $N \rightarrow \infty$,

$$\sum_{n \geq N} \int_{E'_n} |g| \rightarrow 0$$

Yet for any N ,

$$\sum_{n \geq N} \int_{E'_n} |g| = \int_{E_N} |g| \geq \varepsilon$$

which is an impossibility. Thus such a δ exists for every ε , and so if we have disjoint intervals (a_n, b_n) with $\sum(b_n - a_n) < \delta$, then

$$\sum |f(b_n) - f(a_n)| = \sum \left| \int_{a_n}^{b_n} g(t) dt \right| \leq \sum \int_{a_n}^{b_n} |g| = \int_{\bigcup (a_n, b_n)} |g| < \varepsilon$$

which shows the function is absolutely continuous. \square

To prove the differentiation theorem, we require a covering estimate not unlike that used to prove the Lebesgue differentiation theorem. We say a collection of balls is a *Vitali covering* of a set E if for every $x \in E$ and every $\eta > 0$, there is a ball B in the cover containing x with radius smaller than η . Thus every point is covered by an arbitrary small ball.

Lemma 21.8. *If E is a set of finite measure, and $\{B_\alpha\}$ is a Vitali covering of E , then there exists a disjoint family of cubes $\{B_\beta\}$ in the covering such that*

$$\left| E - \bigcup_{\beta} B_{\beta} \right| = 0.$$

Proof. Fix $\delta > 0$. We claim we can find a disjoint family of cubes B_1, \dots, B_N in the Vitali cover such that

$$\left| E - \bigcup_{i=1}^N B_i \right| \leq \delta.$$

By inner regularity, pick a compact set $K \subset E$ with $|K| \geq |E| - \delta/2$. Then K is covered by finitely many balls of radius less than η in the covering $\{B_\alpha\}$, and the elementary Vitali covering lemma gives a disjoint subcollection of balls B_1, \dots, B_{n_0} with

$$|K| \leq \left| \bigcup B_\alpha \right| \leq 3^d \sum |B_k|$$

so $\sum |B_k| \geq 3^{-d} |K|$. If $\sum |B_k| \geq |K| - \delta/2$, we're done. Otherwise, define $E_1 = K - \bigcup \overline{B_k}$. Then

$$|E_1| \geq |K| - \sum |\overline{B_k}| = |K| - \sum |B_k| > \delta/2$$

If we pick a compact set $K_1 \subset E_1$ with $|K_1| \geq \delta/2$, then if we remove all sets in the Vitali covering which intersect B_1, \dots, B_{n_0} , then we still obtain a Vitali covering for K_1 , and we can repeat the argument above to find a disjoint collection of open sets $B_1^1, \dots, B_{n_1}^1$ with $\sum |B_k^1| \geq 3^{-d} |K_1|$. Then $\sum |B_k| + \sum |B_k^1| \geq 2(3^{-d} \delta)$. If $\sum |B_k| + \sum |B_k^1| < |K| - \delta/2$, we repeat the same process, finding a disjoint family for $K_2 \subset E_2$, where $E_2 = K_1 - \bigcup B_k^1$. If this

process repeats itself k times, then we obtain a family of open sets with total measure greater than or equal to $k(3^{-d}\delta)$. But then if we eventually have $k \geq (|E| - \delta)3^d/\delta$, then we obtain the required bound.

We now construct our final cover inductively. Given E , we can find finitely many balls B_{11}, \dots, B_{1n_1} such that

$$\left| E - \bigcup_{i=1}^{n_1} B_i \right| \leq 1/2.$$

Set $E_1 = E - \bigcup_{i=1}^{n_1} B_i$. If we remove all balls from the Vitali cover that intersect the balls B_1, \dots, B_{n_1} , we still have a Vitali cover of E_1 . Inductively, we can then find a disjoint family of balls B_{k1}, \dots, B_{kn_k} which are disjoint from all previously selected balls such that

$$\left| E - \bigcup_{k=1}^{k_0} \bigcup_{i=1}^{n_k} B_{ki} \right| \leq 1/2^{k_0}.$$

Taking $k_0 \rightarrow \infty$ gives an infinite family of disjoint balls which cover E up to a set of zero Lebesgue measure. \square

Theorem 21.9. *If $f : [a, b] \rightarrow \mathbf{R}$ is absolutely continuous, then f' exists almost everywhere, and if $f'(x) = 0$ almost surely, then f is constant.*

Proof. It suffices to prove that $f(a) = f(b)$, since we can then apply the theorem on any subinterval. Let $E = \{x \in (a, b) : f'(x) = 0\}$. Then $|E| = b - a$. Fix $\varepsilon > 0$. Since for each $x \in E$, we have

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = 0$$

This implies that the family of intervals (x, y) such that the inequality $|f(y) - f(x)| \leq \varepsilon(y - x)$ holds forms a Vitali covering of E , and we may therefore select a family of disjoint intervals $I_i = (x_i, y_i)$ with

$$\sum |I_i| \geq |E| - \delta = (b - a) - \delta$$

But $|f(y_i) - f(x_i)| \leq \varepsilon(y_i - x_i)$, so we conclude

$$\sum |f(y_i) - f(x_i)| \leq \varepsilon(b - a)$$

The complement of I_i is a union of intervals $J_i = (x'_i, y'_i)$ of total length $\leq \delta$. Applying the absolute continuity of f , we conclude

$$\sum |f(y'_i) - f(x'_i)| \leq \varepsilon$$

so applying the triangle inequality,

$$|f(b) - f(a)| \leq \sum |f(y'_i) - f(x'_i)| + \sum |f(y_i) - f(x_i)| \leq \varepsilon(b - a + 1)$$

We can then let $\varepsilon \rightarrow 0$ to obtain equality. \square

Theorem 21.10. *Suppose f is absolutely continuous on $[a, b]$. Then f' exists almost every and is integrable, and*

$$f(b) - f(a) = \int_a^b f'(y) \, dy$$

so the fundamental theorem of calculus holds everywhere. Conversely, if $f \in L^1[a, b]$, then there is an absolutely continuous function g with $g' = f$ almost everywhere.

Proof. Since f is absolutely continuous, we can write f as the difference of two continuous increasing functions on $[a, b]$, and this easily implies f is differentiable almost everywhere and is integrable on $[a, b]$. If $g(x) = \int_a^x f'(x)$, then g is absolutely continuous, hence $g - f$ is also absolutely continuous. But we know that $(g - f)' = g' - f' = 0$ almost everywhere, so the last theorem implies that g differs from f by a constant. Since $g(a) = 0$, $g(x) = f(x) - f(a)$. The converse was proved exactly in our understanding of differentiating integrals. \square

We now dwell slightly longer on the properties of absolutely continuous functions, which enables us to generalize other properties of integrals found in the calculus. We begin by noting that it is easy to verify that if f and g are absolutely continuous functions, then fg is also absolutely continuous. We know f' , g' , and $(fg)'$ exist almost everywhere. But when all three exist simultaneously, the product rule gives $(fg)' = f'g + fg'$. The absolute continuity implies that

$$\int_a^b f'g + fg' = \int_a^b (fg)' = f(b)g(b) - f(a)g(a)$$

Thus one can integrate a pair of absolutely continuous functions by parts. Next, we shall show that monotone absolutely continuous functions are precisely those we can use to change variables. One important thing to note is that even if f is a continuous function, and g is measurable, $g \circ f$ need not be measurable. The easy reason to see this is that the inverse image of every open set in g is measurable, so in order to guarantee $g \circ f$ is measurable we need the inverse image of every measurable set under f be measurable.

Example. Consider the function $f(x) = \int_0^x \chi_E(x) dx$, where E is a thick Cantor set. Then f is absolutely continuous, strictly increasing on $[0, 1]$, and maps E to a set of measure zero. This is because $E = \lim E_n$, where E_n is a family of intervals with $|E_n| \downarrow |E|$. Then $f(E_n)$ has total length $|E_n - E|$, so as $n \rightarrow \infty$, we see $\lim f(E_n) = f(E)$ has measure zero. This means that $f(X)$ is measurable for any subset X of E , and in particular, if X is non-measurable, then $f^{-1}(f(X))$ cannot be measurable, even though $f(X)$ is measurable. Note that f is strictly increasing even though its derivatives vanish on a set of positive measure.

The next lemmas will show that even though $g \circ f$ may not be Lebesgue measurable when f is absolutely continuous and g is Lebesgue measurable, this does not really bother us too much when changing variables.

Lemma 21.11. *If f is absolutely continuous, then it maps sets of measure zero to sets of measure zero.*

Proof. Let E be a set of measure zero. Then for each $\delta > 0$, E is coverable by a family of open intervals with total length δ . But if δ is taken small enough, this means that $f(E)$ is coverable by a family of open intervals with total length bounded by ε , for any ε . \square

This property of absolutely continuous functions is independent of the properties of the Euclidean domain as its domain, and is used in the generalization of absolute continuity to more general domains, or even to measures. If f is absolutely continuous, then the image of every interval is an interval, and since $f(\bigcup K_n) = \bigcup f(K_n)$, this implies that the image of a F_σ set is measurable. But since every measurable set of \mathbf{R} differs from a F_σ set by a set of measure zero, the image of every Lebesgue measurable set is Lebesgue measurable. The reverse is almost true.

Lemma 21.12. *If f is absolutely continuous, and E measurable, then the set*

$$f^{-1}(E) \cap \{x : f'(x) > 0\}$$

is measurable.

Proof. If E is an open set, then

$$|E| = \int_{f^{-1}(E)} f'(x) dx$$

It suffices to prove this when E is an interval, and then this is just the theorem of differentiation for absolutely continuous functions. But then applying the dominated convergence theorem shows that this equation remains true if E is an G_δ set. Furthermore, this means the theorem is true if E is a closed set, and so by applying the monotone convergence theorem, the theorem is true if E is an F_σ set. But if E is an arbitrary measurable set, then for every ε there are F_σ and G_δ sets $K \subset E \subset U$ with $|U - K| = 0$. But

$$\alpha |f^{-1}(U - K) \cap \{f' \geq \alpha\}| \leq \int_{f^{-1}(U - K)} f'(x) dx = |U - K| = 0$$

Thus $f^{-1}(U - K) \cap \{f' \geq \alpha\}$ is a set of measure zero, and so in particular by completeness, every set contained in this set is measurable, in particular $f^{-1}(U - E) \cap \{f' \geq \alpha\}$ is measurable. But now this means

$$\{f' \geq \alpha\} - f^{-1}(U - E) \cap \{f' \geq \alpha\} = f^{-1}(E) \cap \{f' \geq \alpha\}$$

is measurable. Taking $\alpha \downarrow 0$ completes the proof. \square

Because of this, even though $g \circ f$ is not necessarily measurable, $(g \circ f)f'$ is always measurable if f is absolutely continuous. Thus the expression $\int (g \circ f)f'$ makes sense, and thus we can always interpret the change of variables formula.

Theorem 21.13. *If f is absolutely continuous, and g is integrable, then*

$$\int g(f(x))f'(x) dx = \int g(y) dy$$

Proof. Using the notation in the last proof, if E is measurable, then

$$|K| = \int_{f^{-1}(K)} f'(x) dx \leq \int_{f^{-1}(E)} f'(x) dx \leq \int_{f^{-1}(U)} f'(x) dx = |U|$$

and $|U| = |K| = |E|$, so that for any measurable set E ,

$$|E| = \int_{f^{-1}(E)} f'(x) dx$$

This implies the theorem we need to prove is true whenever g is the characteristic function of any measurable set. But then by linearity, it is true for any simple function. By monotone convergence, it is then true for any non-negative function, and then by partitioning g into the sum of simple functions, we obtain the theorem in general. \square

21.2 Differentiability of Jump Functions

We now consider the differentiability of not necessarily continuous monotonic functions. Set f to be an increasing function on $[a, b]$, which we may assume to be bounded. Then the left and right limits of f exist at every point, which we will denote by $f(x-)$ and $f(x+)$. Of course, we have $f(x-) \leq f(x) \leq f(x+)$. If there is a discontinuity, this means we are forced to have a ‘jump discontinuity’ where f skips an interval. This implies that f can only have countably many such discontinuities, because a family of disjoint intervals on \mathbf{R} is at most countable. Now define the jump function $\Delta f(x) = f(x+) - f(x-)$, with $\theta(x) \in [0, 1]$ defined such that $f(x_n) = f(x_n-) + \theta(x)\Delta f(x)$. If we define the functions

$$j_y(x) = \begin{cases} 0 & : x < y \\ \theta(y) & : x = y \\ 1 & : x > y \end{cases}$$

then we can define the *jump function* associated with f by

$$J(x) = \sum_x \Delta f(x) j_n(x)$$

Since f is bounded on $[a, b]$, we make the final observation that

$$\sum_{x \in [a, b]} \Delta f(x) \leq f(b) - f(a) < \infty$$

so the series defining J converges absolutely and uniformly.

Lemma 21.14. *If f is increasing and bounded on $[a, b]$, then J is discontinuous precisely at the values x with $\Delta f(x) \neq 0$ with $\Delta J(x) = \Delta f(x)$. The function $f - J$ is continuous and increasing.*

Proof. If x is a continuity point of f , then j_y is continuous at x , and hence, because $\sum_y \Delta f(y) j_y(x) \rightarrow J(x)$ uniformly, so we conclude that J is continuous at x . On the other hand, for each y , $j_y(y-) = 0$ and $j_y(y+) = 1$, and if we label the points of discontinuity of f by x_1, x_2, \dots , then

$$J(x) = \sum_{i=1}^k \Delta f(x_i) j_{x_i} + \sum_{i=k+1}^{\infty} \Delta f(x_i) j_{x_i}$$

The right hand partial sums are continuous at x_k , whereas the left hand sum has a jump discontinuity of the same order as f at x_k , we conclude J also has this discontinuity. But this means that

$$(f - J)(x_k+) - (f - J)(x_k-) = 0$$

so $f - J$ is continuous at every point. $f - J$ is increasing because of the inequality

$$J(y) - J(x) \leq \sum_{x < x_n \leq y} \alpha_n \leq f(y) - f(x)$$

which follows because J is just the sum of jump discontinuities, and the right hand side because f can decrease and increase outside of the jump discontinuities. \square

Since $f - J$ is continuous and increasing, it is differentiable almost everywhere. It therefore remains to analyze the differentiability of the jump function J .

Theorem 21.15. *J' exists and vanishes almost everywhere.*

Proof. Fix $\varepsilon > 0$, and consider

$$E = \left\{ x \in [a, b] : \limsup_{h \rightarrow 0} \frac{J(x+h) - J(x)}{h} > \varepsilon \right\}$$

Then E is measurable, because we can take the limsup over rational numbers because J is increasing. We want to show it has measure zero. Suppose $\delta = |E|$. Consider $\eta > 0$ to be chosen later, and find n such that $\sum_{k=n}^{\infty} \alpha_k < \eta$. Write

$$J_0(x) = \sum_{n > N} \alpha_n j_n$$

Then $J_0(b) - J_0(a) < \eta$. Now E differs from the set

$$E' = \left\{ x \in [a, b] : \limsup_{h \rightarrow 0} \frac{J_0(x+h) - J_0(x)}{h} > \varepsilon \right\}$$

by finitely many points. Using inner regularity, find a compact set $K \subset E'$ with $|K| \geq \delta/2$. For each $x \in K$, we can find intervals (α_x, β_x) upon which $J_0(\beta_x) - J_0(\alpha_x) \geq \varepsilon |\beta_x - \alpha_x|$. But applying the elementary Vitali covering lemma, we can find a disjoint family of such intervals with $\sum (\beta_{x_i} - \alpha_{x_i}) \geq |K|/3 \geq \delta/6$. But now we find

$$J_0(b) - J_0(a) \geq \sum J_0(\beta_{x_i}) - J_0(\alpha_{x_i}) \geq \varepsilon \delta / 6$$

This means $\delta \leq 6\eta/\varepsilon$, and by letting $\eta \rightarrow 0$, we can conclude $\delta = 0$. \square

This concludes our argument that *every* function of bounded variation has a derivative almost everywhere, because every such function can be uniquely written (up to a shift in the range of the functions) as the sum of a continuous function and a jump function. If f is a function with bounded variation, then the function

$$F(x) = \int_0^x f'(x)$$

is absolutely continuous, and $f - F$ is a continuous function with derivative zero almost everywhere. The fact that this decomposition is unique up to a shift as well (which can easily be seen in the case of an increasing function, from which the general case follows) leads us to refer to this as the *Lebesgue decomposition* of a function of bounded variation on the real line.

21.3 Rectifiable Curves

We now consider the validity of the length formula

$$L = \int_a^b (x'(t)^2 + y'(t)^2)^{1/2} dt$$

where L is the length of the curve parameterized by (x, y) on $[a, b]$. We cannot always expect this formula to hold, because if x and y are both the Cantor devil staircase function, then the formula above gives a length of zero, whereas we know the curve traces a line between 0 and 1, hence has length at least $\sqrt{2}$.

Theorem 21.16. *If a curve is parameterized by absolutely continuous functions x and y on $[a, b]$, then the curve is rectifiable, and has length*

$$\int_a^b (x'(t)^2 + y'(t)^2)^{1/2} dt$$

Proof. This proof can be reworded as saying if f is complex-valued and absolutely continuous, then its total variation can be expressed as

$$V(f, a, b) = \int_a^b |f'(t)| dt$$

If $P = \{a \leq t_1 < \cdots < t_n \leq b\}$ is a partition, then

$$\sum |f(t_{n+1}) - f(t_n)| = \sum \left| \int_{t_n}^{t_{n+1}} f'(t) dt \right| \leq \sum \int_{t_n}^{t_{n+1}} |f'(t)| dt \leq \int_a^b |f'(t)| dt$$

so $V(f, a, b) \leq \int_a^b |f'(t)| dt$. To prove the converse inequality, fix $\varepsilon > 0$, and find a step function g with $f' = g + h$, with $\|h\|_1 \leq \varepsilon$. If $G(x) = \int_a^x g(t) dt$ and $H(x) = \int_a^x h(t) dt$, then $F = G + H$, and $V(f, a, b) \geq V(G, a, b) - V(H, a, b) \geq V(G, a, b) - \varepsilon$, and if we partition $[a, b]$ into $a = t_0 < \cdots < t_N$, where G is constant on each (t_n, t_{n+1}) , then

$$\begin{aligned} V(G, a, b) &\geq \sum |G(t_n) - G(t_{n-1})| = \sum \left| \int_{t_{n-1}}^{t_n} g(t) dt \right| \\ &= \sum \int_{t_{n-1}}^{t_n} |g(t)| dt = \int_a^b |g(t)| dt \geq \|f'\|_1 - \varepsilon \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ now gives the result. □

It is interesting to note that any rectifiable curve has a special *parameterization by arclength*, i.e. a parameterization $(x(t), y(t))$ such that if L is the length function associated to the parameterization, then $L(A, B) = B - A$. This is obtainable by inverting the length function.

Theorem 21.17. *If $z = (x, y)$ is a parameterization of a rectifiable curve by arclength, then x and y are absolutely continuous, and $|z'| = 1$ almost everywhere.*

Proof. For any $s < t$,

$$t - s = L(s, t) = V(f, s, t) \geq |z(t) - z(s)|$$

so it follows immediately that $|z|$ is an absolutely continuous function, and $|z'| \leq 1$ almost surely. But now we know that

$$\int_a^b |z'(t)| = b - a$$

and this equality can now only hold if $|z'(t)| = 1$ almost surely. \square

21.4 Bounded Variation in Higher Dimensions

Since the higher dimensional Euclidean domains do not have an ordering, it is impossible to define their length by partitioning their domain, and the meaning of a jump discontinuity is no longer clear. However, there are properties equivalent to having bounded variation which are more extendable to higher dimensions.

Theorem 21.18. *The following properties of $f : \mathbf{R} \rightarrow \mathbf{R}$ are equivalent, for some fixed finite constant A .*

- *f can be modified on a set of measure zero so that it has bounded variation not exceeding A .*
- *$\int |f(x+h) - f(x)| \leq A|h|$ for all $h \in \mathbf{R}$.*
- *For any C^1 function φ with compact support, $|\int f(x)\varphi'(x)| \leq A\|\varphi\|_\infty$.*

Proof. If $V(f) = A$, where $A < \infty$, then we can write $f = f^+ - f^-$, where f^+ and f^- are both increasing functions, and with $V(f) = V(f^+) + V(f^-)$. It then follows that $|f(x+h) - f(x)| \leq (f^+(x+h) - f^+(x)) + |f^-(x+h) - f^-(x)|$, so it suffices to prove the second property by assuming f is increasing. But then by the monotone convergence theorem, assuming $h > 0$ without loss of generality,

$$\int |f(x+h) - f(x)| = \lim_{y \rightarrow \infty} \int_{-y}^y f(x+h) - f(x) = \lim_{y \rightarrow \infty} \int_y^{y+h} f(x) - \int_{-y-h}^{-y} f(x)$$

The first term of the limit converges to $hV(f)$, and the second to zero, completing the first part of the theorem. Now assuming the second point, we prove the third point. Then using the second point, we find

$$\begin{aligned} \left| \int f(x) \varphi'(x) \right| &= \left| \lim_{h \rightarrow 0} \int f(x) \frac{\varphi(x+h) - \varphi(x)}{h} \right| \\ &= \left| \lim_{h \rightarrow 0} \int \frac{f(x-h) - f(x)}{h} \varphi(x) \right| \leq A \|\varphi\|_\infty \end{aligned}$$

Finally, we consider the third point being true. The set of all partitions with rational points is countable. Suppose that for each rational $P = \{t_0 < \dots < t_N\}$ there is a set E_P of measure zero for each rational partition P such that

$$\sum_{n=1}^N \sup_{\substack{x \in [t_{n-1}, t_n] \\ x \notin E_P}} f(x) - \inf_{\substack{x \in [t_{n-1}, t_n] \\ x \notin E_P}} f(x) \leq A$$

Then the union of E_P over all rational P has measure zero. We can modify f on E_P by setting $f(x) = \liminf_{y \rightarrow 0} f(x+y)$, and then $V(f, P) \leq A$ for all rational partitions P . If Q is now any partition, we can find a rational partition P with $V(f, P) \geq V(f, Q) - \varepsilon$, and so $V(f, P) \leq A - \varepsilon$. Taking $\varepsilon \rightarrow 0$ completes the argument. Thus if f cannot be modified to have finite variation A , there exists a rational partition P such that for any set E of measure zero,

$$\sum_{n=1}^N \sup_{\substack{x \in [t_{n-1}, t_n] \\ x \notin E}} f(x) - \inf_{\substack{x \in [t_{n-1}, t_n] \\ x \notin E}} f(x) > A$$

Thus for any ε , there exists $E_n^+, E_n^- \subset [t_{n-1}, t_n]$ of positive measure such that

$$\sum_{n=1}^N \inf_{x \in E_n^+} f(x) - \sup_{x \in E_n^-} f(x) > A$$

If we consider the polygonal function ϕ which □

21.5 Minkowski Content

Given a set $K \in \mathbf{R}^n$, we let K^δ denote the open set consisting of points x with $d(x, K) < \delta$. The m dimensional *Minkowski content* of K is defined to be

$$\lim_{\delta \rightarrow 0} \frac{1}{\alpha(n-m)} \frac{|K^\delta|}{\delta^m}$$

where $\alpha(d)$ is the volume of the unit ball in d dimensions. When this limit exists, we denote it by $M^m(K)$. In this section, we mainly discuss the one dimensional Minkowski content in two dimensions, i.e. the values of

$$\lim_{\delta \rightarrow 0} \frac{|K^\delta|}{2\delta^m}$$

and it's relation the length of curves. Since we now only care about the one dimensional Minkowski content, we let $M(K)$ denote the one dimension Minkowski content.

Lemma 21.19. *If $\Gamma = \{z(t) : a \leq t \leq b\}$ is a curve, and Δ is the distance between the endpoints of the curve, then $|\Gamma^\delta| \geq 2\delta\Delta$.*

Proof. By rotating, we may assume that both endpoints of the curve lie on the x axis, so $z(a) = (A, 0)$, $z(b) = (B, 0)$ with $A < B$, so $\Delta = B - A$. If $\Delta = 0$, the theorem is obvious. Otherwise, for each point $x \in [A, B]$ there is $t(x)$ such that if $z_1(t(x)) = x$, and so Γ^δ contains $x \times [z_2(t(x)) - \delta, z_2(t(x)) + \delta]$, which has length 2δ . Thus by Fubini's theorem,

$$|\Gamma^\delta| = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \chi_{\Gamma^\delta}(x, y) dx dy \geq \int_A^B 2\delta = 2\delta\Delta$$

so the theorem is proved. □

Theorem 21.20. *If $\Gamma = \{z(t) : a \leq t \leq b\}$ is a quasi-simple curve (simple except at finitely many points), then the Minkowski content of Γ exists if and only if Γ is rectifiable, and in this case $M^1(\Gamma)$ is the length of the curve L .*

Proof. To prove the theorem, we consider the upper and lower Minkowski contents

$$M^*(\Gamma) = \limsup_{\delta \rightarrow 0} \frac{|\Gamma|^\delta}{\alpha(n-1)\delta} \quad M_*(\Gamma) = \liminf_{\delta \rightarrow 0} \frac{|\Gamma|^\delta}{\alpha(n-1)\delta}$$

First, we prove that $M^*(\Gamma) \leq L$. Consider a partition P of $[a, b]$, and let L_P be the length of the polygonal approximation to the curve. By refining the partition, we may assume that Γ is simple, with the repeated points at the boundaries of the intervals. For each interval I_n in the partition, we select a closed subinterval $J_n = [t_n, u_n]$ such that Γ is simple on $\bigcup J_n$, and

$$\sum |z(u_n) - z(t_n)| \geq L_P - \varepsilon$$

Since the intervals J_n are disjoint, for suitably small δ the sets J_n^δ are disjoint. Applying the previous lemma, we conclude that

$$|\Gamma^\delta| \geq \sum |J_n^\delta| \geq 2\delta \sum |z(u_n) - z(t_n)| = 2\delta(L_P - \varepsilon)$$

First, by letting $\varepsilon \rightarrow 0$ and then $\delta \rightarrow 0$, we conclude that $M_*(\Gamma) \geq \lim_P L_P$. In particular, this shows that if Γ has Minkowski content one, then the curve is rectifiable. Conversely, we consider the functions

$$F_n(s) = \sup_{0 < |h| < 1/n} \left| \frac{z(s+h) - z(s)}{h} - z'(s) \right|$$

Because z is continuous, this supremum can be considered over a countable, dense subset, and so each F_n is measurable. Since $F_n(s) \rightarrow 0$ for almost every s , we can apply Egorov's theorem to show that this limit is uniform except on a singular set E with $|E| < \varepsilon$, so that for some large N , for $s \notin E$ and $|h| < 1/N$, $|z(s+h) - z(s) - hz'(s)| < \varepsilon h$. We now split the interval $[a, b]$ into consecutive intervals I_1, \dots, I_{M+1} , with each interval but I_{M+1} having length $1/N$. We let Γ_n denote the section of the curve travelled along the interval I_n . Thus $|\Gamma^\delta| \leq \sum |\Gamma_n^\delta|$. If an interval I_n contains an

element of E^c , we say I_n is a ‘good’ interval. Then we can pick an element $x_n \in I_n$ for which for any $x \in I_n$,

$$|z(x) - z(x_n) - (x - x_n)z'(x_n)| < \varepsilon|x - x_n| < \varepsilon/N$$

Thus Γ_n is covered by a ε/N thickening of a length $1/N$ line J_n in \mathbf{R}^2 through $z(x_n)$ with slope $z'(x_n)$. Thus if $\varepsilon \leq 1$, we conclude

$$\begin{aligned} |\Gamma_n^\delta| &\leq J_n^{\varepsilon/N+\delta} \leq (1/N + 2\varepsilon/N + 2\delta)(2\varepsilon/N + 2\delta) \\ &\leq 2\delta/N + O(\delta\varepsilon/N + \delta^2 + \varepsilon/N^2) \end{aligned}$$

Since $M \leq NL$, if we take the sum of $|\Gamma_n^\delta|$ over all ‘good’ intervals we obtain an upper bound of

$$NL(2\delta/N + O(\delta\varepsilon/N + \delta^2 + \varepsilon/N^2)) = 2\delta L + O(\delta\varepsilon + \delta^2 N + \varepsilon/N)$$

On the other hand, if I_n is contained within E , or if $n = M + 1$, we say I_n is a bad interval. Since E has total measure bounded by ε , there can be at most $\varepsilon N + 1$ bad intervals. On these intervals we use the crude estimate $|z(t) - z(u)| \leq |t - u|$ (true because z is an arclength parameterization) to show Γ_n is contained in a rectangle with sidelengths $1/N$, so we obtain that $|\Gamma_n^\delta| \leq (1/N + 2\delta)^2 = O(1/N^2 + \delta^2)$. Thus the sum of $|\Gamma_n^\delta|$ over the ‘bad intervals’ is bounded by

$$O(\varepsilon/N + 1/N^2 + \varepsilon N \delta^2 + \delta^2)$$

In particular, the sum of the two bounds gives

$$|\Gamma^\delta| \leq 2\delta L + O(\delta\varepsilon + \delta^2 N + \varepsilon/N + 1/N^2)$$

Or

$$\frac{|\Gamma^\delta|}{2\delta} \leq L + O(\varepsilon + \delta N + \varepsilon/N + 1/N^2)$$

If we choose $N \geq 1/\delta$, we get that

$$\frac{|\Gamma^\delta|}{2\delta} \leq L + O(\varepsilon + \delta N + \delta\varepsilon) = L + O(\varepsilon + \delta N)$$

Letting $\delta \downarrow 0$, we conclude that $M^*(\Gamma) \leq L + O(\varepsilon)$, and we can then let $\varepsilon \downarrow 0$ to conclude $M^*(\Gamma) \leq L$. This completes the proof that if Γ is rectifiable, then Γ has one dimensional Minkowski content, and $M(\Gamma) = L$. \square

If Γ is rectifiable, it is parameterizable by a Lipschitz map (the arclength parameterization). If we instead consider a curve parameterizable by a map z which is Lipschitz of order α , which may no longer be absolutely continuous, but still has a decay very similar to the Minkowski dimension decay.

Theorem 21.21. *If z is a planar curve which is Lipschitz of order $\alpha > 1/2$, then its trace Γ satisfies $|\Gamma^\delta| = O(\delta^{2-1/\alpha})$.*

Proof. Since $|z(t) - z(s)| \leq |t - s|^\alpha$, we can cover z by $O(N)$ radius $1/N^\alpha$ balls, so $|\Gamma| \lesssim N^{1-2\alpha}$, and so $|\Gamma^\delta| \lesssim N(1/N^\alpha + \delta)^2$. Setting $N = \delta^{-1/\alpha} + O(1)$ gives $|\Gamma^\delta| \lesssim \delta^{2-\alpha-1/\alpha}$. \square

21.6 The Isoperimetric Inequality

We now use our Minkowski content techniques to prove the isoperimetric inequality, which asks us to find the region in the plane with largest area whose boundary has a bounded length L . We suppose Ω is a bounded region of the plane, whose boundary $\partial\Omega$ is a rectifiable curve with length L . In particular, we shall find the region with the largest area whose boundary has a fixed length are balls. A key inequality used in the proof is the Brun Minkowski inequality, which lowers bounds the measure of $A + B$ in terms of A and B . If we hope for an estimate $|A + B|^\alpha \gtrsim |A|^\alpha + |B|^\alpha$, then taking $B = \alpha A$, where A is convex and, for which $A + \alpha A = (1 + \alpha)A$, we find $(1 + \alpha)^{d\alpha} \gtrsim (1 + \alpha^{d\alpha})$. Thus $\alpha \geq 1/d$.

Lemma 21.22. *If A , B , and $A + B$ are measurable, $|A + B|^{1/d} \geq |A|^{1/d} + |B|^{1/d}$.*

Proof. Suppose first that A and B are rectangles with side lengths x_n and y_n . Then the Minkowski inequality becomes

$$\left(\prod (x_n + y_n)\right)^{1/d} \geq \left(\prod x_n\right)^{1/d} + \left(\prod y_n\right)^{1/d}$$

Replacing x_n with $\lambda_n x_n$ and y_n with $\lambda_n y_n$, we find that we may assume $x_n + y_n = 1$, and so we must prove that for any $x_n \leq 1$,

$$\left(\prod x_n\right)^{1/d} + \left(\prod (1 - x_n)\right)^{1/d} \leq 1$$

But this inequality is an immediate consequence of the arithmetic geometric mean inequality. Thus the case is proved. Next, we suppose A and B are unions of disjoint closed rectangles, and we prove the inequality by induction on the number of rectangles. Without loss of generality, by symmetry in A and B , we may assume that A has at least two rectangles R_1 and R_2 . Since the inequality is translation invariant separately in A and B , and R_1 and R_2 is disjoint, hence separated by a coordinate axis, we may assume there exists an index j such that every element x of R_1 has $x_j < 0$ and every element x of R_2 has $x_j > 0$. Let $A^+ = A \cap \{x_j \leq 0\}$ and $A^- = A \cap \{x_j \geq 0\}$. Next, we translate B such that if B^\pm are defined similarly, then

$$\frac{|B^\pm|}{|B|} = \frac{|A^\pm|}{|A|}$$

Note that $A + B$ contains the union of $A^+ + B^+$ and $A^- + B^-$, and this union is disjoint. Thus by induction,

$$\begin{aligned} |A + B| &\geq |A^+ + B^+| + |A^- + B^-| \\ &\geq (|A^+|^{1/d} + |B^+|^{1/d})^d + (|A^-|^{1/d} + |B^-|^{1/d})^d \\ &= |A^+| \left(1 + \left(\frac{|B^+|}{|A^+|} \right)^{1/d} \right)^d + |A^-| \left(1 + \left(\frac{|B^-|}{|A^-|} \right)^{1/d} \right)^d \\ &= (|A|^{1/d} + |B|^{1/d})^d \end{aligned}$$

Thus the proof is completed for unions of rectangles. The proof then passes to open sets by approximating open sets by closed rectangles contained within. Then we can pass to where A and B are compact sets, since then $A + B$ is compact, and so if we consider the open thickenings A^ε , B^ε , and $(A + B)^\varepsilon$, then

$$|A| = \lim |A^\varepsilon| \quad |B| = \lim |B^\varepsilon| \quad |A + B| = \lim |(A + B)^\varepsilon|$$

and $(A + B)^\varepsilon \subset A^\varepsilon + B^\varepsilon \subset (A + B)^{2\varepsilon}$. Finally, we can use inner regularity to obtain the theorem in full. \square

Theorem 21.23. *For any region Ω , $4\pi|\Omega| \leq L^2$.*

Proof. For $\delta > 0$, consider

$$\Omega_+(\delta) = \{x : d(x, \Omega) < \delta\} \quad \Omega_-(\delta) = \{x : d(x, \Omega^c) \geq \delta\}$$

Then we have a disjoint union $\Omega_+(\delta) = \Omega_-(\delta) + \Gamma^\delta$, where Γ is the boundary curve of Ω . Furthermore, $\Omega_+(\delta)$ contains $\Omega + B_\delta$, and Ω contains $\Omega_-(\delta) + B_\delta$. Applying the Brun Minkowski inequality, we conclude

$$\begin{aligned} |\Omega_+(\delta)| &\geq (|\Omega|^{1/2} + \pi^{1/2}\delta)^2 \geq |\Omega| + 2\pi^{1/2}\delta|\Omega|^{1/2} \\ |\Omega| &\geq (|\Omega_-(\delta)|^{1/2} + \pi^{1/2}\delta)^2 \geq |\Omega_-(\delta)| + 2\pi^{1/2}\delta|\Omega_-(\delta)|^{1/2} \end{aligned}$$

But

$$|\Gamma^\delta| = |\Omega_+(\delta)| - |\Omega_-(\delta)| \geq 2\pi^{1/2}\delta \left(|\Omega|^{1/2} + |\Omega_-(\delta)|^{1/2} \right)$$

Dividing by 2δ and letting $\delta \rightarrow 0$, we conclude $L \geq 2\pi^{1/2}|\Omega|^{1/2}$. This is precisely the inequality we need. \square

Using some Fourier analysis, we can prove that the only smooth curves which make this inequality tight are circles. Indeed, if a closed C^1 curve $\Gamma = \{z(t) : a \leq t \leq b\}$ is given, then Green's theorem implies the area of its interior is given by

$$\frac{1}{2} \left| \int_{\Gamma} x dy - y dx \right| = \frac{1}{2} \left| \int_a^b x(t)y'(t) - y(t)x'(t) dt \right|$$

We then take a Fourier series in x and y .

Theorem 21.24. *The only curves Γ with rectifiable boundary such that $A = \pi(L/2)^2$ are circles.*

Proof. By normalizing, we may assume z is an arcline parameterization, and Γ has length 2π , so $z : [0, 2\pi] \rightarrow \mathbf{R}^2$, and z is absolutely continuous. If $x(t) \sim \sum a_n e^{nit}$ and $y(t) \sim \sum b_n e^{int}$, then $x'(t) \sim \sum i n a_n e^{nit}$ and $y(t) \sim \sum i n b_n e^{nit}$. Parseval's equality implies

$$\int_0^{2\pi} x(t)y'(t) - y(t)x'(t) dt = 2\pi i \sum n(b_n \overline{a_n} - a_n \overline{b_n})$$

Thus the area of the curve is precisely

$$\pi \left| \sum n(b_n \overline{a_n} - a_n \overline{b_n}) \right| \leq \pi \sum 2n |b_n a_n| \leq \pi \sum |n|(|a_n|^2 + |b_n|^2)$$

On the other hand, the length constraint implies that, since $|z'(t)| = 1$,

$$1 = \frac{1}{2\pi} \int_0^{2\pi} x'(t)^2 + y'(t)^2 dt = \sum |n|^2(|a_n|^2 + |b_n|^2)$$

If $A = \pi$, then

$$\sum |n|(|a_n|^2 + |b_n|^2) \geq 1 = \sum |n|^2(|a_n|^2 + |b_n|^2)$$

This means we cannot have $|n| < |n|^2$ whenever a_n or b_n is nonzero. Thus the Fourier support of x and y is precisely $\{-1, 0, 1\}$. Since x is real valued, $a_1 = \overline{a_{-1}} = a$, $b_1 = \overline{b_{-1}}$. We thus have $2(|a_1|^2 + |b_1|^2) = 1$, and since we must have a a scalar multiple of b so the Cauchy Schwarz inequality application becomes an equality, we must have $|a_1| = |b_1| = 1/2$. If $a_1 = e^{i\alpha}/2$ and $b_1 = e^{i\beta}/2$, the fact that $1 = 2|a_1 \overline{b_1} - \overline{a_1} b_1|$ implies $|\sin(\alpha - \beta)| = 1$, hence $\alpha - \beta = k\pi/2$, where k is an odd integer. Thus $x(s) = \cos(\alpha + s)$, and $y(s) = \cos(\beta + s)$, which parameterizes a circle. \square

Chapter 22

Singular Integral Operators

Let us now consider some kernel operators

$$Tf(y) = \int K(x, y)f(x) dx$$

where $K(x, y)$ is singular for $x = y$. The prototypical example is the Hilbert transform on \mathbf{R} , i.e.

$$Hf(y) = \int \frac{f(x - y)}{y} dy.$$

One cannot even interpret the right hand side in the Lebesgue sense because the function $1/y$ is not Lebesgue integrable. We can proceed

22.1 The Calderon-Zygmund Decomposition

The main trick of the Calderon-Zygmund decomposition, to exploit the local cancellation properties of the singular kernel K , is to decompose a general function into a bounded term (the ‘good’ term), and a family of terms which are locally oscillating (the ‘bad’ terms). The resulting decomposition is called the Calderon-Zygmund decomposition. The trick here is to exploit the compatibility of measure and metric that has already been exploited, e.g. in proving Hardy-Littlewood maximal bounds. We begin with a version of the decomposition obtained by dyadic methods.

Theorem 22.1. Let $f \in L^1(\mathbf{R}^d)$ and set $\lambda > 0$. Then there exists a set \mathcal{Q} of disjoint, dyadic cubes, and a decomposition

$$f = g + \sum_{Q \in \mathcal{Q}} b_Q,$$

such that

$$\|g\|_{L^1(\mathbf{R}^d)} \leq \|f\|_{L^1(\mathbf{R}^d)} \quad \text{and} \quad \|g\|_{L^\infty(\mathbf{R}^d)} \lesssim_d \lambda,$$

and for $Q \in \mathcal{Q}$, $\text{supp}(b_Q) \subset Q$, $\int b_Q = 0$, and $\|b_Q\|_{L^1(\mathbf{R}^d)} \lesssim_d \lambda |Q|$. We have

$$\bigcup \mathcal{Q} = \{x : M_\Delta f(x) > \lambda\},$$

where M_Δ is the maximal averaging operator over dyadic cubes. From the weak L^1 bound for this operator we thus obtain that

$$\sum_{Q \in \mathcal{Q}} |Q| \leq \frac{\|f\|_{L^1(\mathbf{R}^d)}}{\lambda}.$$

Proof. Set $c = 2^d$. Call a dyadic cube Q *bad* if $\int_Q |f(x)| dx > \lambda$. Since $f \in L^1(\mathbf{R}^d)$, any cube with sidelength exceeding $\|f\|_{L^1(\mathbf{R}^d)}/\lambda$ is good. Thus every bad cube is contained in a *maximal* bad cube, and the union of these bad cubes is precisely $\{x : M_\Delta f(x) > \lambda\}$. Let \mathcal{Q} be the set of all maximal bad cubes. Define

$$g(x) = \sum_{Q \in \mathcal{Q}} \mathbf{I}(x \in Q) \int_Q f(y) dy + \mathbf{I}(M_\Delta f(y) \leq \lambda) f(y).$$

By the Lebesgue density theorem, $|f| \leq M_\Delta f$ almost everywhere. Thus $|f(y)| \leq \lambda$ for almost every y with $M_\Delta f(y) \leq \lambda$. Conversely, if Q is a maximal bad cube, then the parent cube Q' of Q is good, which means that

$$\int_Q |f(y)| dy \leq 2^d \int_{Q'} |f(y)| dy \leq 2^d \lambda.$$

Thus we conclude that $\|g\|_{L^\infty(\mathbf{R}^d)} \lesssim_d \lambda$. It is also easy to see that $\|g\|_{L^1(\mathbf{R}^d)} \leq \|f\|_{L^1(\mathbf{R}^d)}$. On the other hand, if we set

$$b_Q(x) = \mathbf{I}(x \in Q) [f(x) - \int_Q f(y) dy],$$

then $\int b_Q = 0$, $\text{supp}(b_Q) \subset Q$, and $\int |b_Q(x)| dx \leq 2 \int_Q |f(x)| dx \leq 2^{d+1} \lambda$. □

Remark. On a general measure space X , for $f \in L^1(X)$ one can write $f = f\mathbf{I}_{x \leq \lambda} + f\mathbf{I}_{x > \lambda} = g + b$. We have

$$\|g\|_{L^1(X)} \leq \|f\|_{L^1(X)} \quad \text{and} \quad \|g\|_{L^\infty(X)} \leq \lambda.$$

And for the bad function,

$$\|b\|_{L^1(X)} \leq \|f\|_{L^1(X)} \quad \text{and} \quad |\text{supp}(b)| \leq \|f\|_{L^1(X)}/\lambda.$$

Thus the Calderon-Zygmund decomposition obtains a similar version of this result, but accounting for additional metric structure.

Remark. For each point $x \in \mathbf{R}^d$ contained in a bad cube, let T_x be the integer n such that the maximal bad cube containing x has sidelength 2^{-n} . Otherwise, let $T_x = \infty$. Then in the language of conditional expectations, $g = \mathbf{E}[f|T_x]$. Thus one can view the Calderon-Zygmund decomposition as a kind of non-probabilistic *stopping time* result.

The key idea in the proof above was to first decompose the open set $\Omega = \{x : M_\Delta f(x) > \lambda\}$ into disjoint dyadic cubes, which gives another very useful geometric result, the *Whitney decomposition*.

Theorem 22.2. *Let $\Omega \subset \mathbf{R}^d$ be an open set. For any $K \geq 1$, there exists a decomposition Ω into a disjoint union of dyadic cubes Q , where for any $Q \in \mathcal{Q}$, $\text{diam}(Q) \sim Kd(Q, \Omega^c)$.*

Proof. Let \mathcal{Q}' denote the dyadic cubes Q contained in Ω such that

$$K\text{diam}(Q) \leq d(Q, \Omega^c) \leq 4K\text{diam}(Q).$$

These cubes cover Ω . Indeed, for each $x \in \Omega$, one can find a dyadic cube Q with diameter between $d(x, \Omega^c)/4K$ and $d(x, \Omega^c)/2K$ containing x . Then

$$\begin{aligned} d(Q, \Omega^c) &\geq d(x, \Omega^c) - \text{diam}(Q) \\ &\geq (1 - 1/4K) \cdot d(x, \Omega^c) \\ &\geq (2K - 1/2) \cdot \text{diam}(Q) \\ &\geq K\text{diam}(Q). \end{aligned}$$

Conversely,

$$d(Q, \Omega^c) \leq d(x, \Omega^c) \leq 4K\text{diam}(Q).$$

But now we can take a maximal subfamily \mathcal{Q} of \mathcal{Q}' . □

Remark. Consider the decomposition \mathcal{Q} in the last theorem. If Q_1, Q_2 are two nearby cubes in the decomposition, and

$$d(Q_1, Q_2) \leq \delta(\text{diam}(Q_1) + \text{diam}(Q_2))$$

for some $\delta > 0$, then

$$\begin{aligned} |d(Q_1, \Omega^c) - d(Q_2, \Omega^c)| &\leq d(Q_1, Q_2) + \text{diam}(Q_1) + \text{diam}(Q_2) \\ &\leq (1 + \delta)(\text{diam}(Q_1) + \text{diam}(Q_2)). \end{aligned}$$

Thus

$$\text{diam}(Q_2) \leq \frac{d(Q_2, \Omega^c)}{K} \leq \frac{d(Q_1, \Omega^c)}{K} + \frac{1 + \delta}{K} \text{diam}(Q_1) + \frac{1 + \delta}{K} \text{diam}(Q_2).$$

Rearranging, we find that for $\delta \leq K/2 - 1$,

$$\frac{\text{diam}(Q_2)}{2} \leq \frac{d(Q_1, \Omega^c)}{K} + (1 + \delta) \text{diam}(Q_1) \leq (5 + \delta) \text{diam}(Q_1),$$

and so

$$\text{diam}(Q_2) \leq (5 + \delta) \cdot \text{diam}(Q_1).$$

By symmetry, we automatically obtain the two sided inequality

$$(5 + \delta)^{-1} \cdot \text{diam}(Q_1) \leq \text{diam}(Q_2) \leq (5 + \delta) \cdot \text{diam}(Q_1).$$

Thus balls which are close together also have comparable diameter.

Given the Whitney decomposition, how does Calderon-Zygmund follow? Set $\Omega = \{x : M_\Delta f(y) > \lambda\}$, and let \mathcal{Q} denote the Whitney decomposition. If $Q \in \mathcal{Q}$, then $d(Q, \Omega^c) \lesssim \text{diam}(Q)$. Thus we can find a point x with $d(x, Q) \lesssim \text{diam}(Q)$ and with $M_\Delta f(x) \leq \lambda$. Thus for a universal constant $C > 0$, $x \in C \cdot Q$, and so

$$\int_Q |f(y)| dy \leq C^d \int_{CQ} |f(y)| dy \leq C^d \lambda.$$

Writing $b_Q(x) = \mathbf{I}(x \in Q) \cdot [f(x) - (\int_Q f(y) dy)]$ and $g = f - \sum b_Q$, we obtain an analogous decomposition to the one obtained before.

Certainly one cannot prove an exact analogy of the Whitney decomposition for *balls*, i.e. one cannot cover a general open set by disjoint open balls with radius comparable to their distance to the boundary of the open set. Nonetheless, we can obtain such a decomposition with the *bounded intersection property*, i.e. such that no point is contained in too many of the balls in the particular cover.

Theorem 22.3. *Let $\Omega \subset \mathbf{R}^d$ be an open set. For any $K \geq 1$, Ω is the union of a family of balls \mathcal{B} , where for each ball $B \in \mathcal{B}$, the radius r_B of this ball is comparable to $Kd(Q, \Omega^c)$, and each point in Ω is contained in $O_d(1)$ balls.*

Proof. Applying the last remark, without loss of generality assume $K \geq 2$, and pick $\delta = 1$. Consider the cubes Q in the decomposition above, and let \mathcal{B} be a family of balls obtained by covering each cube Q in \mathcal{Q} by a ball B with $\text{diam}(B) = \text{diam}(Q)$. If a point x is contained in a family of balls B_1, \dots, B_k corresponding to dyadic cubes Q_1, \dots, Q_k , then $d(Q_i, Q_j) \leq \text{diam}(Q_i) + \text{diam}(Q_j)$. Thus we find that for any Q_i and Q_j

$$\text{diam}(Q_i) \leq 50 \cdot \text{diam}(Q_j).$$

Thus there is a common quantity integer n such that for all i , $2^n \leq \text{diam}(Q_i) \leq 50^2 \cdot 2^n$. The number of dyadic lattice points whose vertices have coordinates which can be written as a fraction with denominator 2^n , and which lie a distance at most $50^2 \cdot 2^n$ from x is $O_d(1)$, which shows $k \lesssim_d 1$, and thus we obtain the required bounded intersection property. \square

This decomposition using balls is the one most generalizable to more general situations where one is working in a space without quite as regular a tiling as the family of cubes. To obtain this generalization, we assume the slightly stronger assumptions we made in our analysis of the Hardy-Littlewood maximal function, i.e. we assume we are working on a locally compact topological space X equipped with a locally finite Radon measure such that the technical assumptions made there, and the slightly stronger *engulfing* and *doubling* conditions mentioned, hold true. To generalize the Whitney decomposition, we fix two additional constants c^* and c^{**} with $1 < c^* < c^{**}$ which will depend only on c_1 and c_2 , and for a ball $B = B(x, \delta)$, we let $B^* = B(x, c^* \delta)$ and $B^{**} = B(x, c^{**} \delta)$.

Theorem 22.4. *Let $\Omega \subset X$ be an open set. Then there is a collection of pairwise disjoint balls \mathcal{B} , such that*

$$\bigcup_{B \in \mathcal{B}} B^* = \Omega, \quad \text{and} \quad B^{**} \cap \Omega^c \neq \emptyset \text{ for any } B \in \mathcal{B}.$$

It is simple to construct from these balls a disjoint family of sets Q_B for each $B \in \mathcal{B}$ such that $B \subset Q_B \subset B^$, and $\bigcup Q_B = \Omega$, which can be taken as a substitute for cubes in the original Whitney covering lemma.*

Remark. As with the Euclidean case above, the balls $\{B^* : B \in \mathcal{B}\}$, though not disjoint, have the bounded intersection property. This follows from a similar argument to that of the above argument. First we show intersecting balls have comparable radii. Suppose $B_1, B_2 \in \mathcal{B}$ and $B_1 \cap B_2 \neq \emptyset$, where $B_i = B(x_i, r_i)$, and $r_1 \leq r_2$. Then

$$B_1^{**} = B(x_1, c^{**}r_1) \subset B(x_2, c_2 c^{**}r_1).$$

Since $B_2^* = B(x_2, c^*r_2)$ is disjoint from Ω^c , whereas B_1^{**} is not disjoint from Ω^c it follows that $c^*r_2 \leq c_2 c^{**}r_1$, so that $r_2 \lesssim r_1$, and by symmetry, $r_1 \sim r_2$. Now if B_1^*, \dots, B_N^* are balls whose intersection contains a point x_0 , where $B_i = B(x_i, r_i)$. Then the balls all have comparable radii, i.e. if $r = \max(r_1, \dots, r_N)$ then $r \sim r_i$. Set $B = B(x_0, r)$. The doubling / engulfing property implies that $|B_i| \sim |B|$ for all i . But also $B_i \subset B(x_0, c_1 r)$ for all i , and since the $\{B_i\}$ are disjoint, we find that

$$N|B| \lesssim |B_1| + \dots + |B_N| \leq |B(x_0, c_1 r)| \lesssim |B|.$$

Thus we find $N \lesssim 1$, i.e. so we have the bounded intersection property.

Proof.

□

Chapter 23

Fourier Multiplier Operators

Our aim in this chapter is to study the boundedness of *Fourier multiplier operators*. Given a function $m : \mathbf{R}^d \rightarrow \mathbf{C}$, known as a *symbol*, we want to associate a multiplier operator T , sometimes denoted $m(D)$, which when applied to a function $f : \mathbf{R}^d \rightarrow \mathbf{C}$ should be formally given by the equation

$$Tf(x) = \int_{\mathbf{R}^d} m(\xi) \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

In maximum generality, for any tempered distribution m on \mathbf{R}^d we can define T as a continuous operator from $\mathcal{S}(\mathbf{R}^d)$ to itself. But often times we will consider much more regular symbols m , i.e. those which are locally integrable functions, and our goal will be to obtain stronger continuity statements for the associated multiplier operators. If K is the tempered distribution which is the Fourier transform of m , then $Tf = K * f$. Thus Fourier multiplier operators are precisely the same as the class of convolution operators formed by tempered distributions. In any case, the map $m \mapsto m(D)$ gives an injective *algebra homomorphism* from the family of all tempered distributions to the family of continuous operators on $\mathcal{S}(\mathbf{R}^d)$ (another injective homomorphism is obtained by considering the multiplier operators $m \mapsto m(X)$, where $m(X)$ is the multiplier operator $m(X)f = m \cdot f$, the family of *spatial* multiplier operators, which have a much simpler theory). The main goal, of course, is to determine what properties of the symbol or it's Fourier transform imply boundedness properties of the operator T .

Remark. In engineering these operators are known as *filters*, and occur in a variety of contexts. Due to the presence of error the regularity of these op-

erators are of utmost importance. The function m is known as the *system-transfer function*, *optical-transfer function*, or *frequency response*, depending on the context, and the function K is known as the *point-spread function*.

Example. Over \mathbf{R} , consider a rough cutoff $\mathbf{I}_{[-1,1]}$. Then we calculate explicitly that

$$K(x) = \int_{-1}^1 e^{2\pi i \xi \cdot x} = \frac{\sin(2\pi x)}{\pi x}.$$

Thus convolution by K acts by cutting off higher frequency parts of the function. In engineering this operator is called a low pass filter.

Example. Over \mathbf{R} , we consider the Fourier multiplier

$$m(\xi) = -i \cdot \operatorname{sgn}(\xi).$$

Then $m(D)$ is the Hilbert transform.

Example. In \mathbf{R}^d , we consider the Fourier multiplier

$$m_R(\xi) = \mathbf{I}(|\xi| \leq R).$$

The operator $m_R(D)$ is known as the ball multiplier operator. More generally, given any compact set S we can consider the Fourier multiplier $\mathbf{I}_S(D)$. In the engineering literature these multipliers are called ideal low pass filters.

Example. In this chapter, it is natural to renormalize the differentiation operators $D^\alpha : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}(\mathbf{R}^d)$ so that for $f \in \mathcal{S}(\mathbf{R}^d)$,

$$\widehat{D^\alpha f} = \xi^\alpha \widehat{f}.$$

In particular, this implies that if $m(\xi) = \xi_i^\alpha$, then $m(D) = D^\alpha$. More generally, if $m(\xi) = \sum_{|\alpha| \leq k} c_\alpha \xi^\alpha$, then

$$m(D) = \sum_{|\alpha| \leq k} c_\alpha D^\alpha.$$

Thus the family of Fourier multiplier operators contains all constant coefficient differential operators.

Fourier multiplier operators have proved essential to our study of classical Fourier analysis. In particular, we have used Fourier multiplier operators to prove a great many results; the convolution operator by the Poisson kernel is a Fourier multiplier given by the symbol $e^{-|x|}$, and the heat kernel is a Fourier multiplier with symbol $e^{-\pi|x|^2}$. This is no coincidence. It is a general heuristic that any well-behaved translation invariant operator is given by convolution with an appropriate function.

We have already seen in our study of distributions that any translation invariant continuous linear operator $T : C_c^\infty(\mathbf{R}^d) \rightarrow C_{\text{loc}}^\infty(\mathbf{R}^d)$ is given by convolution with a distribution. If the distribution is tempered, we can take the Fourier transform to conclude that the operator is a Fourier multiplier operator. In fact, if $1 \leq p, q \leq \infty$ and a translation invariant operator T satisfies a bound of the form

$$\|Tf\|_{L^q(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

for any $f \in \mathcal{S}(\mathbf{R}^d)$, then T is a Fourier multiplier operator. To prove this, we apply the theory of Sobolev embeddings.

Theorem 23.1. *Suppose $1 \leq p, q \leq \infty$, and $T : \mathcal{S}(\mathbf{R}^d) \rightarrow L^q(\mathbf{R}^d)$ is a linear map commuting with translations and satisfies*

$$\|Tf\|_{L^q(\mathbf{R}^d)} \leq \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in \mathcal{S}(\mathbf{R}^d)$. Then T is a Fourier multiplier operator.

Proof. For any $f \in \mathcal{S}(\mathbf{R}^d)$, $Tf \in W^{q,n}(\mathbf{R}^d)$ for any $n > 0$. To see this, we note that for any $h > 0$ and $k \in \{1, \dots, d\}$, and

$$(\Delta_h f)(x) = \frac{f(x + he_k) - f(x)}{h}.$$

Then $\Delta_h(Tf) = T(\Delta_h f)$ because T is translation invariant. Since f is a Schwartz function, $\Delta_h f$ converges to $D^k f$ in $L^p(\mathbf{R}^d)$. Thus by continuity of f , Tf has a strong derivative $T(D^k f)$ in $L^q(\mathbf{R}^d)$. Induction shows Tf has strong derivatives of all orders. The last lemma shows that $Tf \in C(\mathbf{R}^d)$,

and

$$\begin{aligned}
\|Tf\|_{L^\infty(\mathbf{R}^d)} &\lesssim \sum_{|\alpha| \leq n+1} \|D^\alpha(Tf)\|_{L^q(\mathbf{R}^d)} \\
&= \sum_{|\alpha| \leq n+1} \|T(D^\alpha f)\|_{L^q(\mathbf{R}^d)} \\
&\lesssim \sum_{|\alpha| \leq n+1} \|D^\alpha f\|_{L^q(\mathbf{R}^d)}.
\end{aligned}$$

The map $f \mapsto Tf(0)$ is thus a continuous operator on $\mathcal{S}(\mathbf{R}^d)$, and therefore defines a tempered distribution Λ . Translation invariance shows that $Tf = \Lambda * f$, and setting $m = \hat{\Lambda}$ completes the proof. \square

Remark. It follows from this argument that if $T : \mathcal{S}(\mathbf{R}^d) \rightarrow L^q(\mathbf{R}^d)$ is a linear operator commuting with translations satisfying a bound

$$\|Tf\|_{L^q(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)},$$

then for any $f \in \mathcal{S}(\mathbf{R}^d)$, $Tf \in C_{\text{loc}}^\infty(\mathbf{R}^d)$ and is slowly increasing, as is all of its derivatives.

23.1 Frequency Localization

One use of Fourier multipliers is to localize the support of the Fourier transform of a function to a portion of space. Given a compactly supported function m supported on some set S , and any function f , $\widehat{m(D)f} = m\hat{f}$ is supported on S . Frequency localization has the additional feature that $m(D)f$ is smooth (analytic even), since its Fourier transform is compactly supported. One major advantage of localizing in frequency is that it enables us to control the behaviour of the derivatives of a function; if m is a cutoff function supported ξ_0 , then $\widehat{m(D)f}$ is supported near ξ_0 , and so we might expect that $\partial_\alpha(m(D)f) \approx \xi_0^\alpha f$. This makes frequency localization useful especially useful in problems involving derivatives. The uncertainty principle also gives another heuristic application of this principle; if m is supported on a cube centred at the origin with sidelengths R_1, \dots, R_d , then the function $m(D)f$ will be roughly speaking, locally constant on ‘dual rectangles’ with sidelength $1/R_1, \dots, 1/R_d$. More generally, if

we instead choose m supported on a cube centered at ξ_0 with sidelengths R_1, \dots, R_n , then f acts roughly like a constant multiple of $e^{2\pi i \xi \cdot x}$ on dual rectangles. This trick comes up all over the place since the behaviour of certain operators can be exploited more clearly once inputs or outputs have been localized in frequency.

Let us consider some examples. If we consider the Fourier multiplier $\mathbf{I}_{[-R,R]}$ in \mathbf{R}^1 , then this multiplier corresponds to the kernel

$$K(x) = \int_{-R}^R e^{2\pi i \xi x} = \frac{\sin(2\pi R x)}{\pi x} = 2R \cdot \text{sinc}(2\pi R x),$$

i.e. a sinc function. If f is supported on an interval I , then it follows from a simple estimate that

$$|(K * f)(x)| \lesssim \frac{\|f\|_{L^1(\mathbf{R})}}{d(x, I)}.$$

Thus we see that we can localize in frequency while remaining somewhat localized in space, but with some additional fuzziness that decays away from this interval at a rate of $1/x$. Unfortunately, this is often not enough to obtain useful estimates, e.g. the fuzziness does not even lie in $L^1(\mathbf{R})$. In \mathbf{R}^d , if we write $K = K_1 \otimes \dots \otimes K_d$ be the kernel associated with the rectangle multiplier, then we have a similar $1/x$ decay estimate (though we do get up to $1/x^d$ decay in directions not close to being parallel with any axis, an error that still remains non integrable).

We can do better if we use a smooth cutoff function, i.e. we choose a smooth non-negative function m compactly supported on a cube I of sidelength L which equals one on the interior third of I , and satisfies

$$\|D^\alpha m\|_{L^\infty(\mathbf{R}^d)} \lesssim_\alpha L^{-|\alpha|}$$

for all multi-indices α , and then consider the Fourier multiplier $m(D)$. Then $m(D)$ corresponds to the convolution kernel

$$K(x) = \int e^{2\pi i \xi \cdot x} m(\xi) d\xi.$$

Integrating by parts gives that for all $n > 0$, $|K(x)| \lesssim_n L^{d-n} |x|^{-n}$, which decreases away from the origin much faster than for a rough cutoff. This

implies that we have the decay estimates

$$|(K * f)(x)| \lesssim_n \frac{L^{d-n} \|f\|_{L^1(\mathbf{R}^d)}}{d(x, J)^n}.$$

for f supported on a cube J , and all $n > 0$. Thus the majority of the mass of $K * f$ is concentrated in an $O(1/L)$ -thickening of the interval J , so spatial localization is preserved provided the sidelengths of J are $\Omega(1/L)$. This means the ‘fuzziness’ outside of the interval J here is fast decaying in a quantitative manner, which is often enough that it can be safely ignored. Thus frequency localization in this setting also preserves spatial localization as long as we respect the uncertainty principle, i.e. we do not care about localization in space to an accuracy finer than $\Omega(1/L)$.

23.2 L^p Regularity

We now wish to study conditions on m which guarantee bounds of the form

$$\|m(D)f\|_{L^q(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

for all $f \in \mathcal{S}(\mathbf{R}^d)$. This question was of course prominent throughout the study of Harmonic analysis, but the explicit problem of studying general multipliers was pushed forwards by analysts like Hörmander in the 1960s.

Example. Let $k_s(x) = |x|^{-s}$, for $0 < s < d$ be the Riesz kernel. Then $k_s \in L^{r,\infty}(\mathbf{R}^d)$ for $r = d/s$, so the weak-type variant of Young’s inequality implies that

$$\|k_s * f\|_{L^q(\mathbf{R}^d)} \lesssim_{p,q} \|f\|_{L^p(\mathbf{R}^d)}$$

provided $1/p - 1/q = 1 - s/d$. Analysing the scaling properties of each side of this equation shows these are the only exponents for which this inequality holds. Now it turns out that $\widehat{k_s} = C_{d,s}k_{d-s}$ for some constants $C_{d,s}$, and so it follows that $k_s \in M^{p,q}(\mathbf{R}^d)$ when $1/p - 1/q = s/d$.

In general, a characterization of the tempered distributions which give bounded convolution operators is unknown except in a few very particular situations. For each $1 \leq p \leq q \leq \infty$, we let $\|m\|_{M^{p,q}(\mathbf{R}^d)}$ denote the operator norm of $m(D)$ from $\mathcal{S}(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^d)^*$, via the $L^p(\mathbf{R}^d)$ norm in the input,

and via the $L^q(\mathbf{R}^d)$ norm in the output, i.e. it is the smallest quantity $\|m\|_{M^{p,q}(\mathbf{R}^d)}$ such that

$$\|m(D)f\|_{L^q(\mathbf{R}^d)} \leq \|m\|_{M^{p,q}(\mathbf{R}^d)} \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in \mathcal{S}(\mathbf{R}^d)$. If $1 \leq p < \infty$, then the Hahn-Banach density theorem shows every element of $M^{p,q}(\mathbf{R}^d)$ extends uniquely to an operator from $L^p(\mathbf{R}^d)$ to $L^q(\mathbf{R}^d)$ with the same operator norm. If $p = \infty$, then every such operator extends to a map from $L^\infty(\mathbf{R}^d)$ to $L^q(\mathbf{R}^d)$ with the same operator norm, but this extension need not be unique. We let $M^{p,q}(\mathbf{R}^d)$ be the set of tempered distributions for which the bound is finite. For simplicity, we also let $M^p(\mathbf{R}^d)$ denote $M^{p,p}(\mathbf{R}^d)$. All of these spaces are Banach spaces. By symmetries of the Fourier transform, it is easy to check that translations, modulations, and dilations all preserve the space $M^{p,q}(\mathbf{R}^d)$, though dilations need not preserve the $M^{p,q}$ norm unless $p = q$. Thus we have a family of Banach Algebras $M^p(\mathbf{R}^d)$. On these spaces, translation, dilation, conjugation, and modulation are all isometries, so this space is highly symmetric.

Littlewood's principle tells us that the only interesting Fourier multipliers in $M^{p,q}(\mathbf{R}^d)$ occur with 'the larger exponent on the left'.

Theorem 23.2. Fix $1 \leq q < p \leq \infty$, and suppose m is a tempered distribution on \mathbf{R}^d satisfying a uniform bound

$$\|m(D)f\|_{L^q(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in \mathcal{S}(\mathbf{R}^d)$. Then $m = 0$. In other words, $M^{p,q}(\mathbf{R}^d) = 0$ for $q < p$.

Proof. Suppose $m \neq 0$ and $q < p$. Then there is $f_0 \in \mathcal{S}(\mathbf{R}^d)$ with $m(D)f_0 \neq 0$. Thus $m(D)f_0$ lies in $L^q(\mathbf{R}^d)$. Fix a large integer N and pick $x_1, \dots, x_N \in \mathbf{R}^d$ separated far enough apart that

$$\left\| \sum_{n=1}^N \text{Trans}_{x_n} f_0 \right\|_{L^p(\mathbf{R}^d)} \gtrsim N^{1/p} \|f_0\|_{L^p(\mathbf{R}^d)}$$

and

$$\left\| \sum_{n=1}^N \text{Trans}_{x_n} m(D)f_0 \right\|_{L^q(\mathbf{R}^d)} \sim N^{1/q} \|m(D)f_0\|_{L^q(\mathbf{R}^d)} \lesssim N^{1/q} \|f_0\|_{L^p(\mathbf{R}^d)}.$$

Translation invariance of convolution shows $N^{1/q} \lesssim N^{1/p}$, which is impossible for suitably large N . Thus $m = 0$. \square

Example. We claim that $M^{2,2}(\mathbf{R}^d) = L^\infty(\mathbf{R}^d)$, in the sense that the two spaces consist of the same family of distributions, and the norms on both spaces are equal, i.e. $\|m\|_{M^{2,2}(\mathbf{R}^d)} = \|m\|_{L^\infty(\mathbf{R}^d)}$. To see this, suppose either $m \in M^{2,2}(\mathbf{R}^d)$ or $m \in L^\infty(\mathbf{R}^d)$, and let

$$\Phi(x) = e^{-\pi|x|^2}$$

be the Gaussian distribution. Then Φ is a Schwartz function, and satisfies

$$\widehat{m(D)\Phi} = \Phi \cdot m.$$

Since $\Phi \in L^2(\mathbf{R}^d)$, $\Phi \cdot m \in L^2(\mathbf{R}^d)$. But since $1/\Phi \in L_{loc}^\infty(\mathbf{R}^d)$, this implies that $m \in L_{loc}^1(\mathbf{R}^d)$.

Now we know m is locally integrable, we can obtain the general result by Parseval's inequality, since a bound of the form

$$\|m(D)f\|_{L^2(\mathbf{R}^d)} \leq C\|f\|_{L^2(\mathbf{R}^d)}$$

holds for all $f \in \mathcal{S}(\mathbf{R}^d)$ if and only if

$$\|m \cdot g\|_{L^2(\mathbf{R}^d)} \leq C\|g\|_{L^2(\mathbf{R}^d)}$$

for all $g \in \mathcal{S}(\mathbf{R}^d)$. The former bound is equivalent to $\|m\|_{M^{2,2}(\mathbf{R}^d)} \leq C$, and the second bound is equivalent to $\|m\|_{L^\infty(\mathbf{R}^d)} \leq C$ by Hölder's inequality.

For any tempered distribution m and $f, g \in \mathcal{S}(\mathbf{R}^d)$, the fact that the Fourier transform is self adjoint implies that

$$\begin{aligned} \langle m(D)f, g \rangle &= \langle m \cdot \widehat{f}, \widehat{g} \rangle \\ &= \langle \widehat{f}, \overline{m} \cdot \widehat{g} \rangle \\ &= \langle f, \overline{m}(D)g \rangle. \end{aligned}$$

Thus we have an adjoint relation $m(D)^* = \overline{m}(D)$, which gives a natural duality theory for Fourier multiplier operators. In particular, this means that if R is the reflection operator $[Ru](x) = u(-x)$, then

$$m(D)f = [(R \circ m^*(D) \circ R)f^*]^*.$$

In particular, for any $0 < p \leq \infty$, $\|m(D)f\|_{L^p(\mathbf{R}^d)} = \|m(D)^*Rf^*\|_{L^p(\mathbf{R}^d)}$.

Theorem 23.3. For any $1 \leq p \leq q < \infty$ and any tempered distribution m ,

$$\|m\|_{M^{p,q}(\mathbf{R}^d)} = \|m\|_{M^{q^*,p^*}(\mathbf{R}^d)}.$$

Proof. It will suffice to prove by duality that

$$\|m\|_{M^{q^*,p^*}(\mathbf{R}^d)} \leq \|m\|_{M^{p,q}(\mathbf{R}^d)}.$$

Assume without loss of generality that $\|m\|_{M^{p,q}(\mathbf{R}^d)} < \infty$. Given $f \in \mathcal{S}(\mathbf{R}^d)$, we know $m(D)f \in C_{\text{loc}}^\infty(\mathbf{R}^d) \cap L^q(\mathbf{R}^d)$. If $p \neq \infty$, then

$$\|m(D)f\|_{L^{p^*}(\mathbf{R}^d)} = \sup \left\{ \int m(D)f(x)g(x) dx : g \in \mathcal{S}(\mathbf{R}^d), \|g\|_{L^p(\mathbf{R}^d)} < \infty \right\}.$$

Given any such $g \in \mathcal{S}(\mathbf{R}^d)$, we have by Hölder's inequality,

$$\begin{aligned} \int m(D)f(x)g(x) dx &= \int f(x)m^*(D)g(x) dx \\ &\leq \|f\|_{L^{q^*}(\mathbf{R}^d)} \|m^*(D)g\|_{L^q(\mathbf{R}^d)} \\ &\leq \|m\|_{M^{p,q}(\mathbf{R}^d)} \|f\|_{L^{q^*}(\mathbf{R}^d)}. \end{aligned}$$

Thus we conclude that

$$\|m(D)f\|_{L^{p^*}(\mathbf{R}^d)} \leq \|m\|_{M^{p,q}(\mathbf{R}^d)} \|f\|_{L^{q^*}(\mathbf{R}^d)}$$

and thus $\|m\|_{M^{q^*,p^*}(\mathbf{R}^d)} \leq \|m\|_{M^{p,q}(\mathbf{R}^d)}$. On the other hand, if $p = \infty$, then $q = \infty$ (or else the inequality is trivial). If $\|m\|_{M^{\infty,\infty}(\mathbf{R}^d)} < \infty$, then it follows that for any $f \in \mathcal{S}(\mathbf{R}^d)$, $m(D)f \in C^\infty(\mathbf{R}^d)$. It therefore follows by smoothness that

$$|m(D)f(0)| \leq \|m(D)f\|_{L^\infty(\mathbf{R}^d)} \leq \|m\|_{M^{\infty,\infty}(\mathbf{R}^d)} \|f\|_{L^\infty(\mathbf{R}^d)}.$$

A consequence of this is that for all $f \in \mathcal{S}(\mathbf{R}^d)$,

$$\left| \int \hat{m}(x)f(x) dx \right| \leq \|m\|_{M^{\infty,\infty}(\mathbf{R}^d)} \|f\|_{L^\infty(\mathbf{R}^d)}.$$

This means \hat{m} is a distribution of order zero, and is therefore equal to some Radon measure μ . Moreover, the bound actually implies that μ is a finite Borel measure, with total variation at most $\|m\|_{M^{\infty,\infty}(\mathbf{R}^d)}$. But then Young's inequality implies that

$$\|\hat{m} * f\|_{L^1(\mathbf{R}^d)} \leq \|\hat{m}\|_{M(\mathbf{R}^d)} \|f\|_{L^1(\mathbf{R}^d)} \leq \|m\|_{M^{\infty,\infty}(\mathbf{R}^d)} \|f\|_{L^1(\mathbf{R}^d)}.$$

Thus $\|m\|_{M^{1,1}(\mathbf{R}^d)} \leq \|m\|_{M^{\infty,\infty}(\mathbf{R}^d)}$, which completes the proof of duality. \square

We also have a rescaling law which follows from the standard rescaling properties of the Fourier transform, i.e. for any matrix $A \in \text{GL}(d)$,

$$\|m \circ A\|_{M^{p,q}(\mathbf{R}^d)} = |\det(A)|^{-|1/p-1/q|} \|m\|_{M^{p,q}(\mathbf{R}^d)}.$$

In particular, the quantities $\|m\|_{M^p(\mathbf{R}^d)}$ are invariant under scaling.

In particular, if $1 \leq p \leq \infty$ and $m \in M^p(\mathbf{R}^d)$, then also $m \in M^{p^*}(\mathbf{R}^d)$ and so Riesz-interpolation implies $m \in M^2(\mathbf{R}^d)$. Thus if we are studying L^p to L^p boundedness for any $1 \leq p \leq \infty$, we may restrict our attention to Fourier multipliers with a bounded symbol. More generally, this interpolation approach shows that $M^p(\mathbf{R}^d)$ is a larger space of multipliers the closer p is to two. Thus $M^2(\mathbf{R}^d) = L^\infty(\mathbf{R}^d)$ is the largest family, and $M^1(\mathbf{R}^d) = M^\infty(\mathbf{R}^d)$ is the smallest family of multipliers bounded on some $M^p(\mathbf{R}^d)$. It turns out we can also completely characterize this space of multipliers, it is the space of Fourier transforms of finite Radon measures.

Example. *The only remaining space which we can completely characterize are the spaces $M^{1,q}(\mathbf{R}^d)$ (and thus $M^{q^*,\infty}$), where $1 \leq q \leq \infty$. For $q > 1$, we have*

$$M^{1,q}(\mathbf{R}^d) = \widehat{L^q(\mathbf{R}^d)}$$

and $M^{1,1}(\mathbf{R}^d) = \widehat{M(\mathbf{R}^d)}$, the set of all finite Borel measures. Moreover, in the case $q > 1$ we have

$$\|m\|_{M^{1,q}(\mathbf{R}^d)} = \|\widehat{m}\|_{L^q(\mathbf{R}^d)}$$

and for $q = 1$,

$$\|m\|_{M^{1,1}(\mathbf{R}^d)} = \|\widehat{m}\|_{M(\mathbf{R}^d)}.$$

Given such an $m \in M^{1,q}(\mathbf{R}^d)$, if $K = \widehat{m}$, and $\{\phi_\varepsilon\}$ is a family of standard mollifiers, then

$$\|K * \phi_\varepsilon\|_{L^q(\mathbf{R}^d)} = \|m(D)(\phi_\varepsilon)\|_{L^q(\mathbf{R}^d)} \lesssim \|\phi_\varepsilon\|_{L^1(\mathbf{R}^d)} = 1.$$

Applying the Banach-Alaoglu theorem, the family $\{K * \phi_\varepsilon\}$ has a weak $*$ convergent subsequence in $L^q(\mathbf{R}^d)^{**}$. Thus there is $\lambda \in L^q(\mathbf{R}^d)^{**}$ and $\varepsilon_i \rightarrow 0$ such that $K * \phi_{\varepsilon_i} \rightarrow \lambda$. But this means that $K * \phi_{\varepsilon_i}$ converges to λ distributionally. It also converges to K distributionally, so $K = \lambda \in (L^q(\mathbf{R}^d))^{**}$. If $q > 1$, then $(L^q(\mathbf{R}^d))^{**} = L^q(\mathbf{R}^d)$, with the same norm, and $(L^1(\mathbf{R}^d))^{**} = M(\mathbf{R}^d)$.

Another case where the multiplier problem is trivial is for $m \in L^\infty(\mathbf{R}^d)$ with $\widehat{m} \geq 0$. This is related to the fact that the study of kernels which are non-negative and homogeneously distributed is also trivial, though the fact that we are working over a space with infinite measure complicates the analysis somewhat. To argue more precisely, by Lemma 4.12, $\|m\|_{L^\infty(\mathbf{R}^d)} = \|\widehat{m}\|_{L^1(\mathbf{R}^d)}$, and so for $1 \leq p \leq \infty$,

$$\|\widehat{m}\|_{L^1(\mathbf{R}^d)} = \|m\|_{M^2} \leq \|m\|_{M^p} \leq \|m\|_{M^1} = \|\widehat{m}\|_{L^1(\mathbf{R}^d)},$$

so $\|m\|_{M^p} = \|\widehat{m}\|_{L^1(\mathbf{R}^d)}$ for all $p \in [1, \infty]$.

We have a version of Young's inequality for multipliers. A scaling analysis shows that we can only have an inequality

$$\|m_1 * m_2\|_{M^{p,q}(\mathbf{R}^d)} \leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)}$$

if $1 + (1/p - 1/q) = 1/r_1 + 1/r_2$.

Lemma 23.4. *If $m_1 \in L^{r_1}(\mathbf{R}^d)$ and $m_2 \in L^{r_2}(\mathbf{R}^d)$, then*

$$\|m_1 * m_2\|_{M^{p,q}(\mathbf{R}^d)} \leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)}$$

for $1 \leq p, q \leq \infty$, and $1 \leq r_1, r_2 \leq 2$, and $1 + |1/p - 1/q| = 1/r_1 + 1/r_2$.

Proof. We apply complex interpolation. Young's inequality implies $m_1 * m_2 \in L^r$ for $r = 1/|1/p - 1/q|$. In particular, if $p = q = 2$, then $m_1 * m_2 \in L^\infty(\mathbf{R}^d)$, which means

$$\|m_1 * m_2\|_{M^2(\mathbf{R}^d)} = \|m_1 * m_2\|_{L^\infty(\mathbf{R}^d)} \leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)}.$$

If $p = q = 2$,

If $p = 1$, then $2 - 1/q = 1/r_1 + 1/r_2$ and $1 \leq r_1, r_2 \leq 2$. We may then apply Hölder's inequality and the Hausdorff Young inequality to conclude that

$$\begin{aligned} \|m_1 * m_2\|_{M^{1,q}(\mathbf{R}^d)} &= \|\widehat{m_1 * m_2}\|_{L^q(\mathbf{R}^d)} \\ &= \|\widehat{m_1} \widehat{m_2}\|_{L^q(\mathbf{R}^d)} \\ &\leq \|\widehat{m_1}\|_{L^{r_1^*}(\mathbf{R}^d)} \|\widehat{m_2}\|_{L^{r_2^*}(\mathbf{R}^d)} \\ &\leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)}. \end{aligned}$$

Next, for $p = q = 2$, we have $1/r_1 + 1/r_2 = 1$, and so we can again apply Hausdorff Young to conclude that

$$\begin{aligned}\|m_1 * m_2\|_{M^2(\mathbf{R}^d)} &= \|m_1 * m_2\|_{L^\infty(\mathbf{R}^d)} \\ &\leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)}.\end{aligned}$$

For the remaining cases, given an arbitrary $f \in \mathcal{S}(\mathbf{R}^d)$, and $m_1, m_2 \in C_c^\infty(\mathbf{R}^d)$, the quantity

$$\|(m_1 * m_2)f\|_{L^p(\mathbf{R}^d)} = \left(\left| \int (m_1 * m_2)(\xi) \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi \right|^p dx \right)^{1/p}$$

is holomorphic in p , and so we can apply complex interpolation to prove that

$$\|(m_1 * m_2)f\|_{L^q(\mathbf{R}^d)} \leq \|m_1\|_{L^{r_1}(\mathbf{R}^d)} \|m_2\|_{L^{r_2}(\mathbf{R}^d)} \|f\|_{L^p(\mathbf{R}^d)}. \quad \square$$

Essentially for any other family of multipliers, it is still incredibly difficult to determine conditions when, especially when $d > 1$. For instance, it still remains a major open question in harmonic analysis, for $d > 2$, to determine the values of $p \in [1, \infty]$ and $\delta > 0$ for which the multiplier

$$m^\delta(\xi) = (1 - |\xi|^2)_+^\delta = \max((1 - |\xi|^2)^\delta, 0)$$

lies in $M^p(\mathbf{R}^d)$, a problem known as the *Bochner-Riesz conjecture*.

The difficulty which causes m^δ to be unbounded is that it is singular on the boundary of the unit sphere, which is a large, curved set, upon which m^δ has δ degrees of regularity in the direction tangential to the sphere. Intuition suggests that a smooth Fourier multiplier would have a rapidly decaying Fourier transform, which would therefore be well posed as a convolution operator. One basic instance of this phenomenon occurs if $m \in \mathcal{S}(\mathbf{R}^d)$. Then $\hat{m} \in \mathcal{S}(\mathbf{R}^d)$, hence integrable, and by Young's convolution inequality, it follows that for any $p \leq q$, $\|m\|_{M^{p,q}(\mathbf{R}^d)} \lesssim 1$, and the induced map from $\mathcal{S}(\mathbf{R}^d)$ to $M^{p,q}(\mathbf{R}^d)$ is continuous. Another example is the following calculation: if $\Omega \subset \mathbf{R}^d$ is bounded and open, C is a fixed constant, and $L : \mathbf{R}^d \rightarrow \mathbf{R}^d$ is an invertible linear map, then we say ϕ is a *bump function adapted to $L(\Omega)$* if ϕ is smooth and supported in $L(\Omega)$, and $|D^k(\phi \circ L)(x)| \leq C$ for all $x \in \Omega$. If m is a bump function adapted to $L(\Omega)$, then it follows that $\|m\|_{M^{p,q}(\mathbf{R}^d)} \lesssim_{C,\Omega} 1$.

It is important to note that it is mainly qualitative information that is of interest here. This is because $\|m \circ A\|_{M^p(\mathbf{R}^d)} = \|m\|_{M^p(\mathbf{R}^d)}$ for any multiplier m and invertible linear transformation A . Thus despite the fact that a singularity can become quantitatively more singular as we focus in on it by introducing an appropriate transformation A , the $M^p(\mathbf{R}^d)$ norm stays the same.

Example. Consider the sawtooth function $m(\xi) = \operatorname{sgn}(\xi) \max(1 - |\xi|, 0)$. For each $R > 0$, consider $m_R(\xi) = \operatorname{sgn}(\xi) \max(1 - |\xi/R|, 0)$. Then m_R converges pointwise almost everywhere to $m_\infty(\xi) = \operatorname{sgn}(\xi)$, and actually $m_R(D)$ converges to $m_\infty(D)$ distributinally. Here $m_\infty(D)$ is just the Hilbert transform. Now $\|m_R\|_{M^p(\mathbf{R}^d)} = \|m\|_{M^p(\mathbf{R}^d)}$ for all $p \in [1, \infty]$, so we might expect that $\|m\|_{M^p(\mathbf{R}^d)} = \|m_\infty\|_{M^p(\mathbf{R}^d)}$. Certainly the distributional limiting process implies that

$$\|m_\infty\|_{M^p(\mathbf{R}^d)} \leq \|m\|_{M^p(\mathbf{R}^d)}.$$

Conversely, $m(\xi) = a(\xi)m_\infty(\xi)$, where $a(\xi) = \max(1 - |\xi|, 0)$ is a tent function. Now $\hat{a}(x) = \operatorname{sinc}(x)^2$ is non-negative, so in particular, this means that for any p , $\|a\|_{M^p(\mathbf{R}^d)} = \|a\|_{L^\infty(\mathbf{R}^d)} = 1$. But this means that

$$\|m\|_{M^p(\mathbf{R}^d)} \leq \|a\|_{M^p(\mathbf{R}^d)} \|m_\infty\|_{M^p(\mathbf{R}^d)} \leq \|m_\infty\|_{M^p(\mathbf{R}^d)},$$

which completes the proof.

If the multiplier m is only singular on a small set, we can likely still obtain some interesting estimates. For instance, the Hilbert transform, corresponding to the Fourier multiplier $m(\xi) = i \operatorname{sgn}(\xi)$, is singular at $\xi = 0$, but still satisfies the bounds

$$\|Hf\|_{L^p(\mathbf{R}^d)} \lesssim_p \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in L^p(\mathbf{R}^d)$ and $1 < p < \infty$. One can see this from the Calderon-Zygmund theory of singular integrals, or via the Hörmander-Mikhlin theory we will develop shortly. It follows from translation symmetry that for $1 < p < \infty$ and any (possibly unbounded interval) I ,

$$\|I_I\|_{M^p(\mathbf{R}^d)}, \|I_I\|_{M^p(\mathbf{R}^d)} \lesssim_p 1.$$

Now suppose $m \in L^\infty(\mathbf{R}^d)$ has *bounded variation*, which means the quantity

$$V(m) = \sup_{\xi_1 < \dots < \xi_N} \sum_{i=1}^{N-1} |m(\xi_{i+1}) - m(\xi_i)|.$$

is finite. Then m has countably many discontinuities, and the variation upper bound prevents m from being too nonsmooth. In this case, we can obtain an operator norm bound.

Theorem 23.5. *For any symbol m on \mathbf{R} , and any $1 < p < \infty$,*

$$\|m\|_{M^p(\mathbf{R})} \lesssim_p \|m\|_{L^\infty(\mathbf{R})} + V(m).$$

Proof. For each n , pick ξ_1, \dots, ξ_{N_n} such that

$$\sum_{i=1}^{N-1} |m(\xi_{i+1}) - m(\xi_i)| \geq V(m) - 1/n.$$

If we define

$$m_n = m(\xi_1)\mathbf{I}_{(-\infty, \xi_1)} + \sum_{i=1}^{N_n-1} m(\xi_i)\mathbf{I}_{(\xi_i, \xi_{i+1})} + m(\xi_N)\mathbf{I}_{(\xi_N, \infty)}$$

Then $\|m - m_n\|_{L^1(\mathbf{R})} \leq 1/n$. But this means that $m_n(D)$ converges to $m(D)$ weakly as operators on $\mathcal{S}(\mathbf{R})$, and so

$$\|m\|_{M^p(\mathbf{R})} \leq \limsup_{n \rightarrow \infty} \|m_n\|_{M^p(\mathbf{R})}.$$

Now we can rewrite

$$m_n(\xi) = m(\xi_1)\mathbf{I}_{(-\infty, \xi_1)} + \sum_{i=1}^{N-1} [m(\xi_i) - m(\xi_{i+1})]\mathbf{I}_{(\xi_1, \xi_i)}(\xi) + m(\xi_N)\mathbf{I}_{(\xi_N, \infty)}.$$

Since the multiplier operators corresponding to indicator functions of intervals are uniformly bounded in $M^p(\mathbf{R})$ for $1 < p < \infty$, we conclude that

$$\|m_n\|_{M^p(\mathbf{R})} \lesssim_p |m(\xi_1)| + \sum_{i=1}^{N-1} |m(\xi_i) - m(\xi_{i+1})| + |m(\xi_N)| \leq \|m\|_{L^\infty(\mathbf{R})} + V(m).$$

Taking $n \rightarrow \infty$ gives $\|m\|_{M^p(\mathbf{R})} \lesssim_p \|m\|_{L^\infty(\mathbf{R})} + V(m)$. \square

The theorem of Hörmander-Mikhlin is a more sophisticated instance of the smoothness-regularity phenomenon, giving L^p to L^p bounds to Fourier multipliers which decay smoothly away from the origin. There are several

formulations of the phenomenon, of greater and greater generality. Without loss of generality, we will assume that $m \in L^\infty(\mathbf{R}^d)$, since otherwise m cannot lie in any of the spaces M^p . Let us state them in increasing order of generality, the former being most easily seen as relating to the smoothness of the multipliers:

- The multiplier m lies in $C_{\text{loc}}^\infty(\mathbf{R}^d - \{0\})$, and satisfies

$$|D_\xi^\beta m(\xi)| \lesssim_\beta |\xi|^{-\beta} \quad (23.1)$$

for all multi-indices β . In particular, this is true if m is smooth away from the origin, and homogeneous of degree zero.

- For some integer N with $N > d/2$, m has weak derivatives up to order N which are functions, and satisfy (23.1) for all β with $|\beta| \leq N$.
- For some integer $N > d/2$, m has weak derivatives in $L_{\text{loc}}^2(\mathbf{R}^d)$ up to order N , and satisfies an L^2 average estimate

$$\sup_{R>0} \left(\int_{|\xi| \leq 2R} |\xi|^{2\beta} |D_\xi^\beta m(\xi)|^2 d\xi \right)^{1/2} \lesssim 1.$$

- For some non-zero, smooth radial test function φ , compactly supported away from the origin,

$$\sup_{R>0} \|\varphi \text{Dil}_R m\|_{L_\alpha^2} < \infty \quad (23.2)$$

for some $\alpha > d/2$ (not necessarily an integer).

- For some φ as above, and for some $\varepsilon > 0$,

$$\sup_{R>0} \int |\mathcal{F}(\varphi \text{Dil}_R m)(x)| (1 + |x|)^\varepsilon dx < \infty.$$

If the second last point holds, the ε in the last point can be chosen to be any number smaller than $\alpha - d/2$. This result is closely related to the theory of singular integrals. In particular, under the first two assumptions,

Theorem 23.6. Consider $m \in L^\infty(\mathbf{R}^d)$ and suppose there exists $\varepsilon > 0$ and φ as above with

$$\sup_{R>0} \int |\mathcal{F}(\varphi \text{Dil}_R m)(x)| (1 + |x|)^\varepsilon dx < \infty.$$

Then for any $1 < p < \infty$ and $f \in \mathcal{S}(\mathbf{R}^d)$,

$$\|m(D)f\|_{L^p(\mathbf{R}^d)} \lesssim_p \|f\|_{L^p(\mathbf{R}^d)}.$$

Moreover, we have a bound $\|m(D)f\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$, which, together with the fact that $m \in M^2(\mathbf{R}^d)$ because it is bounded, implies the bounds above by Riesz-Thorin and duality.

Proof. If the assumptions hold for some ε , it is simple to argue they hold for any such φ . In particular, if we take a non-negative φ such that for any $\xi \neq 0$,

$$\sum_{n=-\infty}^{\infty} \varphi(2^n \xi) = 1.$$

Write $m_n(\xi) = \varphi(\xi)m(2^n \xi)$, and $K_n = \widehat{m_n}$. Then our assumption implies that

$$\int K_n(x)(1 + |x|)^\varepsilon dx \lesssim 1,$$

In particular, this means that

$$\int_{|x| \geq R} |K_n(x)| dx \leq (1 + R)^{-\varepsilon}.$$

uniformly in n . Similarly, Bernstein's inequality implies that

$$\|\nabla K_n\|_{L^1(\mathbf{R}^d)} \lesssim \|K_n\|_{L^1(\mathbf{R}^d)} \lesssim 1.$$

This implies that for all $y \in \mathbf{R}^d$, uniformly in n ,

$$\int |K_n(x + y) - K_n(x)| dx \lesssim |y|.$$

For any $f \in \mathcal{S}(\mathbf{R}^d)$, we have

$$m(D)f = \sum_{n=-\infty}^{\infty} (\text{Dil}_{2^n} m_n)(D),$$

where the sum converges absolutely in $L^p(\mathbf{R}^d)$. Since $\widehat{\text{Dil}_{2^n} m_n} = 2^{nd} \text{Dil}_{1/2^n} K_n$, it follows that

$$K * f = \sum_{n=-\infty}^{\infty} 2^{nd} \cdot (\text{Dil}_{1/2^n} K_n) * f.$$

The cancellation bound we obtained for the functions K_n indicate the singular-integral nature of the kernels K_n , i.e. they satisfy a cancellation criterion, and the singularity is quantitatively concentrated near the origin. Indeed, the cancellation bound allows us to use the standard Calderon-Zygmund singular integral results to obtain a uniform bound

$$\|K_n * f\|_{L^p(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

in n . However, these bounds cannot be summed in n as $n \rightarrow \infty$ to yield an L^p bound on $K * f$, since the L^p operator norm of $\text{Dil}_{1/2^n} K_n$ is the same as the L^p operator norm of K_n . To get a better bound, we must utilize the fact that the functions decay away from the origin more strongly. We will still use the Calderon Zygmund decomposition to understand this. So fix an integrable function b supported on a cube Q of some sidelength R , with $\int b(x) dx = 0$. Since convolution commutes with translation, we may assume without loss of generality for the calculation of L^p norms that Q is centred at the origin. Let Q^* denote the cube with the same centre and twice the width. Then

$$\begin{aligned} \int_{(Q^*)^c} 2^{nd} |(\text{Dil}_{1/2^n} K_n) * b| &\leq 2^{nd} \int_{(Q^*)^c} \left| \int_Q K_n(2^n(x-y)) b(y) dy \right| dx \\ &\leq 2^{nd} \int_{(Q^*)^c} \left| \int_Q K_n(2^n(x-y)) b(y) dy \right| dx \\ &= 2^{nd} \int_{(Q^*)^c} \left| \int_Q [K_n(2^n(x-y)) - K_n(2^n x)] b(y) dy \right| dx \\ &\leq 2^{nd} \int_Q \int_{(Q^*)^c} |b(y)| |K_n(2^n(x-y)) - K_n(2^n x)| dx dy \\ &= \int_Q \int_{(Q^*)^c} |b(y)| |K_n(x - 2^n y) - K_n(x)| dx dy \\ &\lesssim 2^n \int_Q |y| |b(y)| dx \lesssim 2^n R \cdot \|b\|_{L^1(\mathbf{R}^d)}. \end{aligned}$$

This is a standard Calderon-Zygmund type calculation, as in the singular integral theory, and gives good bounds for $R \leq 1/2^n$. For $R \geq 1/2^n$, we

apply the decay estimate to get a more optimal bound, writing

$$\begin{aligned}
\int_{(Q^*)^c} 2^{nd} |(\text{Dil}_{1/2^n} K_n) * b| &\leq 2^{nd} \int_{(Q^*)^c} \int_Q |K_n(2^n(x-y))| |b(y)| \, dy \, dx \\
&\leq 2^{nd} \int_{|z| \geq R} \int_Q |K_n(2^n z)| |b(y)| \, dy \, dz \\
&\leq (1 + R/2^n)^{-\varepsilon} \|b\|_{L^1(\mathbf{R}^d)}
\end{aligned}$$

This is a good for $R \geq 2^n$. In particular, we now have good bounds for any value of R , which we can sum up to conclude that

$$\int_{(Q^*)^c} |K * b| \lesssim \|b\|_{L^1(\mathbf{R}^d)}.$$

Now given a general $f \in \mathcal{S}(\mathbf{R}^d)$, we apply a Calderon-Zygmund decomposition, fixing $\alpha > 0$, and writing

$$f = g + \sum_{k=1}^{\infty} b_k$$

where $\|g\|_{L^1(\mathbf{R}^d)} + \sum_{k=1}^{\infty} \|b_k\|_{L^1(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$, $\|g\|_{L^\infty(\mathbf{R}^d)} \lesssim \alpha$, and there exists a family of disjoint cubes $\{Q_k\}$ such that b_k is supported on Q_k , $\int b_k = 0$, and $\sum_{k=1}^{\infty} |Q_k| \lesssim \alpha^{-1} \|f\|_{L^1(\mathbf{R}^d)}$. If $\Omega = \bigcup_{k=1}^{\infty} Q_k$, then

$$\int_{\Omega^c} |K * (\sum_{k=1}^{\infty} b_k)| \lesssim \sum_{k=1}^{\infty} \|b_k\|_{L^1(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}.$$

Thus

$$|\{x \in \Omega^c : |K * (\sum_{k=1}^{\infty} b_k)| \geq \alpha/2\}| \lesssim \|f\|_{L^1(\mathbf{R}^d)}/\alpha,$$

and since $|\Omega| \leq \|f\|_{L^1(\mathbf{R}^d)}/\alpha$,

$$|\{x \in \mathbf{R}^d : |K * (\sum_{k=1}^{\infty} b_k)| \geq \alpha/2\}| \lesssim \|f\|_{L^1(\mathbf{R}^d)}/\alpha.$$

Since α was arbitrary, we conclude that $\|K * (\sum b_k)\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$. Next, we use the fact that K is bounded on L^2 (since m is bounded) to conclude that

$$\|K * g\|_{L^2(\mathbf{R}^d)} \lesssim \|g\|_{L^2(\mathbf{R}^d)} \lesssim \alpha^{1/2} \|g\|_{L^1(\mathbf{R}^d)}^{1/2}$$

and so

$$|\{x \in \mathbf{R}^d : |(K * g)(x)| \geq \alpha/2\}| \lesssim \|K * g\|_{L^2(\mathbf{R}^d)}^2 \alpha^{-2} \lesssim \|g\|_{L^1(\mathbf{R}^d)} \alpha^{-1}.$$

But now we know $\|K * g\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim \|g\|_{L^1(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$. We sum and get $\|K * f\|_{L^{1,\infty}(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$. \square

Remark. A fastidious reader may complain that the Calderon-Zygmund decomposition of a Schwartz function yields a family of functions that are only integrable, and not necessarily even continuous. To remedy this situation, we replace the kernel operator K we are originally studying with the more regularized kernels $K^{\leq N} = \sum_{|n| \leq N} 2^{nd} \cdot \text{Dil}_{1/2^n} K_n$. The operators $K^{\leq N}$ are then Schwartz convolution kernels, and, independently of this proof, are thus easily seen to lie in $M^p(\mathbf{R}^d)$ for all $1 \leq p \leq \infty$, and thus extend to be well defined on the components of Calderon-Zygmund decompositions of functions. The important part of this proof is that it shows that the operators $K^{\leq N}$ are *uniformly* when they induce convolution operators from $L^1(\mathbf{R}^d)$ to $L^{1,\infty}(\mathbf{R}^d)$, and thus imply the resultant bounds for the limiting convolution kernel K .

Since convolution can be seen as a way of smoothing out some of the irregularities of a function, one might ask whether the convolution of a multiplier in $M^{p,q}(\mathbf{R}^d)$ with a function lies in $M^{p,q}(\mathbf{R}^d)$. One can obtain such a result, provided the function is integrable.

Theorem 23.7. *If $u \in L^1(\mathbf{R}^d)$, then $\|m * u\|_{M^{p,q}(\mathbf{R}^d)} \leq \|m\|_{M^{p,q}(\mathbf{R}^d)} \|u\|_{L^1(\mathbf{R}^d)}$.*

Proof. Let $K = \hat{m}$. If $v = \hat{u}$, then

$$\begin{aligned} (m * u)(D)f(x) &= \int K(x-y)v(x-y)f(y) dy \\ &= \int u(\xi) \int K(x-y)f(y)e^{2\pi i(x-y)\cdot\xi} dy d\xi \\ &= \int u(\xi) \cdot (\text{Mod}_\xi K * f)(x) d\xi \\ &= \int u(\xi)(\text{Trans}_\xi m)(D)f(x) d\xi. \end{aligned}$$

Since $\|\text{Trans}_\xi m\|_{M^{p,q}(\mathbf{R}^d)} = \|m\|_{M^{p,q}(\mathbf{R}^d)}$, the result follows. \square

One might ask whether multipliers operators need at least *some* regularity to get a bound at all. Provided that $1 \leq p \leq q \leq 2$, one at least needs some kind of integrability condition.

Theorem 23.8. *If $m \in M^{p,q}(\mathbf{R}^d)$, and $1 \leq p \leq q \leq 2$, $M^{p,q}(\mathbf{R}^d) \subset L^q(\mathbf{R}^d)^*$.*

Proof. TODO □

On the other hand, if $q > 2$, then $M^{p,q}(\mathbf{R}^d)$ contains distributions of positive order. We will show an explicit examples for large $d \geq 4$, and non constructive examples otherwise.

Example. Recall that the surface measure σ on the sphere in \mathbf{R}^d satisfies

$$|\hat{\sigma}(\xi)| \lesssim_d |\xi|^{-\frac{d-1}{2}}.$$

Consider the multiplier corresponding to the distribution $\Lambda = \partial\sigma/\partial r$, which is not a distribution of order zero. On the other hand, $|\hat{\Lambda}(\xi)| \lesssim_d |\xi|^{-\frac{d-3}{2}}$. Provided $d \geq 4$, this implies that $\|\hat{\Lambda}\|_{L^q(\mathbf{R}^d)} < \infty$ for $(d-3)/2q > d$, i.e. for $q > 2d/(d-3) = 2 + 6/(d-3)$, and thus $\|\Lambda\|_{M^{1,q}(\mathbf{R}^d)} < \infty$ for this range. For each $\varepsilon > 0$, if we take d appropriately large, apply duality, and interpolate, for each $\varepsilon > 0$, we obtain a distribution Λ of positive order, which lies in $M^{p,q}(\mathbf{R}^d)$ for all (p,q) with $p < 2 - \varepsilon$ and $q > 2 + \varepsilon$.

Example. Our other, non constructive examples apply Baire category arguments. Fix $\eta \in C_c^\infty(\mathbf{R}^d)$, and for any $\lambda > 0$, let $m_\lambda(\xi) = \eta(\xi)e^{2\pi i\lambda|\xi|^2}$. We claim that if $K_\lambda = \widehat{m_\lambda}$, $\|K_\lambda\|_{L^\infty(\mathbf{R}^d)} \lesssim \lambda^{-d/2}$. This follows from a simple oscillatory integral argument. Also K_λ will be essentially supported on a ball of radius $O(\lambda)$ at the origin, so one can show from this that $\|K_\lambda\|_{L^1(\mathbf{R}^d)} \lesssim \lambda^{d/2}$. Thus $\|m_\lambda\|_{M^{1,\infty}} \lesssim \lambda^{-d/2}$ and $\|m_\lambda\|_{M^{1,1}} = \|m_\lambda\|_{M^{\infty,\infty}} \lesssim \lambda^{d/2}$. Interpolating gives for $q > 2$ and $q \geq p$, a bound of the form $\|m_\lambda\|_{M^{p,q}(\mathbf{R}^d)} \lesssim \lambda^{-\varepsilon}$ for some $\varepsilon > 0$. If $M^{p,q}(\mathbf{R}^d)$ solely contained distributions of order zero, if $\chi \in C_c^\infty(\mathbf{R}^d)$ is a bump function equal to one in a neighborhood of the support of η , then the operator $m \mapsto \chi m$ would be a bounded operator from $M^{p,q}(\mathbf{R}^d)$ to $M(\mathbf{R}^d)$ by the closed graph theorem. But this would imply that

$$\|\eta\|_{L^1(\mathbf{R}^d)} = \|\chi m_\lambda\|_{M(\mathbf{R}^d)} \lesssim \|m_\lambda\|_{M^{1,q}(\mathbf{R}^d)} \lesssim \lambda^{-\varepsilon},$$

which certainly cannot be true. Thus $M^{p,q}(\mathbf{R}^d)$ contains distributions of positive order for $q > 2$ and $p < 2$.

De Leeuw's theorem shows slices of continuous $d+1$ dimensional multipliers are bounded by the original multiplier, so that we might view multipliers in higher dimensions as more complicated than lower dimensional multipliers.

Theorem 23.9. *Let $m \in C(\mathbf{R}^{d+1})$. Fix $\xi_0 \in \mathbf{R}$, and define $m_0 \in C(\mathbf{R}^d)$ by setting*

$$m_0(\xi) = m(\xi, \xi_0).$$

Then for any $1 \leq p \leq \infty$, $\|m_0\|_{MP(\mathbf{R}^d)} \leq \|m\|_{MP(\mathbf{R}^{d+1})}$.

Proof. Without loss of generality, assume $\xi_0 = 0$. For $\lambda > 0$ set

$$L(\xi_1, \dots, \xi_{d+1}) = (\xi_1, \dots, \xi_d, \xi_{d+1}/\lambda).$$

Then

$$\|m \circ L_\lambda\|_{MP(\mathbf{R}^{d+1})} = \|m\|_{MP(\mathbf{R}^{d+1})}.$$

Take $\lambda \rightarrow \infty$. Since m is continuous, $m \circ L_\lambda$ converges to $m \circ L_\infty$ pointwise, where $L_\infty(\xi_1, \dots, \xi_{d+1}) = (\xi_1, \dots, \xi_d, 0)$. On the other hand,

$$\|m \circ L_\infty\|_{MP(\mathbf{R}^d)} = \|m_0\|_{MP(\mathbf{R}^d)}.$$

Indeed, this follows from the simple fact that $(m \circ L_\infty)(D) = m_0(D) \otimes 1$, and $m_0(D) \otimes 1$ has the same operator norm as $m_0(D)$. Thus it suffices to show that

$$\|m \circ L_\infty\|_{MP(\mathbf{R}^d)} \leq \limsup_{\lambda \rightarrow \infty} \|m \circ L_\lambda\|_{MP(\mathbf{R}^d)}.$$

But this follows from a simple weak convergence argument; for any $f, g \in \mathcal{S}(\mathbf{R}^{d+1})$, the dominated convergence theorem implies that

$$\lim_{\lambda \rightarrow \infty} |\langle (m \circ L_\lambda)(D)f, g \rangle| = \langle (m \circ L_\infty)(D)f, g \rangle. \quad \square$$

Example. *If the cone multiplier*

$$m_\lambda(\xi, \eta) = \left(1 - \frac{|\xi|^2}{|\eta|^2}\right)_+^\lambda$$

lies in $MP(\mathbf{R}^{d+1})$, then the corresponding Bochner-Riesz multiplier

$$m_\lambda(\xi) = (1 - |\xi|^2)_+^\lambda$$

lies in $MP(\mathbf{R}^d)$.

If m is no longer continuous, this theorem only holds for *almost all* slices of the function.

Theorem 23.10. Fix $m \in M^p(\mathbf{R}^{d+1})$, and for each λ , let $m_\lambda(\xi) = m(\xi, \lambda)$ be defined on \mathbf{R}^d . Then for almost every $\lambda \in \mathbf{R}$, $\|m_\lambda\|_{M^p(\mathbf{R}^d)} \leq \|m\|_{M^p(\mathbf{R}^{d+1})}$.

Proof. By duality and the multiplication formula for the Fourier transform,

$$\left| \int m(\xi) \widehat{f}(\xi) \widehat{g}(\xi) d\xi \right| \leq \|m\|_{M^p(\mathbf{R}^{d+1})} \|f\|_{L^p(\mathbf{R}^{d+1})} \|g\|_{L^q(\mathbf{R}^{d+1})},$$

where q is the dual of p . If $f = f_1 \otimes f_2$ and $g = g_1 \otimes g_2$ then

$$\left| \int m_\lambda(\xi) \widehat{f}_1(\xi) \widehat{f}_2(\lambda) \widehat{g}_1(\xi) \widehat{g}_2(\lambda) d\xi d\lambda \right| \leq \|m\|_{M^p(\mathbf{R}^d)} \|f\|_{L^p(\mathbf{R}^d)} \|g\|_{L^q(\mathbf{R}^d)}.$$

Write the left hand side as

$$\left| \int J(\lambda) \widehat{f}_2(\lambda) \widehat{g}_2(\lambda) d\lambda \right|,$$

where

$$J(\lambda) = \int m_\lambda(\xi) \widehat{f}_1(\xi) \widehat{f}_2(\xi) d\xi.$$

The inequality above this implies that

$$\|J\|_{L^\infty(\mathbf{R})} \leq \|J\|_{M^p(\mathbf{R})} \leq \|m\|_{M^p(\mathbf{R}^{d+1})} \|f_1\|_{L^p(\mathbf{R}^d)} \|g_1\|_{L^p(\mathbf{R}^d)}.$$

But this implies precisely that for almost every $\lambda \in \mathbf{R}$,

$$|J(\lambda)| \leq \|f_1\|_{L^p(\mathbf{R}^d)} \|g_1\|_{L^p(\mathbf{R}^d)},$$

which completes the proof (modulo a separability argument). \square

23.3 Transference

One can view the last result of the last section as a *transference theorem*, showing that the study of the boundedness of multipliers becomes harder

in higher dimensions (since a higher dimensional result implies a corresponding lower dimensional result for the slices). Our goal now is to analyze a similar correspondence between the theory of multipliers on \mathbf{R}^d and the theory of multipliers on \mathbf{T}^d .

Let us briefly describe the analogous properties of multipliers on \mathbf{T}^d that we have proved for \mathbf{R}^d . There exists a theory of Schwartz and tempered sequences $\{a_m : m \in \mathbf{Z}^d\}$ analogous to that of Schwartz functions and tempered distributions on \mathbf{T}^d and \mathbf{R}^d , such that the Fourier transform induces an isomorphism between Schwartz functions on \mathbf{T}^d and Schwartz sequences on \mathbf{Z}^d , and tempered distributions on \mathbf{T}^d with tempered sequences on \mathbf{Z}^d . One can take the convolution in both spaces. If $T : C^\infty(\mathbf{T}^d) \rightarrow C^\infty(\mathbf{T}^d)$ is any continuous operator commuting with translations, then there exists a distribution Λ on \mathbf{T}^d such that $Tf = \Lambda * f$ for any $f \in C^\infty(\mathbf{T}^d)$. Applying the Fourier transform, for any such operator there is also a tempered sequence $\{a_m : m \in \mathbf{Z}^d\}$ such that

$$\widehat{Tf}(n) = a_n \widehat{f}(n),$$

so we have a theory of Fourier multipliers on \mathbf{T}^d . For any $1 \leq p, q \leq \infty$, we let $M^{p,q}(\mathbf{Z}^d)$ denote the family of all tempered sequences $\{a_m\}$ such that the induced Fourier multiplier operator T is bounded from the $L^p(\mathbf{T}^d)$ norm to the $L^q(\mathbf{T}^d)$ norm on smooth functions, with the associated operator norm giving the space a topology. Similarly, we define $M^p(\mathbf{Z}^d) = M^{p,q}(\mathbf{Z}^d)$. We have $M^{p,q}(\mathbf{Z}^d) = M^{q^*,p^*}(\mathbf{Z}^d)$ for all $1 \leq p, q \leq \infty$. As for multipliers on \mathbf{R}^n , we have $M^2(\mathbf{Z}^d) = l^\infty(\mathbf{Z}^d)$, and $M^{1,q}(\mathbf{Z}^d) = M^{q^*,\infty}(\mathbf{Z}^d) = \widehat{L^q(\mathbf{T}^d)}$ for $q > 1$, and $M^{1,1}(\mathbf{Z}^d) = M^{\infty,\infty}(\mathbf{Z}^d) = \widehat{M(\mathbf{T}^d)}$.

Our goal is to relate the boundedness of multipliers on \mathbf{T}^d to the boundedness of associated multipliers on \mathbf{R}^d , and vice versa. To begin with, we show that bounded multipliers on \mathbf{R}^d induce bounded multipliers on \mathbf{T}^d .

Theorem 23.11. *Suppose $m \in M^p(\mathbf{R}^d)$, and every point in \mathbf{R}^d is a Lebesgue point of m . Then for all $R > 0$, the sequence $a_R(n) = m(n/R)$ lies in $M^p(\mathbf{Z}^d)$, and $\|a_R\|_{M^p(\mathbf{Z}^d)} \leq \|m\|_{M^p(\mathbf{R}^d)}$.*

Remark. Every function can be altered on a set of measure zero in order to satisfy the assumptions of this theorem, as the theory of the Hardy-Littlewood maximal function indicates.

Proof. Without loss of generality, by rescaling we may assume that $R = 1$. The case $p = 1$ and $p = \infty$ follows from Poisson summation and the characterizations of $M^1(\mathbf{R}^d)$ and $M^1(\mathbf{Z}^d)$. Thus we may assume $1 < p < \infty$.

Here the proof follows from a general identity, that for any pair of trigonometric periodic polynomials f and g , if $\Phi_\delta(x) = e^{-\pi\delta|x|^2}$, if T is a Fourier multiplier on \mathbf{R}^d with symbol $m(\xi)$, and S is a Fourier multiplier on \mathbf{Z}^d with symbol $m(n)$, then for any $\alpha, \beta > 0$ with $\alpha + \beta = 1$,

$$\lim_{\delta \rightarrow 0} \delta^{d/2} \int_{\mathbf{R}^d} T\{f\Phi_{\alpha\delta}\} \cdot \overline{g\Phi_{\beta\delta}} = \int_{\mathbf{T}^d} Sf \cdot \bar{g} \, dx.$$

To obtain the identity, it suffices to assume by linearity that $f(x) = e^{2\pi i n_1 \cdot x}$ and $g(x) = e^{2\pi i n_2 \cdot x}$ for two integers n_1 and n_2 . If $n_1 = n_2 = n$, we therefore have to prove that the left hand side is equal to $m(n)$, and if $n_1 \neq n_2$, we have to prove the left hand side is equal to zero. By the multiplication formula, the left hand side is

$$\lim_{\delta \rightarrow 0} \delta^{d/2} (\delta\alpha)^{-d/2} (\delta\beta)^{-d/2} \int_{\mathbf{R}^d} m(\xi) \cdot \Phi_{1/\alpha\delta}(\xi - n_1) \Phi_{1/\beta\delta}(\xi - n_2) = \lim_{\delta \rightarrow 0} (\alpha\beta\delta)^{-d/2} \int_{\mathbf{R}^d} m(\xi) \Phi_{1/\alpha\delta}(\xi - n_1) \Phi_{1/\beta\delta}(\xi - n_2) \, d\xi$$

If $n_1 = n_2 = n$, then

$$\Phi_{1/\alpha\delta}(\xi - n_1) \Phi_{1/\beta\delta}(\xi - n_2) = (\alpha\beta\delta)^{-n/2} \Phi_{1/\alpha\beta\delta}(\xi - n),$$

is an approximation to the identity as $\delta \rightarrow 0$, yielding the result. On the other hand, if $n_1 \neq n_2$, then the two Gaussian functions are primarily supported on disjoint sets, which yields the result in this case as well.

Returning to the general proof, we note that it suffices to prove by density and duality that for any trigonometric polynomials f and g ,

$$\left| \int Sf(x) \overline{g(x)} \, dx \right| \leq \|m\|_{M^p(\mathbf{R}^d)} \|f\|_{L^p(\mathbf{R}^d)} \|g\|_{L^{p^*}(\mathbf{R}^d)}.$$

But the identity above gives that

$$\begin{aligned} \left| \int Sf(x) \overline{g(x)} \, dx \right| &\leq \lim_{\delta \rightarrow 0} \delta^{d/2} \left| \int \int_{\mathbf{R}^d} T\{f\Phi_{\alpha\delta}\} \cdot \overline{g\Phi_{\beta\delta}} \right| \\ &\leq \lim_{\delta \rightarrow 0} \delta^{d/2} \|m\|_{M^p(\mathbf{R}^d)} \|f\Phi_{\alpha\delta}\|_{L^p(\mathbf{R}^d)} \|g\Phi_{\beta\delta}\|_{L^{p^*}(\mathbf{R}^d)} \\ &= \|m\|_{M^p(\mathbf{R}^d)} \|f\|_{L^p(\mathbf{T}^d)} \|g\|_{L^{p^*}(\mathbf{T}^d)}. \end{aligned} \quad \square$$

It is clearly not true that if $\{m(n) : n \in \mathbf{Z}^d\}$ gives an element of $M^p(\mathbf{Z}^d)$, then m is in $M^p(\mathbf{T}^d)$, since m can be poorly behaved away from the integers. But if $\{m(n/R) : n \in \mathbf{Z}^d\}$ is uniformly in $M^p(\mathbf{Z}^d)$, we can prove that m is in $M^p(\mathbf{T}^d)$ given weak continuity assumptions.

Theorem 23.12. *Suppose m is Riemann integrable, $1 \leq p \leq \infty$, and we define $a_R(n) = m(n/R)$. Then*

$$\|m\|_{M^p(\mathbf{R}^d)} \leq \sup_{R>0} \|a_R\|_{M^p(\mathbf{Z}^d)}.$$

Proof. Fix $f, g \in C_c^\infty(\mathbf{R}^d)$. Then for suitably large R , $\text{Dil}_{1/R}f$ and $\text{Dil}_{1/R}g$ are supported in $[-1/2, 1/2]^d$. Define the periodic functions

$$\tilde{f}_R(x) = \sum_{n \in \mathbf{Z}^d} \text{Dil}_{1/R}f(x - n) \quad \text{and} \quad \tilde{g}_R(x) = \sum_{n \in \mathbf{Z}^d} \text{Dil}_{1/R}g(x - n).$$

Then

$$\begin{aligned} \int m(D)f(x)\overline{g}(x) &= \int m(\xi)\widehat{f}(\xi)\overline{\widehat{g}(\xi)} \\ &= \lim_{R \rightarrow \infty} R^{-d} \sum_{n \in \mathbf{Z}^d} b(n/R)\widehat{f}(n/R)\overline{\widehat{g}(n/R)} \\ &= \lim_{R \rightarrow \infty} R^d \sum_{n \in \mathbf{Z}^d} b(n/R)\widehat{f_R}(n)\overline{\widehat{\tilde{g}_R}(n)} \\ &\leq \lim_{R \rightarrow \infty} R^d \|a_R\|_{M^p(\mathbf{Z}^d)} \|\tilde{f}_R\|_{L^p(\mathbf{T}^d)} \|\tilde{g}_R\|_{L^{p^*}(\mathbf{T}^d)} \\ &\leq \lim_{R \rightarrow \infty} \|a_R\|_{M^p(\mathbf{Z}^d)} \|f\|_{L^p(\mathbf{R}^d)} \|g\|_{L^{p^*}(\mathbf{R}^d)}. \end{aligned}$$

□

Example. *The Hilbert transform H is a Fourier multiplier with symbol $i\text{sgn}(\xi)$. Since it is homogeneous, Riemann integrable, and every point is a Lebesgue point, the theory of boundedness from $L^p(\mathbf{R}^d)$ to $L^p(\mathbf{R}^d)$ is equivalent to the boundedness of the periodic version of the Hilbert transform from $L^p(\mathbf{T}^d)$ to $L^p(\mathbf{T}^d)$.*

Similarly, we see that the convergence in $L^p(\mathbf{T}^d)$ of various Fourier series (Dirichlet sums, spherical summation, square summation, and so

on) are equivalent to the associated problems of the convergence of the Fourier transform in $L^p(\mathbf{R}^d)$.

Many problems in harmonic analysis deal not only with the boundedness of Fourier multipliers, but also the boundedness of a maximal function associated with a family of multipliers (e.g. the almost everywhere convergence of Fourier series is connected with the properties of the maximal function $Mf = \sup_{R>0} S_R f$, where S_R is the partial summation operator. Let us prove a transference result for maximal functions associated with dilations of multiplier operators.

Theorem 23.13. *Fix a bounded, Riemann integrable function m on \mathbf{R}^d such that every $\xi \in \mathbf{R}^d$ is a Lebesgue point of m . Define the maximal functions $Mf = \sup_{R>0} m(D/R)f$ on \mathbf{R}^d , and the maximal function $Nf = \sup_{R>0} T_R f$, where T_R is the Fourier multiplier operator on \mathbf{T}^d associated with the sequence $a_R(n) = m(n/R)$. Then M is bounded from the $L^p(\mathbf{R}^d)$ norm to itself if and only if N is bounded from the $L^p(\mathbf{R}^d)$ norm to itself.*

Proof. It suffices to show that a uniform family of estimates of the form

$$\left\| \sum_{i=1}^K c_i m(D/R_i) f \right\|_{L^p(\mathbf{R}^d)} \lesssim \sum |c_i|$$

uniform in K and R_1, \dots, R_K , is equivalent to a uniform family of estimates of the form

$$\left\| \sum_{i=1}^K c_i T_{R_i} f \right\|_{L^p(\mathbf{R}^d)} \lesssim \sum |c_i|,$$

which is uniform in K and R_1, \dots, R_K . In the first case, we may assume without loss of generality that $f \in C_c^\infty(\mathbf{R}^d)$, and in the second, that f is a trigonometric polynomial. The proof then follows by essentially the analogous techniques to above. \square

Remark. One can also show that transference holds between maximal functions bounded merely from $L^p(\mathbf{R}^d)$ to $L^{p,\infty}(\mathbf{R}^d)$, using the duality of Lorentz spaces in much the same way.

Here are some other examples of transference.

Example. *TODO: ELABORATE ON THIS.* For a given family of coefficients $\{a_m : m \in \mathbf{Z}^d\}$, consider the periodization operator

$$Tf = \sum_{m \in \mathbf{Z}^d} a_m \text{Trans}_m f.$$

Then, roughly speaking, T is a Fourier multiplier operator with symbol $m(\xi) = \sum a_m e^{-2\pi i \xi \cdot m}$, where this sum must be interpreted in a non-standard sense if the sum $\{a_m\}$ does not converge, which means we might end up obtaining a tempered distribution m rather than a function. But certainly T is well defined for any compactly supported input, since then the sum is locally finite. Let us assume T is bounded on these inputs in the L^p norm for some $1 < p < \infty$ with operator norm A , and thus extends uniquely to a bounded operator from $L^p(\mathbf{R}^n)$ to itself. If we consider any family of coefficients $\{c_m : m \in \mathbf{Z}^d\}$, and consider, for $\varepsilon < 1$, the function

$$f(x) = \sum_{m \in \mathbf{Z}^d} c_m \varepsilon^{-d/p} \mathbf{I} \left(x \in \prod_{i=1}^d [m_i, m_i + \varepsilon] \right),$$

then $\|f\|_{L^p(\mathbf{R}^d)} = \|c\|_{l^p(\mathbf{Z}^d)}$. If $\|c_m\|_{l^p(\mathbf{Z}^d)}$, it follows by continuity that it must be true that

$$\begin{aligned} Tf(x) &= \sum_{k \in \mathbf{Z}^d} \sum_{m+n=k} c_m a_n \varepsilon^{-d/p} \mathbf{I} \left(x \in \prod_{i=1}^d [k_i, k_i + \varepsilon] \right) \\ &= \sum_{k \in \mathbf{Z}^d} (a * c)(k) \varepsilon^{-d/p} \mathbf{I} \left(x \in \prod_{i=1}^d [k_i, k_i + \varepsilon] \right). \end{aligned}$$

It follows that

$$\|Tf\|_{L^p(\mathbf{R}^d)} = \|a * c\|_{l^p(\mathbf{Z}^d)}.$$

Thus we conclude that $\|a * c\|_{l^p(\mathbf{Z}^d)} \leq A \|c\|_{l^p(\mathbf{Z}^d)}$. Conversely, suppose convolution with a is bounded from $l^p(\mathbf{Z}^d)$ to $l^p(\mathbf{Z}^d)$ with norm A . Then

$$\left(\sum_{k \in \mathbf{Z}^d} |Tf(x+k)|^p \right)^{1/p} = \left(\sum_{k \in \mathbf{Z}^d} \left| \sum_m a_m f(x+k-m) \right|^p \right)^{1/p} \leq A \left(\sum_k |f(x+k)|^p \right)^{1/p}.$$

But

$$\begin{aligned}\|Tf\|_{L^p(\mathbf{R}^d)} &= \left(\int_{[0,1]^d} \left(\sum_{k \in \mathbf{Z}^d} |Tf(x+k)|^p \right) \right)^{1/p} \\ &\leq A \left(\int_{[0,1]^d} \sum_k |f(x+k)|^p \right)^{1/p} \\ &= A \|f\|_{L^p(\mathbf{R}^d)}.\end{aligned}$$

Thus we conclude that T is bounded if and only if the associated discrete convolution operator is bounded. Thus we have shown studying the boundedness of multiplier operators associated to symbols with domain \mathbf{T}^d , e.g. convolution operators from $l^p(\mathbf{Z}^d)$ to $l^q(\mathbf{Z}^d)$, via Fourier multiplier operators with periodic symbols in the domain \mathbf{R}^d . More precisely, we can define $M^{p,q}(\mathbf{Z}^d)$ to be the set of all distributions f on \mathbf{T}^d such that convolution with \hat{f} gives a bound from $L^p(\mathbf{Z}^d)$ to $L^q(\mathbf{Z}^d)$ in a suitable domain so we can interpret this convolution, e.g. for compactly supported sequences. It then follows that $M^p(\mathbf{Z}^d)$ is isomorphic to the family of periodic symbols in $M^p(\mathbf{R}^d)$.

Example. Next, consider $m \in M^p(\mathbf{R}^d)$ with compact support in $[0,1]^d$, and consider the periodization

$$m_P(\xi) = \sum_{n \in \mathbf{Z}^d} m(\xi - n).$$

We claim that $m_P \in M^p(\mathbf{R}^d)$. TODO (Look up Jodeit Paper?).

Chapter 24

Sobolev Spaces

Chapter 25

Time Frequency Analysis

In harmonic analysis, it is often useful to study a function $f : \mathbf{R}^d \rightarrow \mathbf{C}$ via its Fourier transform $\hat{f} : \mathbf{R}^d \rightarrow \mathbf{C}$. The goal of time-frequency analysis is to think of such a function as being a function living simultaneously in both spaces, i.e. a function on the domain $\mathbf{R}^d \times \mathbf{R}^d$, which can be either written in temporal coordinates, or frequential coordinates, and in certain special cases, a combination of the two. We call this space the *phase plane*, combining both time and frequency information together. There are several difficulties with rigorously incorporating this approach, for instance, resulting from the uncertainty principle, but the utility makes this. Given a function f , we define a *phase portrait* for f to be a subset of $\mathbf{R}^d \times \mathbf{R}^d$ where the majority of the ‘mass’ of f and \hat{f} are concentrated (for an arbitrary locally compact abelian group G , phase space is $G^* \times G$). Let us consider a simple example.

Example. Consider the Gaussian $f(t) = e^{-\pi t^2}$. Then 70% of the mass of f is concentrated on $[-1, 1]$, and the mass decays exponentially away from this interval. Thus the function f is concentrated in $[-1, 1]$. We have $\hat{f}_\delta(\omega) = e^{-\pi \omega^2}$, which similarly, is concentrated in $[-1, 1]$. A natural choice of the phase portrait of f is therefore $[-1, 1] \times [-1, 1]$.

Example. The Fourier transform of the Dirac delta function δ at a point x is the plane wave $\xi \mapsto e^{-2\pi i \xi \cdot x}$. Thus a natural phase portrait for the Dirac delta function is $\{x\} \times \mathbf{R}^d$. Similarly, the phase portrait of a pure plane wave $x \mapsto e^{2\pi i \xi \cdot x}$ is $\mathbf{R}^d \times \{\xi\}$, since the Fourier transform is the Dirac delta function at ξ .

The symmetries of the Fourier transform have natural effects on the phase portrait of a function f , which has phase portrait S .

- The phase portrait of $\text{Trans}_x f$ is obtained by translating S horizontally by x units, since on the Fourier side the translation acts as a modulation, and does not move mass. Similarly, the phase portrait of the modulation $\text{Mod}_\xi f$ is obtained by translating S vertically by ξ units, since modulation does not affect the position of mass on the spatial side of things.
- Scaling in physical space has a ‘dual’ effect in phase space. More precisely, $\text{Dil}_\delta f$ has phase portrait

$$\{(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d : (x/\lambda, \lambda\xi) \in S\}$$

More generally, given a linear transformation T , the phase portrait of $f \circ T^{-1}$ is equal to

$$\{(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d : (T(x), T^{-t}(\xi)) \in S\}.$$

The rescaling above is a special case. In particular, if $T \in O(d)$, then $f \circ T^{-1}$ has phase portrait

$$\{(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d : (T(x), T(\xi)) \in S\}.$$

- The phase portrait of \hat{f} is equal to

$$\{(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d : (\xi, -x) \in S\},$$

i.e. a clockwise rotation by ninety degrees.

Notice that all the transformations above preserve area, which is where symplectic geometry enters the picture.

We note that the phase portrait of a rescaled, translated, and modulated Gaussian has phase portrait consisting of an axis-oriented rectangle with sidelengths δx and $\delta \xi$, where $\delta x \cdot \delta \xi \sim 1$. Such a rectangle in phase space is called a *Heisenberg tile*, and functions whose phase portraits consist of Heisenberg tiles are called *wave packets*. In light of the uncertainty principle, these functions are the best alternatives to a function which is compactly supported in a sidelength one interval, and whose Fourier transform is also supported in a sidelength one interval.

25.1 Localization in Time and Space

Physical space localization is easy, we just multiply by a function, either a rough cutoff, or a smooth cutoff. Frequency localization is only slightly harder, where we can apply a basic Fourier multiplier. To localize in both time and frequency, one approach is to first smoothly localize in space, then time, or vice versa. This works out fine, as long as we do not localize too finely in both time and space, i.e. breaking the uncertainty principle. Here is a characteristic result in this setting

Lemma 25.1. *Fix two cubes I and J , and two smooth functions ψ_I and ψ_J adapted to I and J , and consider the localization operator $\pi_{I \times J} = \psi_I(D) \circ \psi_J(X)$. Then $\pi_{I \times J} f$ has Fourier support in I , and is localized in J in the sense that for $|x| \geq$*

$$(\pi_{I \times J} f)(x) \lesssim_n |I|^{1-n} |J|^{1/2} \|f\|_{L^2(\mathbf{R}^d)} d(x, J)^{-n}$$

Proof. If we let K_I be the inverse Fourier transform of ψ_I , then for all $n > 0$,

$$|K_I(x)| \lesssim_n |I|^{1-n} |x|^{-n}.$$

The proof then follows from Cauchy-Schwartz applied to the representation $\pi_{I \times J} f = K_I * (\psi_J \cdot f)$.

$$\left(\int |K_I(x-y)|^2 |\psi_J(y)|^2 dy \right)^{1/2} \|f\|_{L^2(\mathbf{R}^d)}$$

□

25.2 Convergence in L^p and the Hilbert Transform

We now move onto a more 20th century viewpoint on Fourier series, namely, those to do with operator theory. Under this viewpoint, the properties of convergence are captured under the boundedness of certain operators on function spaces, allowing us to use the modern theory of functional analysis to it's full extent on our problems. However, unlike in most of basic functional analysis, where we assume all operators we encounter are

bounded to begin with, in harmonic analysis we more often than not are given an operator defined only on a subset of spaces, and must prove the continuity of such an operator to show it is well defined on all of space. We will illustrate this concept through the theory of the circular Hilbert transform, and its relation to the norm convergence of Fourier series.

A *Fourier multiplier* is a linear transform T associated with a given sequence of scalars λ_n , for $n \in \mathbf{Z}$. It is defined for any trigonometric polynomial $f = \sum_{|n| \leq N} c_n e_n$ as $Tf = \sum_{|n| \leq N} \lambda_n c_n e_n$. The trigonometric polynomials are dense in $L^p(\mathbf{T})$, for each $p < \infty$. An important problem is determining whether T is therefore figuring out whether the operator can be extended to a *continuous operator* on the entirety of L^p . Because the trigonometric polynomials are dense in L^p , in the light of the Hahn Banach theorem it suffices to prove an inequality of the form $\|Tf\| \lesssim \|f\|$. Here are some examples of Fourier operators we have already seen.

Example. The truncation operator S_N is the transform associated with the scalars $\lambda_n = [|n| \leq N]$. The truncation is continuous, since for any integrable function f , the Fourier coefficients are uniformly bounded by $\|f\|_1$, so $\|S_N f\|_1 \leq N \|f\|_1$. Similarly, the Féjer truncation σ_N associated to the multipliers $\lambda_N = [|n| \leq N](1 - |n|/N)$ is continuous on all integrable functions. These operators are easy to extend precisely because the nonzero multipliers have finite support.

Example. In the case of the Abel sum, A_r , associated with $\lambda_n = r^{|n|}$, A_r extends in a continuous way to all integrable functions, since

$$|A_r f| = \left| \sum r^{|n|} \hat{f}(n) e_n(t) \right| \leq \|f\|_1 \sum r^{|n|} = \|f\|_1 \left(1 + \frac{2}{1-r} \right)$$

Thus the map is bounded.

To understand whether the truncations $S_N f$ of f converge to f in the L^p norms, rather than pointwise, we turn to the analysis of an operator which is the core of the divergence issue, known as the *Hilbert transform*. It is a Fourier multiplier operator H associated with the coefficients

$$\lambda_n = \frac{\text{sgn}(n)}{i} = \begin{cases} +1/i & n > 0 \\ 0 & n = 0 \\ -1/i & n < 0 \end{cases}$$

Because

$$[|n| \leq N] = \frac{\operatorname{sgn}(n+N) - \operatorname{sgn}(n-N)}{2} + \frac{[n=N] + [n=-N]}{2}$$

we conclude

$$S_n f = \frac{i(e_{-n}H(e_n f) - e_n H(e_{-n} f))}{2} + \frac{\widehat{f}(n)e_n + \widehat{f}(-n)e_{-n}}{2}$$

Since the operators $f \mapsto \widehat{f}(n)e_n$ are bounded in all the L^p spaces since they are continuous in $L^1(\mathbf{T})$, we conclude that the operators S_n are uniformly bounded as endomorphisms on $L^p(\mathbf{T})$ provided that H is bounded as an operator from $L^p(\mathbf{T})$ to $L^q(\mathbf{T})$. Since $S_n f$ converges to f in L^p whenever f is a trigonometric polynomial, this would establish that $S_n f$ converges to f in the L^p norm for any function f in $L^p(\mathbf{T})$. Later on, as a special case of the Hilbert transform on the real line, we will be able to prove that H is a bounded operator on $L^p(\mathbf{T})$ for all $1 < p < \infty$, and as a result, we find that $S_N f \rightarrow f$ in L^p for all such p . Unfortunately, H is not bounded from $L^1(\mathbf{T})$ to itself, and correspondingly, $S_N f$ does not necessarily converge to f in the L^1 norm for all integrable f .

For now, we explore some more ideas in how we can analyze the Hilbert transform via convolution, the dual of Fourier multipliers. The fact that $\widehat{f * g} = \widehat{f} \widehat{g}$ implies that if there is an integrable function g whose Fourier coefficients corresponds to the multipliers of an operator T , then $f * g = Tf$ for any trigonometric polynomial f , and by the continuity of convolution, this is the unique extension of the Fourier multiplier operator. In the theory of distributions, one generalizes the family of objects one can take the Fourier series from integrable functions to a more general family of objects, such that every sequence of Fourier coefficients is the Fourier series of some *distribution*. One can take the convolution of any such distribution Λ with a C^∞ function f , and so one finds that $\Lambda * f = Tf$ for any trigonometric polynomial f . There is a theorem saying that *all* continuous translation invariant operators from $L^p(\mathbf{T})$ to $L^q(\mathbf{T})$ are given by convolution with a Fourier multiplier operator. In practice, we just compute the convolution kernel which defines the Fourier multiplier, but it is certainly a satisfying reason to justify the study of Fourier multipliers. For instance, a natural question is to ask which Fourier multipliers result in bounded operations in space.

Theorem 25.2. *A Fourier multiplier is bounded from $L^2(\mathbf{T})$ to itself if and only if the coefficients are bounded.*

Proof. If a Fourier multiplier is given by λ_n , then for some trigonometric polynomial f ,

$$\|Tf\|_2^2 = \sum |\widehat{Tf}(n)|^2 = \sum |\lambda_n|^2 |\widehat{f}(n)|^2$$

If the λ_n are bounded, then we can obtain from this formula the bound

$$\|Tf\|_2^2 \leq \max |\lambda_n| \|f\|_2^2$$

Conversely, if Tf is bounded, then

$$|\lambda_n|^2 = \|T(e_n)\|_2^2 \leq \|T\|^2$$

so the λ_n are bounded. □

Corollary 25.3. *The Hilbert transform is a bounded endomorphism on $L^2(\mathbf{T})$. Note that we already know that $S_N f \rightarrow f$ in the L^2 norm.*

The terms of the Hilbert transform cannot be considered the Fourier coefficients of any integrable function. Indeed, they don't vanish as $n \rightarrow \infty$. Nonetheless, we can use Abel summation to treat the Hilbert transform as convolution with an appropriate operator. For $0 < r < 1$, consider, for $z = e^{it}$,

$$K_r(z) = \sum_{n \in \mathbf{Z}} \frac{\operatorname{sgn}(n)}{i} r^{|n|} z^n = K * P_r$$

Since we know the Hilbert transform is continuous in $L^2(\mathbf{T})$, we can conclude that, in particular, for any C^∞ function f ,

$$Hf = \lim_{r \rightarrow 1} K * (P_r * f) = \lim_{r \rightarrow 1} (K * P_r) * f = \lim_{r \rightarrow 1} K_r * f$$

So it suffices to determine the limit of the K_r . We find that

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(rz)^n - (r\bar{z})^n}{i} &= \frac{r}{i} \left(\frac{1}{\bar{z} - r} - \frac{1}{z - r} \right) = \frac{r}{i} \frac{z - \bar{z}}{|z|^2 - 2r\operatorname{Re}(z) + r^2} \\ &= \frac{2r \sin(t)}{1 - 2r \cos(t) + r^2} = \frac{4r \sin(t/2) \cos(t/2)}{(1-r)^2 + 4r \sin^2(t/2)} \\ &= \cot(t/2) + O\left(\frac{(1-r)^2}{t^3}\right) \end{aligned}$$

Thus $K_r(t)$ tends to $\cot(t/2)$ locally uniformly away from the origin. But

$$K_r(t) = \frac{4r \sin(t/2) \cos(t/2)}{(1-r)^2 + 4r \sin^2(t/2)} = O\left(\frac{t}{(1-r)^2}\right)$$

If f is any C^∞ function on \mathbf{T} , then

$$\left| \int_{|t| \geq \varepsilon} [K_r(t) - \cot(t/2)] f(t) dt \right| \lesssim (1-r)^2 \|f\|_\infty \int_{|t| \geq \varepsilon} \frac{dt}{|t|^3} \lesssim \frac{(1-r)^2 \|f\|_\infty}{\varepsilon^2}$$

$$\begin{aligned} \left| \int_{|t| < \varepsilon} K_r(t) f(t) dt \right| &\leq \int_0^\varepsilon |K_r(t)| |f(t) - f(-t)| dt \\ &\lesssim \int_0^\varepsilon |t K_r(t)| |f'(0)| dt \lesssim \frac{|f'(0)|}{(1-r)^2} \int_0^\varepsilon t^2 dt \lesssim \|f'\|_\infty \frac{\varepsilon^3}{(1-r)^2} \end{aligned}$$

$$\left| \int_{|t| < \varepsilon} \cot(t/2) f(t) dt \right| \lesssim \int_0^\varepsilon \frac{|f(t) - f(-t)|}{t} dt \lesssim \varepsilon f'(0)$$

Thus

$$\left| \int K_r(t) f(t) dt - \int \cot(t/2) f(t) dt \right| \lesssim \frac{(1-r)^2}{\varepsilon^2} \|f\|_\infty + \left(\frac{\varepsilon^3}{(1-r)^2} + \varepsilon \right) \|f'\|_\infty$$

Choosing $\varepsilon = (1-r)^\alpha$ for some $2/3 < \alpha < 1$ shows that for sufficiently smooth f ,

$$(Hf)(x) = \lim_{r \rightarrow 1} \int \cot(t/2) f(x-t) dt$$

25.3 A Divergent Fourier Series

Analysis was built to analyze continuous functions, so we would hope the method of fourier expansion would work for all continuous functions. Unfortunately, this is not so. The behaviour of the Dirichlet kernel away from the origin already tells us that the convergence of Fourier series is subtle. We shall take advantage of this to construct a continuous function with divergent fourier series at a point.

To start with, we shall consider the series

$$f(t) \sim \sum_{n \neq 0} \frac{e_n(t)}{n}$$

where f is an odd function equaling $i(\pi - t)$ for $t \in (0, \pi]$. Such a function is nice to use, because its Fourier representation is simple, yet very close to diverging. Indeed, if we break the series into the pair

$$\sum_{n=1}^{\infty} \frac{e_n(t)}{n} \quad \sum_{n=-\infty}^{-1} \frac{e_n(t)}{n}$$

Then these series no longer are the Fourier representations of a Riemann integrable function. For instance, if $g(t) \sim \sum_{n=1}^{\infty} \frac{e_n(t)}{n}$, then the Abel means $A_r(f)(t) =$

25.4 Conjugate Fourier Series

When f is a real-valued integrable function, then $\overline{\hat{f}(-n)} = \hat{f}(n)$. Thus we formally calculate that

$$\sum_{n=-\infty}^{\infty} \hat{f}(n) e_n(t) = \operatorname{Re} \left(\hat{f}(0) + 2 \sum_{n=1}^{\infty} \hat{f}(n) e_n(t) \right)$$

This series defines an analytic function in the interior of the unit circle since the coefficients are bounded. Thus the sum is a harmonic function in the interior of the unit circle. The imaginary part of this sum is

$$\operatorname{Im} \left(\hat{f}(0) + 2 \sum_{n=1}^{\infty} \hat{f}(n) e_n(t) \right) = \Re \left(-i \sum_{n=-\infty}^{\infty} \operatorname{sgn}(n) \hat{f}(n) e_n(t) \right)$$

The right hand side is known as the conjugate series to the Fourier series $\hat{f}(n)$. It is closely related to the study of a function \tilde{f} known as the *conjugate function*.

Chapter 26

Higher Dimensional Fourier Analysis

The Fourier transform in dimensions greater than one have strange phenomena, which is not so obvious from the perspective given by the one dimensional theory. For instance, consider the Theorem of Marcel Riesz, which says that if we define the Dirichlet summation operator

$$S_R f(x) = \int_{-R}^R \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi,$$

then for $1 < p < 2$, if $f \in L^p(\mathbf{R}^d)$, $S_R f \rightarrow f$ in $L^p(\mathbf{R}^d)$. The result follows because the operators $\{S_R\}$ are uniformly bounded, and dilation symmetries show this is equivalent to showing that S_1 is a bounded operator on $L^p(\mathbf{R}^d)$. Tensorization shows that Riesz's theorem continues to work in higher dimensions for the *square summation operators*

$$S_R^{\text{sq}} f(x) = \int_{-R}^R \cdots \int_{-R}^R \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi,$$

i.e. the operator S_1^{sq} is bounded on $L^p(\mathbf{R}^d)$ for $1 < p < \infty$. On the other hand, the *spherical summation operators*

$$S_R f(x) = \int_{|\xi| \leq R} \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi$$

act very differently than their one dimensional counterpart. In particular, Riesz's theorem completely fails in this situation. The *ball multiplier* $S = S_1$

is *only* bounded on $L^p(\mathbf{R}^d)$ for $p = 2$ (in which case the bound follows trivially from Plancherel). The unboundedness in $L^p(\mathbf{R}^d)$ for $1 < p < \infty$ follows from a clever result of Fefferman (1971).

Theorem 26.1. *The ball multiplier S is only bounded on $L^2(\mathbf{R}^d)$.*

Proof. We rely on a geometric construction, together with the uncertainty principle. Duality implies that it suffices to show it is unbounded on $L^p(\mathbf{R}^d)$ for $p > 2$. Thus we must find functions f which are 'broad', or 'spread out', and show that Sf can have parts which are large on a 'narrow', or 'concentrated' set. Our first observation is that the bad part of the multiplier occurs on the boundary of the unit ball B ; the multiplier is C^∞ away from this boundary, so behaves very nicely with functions with Fourier support away from the boundary of the unit ball. Thus, without loss of generality, we might as well try and find functions whose Fourier support is concentrated near ∂B .

For any $R > 0$, we construct a function f_R supported on the annulus given by the equation $1 - 1/10R \leq |\xi| \leq 1 + 1/R$. To apply the uncertainty principle, we find $\sim R^{d-1}$ rectangles $\{\Theta\}$, which have dimensions R^{-1} in the direction tangent to ∂B , but have dimension $R^{-1/2}$ in the $d - 1$ directions tangent to the boundary of the sphere, and such that Θ has one tenth of it's mass in B , and half it's mass outside of B . We fix a compactly supported density function $\phi \in C_c^\infty(\mathbf{R}^d)$ on the unit ball, and rescale it to density functions $\{\phi_\Theta\}$ supported on Θ with magnitude roughly $R^{(d+1)/2}$ on Θ . For any x_1, \dots, x_N , the uncertainty principle tells us that the function χ_{T_Θ} , with Fourier transform equal to

$$e^{-2\pi i x_\Theta \cdot \xi} \phi_\Theta(\xi),$$

has mass concentrated on the dual rectangle T_Θ centered at x_Θ and with magnitude roughly $R^{(d+1)/2}$ on T_Θ . The action of S on χ_{T_Θ} is to cut the Fourier support of Θ by a tenth. Thus we lose the 90% of the Fourier mass of Θ , but our support is also made ten times thinner in the tangential direction. As a result, $S\chi_{T_\Theta}$ will now have mass concentrated on a tube T_Θ^* , which is ten times longer than T_Θ .

Let T_Θ^+ be the portion of T_Θ^* which lies at the opposite end of T_Θ^* to T_Θ . Our construction now relies on *Kakeya* like phenomena. For any $\varepsilon > 0$, there exists a large R_0 such that for $R \geq R_0$, we can pick $\{x_\Theta\}$ such that

tubes $\{T_\Theta\}$ are disjoint from one another, but such that the tubes $\{T_\Theta^+\}$ have large overlap, in the sense that

$$|\bigcup_{\Theta} T_\Theta^+| \leq \varepsilon |\bigcup_{\Theta} T_\Theta|.$$

The supports of $S\chi_\Theta$ thus have lots of overlap on this set, but it is difficult to control the sum $\sum S\chi_\Theta$ since these functions might have different signs where they overlap. To fix this, we define

$$f_R = \sum_{\Theta} \varepsilon_\Theta \chi_\Theta$$

where $\{\varepsilon_\Theta\}$ are independent $\{-1, +1\}$ valued Bernoulli random variables, because Khintchine's inequality implies that we have the pointwise inequality, for any $1 < p < \infty$, of the form

$$\left(\mathbf{E} \left| \sum_{\Theta} \varepsilon_\Theta \chi_\Theta \right|^p \right)^{1/p} \sim \left(\sum_{\Theta} |\chi_\Theta|^2 \right)^{1/2}.$$

It follows that, since the tubes T_Θ are disjoint,

$$\begin{aligned} \mathbf{E} \|f_R\|_{L^p(\mathbf{R}^d)}^p &\sim \int \left(\sum_{\Theta} |\chi_\Theta(x)|^2 \right)^{p/2} dx \\ &\sim \left(R^{\frac{d+1}{2}} \right)^p \left| \bigcup_{\Theta} T_\Theta \right|. \end{aligned}$$

Similarly, Khintchine's inequality can again be applied to

$$Sf_R = \sum_{\Theta} \varepsilon_\Theta S\chi_\Theta.$$

to conclude that

$$\begin{aligned} \mathbf{E} \|Sf_R\|_{L^p(\mathbf{R}^d)}^p &\sim \int \left(\sum_{\Theta} |S\chi_\Theta(x)|^2 \right)^{p/2} dx \\ &\gtrsim \left(R^{\frac{d+1}{2}} \right)^p \int \left(\sum_{\Theta} \mathbf{I}_{T_\Theta^+}(x) \right)^{p/2} dx. \end{aligned}$$

If S was bounded on $L^p(\mathbf{R}^d)$, we would conclude that

$$\int \left(\sum_{\Theta} \mathbf{I}_{T_{\Theta}^+}(x) \right)^{p/2} dx \lesssim \left| \bigcup_{\Theta} T_{\Theta} \right|.$$

The trivial bound

$$\int \left(\sum_{\Theta} \mathbf{I}_{T_{\Theta}^+}(x) \right) dx \sim \left| \bigcup_{\Theta} T_{\Theta} \right|$$

combined with the fact that the sum is supported on the finite measure set $\bigcup_{\Theta} T_{\Theta}^+$ yields that

$$\left| \bigcup_{\Theta} T_{\Theta}^+ \right|^{-(p/2-1)} \left| \bigcup_{\Theta} T_{\Theta} \right|^{p/2} \lesssim \int \left(\sum_{\Theta} \mathbf{I}_{T_{\Theta}^+}(x) \right)^{p/2} dx.$$

Thus we find that

$$\left| \bigcup_{\Theta} T_{\Theta} \right|^{p/2-1} \lesssim \left| \bigcup_{\Theta} T_{\Theta}^+ \right|^{p/2-1}.$$

Because of the Kakeya like specification of these sets, this is impossible in the range $p > 2$ specified. \square

We thus see that understanding the behaviour of S is closely connected to Kakeya phenomena. Indeed, one can generalize this problem to determining the behaviour of the *Bochner-Riesz multipliers* S_{δ} , which are the Fourier multipliers with symbol

$$m^{\delta}(\xi) = (1 - |\xi|)_{+}^{\delta}.$$

For $\delta = 0$, S_{δ} is just the ball multiplier above. As δ increases, the singularity on the boundary of the unit ball becomes less and less bad, so it might be expected that S_{δ} becomes bounded on a larger range of L^p spaces. Determining the precise range under which these operators are bounded is a major open problem of harmonic analysis, and is closely connected to a deeper understanding of Kakeya like phenomena.

Chapter 27

Oscillatory Integrals

The goal of the theory of oscillatory integrals is to obtain estimates of integrals with highly oscillatory integrands, where standard techniques such as taking in absolute values, or various spatial decomposition strategies, fail completely to give tight estimates. A typical oscillatory integral is of the form

$$I(\lambda) = \int a(x) e^{2\pi i \lambda \phi(x)} dx,$$

where a is the *amplitude function*, and ϕ is the *phase*. The value λ is a parameter measuring the degree of oscillation. As λ increases, the degree of the oscillatory factor increases, which implies more cancellation should occur on average. Thus we should expect $I(\lambda)$ to decay as $\lambda \rightarrow \infty$. One of the main problems in the study of oscillatory integrals is to measure the asymptotic decay more precisely.

Example. The most basic example of an oscillatory integral is the Fourier transform, where for each function $f \in L^1(\mathbf{R})$, and each $\lambda \in \mathbf{R}$, we consider the quantity

$$\hat{f}(\lambda) = \int_{-\infty}^{\infty} e^{-2\pi i \lambda x} f(x) dx.$$

Thus in the oscillatory integral defining the Fourier transform of f , the function f plays the role of the amplitude, the phase function is $\phi(x) = x$. The basic theory of the Fourier transform hints that the decay of the oscillatory integral is related to the smoothness of the amplitude function f .

There are two main tools to estimate oscillatory integrals. Most classically, the method of steepest descent uses contour integration techniques

from complex analysis to shift the integral to a domain where less oscillation occurs, so we can apply standard estimation strategies. However, it is difficult to apply this method to oscillatory integrals over multivariate domains. The second tool, known as the method of stationary phase, determines the behaviour of the decay of an oscillatory integral by isolating the oscillation of a smooth phase ϕ to points where $\nabla\phi$ is smooth. If the zeroes of $\nabla\phi$ are isolated, then the oscillatory integrals can be localized near these values. Roughly speaking, each zero x_0 contributes $\psi(x_0)e^{2\pi i\lambda\phi(x_0)}$, times the volume of the region around x_0 where ϕ deviates by $\approx 1/\lambda$, to the asymptotic decay of $I(\lambda)$ as $\lambda \rightarrow \infty$.

27.1 One Dimensional Theory

Let us begin with a simple example of an oscillatory integral, i.e.

$$I(\lambda) = \int_J e^{2\pi i\lambda\phi(x)} dx,$$

where J is a closed interval, and $\phi : J \rightarrow \mathbf{R}$ is Borel measurable. Taking in absolute values shows that $|I(\lambda)| \leq |J|$ for all λ . If ϕ is a constant, then $I(\lambda) = |J|e^{i\lambda\phi}$, so in this case the estimate is sharp. But if ϕ varies, we expect $I(\lambda)$ to decay as $\lambda \rightarrow \infty$. For instance, there are various results, such as the Esseen concentration inequality, which show that if we are to expect *average* decay in the integral I over a range of λ , then ϕ must not be concentrated around any point.

Theorem 27.1 (Esseen Concentration Inequality). *Let $\phi : J \rightarrow \mathbf{R}$ be Borel measurable, and for each $\lambda \in \mathbf{R}$, set*

$$I(\lambda) = \int_J e^{2\pi i\lambda\phi(x)} dx.$$

Then for any $\varepsilon > 0$,

$$\sup_{a \in \mathbf{R}} |\{x \in J : |\phi(x) - a| \leq \varepsilon\}| \lesssim \varepsilon \int_0^{1/\varepsilon} |I(\lambda)| d\lambda,$$

where the implicit constant is independent of ϕ .

Proof. By rescaling, we may assume that $J = [0, 1]$. Moreover, for any choice of a , we may replace ϕ with $\phi - a$, reducing the analysis to the case where $a = 0$. Similarly, replacing ϕ with ϕ/ε reduces us to the situation where $\varepsilon = 1$. Thus we must show

$$|\{x \in [0, 1] : |\phi(x)| \leq 1\}| \lesssim \int_0^1 |I(\lambda)| d\lambda,$$

where the implicit constant is independent of the function ϕ . If ψ is an integrable function supported on $[0, 1]$, then Fubini's theorem implies

$$\begin{aligned} \int_0^1 \psi(\lambda) I(\lambda) d\lambda &= \int_0^1 \int_0^1 \psi(\lambda) e^{2\pi i \lambda \phi(x)} d\lambda dx \\ &= \int_0^1 \hat{\psi}(-\phi(x)) dx. \end{aligned}$$

In particular, this means that

$$\left| \int_0^1 \hat{\psi}(-\phi(x)) dx \right| \leq \|\psi\|_{L^\infty[0,1]} \int_0^1 |I(\lambda)| d\lambda.$$

If we choose a bounded function ψ such that $\hat{\psi}$ is non-negative, and bounded below on $[-1, 1]$, then

$$\left| \int_0^1 \hat{\psi}(-\phi(x)) dx \right| \gtrsim |\{x \in [0, 1] : |\phi(x)| \leq 1\}|,$$

and so the claim follows easily. \square

Remark. Recently, I have seen results that *reverse* the concentration bound, for instance, a paper of Basu, Guo, Zhang, and Zorin-Kranich entitled “A Stationary Set Method For Estimating Oscillatory Integrals”, and Wright’s paper “A Theory of Complex Oscillatory Integrals: A Case Study”. I do not know how general these techniques can be developed, but given an oscillatory integral in which we are able to prove stationary set estimates, but do not know asymptotics, it might be useful to look into these papers more.

Thus if large cancellation happens in $I(\lambda)$ for the average λ , this automatically implies that ϕ cannot be concentrated around any particular

point. Conversely, we want to show that if ϕ varies significantly, then I exhibits cancellation as $\lambda \rightarrow \infty$. One way to quantify this rate is through the derivative of the function ϕ , i.e. if the derivative has a large magnitude, the phase is varying fast. A bound on $|\phi'|$ from below is not sufficient to guarantee cancellation independent of the function ϕ , as the next example shows, if the integrand oscillates at a wavelength $1/\lambda$.

Example. Fix a positive integer λ_0 , and let $\phi_{\lambda_0}(x) = x + f(\lambda_0 x)/\lambda_0$, where f is smooth and 1-periodic, $\|f'\|_{L^\infty(\mathbf{R})} \leq 1/2$, and

$$\int_0^1 e^{2\pi i(x+f(x))} dx \neq 0.$$

Then for each $x \in \mathbf{R}$, $1/2 \leq |\phi'_{\lambda_0}(x)| \leq 2$, and in particular, is bounded independently of λ_0 . Since $\phi_{\lambda_0}(x + 1/\lambda_0) = \phi_{\lambda_0}(x) + 1/\lambda_0$, we find $e^{2\pi i \lambda_0 \phi_{\lambda_0}(x)}$ is $1/\lambda_0$ periodic. In particular, this means

$$\begin{aligned} I(\lambda_0) &= \int_0^1 e^{2\pi i \lambda_0 \phi_{\lambda_0}(x)} dx \\ &= \lambda_0 \int_0^{1/\lambda_0} e^{2\pi i(\lambda_0 x + f(\lambda_0 x))} dx \\ &= \int_0^1 e^{2\pi i(x+f(x))} dx. \end{aligned}$$

Thus $I(\lambda_0; \phi_{\lambda_0}) \sim 1$, independent of ϕ_0 , despite a uniform lower bound on the derivatives of the family of functions $\{\phi_{\lambda_0}\}$.

One option is a uniform upper bound on ϕ'' , in addition to lower bounding ϕ' , which eliminates the counterexample above, and yields a positive result.

Theorem 27.2. Let $\phi : J \rightarrow \mathbf{R}$ be twice continuously differentiable, and suppose there exists constants $A, B > 0$ with $|\phi'(x)| \geq A$ and $|\phi''(x)| \leq B$ for all $x \in J$. Then for all $\lambda > 0$, we find

$$|I(\lambda)| \lesssim \frac{1}{\lambda} \left(\frac{1}{A} + \frac{B}{A^2} |J| \right).$$

Proof. A dimensional analysis shows that the inequality is invariant under rescalings in x and λ , so we may assume that $J = [0, 1]$, and $\lambda = 1$. An integration by parts shows that

$$\begin{aligned} \int_0^1 e^{2\pi i \phi(x)} dx &= \int_0^1 \frac{1}{2\pi i \phi'(x)} \frac{d}{dx} \left(e^{2\pi i \phi(x)} \right) dx \\ &= \left(\frac{e^{2\pi i \phi(1)}}{2\pi i \phi'(1)} - \frac{e^{2\pi i \phi(0)}}{2\pi i \phi'(0)} \right) - \int_0^1 \frac{d}{dx} \left(\frac{1}{2\pi i \phi'(x)} \right) e^{2\pi i \phi(x)}. \end{aligned}$$

Now

$$\frac{d}{dx} \left(\frac{1}{\phi'(x)} \right) = -\frac{\phi''(x)}{\phi'(x)^2},$$

so taking in absolute values to the tree quantities in the sum above completes the proof. \square

One can keep applying absolute values to obtain further bounds in terms of higher order derivatives of ϕ . For instance, another integration by parts shows that if there is $A, B, C > 0$ such that for $x \in J$, if $\phi'(x) \geq A$, $\phi''(x) \leq B$, and $\phi'''(x) \leq C$, then

$$|I(\lambda)| \lesssim \frac{1}{\lambda} \left(\frac{1}{A} \right) + \frac{1}{\lambda^2} \left(\frac{B}{A^3} + \frac{C}{A^3} |J| + \frac{B^2}{A^4} |J| \right).$$

One can keep taking in absolute values, but the $1/\lambda$ decay will still remain. This is to be expected; for instance, if $\phi(x) = x$, and $J = [0, 1]$, then as $\lambda \rightarrow \infty$,

$$I(\lambda) = \int_0^1 e^{2\pi i \lambda x} dx = \frac{e^{2\pi i \lambda} - 1}{2\pi i \lambda}$$

and so

$$\limsup_{\lambda \rightarrow \infty} |I(\lambda) \cdot \lambda| = 2,$$

so we cannot obtain any better decay than $1/\lambda$ here.

Another option is to not require control on the second derivative of the phase, but instead to assume that ϕ' is monotone, which prevents the kind of oscillation present in our counterexample.

Lemma 27.3 (Van der Corput). *Let $\phi : \mathbf{R} \rightarrow \mathbf{R}$ be a smooth phase such that $|\phi'(x)| \geq A$ for all $x \in J$, and ϕ' is monotone. Then for all $\lambda > 0$ we have*

$$|I(\lambda)| \lesssim \frac{1}{A\lambda},$$

where the implicit constant is independent of J .

Proof. We may rescale so that $\lambda = 1$ and $J = [0, 1]$. Then the same integration by parts shows that

$$\int_0^1 e^{2\pi i \phi(x)} dx = \left(\frac{e^{2\pi i \phi(1)}}{2\pi i \phi'(1)} - \frac{e^{2\pi i \phi(0)}}{2\pi i \phi'(0)} \right) + \frac{1}{2\pi i} \int_0^1 \frac{d}{dx} \left(\frac{1}{2\pi i \phi'(x)} \right) e^{2\pi i \phi(x)} dx.$$

The two boundary terms are $O(1/A)$. For the integral, we apply a simple trick. Since ϕ' is monotone, so too is $1/\phi'$, so in particular, its derivative has a constant sign. Thus by the fundamental theorem of calculus,

$$\begin{aligned} \left| \int_J \frac{d}{dx} \left(\frac{1}{2\pi i \phi'(x)} \right) e^{2\pi i \phi(x)} dx \right| &\leq \int_J \left| \frac{d}{dx} \left(\frac{1}{2\pi i \phi'(x)} \right) \right| dx \\ &= \left| \int_J \frac{d}{dx} \left(\frac{1}{2\pi i \phi'(x)} \right) dx \right| \\ &= \frac{1}{2\pi \phi'(b)} - \frac{1}{2\pi \phi'(a)}. \end{aligned}$$

Again, this term is $O(1/A)$. □

Since the Van der Corput bound does not depend on $|J|$, it can be easily iterated using a decomposition strategy to give a theorem about higher derivatives of a function ϕ .

Lemma 27.4. *Let $\phi : \mathbf{R} \rightarrow \mathbf{R}$ be smooth, and suppose there is some $k \geq 2$ such that $|\phi^{(k)}(x)| \geq A$ for all $x \in J$. Then for all $\lambda > 0$, we find*

$$|I(\lambda)| \lesssim_k \frac{1}{(A\lambda)^{1/k}},$$

where the implicit constant is independent of J .

Proof. We perform an induction on k , the case $k = 1$ already proven. By scale invariance, we may assume $\lambda = 1$. Now $\phi^{(k-1)}$ is monotone, so for each $\alpha > 0$, outside an interval of length at most $O(\alpha/A)$, $|\phi^{(k-1)}(x)| \geq \alpha$. Thus applying the trivial bound in the excess region, and the case $k - 1$ on the other intervals, we conclude

$$|I(\lambda)| \lesssim_k \frac{\alpha}{A} + \alpha^{-1/(k-1)}$$

Optimizing over α , we find $|I(\lambda)| \lesssim_k A^{-1/k}$. \square

Remark. If $\phi^{(k-1)}$ vanishes at some point x_0 in J , then Taylor expansion shows that

$$|\{x \in J : |\phi(x) - \phi(x_0)| \leq \varepsilon\}| \gtrsim \varepsilon^{1/k}.$$

The Berry-Esseen theorem thus implies that the estimate $I(\lambda) \lesssim_k \lambda^{-1/k}$ is tight; we cannot have $I(\lambda) \lesssim \lambda^{-\alpha}$ for any $\alpha > 1/k$.

Let us now consider a one dimensional oscillatory integral with a varying amplitude function a , i.e.

$$I(\lambda) = \int_{-\infty}^{\infty} a(x) e^{2\pi i \lambda \phi(x)} dx.$$

The Van der Corput lemma also applies here.

Lemma 27.5. *Fix $k \geq 1$. Suppose a is supported on J , $|\phi^{(k)}(x)| \geq A$ for all $x \in J$, with ϕ' monotone if $k = 1$. Then*

$$|I(\lambda)| \lesssim_k \frac{\|a\|_{L^\infty(\mathbf{R})} + \|a'\|_{L^1(\mathbf{R})}}{(A\lambda)^{1/k}}.$$

Proof. Again, without loss of generality, we may assume $\lambda = 1$. Rescaling x means we can assume $J = [0, 1]$. For $x \in [0, 1]$, define

$$I_0(x) = \int_0^x e^{2\pi i \lambda \phi(t)} dt.$$

The standard Van-der Corput lemma implies that for all x ,

$$|I_0(x)| \lesssim_k \frac{1}{(A\lambda)^{1/k}}.$$

Integrating by parts, we find that

$$\begin{aligned} \int_0^1 a(x) e^{2\pi i \lambda \phi(x)} dx &= \int_0^1 a(x) I_0'(x) dx \\ &= [a(1)I_0(1) - a(0)I_0(0)] - \int_0^1 a'(x) I_0(x) dx. \end{aligned}$$

Now

$$|a(1)I_0(1) - a(0)I_0(0)| \lesssim \frac{\|a\|_{L^\infty(\mathbf{R})}}{(A\lambda)^{1/k}}$$

and

$$\left| \int_0^1 a'(x) I_0(x) dx \right| \lesssim_k \frac{\|a'\|_{L^1(\mathbf{R})}}{(A\lambda)^{1/k}}.$$

Putting these two estimates together completes the proof. \square

Remark. Since the bound doesn't depend on the interval, if a is not compactly supported, but a' is integrable, and the other assumptions of the last result holds, then we still have a bound

$$\sup_{a < b} \left| \int_a^b a(x) e^{2\pi i \lambda \phi(x)} dx \right| \lesssim_k \frac{\|a\|_{L^\infty(\mathbf{R})} + \|a'\|_{L^1(\mathbf{R})}}{(A\lambda)^{1/k}}$$

If a is smooth and compactly supported, repeated integration by parts is very successful because there are no boundary terms, and so we get very fast decay as $\lambda \rightarrow \infty$.

Theorem 27.6. *Let J be an interval, and fix $n > 0$. Consider a real-valued phase function $\phi \in C^{n+1}(\mathbf{R})$ with $\phi'(x) \neq 0$ on J , and an amplitude $a \in C^{n+1}(\mathbf{R})$ with $\text{supp}(a) \subset J$. Then*

$$|I(\lambda)| \lesssim_{n,K,\phi} \lambda^{-n} \sum_{k=0}^n \|a^{(k)}\|_{L^\infty(J)} \|\phi'\|^{k-2n}_{L^\infty(J)},$$

where the implicit constant is uniformly bounded over a family of phases ϕ which are bounded in $C^{n+1}(J)$.

Proof. When $n = 0$, the result is trivial. The remaining cases we shall prove by induction. A single integration by parts gives

$$\begin{aligned} I(\lambda) &= \frac{1}{\lambda} \int \frac{a(x)}{2\pi i \phi'(x)} \frac{d}{dx} \left(e^{2\pi i \lambda \phi(x)} \right) \\ &= -\frac{1}{2\pi i \lambda} \int \frac{d}{dx} \left(\frac{a(x)}{\phi'(x)} \right) e^{2\pi i \lambda \phi(x)} \\ &= \frac{1}{2\pi i \lambda} \int \frac{\phi'(x) a'(x) - a(x) \phi''(x)}{\phi'(x)^2} e^{2\pi i \lambda \phi(x)}. \end{aligned}$$

Applying induction, this quantity is bounded up to a universal constant depending solely on the derivatives of ϕ up to order $n - 1$ by

$$\lambda^{-n} \sum_{k=0}^{n-1} \left\| \left(\frac{\phi' a' - a \phi''}{(\phi')^2} \right)^{(k)} |\phi'|^{k-2(n-1)} \right\|_{L^\infty(J)}$$

and applying the product rule, we see this term is bounded, up to a universal constant depending solely on derivatives of ϕ up to order n , by the required quantity. \square

Remark. Differentiation under the integration sign shows

$$\left(\frac{d}{d\lambda} \right)^n I(\lambda) = (2\pi i)^n \int \phi(x)^n a(x) e^{2\pi i \lambda \phi(x)} dx.$$

Since $\phi^n a$ satisfies the same assumptions that a does, it follows that for any N , we have

$$\left| \left(\frac{d}{d\lambda} \right)^n I(\lambda) \right| \lesssim_{a, \phi, N, n} \lambda^{-N}.$$

Thus I is actually a Schwartz function on the real line.

Let us now move onto a ‘stationary phase’, i.e. a phase ϕ whose derivative vanishes at a point. The simplest example of such a phase is the integral

$$I(\lambda) = \int_{-\infty}^{\infty} a(x) e^{2\pi i \lambda x^2} dx,$$

where $a \in C_c^\infty(\mathbf{R})$. If a is nonzero in a neighborhood of the origin, and if $\phi(x) = x^2$, then

$$|\{x \in \mathbf{R} : |\phi(x)| \leq 1/\lambda\}| \lesssim \lambda^{-1/2}.$$

Thus our heuristics tell us that we should expect $I(\lambda)$ decays on the order of $\lambda^{-1/2}$, which agrees with the asymptotics we now find. A dyadic decomposition about the origin, combined with a single integration by parts at each scale yields a bound $|I(\lambda)| \lesssim \lambda^{-1/2}$.

Theorem 27.7. *Let $a \in \mathcal{S}(\mathbf{R})$. Then*

$$I(\lambda) \sim e^{i\pi/4} (2\lambda)^{-1/2} \sum_{n=0}^{\infty} (i/8\pi)^n \frac{a^{(2n)}(0)}{n!} \cdot \frac{1}{\lambda^n}$$

That is, for each pair of nonnegative integers N and k

$$\begin{aligned} \left(\frac{d}{d\lambda}\right)^k I(\lambda) &= \left(\frac{d}{d\lambda}\right)^k \left\{ e^{i\pi/4} \cdot (2\lambda)^{-1/2} \sum_{n=0}^N (i/8\pi)^n \frac{a^{(2n)}(0)}{n!} \cdot \frac{1}{\lambda^n} \right\} \\ &\quad + O_{N,m,a}(1/\lambda^{N+m+3/2}). \end{aligned}$$

In particular, if a equals one near the origin, then $I(\lambda) \sim e^{i\pi/4} (2\lambda)^{-1/2}$.

Proof. Applying the multiplication formula for the Fourier transform, noting that the distributional Fourier transform of $e^{2\pi i \lambda x^2}$ is

$$(2\lambda)^{-1/2} e^{i\pi/4} e^{-i\pi \xi^2 / 2\lambda}$$

Thus

$$I(\lambda) = (2\lambda)^{-1/2} e^{i\pi/4} \int_{-\infty}^{\infty} e^{-i\pi \xi^2 / 2\lambda} \widehat{a}(\xi) d\xi.$$

Now for any N , we can write

$$e^{-i\pi \xi^2 / 2\lambda} = \sum_{n=0}^N \frac{1}{n!} \left(\frac{-i\pi \xi^2}{2\lambda} \right)^n + O_N((\xi^2/\lambda)^{N+1})$$

Thus substituting in the Taylor series, and then applying the Fourier inversion formula, we find

$$\begin{aligned} I(\lambda) &= (2\lambda)^{-1/2} e^{i\pi/4} \sum_{n=0}^N \frac{1}{n!} \int_{-\infty}^{\infty} \left(\frac{-i\pi \xi^2}{2\lambda} \right)^n \widehat{a}(\xi) d\xi + O_{a,N}(1/\lambda^{N+3/2}) \\ &= (2\lambda)^{-1/2} e^{i\pi/4} \sum_{n=0}^N (i/8\pi)^n \frac{1}{n!} \frac{1}{\lambda^n} \int_{-\infty}^{\infty} (2\pi i \xi)^{2n} \widehat{a}(\xi) d\xi + O_{a,N}(1/\lambda^{N+3/2}) \\ &= (2\lambda)^{-1/2} e^{i\pi/4} \sum_{n=0}^N (i/8\pi)^n \frac{a^{(2n)}(0)}{n!} \frac{1}{\lambda^n} + O_{a,N}(1/\lambda^{N+3/2}). \quad \square \end{aligned}$$

One obtains the asymptotic formula for the derivative of I by noting that

$$I^{(k)}(\lambda) = \int (-2\pi i x^2)^k a(x) e^{-2\pi i \lambda x^2} dx,$$

which reduces to the case where $k = 0$.

It requires only a simple change of variables to extend this theorem to arbitrary quadratic phases. We say a critical point of a function is *non-degenerate* if the second derivative at that point is nonzero.

Theorem 27.8. *Let ϕ be a smooth phase with a single, non-degenerate critical point x_0 , and let a be a smooth compactly supported amplitude function. Then there exists a sequence of constants $\{c_n\}$, depending solely on the derivatives of a and ϕ at x_0 , such that*

$$I(\lambda) \sim e^{2\pi i \lambda \phi(x_0)} \lambda^{-1/2} \sum_{n=0}^{\infty} c_n \lambda^{-n}.$$

The implicit constants in the asymptotic formula are uniform over a family of amplitude functions a given uniform bounds on the support of a , and upper bounds on $2N$ derivatives of a . We can explicitly calculate that

$$c_0 = \sqrt{\frac{i}{\phi''(x_0)}} \cdot a(x_0).$$

More generally, in this case there exists a sequence of linear differential operators $\{L_n\}$, with L_n order $2n$, with coefficients depending on the derivatives of ϕ at x_0 , such that $c_n = (L_n a)(x_0)$.

Proof. Translating our integration variables if necessary, we may assume that the critical point x_0 lies at the origin. Partitioning the support of a , applying the principle of nonstationary phase away from the critical point, we may assume without loss of generality that a is supported on an arbitrarily small neighborhood of the critical point. In particular, we may assume that $\phi(x) \neq 0$ for all nonzero x in the support of a . A coordinate change $x \mapsto -x$ means we may assume that $\phi''(0) > 0$. We can then define a function

$$y(x) = \operatorname{sgn}(x) \cdot \phi(x)^{1/2}.$$

It follows from our assumptions that y is a smooth diffeomorphism on the support of a . By the change of variables formula, there exists a smooth, compactly supported function $a_0(y)$ such that

$$I(\lambda) = \int a(x) e^{2\pi i \lambda \phi(x)} dx = \int a_0(y) e^{2\pi i \lambda y^2} dy.$$

Thus we can apply the previous theorem to conclude that there exists a sequence of constants $\{c_n\}$ such that for each N ,

$$\left(\frac{d}{d\lambda}\right)^k I(\lambda) = \left(\frac{d}{d\lambda}\right)^k \left\{ \lambda^{-1/2} \sum_{n=0}^N c_n \lambda^{-n} \right\} + O_{\phi,a,N,k}(1/\lambda^{N+m+3/2}).$$

The existence in this theorem is a *constructive* existence statement. The proof gives an effective algorithm to produce as many constants c_n as required for any particular phase ϕ , provided one can explicitly write out the function $y(x)$. In particular, if the phase has only a single stationary point at the origin,

$$c_0 = 2^{-1/2} e^{i\pi/4} a_0(0) = 2^{-1/2} e^{i\pi/4} a(0)/y'(0).$$

Since

$$\begin{aligned} y'(0) &= \lim_{x \rightarrow 0^+} y'(x) = \lim_{x \rightarrow 0^+} \frac{\phi'(x)}{2\phi(x)^{1/2}} \\ &= \lim_{x \rightarrow 0^+} (1/2) \frac{\phi'(x)}{x} \left(\frac{x^2}{\phi(x)} \right)^{1/2} = (\phi''(0)/2)^{1/2}. \end{aligned}$$

This means that $c_0 = e^{i\pi/4} a(0) \phi''(0)^{-1/2}$. □

If the phase ϕ has a critical point of order greater than two, then the asymptotics of the oscillatory integral get worse. In particular, if ϕ has a zero of order k , then around this region ϕ differs by $1/\lambda$ on an interval of length $1/\lambda^{1/k}$, so we might expect $I(\lambda)$ to be proportional to $\lambda^{1/k}$. This is precisely what happens, but our proof will not rely on the Fourier transform since the computation of the Fourier transform of $e^{\lambda i x^k}$ is quite difficult to calculate when $k > 2$. The next proof also works for the case $k = 2$, but the proof involves some contour shifting. Since the large majority of the examples we consider will have nondegenerate critical points (this is the generic behaviour of critical points), these complicated asymptotics can be safely skipped on a first reading.

Lemma 27.9. *For any non-negative integers l and k , there is a positive constant $A_{kl} > 0$ such that for any $\lambda \in \mathbf{R}$ and $\varepsilon > 0$,*

$$\int_0^\infty e^{2\pi i \lambda x^k} e^{-\varepsilon x^k} x^l dx = A_{kl} (\varepsilon - 2\pi i \lambda)^{-(l+1)/k},$$

where the k th root is the principal root for non-negative complex numbers.

Proof. If $z = (\varepsilon - 2\pi i \lambda)^{1/k} x$, and if α_N is the ray between the origin and the point $N(\varepsilon - 2\pi i \lambda)^{1/k}$, then

$$\int_0^N e^{2\pi i \lambda x^k} e^{-\varepsilon x^k} x^l dx = (\varepsilon - 2\pi i \lambda)^{-(l+1)/k} \int_{\alpha_N} e^{-z^k} z^l dz.$$

Let $\theta \in (-\pi/2, 0]$ be the argument of $(\varepsilon - 2\pi i \lambda)^{1/k}$, and set β_N to be the arc between $N(\varepsilon - 2\pi i \lambda)^{1/k}$ and $N(\varepsilon^2 + \lambda^2)^{1/2}$. Then β_N has length $O(N)$, with implicit constant depending on λ and ε . Moreover, any point z on β_N has modulus $N(\varepsilon^2 + \lambda^2)^{1/2}$ and argument less than or equal to θ/k . But this implies that $\operatorname{Re}(z^k) \geq N^k (\varepsilon^2 + \lambda^2)^{k/2} \cos(\theta)$, and so there exists a constant c depending on ε and λ such that $|e^{-z^k}| \leq e^{cN^k}$. But this means that $|z^l e^{-z^k}| \leq N^l e^{-cN^k}$. Thus taking in absolute values gives that

$$\lim_{N \rightarrow \infty} \int_{\beta_N} e^{-z^k} z^l dz = 0.$$

In particular, applying Cauchy's theorem, we conclude that

$$\lim_{N \rightarrow \infty} \int_{\gamma_N} e^{-z^k} z^l dz = \int_0^\infty e^{-x^k} x^l dx.$$

If we denote the latter integral by $A_{kl} > 0$, then we have shown that

$$\int_0^\infty e^{2\pi i \lambda x^k} e^{-\varepsilon x^k} x^l dx = A_{kl} \cdot (\varepsilon - 2\pi i \lambda)^{-(l+1)/k},$$

as was required to be shown. □

Remark. In particular, this implies that for each ε , there exists constants A_{kln} such that

$$\int_0^\infty e^{2\pi i \lambda x^k} e^{-x^k} x^l dx = (2\pi \lambda)^{-(l+1)/k} \sum_{n=0}^\infty A_{kln} (2\pi \lambda)^{-n}.$$

This is obtained by taking the Laurent series of

$$(1 - 2\pi i \lambda)^{-(l+1)/k} = (2\pi \lambda)^{-(l+1)/k} (\lambda^{-1} - 2\pi i)^{-(l+1)/k},$$

In particular, for each N and for each λ , we conclude

$$\int_0^\infty e^{2\pi i \lambda x^k} e^{-x^k} x^l dx = \lambda^{-(l+1)/k} \sum_{n=0}^N A_{kln} (2\pi \lambda)^{-n} + O_N \left(1/\lambda^{n+1+1/k} \right).$$

Lemma 27.10. *If η is compactly supported and smooth, then*

$$\left| \int_{-\infty}^\infty e^{2\pi i \lambda x^k} x^l \eta(x) dx \right| \lesssim_{l,k,\eta} \lambda^{-(l+1)/k}.$$

Proof. Let α be a bump function supported on $[-2, 2]$ with $\alpha(x) = 1$ for $|x| \leq 1$. For each $\varepsilon > 0$, write

$$\begin{aligned} \int_{-\infty}^\infty e^{2\pi i \lambda x^k} x^l \eta(x) dx &= \int_{-\infty}^\infty e^{2\pi i \lambda x^k} x^l \eta(x) \alpha(x/\varepsilon) dx \\ &\quad + \int_{-\infty}^\infty e^{2\pi i \lambda x^k} x^l \eta(x) (1 - \alpha(x/\varepsilon)) dx, \end{aligned}$$

where we will bound each term and optimize for a small ε . We trivially have

$$\left| \int_{-\infty}^\infty e^{2\pi i \lambda x^k} x^l \eta(x) \alpha(x/\varepsilon) dx \right| \lesssim_\eta \varepsilon^{l+1},$$

We apply an integration by parts to the second integral, noting that $e^{2\pi i \lambda x^k}$ is a fixed point of the differential operator

$$Df = \frac{1}{2\pi i \lambda k x^{k-1}} \frac{df}{dx}.$$

If we consider the differential operator

$$D^*g = \frac{d}{dx} \left(\frac{-f}{2\pi i \lambda k x^{k-1}} \right) = \left(\frac{i}{2\pi \lambda k} \right) \left(\frac{f'(x)}{x^{k-1}} - \frac{(k-1)f(x)}{x^k} \right),$$

then for any smooth f and compactly supported g ,

$$\int_{-\infty}^\infty (Df)(x)g(x) dx = \int_{-\infty}^\infty f(x)(D^*g)(x) dx.$$

In particular,

$$\begin{aligned} \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} x^l \eta(x) (1 - \alpha(x/\varepsilon)) dx &= \int_{-\infty}^{\infty} D^N(e^{2\pi i \lambda x^k}) x^l \eta(x) (1 - \alpha(x/\varepsilon)) dx \\ &= \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} (D^*)^N \{x^l \eta(x) (1 - \alpha(x/\varepsilon))\} dx. \end{aligned}$$

Write $g_N(x) = (D^*)^N \{x^l \eta(x) (1 - \alpha(x/\varepsilon))\}$. Since $x^l \eta(x) (1 - \alpha(x/\varepsilon))$ vanishes for $|x| \leq \varepsilon$, so too does $g_N(x)$. For $N \geq l/(k-1)$, and $|x| \geq \varepsilon$, we have

$$|g_N(x)| \lesssim_{N,\eta} \lambda^{-N} \varepsilon^{-N} |x|^{l-N(k-1)},$$

where the implicit constant depends on upper bounds for the derivatives of η of order to N . We can thus take in absolute values after integrating by parts to conclude that if $N > (l+1)/(k-1)$, then

$$\left| \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} x^l \eta(x) (1 - \alpha(x/\varepsilon)) dx \right| \lesssim_{N,\eta} \lambda^{-N} \varepsilon^{l+1-Nk}$$

Thus we can put the two bounds together to conclude that

$$\left| \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} x^l \eta(x) dx \right| \lesssim_{N,\eta} \varepsilon^{l+1} + \lambda^{-N} \varepsilon^{l+1-Nk}.$$

Picking $\varepsilon = \lambda^{-1/k}$ gives

$$\left| \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} x^l \eta(x) dx \right| \lesssim_{N,\psi} \lambda^{-(l+1)/k}. \quad \square$$

But N was chosen depending only on k and l , so the implicit constants depend on the correct variables.

We can now prove the asymptotics for the model case $\phi(x) = x^k$.

Theorem 27.11. *Suppose a is a smooth compactly supported amplitude, and ϕ is a smooth phase with $\phi'(x) \neq 0$ on the support of a except at some point x_0 , where $\phi'(x_0) = \dots = \phi^{(k-1)}(x_0) = 0$, and $\phi^{(k)}(x_0) \neq 0$. Then there is a sequence $\{c_n\}$ such that*

$$I(\lambda) \sim \lambda^{-1/k} \sum_{n=0}^{\infty} c_n \lambda^{-n/k}.$$

Proof. Let us begin with the model case $\phi(x) = x^k$. Let \tilde{a} be a bump function with $\tilde{a}(x) = 1$ for all x with $a(x) > 0$. Then

$$I(\lambda) = \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} e^{-x^k} [e^{x^k} a(x)] \tilde{a}(x) dx.$$

For each N , perform a Taylor expansion, writing

$$e^{x^k} a(x) = \sum_{n=0}^N c_n x^n + x^{N+1} R_N(x).$$

Thus if $P_N(x) = \sum_{n=0}^N c_n x^n$,

$$\begin{aligned} & \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} e^{-x^k} [e^{x^k} a(x)] \tilde{a}(x) dx \\ &= \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} e^{-x^k} P_N(x) dx \\ &+ \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} e^{-x^k} P_N(x) (\tilde{a}(x) - 1) dx \\ &+ \int_{-\infty}^{\infty} e^{2\pi i \lambda x^k} e^{-x^k} x^{N+1} R_N(x) \tilde{a}(x) dx. \end{aligned}$$

The first integral can be expanded in the required power series. The second integral, since it is supported away from the origin, is $O_M(\lambda^{-M})$ for any $M > 0$. And in the last lemma we showed the third integral is $O(\lambda^{-(N+2)/k})$, so combining these three terms gives the required result. The general case follows from a change of variables. \square

Remark. As we saw in the case $k = 2$, if k is even and n is odd then

$$\int_{-\infty}^{\infty} e^{\lambda i x^k} e^{-x^k} x^n = 0.$$

Thus we can actually improve the asymptotics to the existence of a sequence $\{c_n\}$ such that

$$I(\lambda) = \lambda^{-1/k} \sum_{n=0}^N c_n \lambda^{-2n/k} + O_{\phi, a, N} \left(1/\lambda^{(2N+3)/k} \right).$$

Let us now consider some examples of the method of stationary phase in one dimension.

Example. The Bessel function of order m , denoted $J_m(r)$, is defined to be the oscillatory integral

$$J_m(r) = \int_0^1 e^{2\pi i r \sin(2\pi\theta)} e^{-2\pi i m\theta} d\theta.$$

We want to use the method of stationary phase to determine the decay of $J_m(r)$ as $r \rightarrow \infty$. The amplitude is $a(\theta) = e^{-2\pi i m\theta}$, and the phase is $\phi(\theta) = \sin(2\pi\theta)$. We note that the phase $\phi(\theta) = \sin(2\pi\theta)$ is stationary when $\theta = 1/4$ and $\theta = 3/4$, and that these stationary points are nondegenerate. Thus we might expect $|J_m(r)| = O_m(r^{-1/2})$. More precisely, we write $a = a_1 + a_2 + a_3$, where a_1 is supported in a small neighbourhood of $1/4$, a_2 in a neighbourhood of $3/4$, and a_3 is supported away from $1/4$ and $3/4$. Then $I_{a_1}(\lambda)$ and $I_{a_2}(\lambda)$ are oscillatory integrals with a unique nondegenerate stationary point. In fact, we find that

$$I_{a_1}(r) = (1/2\pi) e^{2\pi i(r-1/8-m/4)} + O(r^{-3/2}).$$

$$I_{a_2}(r) = (1/2\pi) e^{-2\pi i(r-1/8-m/4)} + O(r^{-3/2}).$$

The integral $I_{a_3}(\lambda)$ can be shifted (using periodicity) into a compactly supported integral with smooth amplitude and no stationary points, and thus decays arbitrarily fast, i.e. $|I_{a_3}(\lambda)| = O(\lambda^{-3/2})$. Thus summing up these estimates gives

$$I_a(r) = (1/\pi) \cos(2\pi r - \pi/4 - \pi m/2) + O(r^{-3/2}).$$

Example. Consider the Airy function

$$Ai(x) = \int_{-\infty}^{\infty} e^{2\pi i(x\xi + \xi^3/3)} d\xi,$$

which arises as a solution to the differential equation $y'' = xy$. Again, this integral is not defined absolutely. Nonetheless, for large N , an application of the Van der Corput lemma implies that for any finite interval I containing only points x with $|x| \geq N$,

$$\int_I e^{2\pi i(x\xi + \xi^3/3)} d\xi = O(1/N),$$

where the implicit constant is independent of I . Thus we can interpret the integral as

$$\lim_{n \rightarrow \infty} \int_{a_n}^{b_n} e^{2\pi i(x\xi + \xi^3/3)} d\xi,$$

where $\{a_n\}$ and $\{b_n\}$ are any sequences with $a_n \rightarrow -\infty$, $b_n \rightarrow \infty$.

Now consider the phase $\phi(\xi) = x\xi + \xi^3/3$. Then $\phi'(\xi) = x + \xi^2$. When x is negative, there are two stationary points. Thus we can rescale the integral, writing $v = x^{-1/2}\xi$, so that for $x > 0$,

$$\text{Ai}(-x) = x^{1/2} \int_{-\infty}^{\infty} e^{2\pi i x^{3/2}(v^3/3 - v)} dv.$$

If we write $\phi_0(v) = v^3/3 - v$, then ϕ_0 has two stationary points, at $v = \pm 1$. These stationary points are non-degenerate, so if we write $1 = a_1 + a_2 + a_3 + a_4$, where a_1 equal to one in a neighbourhood of 1, a_2 equal to one in a neighbourhood of -1 , and a_3 is supported in the region between -1 and 1 , and a_4 vanishes in all such regions, then we decompose $\text{Ai}(-x)$ as $I_1 + I_2 + I_3 + I_4$. Now the principle of stationary phase tells us that

$$I_1 = (1/\sqrt{2})e^{2\pi i(-2/3x^{3/2}+1/8)}x^{-1/4} + O(x^{-7/4})$$

$$I_2 = (1/\sqrt{2})e^{2\pi i(2/3x^{3/2}-1/8)}x^{-1/4} + O(x^{-7/4})$$

Moreover, $I_3 = O_N(x^{-N})$ for all $N \geq 0$. It remains to show $I_4 = O(x^{-1})$. Indeed, an integration by parts shows that

$$\begin{aligned} I_4 &= x^{1/2} \int_{-\infty}^{\infty} e^{2\pi i x^{3/2} \phi_0(v)} a_4(v) dv \\ &= \frac{-1}{2\pi i x} \int_{-\infty}^{\infty} e^{2\pi i x^{3/2} \phi_0(v)} \frac{d}{dv} \left(\frac{a_4(v)}{v^2 - 1} \right) dv. \end{aligned}$$

Taking in absolute values shows $|I_4| \lesssim 1/x$. Thus as $x \rightarrow \infty$,

$$\text{Ai}(-x) = \sqrt{2}x^{-1/4} \cos((-4\pi/3)x^{3/2} + \pi/4) + O(1/x),$$

which gives the first order asymptotics of the integral.

On the other hand, let us consider large positive x . Then the phase ϕ has no critical points, and we therefore expect very fast decay. To achieve this decay, we employ a contour shift, replacing the oscillatory integral with a different

oscillatory integral which has a stationary point, so we can obtain asymptotics here. If we write $\phi(z) = xz + z^3/3$, then $\phi'(z) = 0$ when $z = \pm ix^{1/2}$. A simple contour shift argument to the line $\mathbf{R} + ix^{1/2}$ gives

$$Ai(x) = \int_{-\infty}^{\infty} e^{2\pi i \phi(\xi + ix^{1/2})} d\xi = e^{-(4\pi/3)x^{3/2}} \int_{-\infty}^{\infty} e^{-2\pi \xi^2 x^{1/2}} e^{2\pi i \xi^3/3} d\xi.$$

We have

$$\int_{-\infty}^{\infty} e^{-2\pi \xi^2 x^{1/2}} e^{2\pi i \xi^3/3} d\xi \approx x^{-1/4} \int_{-\infty}^{\infty} e^{-2\pi \xi^2} e^{2\pi i x^{-3/4} \xi^3/3} d\xi.$$

Now a Taylor series shows

$$e^{2\pi i x^{-3/4} \xi^3/3} = 1 + O(x^{-3/4} \xi^3/3),$$

so, plugging in, we conclude

$$Ai(x) = 2^{-1/2} x^{-1/4} e^{-(4\pi/3)x^{3/2}} + O(x^{-3/4} e^{-(2/3)x^{3/2}}).$$

Thus Airy's function decreases exponentially as $x \rightarrow \infty$.

Example. Let us consider the integral quantities

$$\int_0^1 e^{2\pi i x \xi} e^{2\pi i/x} x^{-\gamma} dx$$

where to avoid technicalities we assume $0 \leq \gamma < 2$. These integral quantities are not defined absolutely, so we actually interpret this integral as

$$\lim_{\varepsilon \rightarrow 0} \int_0^1 e^{2\pi i x \xi} e^{2\pi i/x} x^{-\gamma} dx$$

If we write $\phi(x) = x\xi + 1/x$, then

$$\int_0^1 e^{2\pi i x \xi} e^{2\pi i/x} x^{-\gamma} dx = \int_0^1 e^{2\pi i \phi(x)} x^{-\gamma} dx.$$

For $0 < \varepsilon_1 < \varepsilon_2 < \varepsilon$, since $\phi'(x) = \xi - 1/x^2$, an easy integration by parts shows that for $\varepsilon \leq \xi^{-1/2}/2$,

$$\begin{aligned} \int_{\varepsilon_1}^{\varepsilon_2} e^{2\pi i \phi(x)} x^{-\gamma} dx &= \frac{1}{2\pi i \xi} \int_{\varepsilon_1}^{\varepsilon_2} \frac{d}{dx} \left(e^{2\pi i \phi(x)} \right) \frac{x^{2-\gamma}}{x^2 - 1/\xi} dx \\ &= \frac{-1}{2\pi i \xi} \int_{\varepsilon_1}^{\varepsilon_2} e^{2\pi i \phi(x)} \frac{d}{dx} \left(\frac{x^{2-\gamma}}{x^2 - 1/\xi} \right) + O(\varepsilon^{2-\gamma}) \quad (27.1) \\ &= O(\varepsilon^{2-\gamma}), \end{aligned}$$

where the constant is independent of ξ . This implies the limit we study exists. We wish to prove an asymptotic formula for this integral as $\xi \rightarrow \infty$. If we write $\phi(x) = x\xi + 1/x$, then

$$\int_0^1 e^{2\pi i x \xi} e^{2\pi i/x} x^{-\gamma} dx = \int_0^1 e^{2\pi i \phi(x)} x^{-\gamma} dx.$$

Since ϕ has a nondegenerate stationary point when $x = \xi^{-1/2}$, our heuristics might suggest that if the phase and amplitude were smooth at the origin, then

$$\int_0^1 e^{2\pi i \phi(x)} x^{-\gamma} \approx (1/2)^{1/2} e^{4\pi i \xi^{1/2}} e^{-i\pi/4} \xi^{\gamma/2-3/4}.$$

We shall show that these heuristics continue to hold, up to an error of $O(\xi^{\gamma/2-1})$.

In an attempt to isolate the critical point, we split the interval $[0, 1]$ into three parts, $[0, 0.5\xi^{-1/2}]$, $[0.5\xi^{-1/2}, 1.5\xi^{-1/2}]$, and $[1.5\xi^{-1/2}, 1]$, obtaining three integrals I_1 , I_2 , and I_3 . The calculation (27.1) shows that $|I_1| \lesssim \xi^{\gamma/2-1}$, and thus is negligible to our asymptotic formula. To obtain a bound on I_3 , we use the Van der Corput lemma, noting that $\phi'(x) = \xi - 1/x^2$ is monotone, and $|\phi'(x)| \gtrsim \xi$ for $x \geq 1.5\xi^{-1/2}$. Thus we find $|I_3| \lesssim \xi^{-1}$, and thus is also negligible to our formula. Thus we are left with the trick part of calculating I_2 accurately. It will be easiest to do this by renormalizing the integral, i.e. writing $y = \xi^{1/2}x$, and calculating

$$I_2 = \int_{0.5\xi^{-1/2}}^{1.5\xi^{-1/2}} e^{2\pi i \phi(x)} x^{-\gamma} dx = \xi^{\gamma/2-1/2} \int_{0.5}^{1.5} e^{2\pi i \xi^{1/2}(y+1/y)} y^{-\gamma} dy.$$

We consider a smooth amplitude function $\psi(y)$ supported on the interior of $[0.5, 1.5]$. Then since $y + 1/y$ is stationary at $y = 1$, but non-degenerate, we can write

$$\int e^{2\pi i \xi^{1/2}(y+1/y)} y^{-\gamma} \psi(y) dy = \xi^{-1/4} \pi^{1/2} e^{2\pi i (2\xi^{1/2}+1/4)} + O(\xi^{-1/2}),$$

from which we obtain our main term. On the other hand, we can apply the Van der Corput lemma to show that

$$\int_{0.5}^{1.5} e^{2\pi i \xi^{1/2}(y+1/y)} y^{-\gamma} (1-\psi(y)) dy = \int e^{2\pi i \xi^{1/2}(y+1/y)} y^{-\gamma} \psi(y) dy = O(\xi^{-1/2}).$$

Combining all these estimates gives the theorem.

On the other hand, consider the integral

$$I(\xi) = \int_0^1 e^{-2\pi i \xi x} e^{2\pi i/x} x^{-\gamma} dx = \int_0^1 e^{2\pi i \phi(x)} x^{-\gamma} dx,$$

where $\phi(x) = 1/x - \xi x$ is the phase. Then the phase has no critical points so we can assume that we can large decay for large ξ . We decompose the integral onto the intervals $[0, \xi^{-1/2}]$ and $[\xi^{-1/2}, 1]$, inducing the two quantities I_1 and I_2 . Now applying the Van der Corput lemma to I_2 with $|\phi'(x)| = |1/x^2 + \xi| \geq \xi$ for $x \geq 0$, gives $|I_2| \lesssim \xi^{\gamma/2-1}$. On the other hand, renormalizing with $y = \xi^{1/2}x$, we have

$$I_1 = \xi^{\gamma/2-1/2} \int_0^1 e^{2\pi i \xi^{1/2}(1/y-y)} y^{-\gamma} dy.$$

For each n , we note that for the phase $\phi_0(x) = 1/y - y$, for $1/2^{n+1} \leq y \leq 1/2^n$, we have $|\phi'_0(x)| \gtrsim 4^n$. Thus we can apply the Van der Corput lemma to conclude

$$\left| \int_{1/2^{n+1}}^{1/2^n} e^{2\pi i \phi_0(x)} y^{-\gamma} dy \right| \lesssim \frac{2^{\gamma n}}{4^n \xi^{1/2}}.$$

Summing up over all $n \geq 0$, we conclude $|I_1| \lesssim \xi^{\gamma/2-1}$. Thus $|I(\xi)| \lesssim \xi^{\gamma/2-1}$.

One way to interpret this asymptotic formula is through a Riemann singularity, i.e. a tempered distribution Λ supported on the half-line $x \geq 0$, that agrees with the oscillatory function $e^{2\pi i/x} x^{-\gamma}$ for small x , but is compactly supported and smooth away from the origin. We consider the case $0 \leq \gamma < 2$ for simplicity. Thus for Schwartz $f \in \mathcal{S}(\mathbf{R})$, we have

$$\Lambda(f) = \lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon}^{\infty} f(x) e^{2\pi i/x} x^{-\gamma} \psi(x) dx,$$

where ψ is smooth and compactly supported, and equals one in a neighbourhood of the origin. An easy integration by parts shows that for a fixed Schwartz f , and for $0 < \varepsilon_1 < \varepsilon_2 < \varepsilon$,

$$\left| \int_{\varepsilon_1}^{\varepsilon_2} f(x) e^{2\pi i/x} x^{-\gamma} dx \right| = O(\varepsilon^{2-\gamma}),$$

where the implicit constants depend on upper bounds for f and f' in a neighbourhood of the origin. Thus we find $\Lambda(f)$ is well defined, and moreover, Λ is

a distribution of order one. Since Λ is compactly supported, the Paley-Weiner theorem implies that $\hat{\Lambda}$ is a distribution represented by a locally integrable function, and

$$\hat{\Lambda}(\xi) = \int_0^\infty e^{2\pi i/x} x^{-\gamma} e^{-2\pi \xi i x} dx.$$

The calculations above give asymptotic formulas for this Fourier transform. In particular, we see that the Fourier transform of Λ decays much faster to the right than to the left.

27.2 Stationary Phase in Multiple Variables

When we move from a single variable oscillatory integral to a multivariable oscillatory integrals. Thus we consider the oscillatory integral

$$I(\lambda) = \int_{\mathbf{R}^d} a(x) e^{2\pi i \lambda \phi(x)} dx.$$

for large λ . The method of stationary phase becomes significantly more complicated in this setting because the stationary points of the phase function are no longer necessarily isolated. In certain basic situations, such as when the stationary points are isolated and satisfy a nondegeneracy condition, we can obtain asymptotic formulae.

Theorem 27.12. *Fix a compact set K and a positive integer $n > 0$, and consider an amplitude $a \in C^n(\mathbf{R}^d)$ with $\text{supp}(a) \subset K$, and a real-valued phase $\phi \in C^{n+1}(\mathbf{R}^d)$, with $\nabla \phi(x) \neq 0$ on K . Then*

$$|I(\lambda)| \lesssim_{n,K,\phi} \sum_{|\alpha| \leq n} \|(D^\alpha a) |\nabla \phi|^{|\alpha|-2n}\|_{L^\infty(\mathbf{R}^d)}.$$

where the implicit constant can be made independent of ϕ for a family of phases if the partial derivatives of ϕ up to order $n+1$ are uniformly bounded in a neighborhood of K .

Proof. Set $v = (\nabla \phi)/|\nabla \phi|^2$. Note that the phase ϕ is an eigenfunction of the differential operator D defined such that

$$Df(x) = \frac{v \cdot \nabla f}{2\pi i \lambda}.$$

The adjoint operator of D is the operator D^* defined by setting

$$D^*f(x) = \frac{\nabla \cdot (vf)}{-2\pi i \lambda},$$

i.e. for any smooth f and g , with one of these functions compactly supported,

$$\int Df(x)g(x) dx = \int f(x)(D^*g)(x) dx.$$

Thus

$$I(\lambda) = \int D^n(e^{2\pi i \lambda \phi(x)})a(x) dx = \int e^{2\pi i \lambda \phi(x)}((D^*)^n a)(x) dx.$$

Taking absolute values in the last integral gives that

$$|I(\lambda)| \leq \int |(D^*)^n a(x)| dx \lesssim_{\phi, a, n} \frac{1}{\lambda^n},$$

and a more careful analysis of this gives the more precise estimates in the lemma. \square

A tensorization argument establishes stationary phase asymptotics for a quadratic phase.

Theorem 27.13. *Let A be an invertible $d \times d$ matrix, fix $x_0 \in \mathbf{R}^d$, and consider the phase $\phi(x) = A(x - x_0) \cdot (x - x_0)$. Then for any compactly supported smooth amplitude a , there exists constants $\{c_n\}$ depending only on the derivatives of a at the origin, such that*

$$I(\lambda) \sim \lambda^{-d/2} \sum_{n=0}^{\infty} c_n \lambda^{-n}.$$

Moreover,

$$c_0 = a(x_0) \prod_{k=1}^d (i/\mu_k)^{1/2},$$

where μ_1, \dots, μ_d are the eigenvalues of A .

Proof. Suppose first that a is a tensor product of d compactly supported functions in \mathbf{R} . The constant c_0 is invariant under affine changes of coordinates. Thus we may assume that A is a diagonal matrix. But then the oscillatory integral splits into the product of single variable integrals, to which we can apply our one-dimensional asymptotics. A density argument then shows the argument generalizes to any smooth a . \square

Morse's theorem says that if x_0 is a non-degenerate critical point of a smooth function ϕ , then there exists a coordinate system around x_0 and $a_1, \dots, a_d \in \{\pm 1\}$ such that, in this coordinate system,

$$\phi(x_0 + t) = a_1 t_1^2 + \dots + a_d t_d^2.$$

In one dimension, the same is true if x_0 has a higher order critical point, but this does not generalize to higher dimensions, which reflects the lack of as nice a theory in this case. But in the case of functions with finitely many non-degenerate critical points, we can obtain nice asymptotics. Applying Morse's theorem gives the following theorem.

Theorem 27.14. *Let ϕ and a be smooth functions, with a compactly supported. Suppose ϕ has a single critical point x_0 on the support of a , which is nondegenerate. Then there exists constants $\{c_n\}$ depending only on finitely many derivatives of Φ and ψ at x_0 , such that*

$$I(\lambda) \sim e^{2\pi i \lambda \phi(x_0)} \lambda^{-d/2} \sum_{n=0}^{\infty} c_n \lambda^{-n}.$$

Moreover,

$$c_0 = a(x_0) \cdot \prod_{k=1}^d (i/\mu_k)^{1/2},$$

where μ_1, \dots, μ_d are the Eigenvalues of the Hessian of ϕ at x_0 .

27.3 Variable Coefficient Results

We can also obtain results given amplitude functions that depend on λ , and also vary in a third variable, under suitable conditions. Consider a symbol $a(x, y, \lambda)$, i.e. a smooth function such that $\text{supp}_y(a)$ is compact, and

$$\left| \nabla_x^n \nabla_y^m \nabla_\lambda^k a(x, y, \lambda) \right| \lesssim_{n,m,k} \lambda^{\alpha + \delta k},$$

for some $\alpha \in \mathbf{R}$ and $\delta > 0$, where the implicit constant is *locally uniform* in x . We can then consider the oscillatory integrals

$$I(x, \lambda) = \int a(x, y, \lambda) e^{2\pi i \lambda \phi(x, y)} dy.$$

Provided that $\delta < 1$, and $\nabla_y \phi \neq 0$ on the support of a , integration by parts gives that for any n and N ,

$$\nabla_x^n \nabla_\lambda^m I(x, \lambda) \lesssim_{N, n, m} \lambda^{-N},$$

where the implicit constant is locally uniform in x .

On the other hand, suppose that we have a nondegenerate critical point, i.e. there exists (x_0, y_0) with $\nabla_y \phi(x_0, y_0) = 0$, but with $H_y \phi(x_0, y_0)$ invertible. Then by the implicit function theorem, there exists a unique solution $(x, y(x))$ to $\nabla_y \phi(x, y)$ in a small neighborhood of y_0 , for x in a small neighborhood of x_0 . Intuitively, if a is supported on the small neighborhood of x_0 and y_0 , we should have

$$\begin{aligned} I(x, \lambda) &= \int a(x, y, \lambda) e^{2\pi i \lambda \phi(x, y)} dy \\ &\approx \lambda^{-d/2} e^{2\pi i \lambda \phi(x, y(x))} a(x, y(x), \lambda) \prod_{k=1}^d (i/\mu_k)^{1/2}, \end{aligned}$$

where μ_1, \dots, μ_d are the eigenvalues of $H_y \phi(x, y(x))$. Ignoring the $e^{2\pi i \lambda \phi(x, y(x))}$ factor (which introduces powers of λ), differentiating by x should introduce negligible effects on the value of the oscillatory integral. In fact, it is not difficult to prove that if

$$\left| \nabla_x^n \nabla_y^m \nabla_\lambda^k a(x, y, \lambda) \right| \lesssim_{n, m, k} \lambda^{-k},$$

then, under the support assumptions on a ,

$$\nabla_x^n \nabla_\lambda^m \left\{ e^{-2\pi i \lambda \phi(x, y(x))} I(x, \lambda) \right\} \lesssim_{N, n, m} \lambda^{-1/2-m}.$$

The result is a simple calculation, left as an exercise. A discussion can be found in Sogge's book 'Fourier Integrals in Classical Analysis'. On the other hand, if we have bounds of the form

$$\left| \nabla_x^n \nabla_y^m \nabla_\lambda^k a(x, y, \lambda) \right| \lesssim_{n, m, k} \lambda^{\alpha + \delta k},$$

for $\delta < 1/2$, then we can still obtain an asymptotic expansion of the form

$$I(x, \lambda) \sim e^{2\pi i \lambda \phi(x, y(x))} \lambda^{-d/2} \sum_{n=0}^{\infty} c_n(x, \lambda) \lambda^{-n},$$

locally uniform in x , where the coefficients c_n are symbols of order TODO , equal to a linear differential operator applied to a at $(x, y(x), \lambda)$, with coefficients depending solely on the behaviour of the Hessian matrix of ϕ in y , at $(x, y(x))$. In particular,

$$c_0 = a(x, y(x), \lambda) \prod_{k=1}^d (i/\mu_k(x))^{1/2},$$

where $\mu_1(x), \dots, \mu_d(x)$ are the eigenvalues of the Hessian of ϕ at $(x, y(x))$. Duistermaat's book on Fourier Integral Operators includes an explicit description of these differential operators.

27.4 Surface Carried Measures

Let us consider oscillatory integrals on a 'curved' version of Euclidean space. One most basic example is the Fourier transform of the surface measure of the sphere, i.e.

$$\hat{\sigma}(\xi) = \int_{S^{d-1}} e^{-2\pi i \xi x} d\sigma(x).$$

Studying the decay of this surface measure is of much interest to many problems in analysis. One can reduce the study of this Fourier transform to the study of Bessel functions, to which we have already developed an asymptotic theory.

Theorem 27.15. *If σ is the surface measure on the sphere S^{d-1} , then*

$$\hat{\sigma}(\xi) = \frac{2\pi \cdot J_{d/2-1}(2\pi|\xi|)}{|\xi|^{d/2-1}}.$$

In particular,

$$\hat{\sigma}(\xi) = \frac{2 \cos(2\pi|\xi| - (d/2 - 1)(\pi/2) - \pi/4)}{|\xi|^{(d-1)/2}} + O_d(1/|\xi|^{(d+1)/2}).$$

Proof. Since σ is rotationally symmetric, so too is $\hat{\sigma}$. In particular, we can apply Fubini's theorem to conclude that if V_{d-2} is the surface area of the unit sphere in \mathbf{R}^{d-2} , then

$$\begin{aligned}\hat{\sigma}(\xi) &= \int_{S^{d-1}} e^{-2\pi|\xi||x_1|} d\sigma(x) \\ &= V_{d-2} \int_{-1}^1 e^{-2\pi|\xi|t} (1-t^2)^{d/2-1} dt.\end{aligned}$$

Setting $r = 2\pi|\xi|$ completes the argument. \square

Since the multivariate stationary phase approach is essentially ‘coordinate independent’, we can also generalize the approach to manifolds. If M is a d dimensional Riemannian manifold, and ϕ and a are complex-valued functions on the manifold, we can consider the oscillatory integral

$$I(\lambda) = \int_M a(x) e^{2\pi i \lambda \phi(x)} d\sigma(x),$$

where σ is the surface measure induced by the metric on M . If ϕ and a are compactly supported, then this integral is well defined in the Lebesgue sense.

Theorem 27.16. *Suppose that a is a compactly supported smooth amplitude on a Riemannian manifold M , ϕ is a smooth phase, and $\nabla\phi$ vanishes at a single critical point x_0 on the support of ψ , upon each of which the Hessian $H\phi$ is non-degenerate at each point. Then there exists constants $\{c_n\}$ such that*

$$I(\lambda) \sim e^{2\pi i \lambda \phi(x_0)} \lambda^{-d/2} \sum_{n=0}^{\infty} c_n \lambda^{-n} + O(1/\lambda^{N+d/2+1}).$$

Moreover,

$$c_0 = a(x_0) \prod_{k=1}^d (i/\mu_k)^{1/2},$$

where μ_1, \dots, μ_d are the eigenvalues of the Hessian $H\phi$ at x_0 .

The theorem is proved by a simple partition of unity approach which reduces to the Euclidean case. It has the following important corollary.

Theorem 27.17. *If a d dimensional surface Σ is a smooth submanifold of \mathbf{R}^{d+1} , and has non-vanishing Gauss curvature, and if a is a smooth, compactly supported function on Σ , then*

$$|\widehat{a\sigma}(\xi)| \lesssim_{a,\sigma} \frac{1}{|\xi|^{d/2}},$$

where σ is the surface measure of Σ .

Proof. For each $\xi \in S^d$,

$$\widehat{a\sigma}(\xi) = \int_M a(x) e^{-2\pi i \xi \cdot x} d\sigma(x)$$

If $\lambda = |\xi|$, for $|\xi| = 1$, we can write

$$\widehat{a\sigma}(\lambda\xi) = \int_M a(x) e^{2\pi i \lambda \phi(\xi, x)} d\sigma(x),$$

where $\phi(\xi, x) = -\xi \cdot x$. Then the gradient of ϕ on the surface M vanishes at a point (ξ, x) precisely when ξ is normal to Σ . There are precisely two different smooth unit normal vector fields n_1, n_2 defined in a neighborhood of any given point x_0 .

We can find two smooth vector fields n_1 and n_2

$$I_\xi(\lambda) = \int_M a(x) e^{2\pi i \lambda \phi(\xi, x)} d\sigma,$$

where $\phi(\xi, x) = -2\pi i \xi \cdot x$. The derivatives of ϕ_ξ of order $\leq N$ on M are $O_N(1)$, independently of ξ . Similarly, $H_M \phi_\xi$ is uniformly non-degenerate, in the sense that the operator norm of $(H_M \phi_\xi)^{-1}(x)$ is $O(1)$, independently of ξ . Working with Σ as a local graph, and then applying the curvature condition on Σ implies that for each $\xi \in S^d$, ϕ_ξ has $O(1)$ stationary points on the support of ψ . There also exists a constant r such that if x does not lie in any ball of radius r around a stationary point, then $|\nabla_M \phi_\xi| \gtrsim 1$. Moreover, the Hessian $H_M \phi_\xi$ is uniformly non-degenerate in the radius r balls around the critical point, independently of ξ . Thus we can apply the last result to conclude

$$I_\xi(\lambda) \lesssim \lambda^{-d/2},$$

where the implicit constant is independent of ξ , because all the required derivatives are uniformly bounded. \square

If Ω is a bounded open subset of \mathbf{R}^{d+1} whose boundary is a smooth manifold with non-zero Gaussian curvature at each point, then its Fourier transform has decay one order better than the Fourier transform of its boundary.

Corollary 27.18. *If Ω is a bounded open subset of \mathbf{R}^d whose boundary is a smooth manifold Σ with non-zero Gaussian curvature at each point. If I_Ω is the indicator function on Ω , then*

$$|\widehat{I_\Omega}(\xi)| \lesssim_\Omega |\xi|^{-(d+1)/2}.$$

Proof. We have

$$\widehat{I_\Omega}(\xi) = \int_{\Omega} e^{-2\pi i \xi \cdot x} dx$$

Then we can apply Stoke's theorem for each $1 \leq k \leq d+1$ to conclude

$$\int_{\Omega} e^{-2\pi i \xi \cdot x} dx = \frac{(-1)^k}{2\pi i \xi_k} \int_{\Sigma} e^{-2\pi i \xi \cdot x} (dx^1 \wedge \cdots \wedge \widehat{dx^k} \wedge \cdots \wedge dx^n).$$

For each k , there is a smooth function ψ_k such that

$$dx^1 \wedge \cdots \wedge \widehat{dx^k} \wedge \cdots \wedge dx^n = \psi_k d\sigma.$$

Thus applying the last case, we find

$$\left| \frac{(-1)^k}{2\pi i \xi_k} \int_{\Sigma} e^{-2\pi i \xi \cdot x} (dx^1 \wedge \cdots \wedge \widehat{dx^k} \wedge \cdots \wedge dx^n) \right| \lesssim \xi_k^{-1} |\xi|^{-(d-1)/2}.$$

At each point ξ , if we choose ξ_k with the largest value, then $|\xi_k| \sim |\xi|$, so

$$\left| \int_{\Omega} e^{-2\pi i \xi \cdot x} dx \right| \lesssim |\xi|^{-(d+1)/2}. \quad \square$$

The fact that curved surfaces have Fourier decay is of supreme importance to much of modern harmonic analysis. Let us see some basic consequences, which is a classical result due to Hardy, Littlewood, and Hlawka.

Theorem 27.19. *Let B be the unit ball in \mathbf{R}^d , and for each $\lambda > 0$, let $N(\lambda)$ denote the number of integer lattice points in λB . Then*

$$N(\lambda) = |B| \cdot \lambda^d + O(\lambda^{n-2+2/(n+1)}).$$

Proof. We have

$$N(\lambda) = \widehat{f}(0),$$

where $f = \mathbf{I}_{\lambda B} \cdot \sum_{n \in \mathbf{Z}^d} \delta_n$. The Poisson summation formula implies that

$$\widehat{f} = \widehat{\mathbf{I}_{\lambda B}} * \left(\sum_{n \in \mathbf{Z}^d} \delta_n \right)$$

Thus

$$\widehat{f}(0) = \sum_{n \in \mathbf{Z}^d} \widehat{\mathbf{I}_{\lambda B}}(n) = \lambda^d \sum_{n \in \mathbf{Z}^d} \widehat{\mathbf{I}_B}(\lambda n) = |B| \lambda^d + \sum_{n \neq 0} \widehat{\mathbf{I}_B}(\lambda n).$$

We want to apply the estimate $\widehat{\mathbf{I}_B}(\lambda n) \lesssim (\lambda n)^{-(d+1)/2}$, but the decay here isn't enough for the sum in the Poisson summation formula to converge. To fix this, we mollify \mathbf{I}_B slightly. Let β be a non-negative bump function supported on $|x| \leq 1/2$ with $\int \beta(x) dx = 1$. Fix $\varepsilon > 0$, and define

$$\chi_\lambda(x) = \varepsilon^{-d} \text{Dil}_\varepsilon \beta * \mathbf{I}_{\lambda B}.$$

Then set $\tilde{N}(\lambda) = \sum_j \chi_\lambda(j)$. Then $\mathbf{I}_{\lambda B}$ is equal to χ_λ everywhere except on $\lambda - \varepsilon \leq |x| \leq \lambda + \varepsilon$, and since both functions are positive,

$$\tilde{N}(\lambda - \varepsilon) \leq N(\lambda) \leq \tilde{N}(\lambda + \varepsilon).$$

We can apply the Poisson summation formula now as above to obtain that

$$\tilde{N}(\lambda) = \lambda^d |B| + \sum_{n \neq 0} \widehat{\mathbf{I}_B}(\lambda n) \widehat{\beta}(\varepsilon n).$$

The first term is bounded by $(\lambda n)^{-(d+1)/2}$, and the second by $(\varepsilon n)^{-K}$, where $n \geq 1/\varepsilon$ and K is large, and bounded by $O(1)$ for $n \leq 1/\varepsilon$. Applying these bounds thus gives

$$\tilde{N}(\lambda) = \lambda^d |B| + O(\lambda^{-(d+1)/2} \varepsilon^{-(d-1)/2}).$$

Thus

$$N(\lambda) = \lambda^d |B| + O(\varepsilon \lambda^{d-1} + \lambda^{(d-1)/2} \varepsilon^{-(d-1)/2})$$

Choosing $\varepsilon = \lambda^{-(d-1)/(d+1)}$ completes the proof. \square

Remark. One can never have $N(\lambda) = |B|\lambda^d + O(\lambda^{d-2})$, because TODO

Example. If M is a hypersurface in \mathbf{R}^d , and ψ is a smooth, compactly supported function on M , and f is a smooth, compactly supported function on \mathbf{R}^d , we can define a function Af on \mathbf{R}^d by defining

$$(Af)(y) = \int_M f(y-x)\psi(x) d\sigma(x).$$

We note that Af is really the convolution of f with $\psi\sigma$. Thus

$$\widehat{Af}(\xi) = \widehat{f}(\xi)\widehat{\psi\sigma}(\xi).$$

For each multi-index α , the derivative $(Af)_\alpha$ is equal to

$$\int_M f_\alpha(y-x)\psi(x) d\sigma(x) = f_\alpha * (\psi\sigma).$$

In particular, we have

$$\widehat{(Af)_\alpha} = (2\pi i\xi)^\alpha \widehat{f}(\xi)\widehat{\psi\sigma}(\xi).$$

Since we have shown

$$|\widehat{\psi\sigma}(\xi)| \lesssim |\xi|^{-(d-1)/2},$$

we conclude that if $|\alpha| \leq k$, where $k = (d-1)/2$,

$$\|(Af)_\alpha\|_{L^2(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^d)}.$$

In particular, this implies that A extends to a unique bounded operator from $L^2(\mathbf{R}^d)$ to $L^2_k(\mathbf{R}^d)$, i.e. to a map such that for each $f \in L^2(\mathbf{R}^d)$, Af is a square integrable function which has square integrable weak derivatives of all orders less than or equal to k , and moreover, $\|(Af)_\alpha\|_{L^2(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^d)}$ for all $|\alpha| \leq k$. Thus the operator A is ‘smoothing’, in a certain sense.

The operator A is obviously bounded from $L^1(\mathbf{R}^d)$ to $L^1(\mathbf{R}^d)$ and $L^\infty(\mathbf{R}^d)$ to $L^\infty(\mathbf{R}^d)$, purely from the fact that $\psi\sigma$ is a finite measure. Using curvature and some analytic interpolation, we will now also show that A is bounded from $L^p(\mathbf{R}^d)$ to $L^q(\mathbf{R}^d)$, where $p = (d+1)/d$, and $q = d+1$. Interpolation thus yields a number of intermediate estimates. The trick here is to obtain an (L^1, L^∞) bound for an ‘improved’ version of A , and an (L^2, L^2) bound for a ‘worsened’ version of A . Interpolating between these two results gives a bound

for precisely A . It suffices to prove this bound ‘locally’ on M , since we can then sum up these bounds, so we may assume that M is given as the graph of some function, i.e. there exists u such that

$$M = \{(x, u(x)) : \}$$

For each s , we write $A_s f = K_s * f$, where

$$K_s(x) = \gamma_s |x_d - \phi(x')|_+^{s-1} \psi_0(x).$$

Here $\gamma_s = s \dots (s + N) e^{s^2}$, where N is some large parameter to be fixed in a moment. The e^{s^2} parameter is to mitigate the growth of γ_s as $|Im(s)| \rightarrow \infty$, which allows us to interpolate. The quantity $|u|_+^{s-1}$ is equal to u^{s-1} where $u > 0$, and is equal to 0 when $u \leq 0$. And $\psi_0(x) = \psi(x)(1 + |\nabla_{x'} \phi(x')|^2)^{1/2}$.

27.5 Oscillatory Integral Operators

An oscillatory integral takes as input a particular frequency scale, and gives back a scalar value. The goal of the theory of stationary phase is to determine the decay of the outputs as we increase the value of the frequency scale. Oscillatory integral operators are a generalization of this property. Most often, these oscillatory operators are a family of operators $\{T_\lambda\}$, where λ parameterizes the frequency scale of some oscillatory factor. The goal is to understand how the properties of these operators is affected asymptotically as we increase the frequency factor.

Let us begin with a family of such operators given by the formula

$$T_\lambda f(x) = \int_{\mathbf{R}^n} a(x, y) e^{2\pi i \lambda \phi(x, y)} f(y) dy,$$

where $a \in C_c^\infty(\mathbf{R}^m \times \mathbf{R}^n)$ is real-valued and $\phi \in C^\infty(\mathbf{R}^m \times \mathbf{R}^n)$. The regularity of all these parameters means that the operators T_λ are defined even for a general distributional input f . For intuition, we recall the theory of oscillatory integral distributions, and in particular, canonical relations and wave front sets. The kernel $K_\lambda(x, y) = a(x, y) e^{2\pi i \lambda \phi(x, y)}$ is an ‘oscillatory integral distribution’ of ‘dimension zero’. Applying the theory of wave front sets for such distributions blindly, we obtain the formula

$$\text{WF}(K_\lambda) \subset \{(x, y; \lambda \nabla_x \phi(x, y), \lambda \nabla_y \phi(x, y)) : x \in \mathbf{R}^m, y \in \mathbf{R}^n\}.$$

Since K_λ is a smooth function, the wave front set of K_λ is actually empty, so this formula is pretty redundant. Nonetheless, this formula does give us some information about where the operator T_λ may be singular in a quantitative sense. Consider the *canonical relation*

$$C_\phi = \{(x, y; \nabla_x \phi(x, y), -\nabla_y \phi(x, y)) : x \in \mathbf{R}^m, y \in \mathbf{R}^n\}.$$

A rudimentary Taylor expansion calculation shows that if $f(y) = \psi((y - y_0)/\delta_y)e^{2\pi i \eta_0 \cdot y}$ for some y_0 and ξ_0 , then for any particular x_0 , if $|x - x_0| \leq \delta_x$,

$$T_\lambda f_\delta(x) \approx \delta_y^n a(x_0, y_0) e^{2\pi i \lambda (\phi(x_0, y_0) + \nabla_x \phi(x_0, y_0) \cdot (x - x_0))} \hat{\psi}(-\delta_y [\lambda \nabla_y \phi(x_0, y_0) + \eta_0]),$$

where the left and right differ by $\delta_y^n \cdot O(\delta_x + \lambda \delta_x^2 + \lambda \delta_y^2)$. In particular, this approximation is good provided that $\delta_x \lesssim \lambda^{-1/2}$ and for $\delta_y \lesssim \lambda^{-1/2}$. We intuit from this that if f is a wave packet with unit amplitude, localized in space near y_0 with an uncertainty of $\lambda^{-1/2}$, and localized near a frequency η_0 , then, when localized in space near x_0 , with an uncertainty of $\lambda^{-1/2}$, $T_\lambda f$ looks like a wave packet with amplitude

$$O_N(\lambda^{-n/2}(1 + \lambda^{1/2}|\lambda \nabla_y \phi(x_0, y_0) + \eta_0|)^{-N})$$

for all $N > 0$, and with frequency localized near $\lambda \nabla_x \phi(x_0, y_0)$. In particular, if $\eta_0 = -\lambda \nabla_y \phi(x_0, y_0)$, then we obtain a wave packet in the output with an amplitude of $\approx \lambda^{-n/2}$ when localized in a $\Theta(\lambda^{-1/2})$ radius neighbourhood of x_0 . And if η_0 is chosen away from this value, the amplitude becomes very small as $\lambda \rightarrow \infty$. This indicates the canonical relation determines which wave packets induce a noticeable effect on the output, and where this effect occurs. In particular, if we take f with the former properties, then $\|T_\lambda f\|_{L^q(\mathbf{R}^m)} \gtrsim \lambda^{-n/2} \lambda^{-m/2q}$, whereas $\|f\|_{L^p(\mathbf{R}^n)} \sim \lambda^{-n/2p}$, which implies that the operator norm of T_λ from $L^p(\mathbf{R}^n)$ to $L^q(\mathbf{R}^m)$ is at least $\Omega(\lambda^{(n/2)(1/p-1)-(m/2)(1/q)})$. In particular, we cannot expect to do any better than the trivial bound $\|T_\lambda f\|_{L^\infty(\mathbf{R}^m)} \lesssim \|f\|_{L^1(\mathbf{R}^n)}$ when $p = 1$ and $q = \infty$. In certain situations, we will be able to conclude that T_λ actually does have an operator norm achieving this decay bound, provided we assume suitable geometric conditions on the canonical relation C_ϕ .

To begin with, let us consider the case where the matrices $D_y \nabla_x \phi = (D_x \nabla_y \phi)^T$ always has full rank on the support of the function a . If $m \geq n$, this information can be expressed by saying that $C_\phi \rightarrow T^*\mathbf{R}^n$ is a submersion, and $C_\phi \rightarrow T^*\mathbf{R}^m$ is an immersion, with these properties swapped if

$m \leq n$. Thus, locally speaking, we have a smooth map $T^*\mathbf{R}^m \rightarrow C_\phi$ which is a left inverse to the corresponding projection map. For intuitions sake, let us assume this smooth map is globally defined, so we can obtain a smooth map $\alpha = (y, \eta) : T^*\mathbf{R}^m \rightarrow T^*\mathbf{R}^n$ such that if $(x_0, y_0; \xi_0, \eta_0) \in C_\phi$, then $y_0 = y(x_0, \xi_0)$ and $\eta_0 = \eta(x_0, \xi_0)$. Thus, roughly speaking, the only possible wave packet f localized spatially at a scale $\lambda^{-1/2}$ whose corresponding output $T_\lambda f$ has significant amplitude when localized at a scale $\lambda^{-1/2}$ near x_0 and oscillates at a frequency ξ_0 is a wave packet at position $y(x_0, \xi_0)$ and with frequency $\eta(x_0, \xi_0)$. The fact that $|\alpha(x_0, \xi_0) - \alpha(x_1, \xi_1)| \gtrsim |(x_0, \xi_0) - (x_1, \xi_1)|$ reflects a kind of ‘orthogonality property’ of the operator T_λ . To understand this orthogonality, we rely on L^2 techniques, namely a T^*T argument.

Theorem 27.20. *Consider a family of oscillatory integral operators*

$$\{T_\lambda : L^2(\mathbf{R}^n) \rightarrow L^2(\mathbf{R}^m)\}$$

with associated phase ϕ . If $D_y \nabla_x \phi$ is injective, i.e. the map $C_\phi \rightarrow T^\mathbf{R}^m$ is an immersion, then*

$$\|T_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{-n/2} \|f\|_{L^2(\mathbf{R}^n)}.$$

Proof. Once sufficiently localized, the assumption above implies that

$$|\nabla_x \phi(x, y_1) - \nabla_x \phi(x, y_2)| \gtrsim |y_1 - y_2|.$$

We calculate that

$$T_\lambda^* g(y) = \int \overline{a(x, y)} e^{-2\pi i \lambda \phi(x, y)} dx.$$

Thus the kernel of the operator $T_\lambda^* T_\lambda$ is equal to

$$K(y_1, y_2) = \int \overline{a(x, y_1)} a(x, y_2) e^{2\pi i \lambda [\phi(x, y_1) - \phi(x, y_2)]} dx.$$

The principle of nonstationary phase thus tells us that

$$|K(y_1, y_2)| \lesssim (1 + \lambda |y_1 - y_2|)^{-N}.$$

Thus for $|K(y_1, y_2)| \lesssim 1$ for $|y_1 - y_2| \lesssim 1/\lambda$, and $|K(y_1, y_2)| \lesssim_N \lambda^{-N} |y_1 - y_2|^{-N}$ for $|y_1 - y_2| \gtrsim 1/\lambda$. Schur’s test thus gives that $\|T_\lambda^* T_\lambda f\|_{L^1(\mathbf{R}^n)} \lesssim \lambda^{-n} \|f\|_{L^1(\mathbf{R}^n)}$, and that $\|T_\lambda^* T_\lambda f\|_{L^\infty(\mathbf{R}^n)} \lesssim \lambda^{-n} \|f\|_{L^\infty(\mathbf{R}^n)}$, and interpolation that $\|T_\lambda^* T_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{-n} \|f\|_{L^2(\mathbf{R}^n)}$, and so $\|T_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{-n/2} \|f\|_{L^2(\mathbf{R}^n)}$. \square

Remark. The best possible decay bound for the operator norm of T_λ for general n and m is $\lambda^{-(n+m)/4}$, and since the theorem above can only be applied when $n \leq m$, the result is only optimal when $n = m$.

Interpolating between the trivial bound $\|T_\lambda f\|_{L^\infty(\mathbf{R}^n)} \lesssim \|f\|_{L^1(\mathbf{R}^n)}$ yields that for $2 \leq q \leq \infty$, if p is the conjugate dual of q , then

$$\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}.$$

More generally, using the fact that $T_\lambda f$ only depends on the behaviour of f on $\text{supp}_y(a)$, we conclude that we have a bound $\|T_\lambda f\|_{L^\infty(\mathbf{R}^n)} \lesssim \|f\|_{L^p(\mathbf{R}^n)}$ for any $1 \leq p \leq \infty$. Interpolating this bound with $\|T_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{-n/2} \|f\|_{L^2(\mathbf{R}^n)}$ then gives $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}$ whenever $1/p + 1/q \leq 1$ and $1 \leq p \leq 2$ (equivalently, $2 \leq q \leq \infty$).

One might see an analogy with this family of inequalities to the Hausdorff-Young inequality $\|\hat{f}\|_{L^q(\mathbf{R}^n)} \lesssim \|f\|_{L^p(\mathbf{R}^n)}$, which holds when $1/p + 1/q = 1$. Indeed, the Hausdorff-Young inequality actually follows from these bounds, since one can consider the truncated Fourier transforms

$$F_\lambda f(\xi) = \int a(x/\lambda, \xi/\lambda) e^{2\pi i \xi \cdot x} f(x) dx = \lambda^n \int a(x, \xi) e^{2\pi i \lambda \xi \cdot x} f(\lambda x) dx$$

for some bump function $a \in C_c^\infty(\mathbf{R})$, which equal the normal Fourier transform in the limit as $\lambda \rightarrow \infty$ for suitably regular functions f . The operators $T_\lambda = \lambda^{-n} F_\lambda \circ \text{Dil}_\lambda$ are given by

$$T_\lambda f(\xi) = \int a(x, \xi) e^{2\pi i \lambda \xi \cdot x} f(x) dx,$$

and $D_\xi \nabla_x(\xi \cdot x)$ is the identity matrix, so the theory above implies that $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}$. Rescaling this inequality shows that $\|F_\lambda f\|_{L^q} \lesssim \|f\|_{L^p(\mathbf{R}^n)}$, and taking limits as $\lambda \rightarrow \infty$ then yields Hausdorff Young. Thus we see obtaining results for oscillatory integral operators without compact support can be obtained from this theory if we can obtain oscillatory integral bounds with a decay of the form $\lambda^{-n/q}$.

One phase to which the last result does not imply is that phase $\phi(x, y) = |x - y|$, since, for $x \neq y$,

$$\nabla_x \phi = \frac{x - y}{|x - y|} \quad \text{and} \quad \nabla_y \phi = \frac{y - x}{|y - x|}.$$

This implies that $D_y \nabla_x \phi$ cannot possibly be invertible, since the map $y \mapsto \nabla_x \phi(x, y)$ is not an open map (it's image lies in S^{n-1}). The canonical relation C_ϕ is a cone of points of the form

$$\left\{ \left(x, y; \frac{x-y}{|x-y|}, \frac{x-y}{|x-y|} \right) \right\},$$

and so each pair (x_0, ξ_0) is related to a one dimensional family of points (y, ξ_0) , where y lies on the half-line of points through x_0 pointing in the direction $-\xi_0$. Similarly, each pair (y_0, η_0) is related to a one dimensional family of points (x, ξ_0) , where x lies on a half-line of points pointing in the direction ξ_0 . In particular, this means that obtaining a bound of the form $\|T_\lambda f\|_{L^2(\mathbf{R}^n)} \lesssim \lambda^{-n/2} \|f\|_{L^2(\mathbf{R}^n)}$ is impossible for such a phase. More precisely, fix $x_0 \in \text{supp}_x(a)$ and a half-line $l = \{x \in \mathbf{R}^n : (x-x_0)/|x-x_0| = \xi_0\}$ starting at x_0 for some $|\xi_0| = 1$ whose intersection with $\text{supp}_y(a)$ has non-empty interior in l . Fix $N \sim \lambda^{1/2}$, and consider a family of N wave packets f_1, \dots, f_N , each with pairwise disjoint support, but supported spatially near a family of $O(\lambda^{-1/2})$ separated points lying on $l \cap \text{supp}_y(a)$, with spatial uncertainty $\lambda^{-1/2}$, and with frequency concentrated near ξ_0 . If $f = f_1 + \dots + f_N$, then the formula above implies that $T_\lambda f$ will have the majority of it's spatial support in the set

$$\{x_0 \in \mathbf{R}^n : |\lambda(x_0 - y_0)/|x_0 - y_0| - \xi_0| \lesssim \lambda^{-1/2}\},$$

and have amplitude $\lambda^{-(n-1)/2}$ over there. Roughly speaking, this is a tube centered at y_0 , pointing in the direction ξ_0 , and with thickness $O(\lambda^{-1/2})$, and thus has volume $\Theta(\lambda^{-(n-1)/2})$. But this means that $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \gtrsim \lambda^{-(n-1)/2} \lambda^{-(n-1)/2q}$, and so combined with the fact that $\|T_\lambda f\|_{L^p(\mathbf{R}^n)} \sim \lambda^{-(n-1)/2p}$, this means the operator norm of T_λ must be $\Omega(\lambda^{(n-1)/2(1/p-1/q-1)})$, which for $p = q = 2$ gives $\Omega(\lambda^{-(n-1)/2})$. On the other hand, this does not preclude a bound of the form $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}$ from holding when $q \geq (n+1)/(n-1) \cdot p^*$, which forms a subset of the family of all of the cases we proved above when, in addition, $1 \leq p \leq 2$. We will show, using the fact that the projections of C_ϕ onto $T^*\mathbf{R}^n$ are surfaces of constant curvature, that we can obtain the bound under this assumption when $1 \leq p \leq 2$, and even when $1 \leq p \leq 4$ when $n = 2$.

To analyze the operator, we consider a cutoff ψ with $\psi(z) = 0$ for $|z| \lesssim 1$,

and with $\psi(z) = 1$ for $|z| \gtrsim 1$. Write $T_\lambda = \tilde{T}_\lambda + R_\lambda$, where

$$\tilde{T}_\lambda f(x) = \int a(x, y) \psi(x - y) e^{2\pi i \lambda |x - y|} f(y) dy.$$

If we can establish that $\|\tilde{T}_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}$, then we know that

$$R_\lambda f(x) = \int a(x, y) (1 - \psi(x - y)) e^{2\pi i \lambda |x - y|}$$

TODO: COMPLETE ARGUMENT.

To study \tilde{T}_λ , we can cover the support of a by finitely many open sets $\{U_\alpha \times V_\alpha\}$, upon each of which we may find a diffeomorphism $y_\alpha : W_\alpha \times I_\alpha \rightarrow V_\alpha$, where $W_\alpha \subset \mathbf{R}^{n-1}$, I_α is an interval, and for any $x \in U_\alpha$ and $t \in I_\alpha$, the map

$$v \mapsto \frac{y_\alpha(v, t) - x}{|y_\alpha(v, t) - x|}$$

is an immersion from W_α to \mathbf{R}^n . Applying a partition of unity $\{\psi_\alpha\}$ over these open sets, we can write $\tilde{T}_\lambda = \sum_\alpha T_\lambda^\alpha$. Moreover, we can write

$$\begin{aligned} T_\lambda^\alpha &= \int a(x, y) \psi_\alpha(x, y) e^{2\pi i \lambda |x - y|} dy \\ &= \int \frac{a(x, y_\alpha(v, t)) \psi_\alpha(x, y_\alpha(v, t))}{|\det(Dy_\alpha(v, t))|} e^{2\pi i \lambda |x - y_\alpha(v, t)|} f(y_\alpha(v, t)) dv dt \\ &= \int_{I_\alpha} \int_{W_\alpha} a_\alpha(x, v; t) e^{2\pi i \lambda |x - y_\alpha(v, t)|} f(y_\alpha(v, t)) dv dt \\ &= \int_{I_\alpha} T_\lambda^{\alpha, t} f_{t, \alpha}(x), \end{aligned}$$

where $f_{t, \alpha}(v) = f(y_\alpha(v, t))$, and if we set $\phi(x, v; t) = |x - y_\alpha(v, t)|$, then

$$T_\lambda^{\alpha, t} f(x) = \int_{W_\alpha} a_\alpha(x, v; t) e^{2\pi i \lambda \phi(x, v; t)} f(v) dv.$$

The next Lemma will justify that $\|T_\lambda^{\alpha, t} f_{t, \alpha}\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f_{t, \alpha}\|_{L^p(W_\alpha)}$, from

which, together with Hölder's inequality, it follows that

$$\begin{aligned}
\|T_\lambda^\alpha f\|_{L^q(\mathbf{R}^n)} &\lesssim \int_{I_\alpha} \|T_\lambda^{\alpha,t} f_{t,\alpha}\|_{L^q(\mathbf{R}^n)} \\
&\lesssim \lambda^{-n/q} \int_{I_\alpha} \|f_{t,\alpha}\|_{L^p(W_\alpha)} dt \\
&\lesssim \lambda^{-n/q} |I_\alpha|^{1-1/p} \left(\int \|f_{t,\alpha}\|_{L^p(W_\alpha)}^p \right)^{1/p} \\
&\lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^n)}.
\end{aligned}$$

Let us now prove this lemma, we note that the canonical relation associated with the family of operators $\{T_\lambda^{\alpha,t}\}$ is

$$C^{\alpha,t} = \left\{ \left(x, v; \frac{|x - y_\alpha(v, t)|}{|x - y_\alpha(v, t)|}, D_v(y_\alpha)(v, t)^T \cdot \frac{x - y_\alpha(v, t)}{|x - y_\alpha(v, t)|} \right) \right\}.$$

Our assumptions on the coordinate system y_α implies that the projection map $C^{\alpha,t} \rightarrow T^*W_\alpha$ is an immersion, i.e. the matrix $D_v \nabla_x \phi_{t,\alpha}$ has full rank $n - 1$ on the domain of the integral. And the image of the projection $C^{\alpha,t} \rightarrow T^*\mathbf{R}^n$ is an open subset of the submanifold of unit vectors in $T^*\mathbf{R}^n$, which, at each point x , is a $n - 1$ dimensional manifold with non-vanishing curvature. These assumptions justify the application of the next Lemma.

Lemma 27.21. *Suppose $\phi : \mathbf{R}^n \times \mathbf{R}^{n-1}$ is a C^∞ phase, that $\nabla_x D_y \phi$ has rank $n - 1$ on the support of $a \in C_c^\infty(\mathbf{R}^n \times \mathbf{R}^{n-1})$, and that the image of the projection $C_\phi \rightarrow T^*\mathbf{R}^n$ is on each fibre a hypersurface of non-vanishing curvature. Then if $1/p + 1/q \leq 1$, $1 \leq p \leq 2$, and $q \geq [(n + 1)/(n - 1)] \cdot p^*$, then*

$$\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^p(\mathbf{R}^{n-1})}.$$

Remark. Using a similar construction as for the operator associated with the phase $\phi(x, y) = |x - y|$, one can show from the assumption that there exists f such that $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \geq \lambda^{(n-2)/2p - (n-2)/2 - n/2q} \|f\|_{L^p(\mathbf{R}^{n-1})}$, which shows that one can only obtain a bound of the form above when $q \geq [(n + 1)/(n - 1)] \cdot p^*$. We also note that the theorem above, once rescale, implies extension estimates of the form

$$\left\| \int_{\mathbf{R}^{n-1}} e^{2\pi i x \cdot \phi(\xi)} f(\xi) d\xi \right\|_{L^q(\mathbf{R}^n)} \lesssim \|f\|_{L^p(\mathbf{R}^{n-1})}$$

for the same range of p and q , which are dual to the corresponding restriction estimates for the hypersurface $H = \{(\xi, \phi(\xi))\}$.

Proof. To prove the result, we may assume that $q = [(n+1)/(n-1)] \cdot p^*$. Since the result is proved when $p = 1$, it suffices to show that

$$\|T_\lambda f\|_{L^{(n+1)/2(n-1)}(\mathbf{R}^n)} \lesssim \lambda^{-n(n-1)/2(n+1)} \|f\|_{L^2(\mathbf{R}^{n-1})}.$$

The assumption that the surface $y \mapsto \nabla_x \phi(x, y)$ has non-vanishing curvature can be expressed in the following manner: I can, locally at least, for each $x \in \mathbf{R}^n$ and $y \in \mathbf{R}^{n-1}$, pick a unit normal vector $\nu(x, y) \in S^{n-1}$ to the surface $y \mapsto \nabla_x \phi(x, y)$ at the point $\nabla_x \phi(x, y)$. This means precisely that for each fixed x_0 and y_0 ,

$$\nabla_y (\nabla_x \phi(x_0, y) \cdot \nu(x_0, y_0)) = 0$$

when $y = y_0$. If $H(x_0, y_0)$ is the Hessian matrix of the operator $y \mapsto \nabla_x \phi(x_0, y) \cdot \nu(x_0, y_0)$, then this means precisely that

$$\nabla_x \phi(x_0, y) \cdot \nu(x_0, y_0) = (y - y_0)^T H(x_0, y_0) (y - y_0) + O(|y - y_0|^3).$$

The statement that the surface does not have vanishing curvature at (x_0, y_0) is then equivalent to the fact that $H(x_0, y_0)$ is invertible.

To prove the theorem, perhaps after localizing and taking a change of x -variables, we may assume that $x = (z, t)$, where $D_z \nabla_y \phi$ is invertible. Introduce the operators T_λ^t , mapping functions from \mathbf{R}^{n-1} to \mathbf{R}^{n-1} by the formula

$$T_\lambda^t f(z) = \int a(z, y; t) e^{2\pi i \lambda \phi(z, y; t)} f(y) dy.$$

Then

$$\|T_\lambda f\|_{L^q(\mathbf{R}^d)} \sim \left(\int \|T_\lambda^t f\|_{L^q(\mathbf{R}^{n-1})}^q dt \right)^{1/q},$$

and it suffices to bound this integral. The advantage of rephrasing the problem in terms of the oscillatory integral operators T_λ^t is because we already have some tight estimates for such operators, namely

$$\|T_\lambda^t f\|_{L^q(\mathbf{R}^{n-1})} \lesssim \lambda^{-(n-1)/q} \|f\|_{L^p(\mathbf{R}^{n-1})}.$$

However, directly plugging this into the inequality only gives a bound $\|T_\lambda f\|_{L^q(\mathbf{R}^n)} \lesssim \lambda^{-n/q} \|f\|_{L^q(\mathbf{R}^{n-1})}$. To gain a boost on what is essentially a

trivial application of the triangle inequality, we need to know more about the interactions between the family of operators $\{T_\lambda^t\}$. This is where the curvature of the canonical relation comes into play, since it implies a kind of orthogonality of the operators.

To utilize this kind of orthogonality, we again apply adjoint techniques. There is $b \in C_c^\infty$ and $\Phi(z_1, z_2, y; t_1, t_2) = \phi(z_1, y, t_1) - \phi(z_2, y, t_2)$ such that the kernel of the operator $T_\lambda^{t_1} (T_\lambda^{t_2})^*$ is

$$K_{t_1, t_2}(z_1, z_2) = \int a(z_1, z_2, y; t_1, t_2) e^{2\pi i \lambda (\phi(z_1, y, t_1) - \phi(z_2, y, t_2))} dy.$$

Let's analyze this using stationary phase. We have $\nabla_y \Phi(z_0, z_0, y_0, t_0, t_0) = 0$ for any choice of z_0, y_0 , and t_0 , yet we have $D_{z_1} \nabla_y \Phi(z, z, y, t, t) = D_{z_1} \nabla_y \phi(z_1, y, t_1)$, which is an invertible matrix by assumption. Thus by the implicit function theorem, there exists a function $z_1 = z_1(z_2, y, t_1, t_2)$ defined in a neighborhood of (z_0, y_0, t_0, t_0) which is a unique solution to the equation $\nabla_y \Phi(z_1, z_2, y, t_1, t_2) = 0$. By localizing in these variables, we may assume these are the only such stationary points. A Taylor expansion gives

$$\begin{aligned} \nabla_y \Phi(z_1, z_2, y, t_1, t_2) &= D_z \nabla_y \phi(z_2, y, t_2) \cdot (z_1 - z_2) \\ &\quad + D_t \nabla_y \phi(z_2, y, t_2) \cdot (t_1 - t_2) \\ &\quad + O(|z_1 - z_2|^2 + |t_1 - t_2|^2). \end{aligned}$$

Thus

$$|D_z \nabla_y \phi(z_2, y, t_2) \cdot (z_1 - z_2) + D_t \nabla_y \phi(z_2, y, t_2) \cdot (t_1 - t_2)| \lesssim O(|z_1 - z_2|^2 + |t_1 - t_2|^2).$$

It follows from this that $(z_1 - z_2, t_1 - t_2)$ must be in a small, conical neighborhood of $\pm \nu(z_2, y, t_2)$ (For any matrix A , $|Ax| \gtrsim |x|$ unless x is close to the kernel of A). If we write $x_i = (z_i, t_i)$ again, this means that for the appropriate sign $|(x_1 - x_2) \pm |x_1 - x_2| \nu(x_2, y)| \lesssim |x_1 - x_2|$. But then

$$\begin{aligned} D_y \nabla_y \Phi(x_1, x_2, y) &= D_y \nabla_y \{D_x \phi(x_2, y) \cdot (x_1 - x_2)\} + O(|x_1 - x_2|^2) \\ &= |x_1 - x_2| D_y \nabla_y \{D_x \phi(x_2, y) \cdot \nu(x_2, y)\} + O(|x_1 - x_2|), \end{aligned}$$

If we choose all the localization values above carefully enough, our curvature assumption implies that this matrix is invertible with determinant $|x_1 - x_2|^{n-1}$, so we have nondegenerate stationary points. Thus isolating near these critical points, we obtain a decay in the oscillatory integral of

the form $\lesssim (1 + \lambda|x_1 - x_2|)^{-(n-1)/2}$. Away from these critical points, we have $|\nabla_y \Phi(x_1, x_2, y)| \gtrsim |x_1 - x_2|$ (since if $|\nabla_y \Phi(x_1, x_2, y)| \ll |x_1 - x_2|$, then $x_1 - x_2$ will be in one of the neighborhoods we considered above), and so when we integrate here, nonstationary phase implies that we have a decay of the form $O_N((1 + \lambda|x_1 - x_2|)^{-N})$ for all N . Putting together these estimates gives that

$$|K^{t_1, t_2}(z_1, z_2)| \lesssim (1 + \lambda|t_1 - t_2| + \lambda|z_1 - z_2|)^{-(n-1)/2}.$$

This implies the estimates

$$\|T_\lambda^{t_1}(T_\lambda^{t_2})^* f\|_{L^\infty(\mathbf{R}^{n-1})} \lesssim \lambda^{-(n-1)/2} |t_1 - t_2|^{-(n-1)/2} \|f\|_{L^1(\mathbf{R}^{n-1})}.$$

This looks like something we could apply the theory of fractional integrals to understand. And indeed we can, as the next result shows: TODO. \square

Chapter 28

Restriction Theorems

If a function f lies in $L^p(\mathbf{R}^d)$, there does not exist a numerically meaningful way to restrict f to a set of measure zero, since f is only defined up to measure zero. More precisely, for any $0 < p < \infty$, $0 < q \leq \infty$, and any Radon measure σ supported on a set S with Lebesgue measure zero, there does not exist a bound

$$\|f\|_{L^q(S, \mu)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in C(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$, and thus there does not exist a bounded operator $T : L^p(\mathbf{R}^d) \rightarrow L^q(S, \mu)$ which agrees with the standard restriction of continuous functions to sets of measure zero. We can consider a very similar problem for the Fourier transform; the Fourier transform of a function $f \in L^1(\mathbf{R}^d)$ is continuous, and thus it is meaningful to consider the restriction $\hat{f}|_S$. If there exists a bound of the form

$$\|\hat{f}\|_{L^q(S, \mu)} \lesssim \|f\|_{L^p(\mathbf{R}^d)} \quad (28.1)$$

for all $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$, then the Hahn-Banach theorem ensures that there exists a bounded operator $R : L^p(\mathbf{R}^d) \rightarrow L^q(S, \mu)$, such that $Rf = \hat{f}|_S$ for $f \in L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$; for $p < \infty$, R will be unique since $L^1(\mathbf{R}^d) \cap L^p(\mathbf{R}^d)$ will be dense in $L^\infty(\mathbf{R}^d)$. Thus we have a meaningful way of restricting the Fourier transforms of functions in $L^p(\mathbf{R}^d)$ to S . In light of the failure to restrict elements of $L^p(\mathbf{R}^d)$ to $L^q(S, \mu)$ without taking the Fourier transform, this indicates that the Fourier transform of $L^p(\mathbf{R}^d)$ is a fairly special element of $L^q(\mathbf{R}^d)$, in particular, taking values in a subclass of $L^q(\mathbf{R}^d)$ in

which it is meaningful to restrict to the set S . We refer to an estimate of the form above as a *restriction estimate*.

The approximate translation invariance of restriction bounds implies, from Littlewood's principle, that $p \leq q$ holds for any meaningful restriction estimate. For $p = 1$, a restriction bound from $L^1(\mathbf{R}^d)$ to $L^\infty(S, \mu)$ is trivial for any set S and measure μ . On the other hand, for $p = 2$ and $q > 0$, *no such bound is possible* unless μ is absolutely continuous with respect to the Lebesgue measure. This follows from a simple application of Parseval's theorem, i.e. because the Fourier transform of an element of $L^2(\mathbf{R}_x^d)$ in general behaves like an arbitrary element of $L^2(\mathbf{R}_\xi^d)$, and thus cannot be restricted to singular sets. Interpolation shows we also should not expect any bounds when $p > 2$, and indeed such restriction estimates even fail when μ is absolutely continuous with respect to the Lebesgue measure, for the Hausdorff-Young inequality fails in this setting. Thus the interesting bounds occur when $1 < p < 2$. For a particular pair (S, μ) and exponent q , the goal is to push the value of p as close as possible to $\max(2, q)$.

There is a dual problem associated with the pair (S, μ) . For $g \in L^1(S, \mu)$, we consider the *extension operator*

$$E_S g(x) = \int_S g(\xi) e^{2\pi i \xi \cdot x} d\mu(\xi).$$

In other words, $E_S g$ is the inverse Fourier transform of the finite Borel measure $g \cdot \mu$. Analogous to the restriction operator, it is simple to verify the estimate

$$\|E_S g\|_{L^\infty(\mathbf{R}^d)} \leq \|g\|_{L^1(S, \mu)},$$

and that unless μ is absolutely continuous with respect to the Lebesgue measure, no bound of the form

$$\|E_S g\|_{L^2(\mathbf{R}^d)} \lesssim \|g\|_{L^p(S, \mu)}$$

exists. The restriction and extension operators are connected by adjointness for all cases of interest, i.e. using Fubini's theorem that if $f \in L^1(\mathbf{R}^d)$ and $g \in L^1(S, \mu)$, then

$$\int_S (R_S f)(\xi) \overline{g(\xi)} d\mu(\xi) = \int_{\mathbf{R}^d} f(x) \overline{(E_S g)(x)} dx.$$

A density argument allows us to conclude that for $1 \leq p < \infty$ and $1 < q \leq \infty$, there exists a bound of the form

$$\|R_S f\|_{L^q(S, \mu)} \leq C \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in L^p(\mathbf{R}^d)$ if and only if there exists a bound of the form

$$\|E_S f\|_{L^{p^*}(\mathbf{R}^d)} \leq C \|f\|_{L^{q^*}(\mathbf{R}^d)}.$$

But this covers all the cases of interest anyway, since interesting results only hold in the regime $1 < p < 2$ and $p \leq q$.

Restriction bounds for product sets reduce to bounds on their projections. Thus the interesting sets S we consider will never be product sets.

Theorem 28.1. *Consider two pairs (S_1, μ_1) and (S_2, μ_2) in \mathbf{R}^n and \mathbf{R}^m , and let $S = S_1 \times S_2$, $\mu = \mu_1 \times \mu_2$. Then for $0 < p \leq q \leq \infty$, a restriction bound of the form*

$$\|\widehat{f}\|_{L^q(S, \mu)} \lesssim \|f\|_{L^p(\mathbf{R}^{n+m})}$$

for all $f \in \mathcal{S}(\mathbf{R}^{n+m})$ is equivalent to a pair of bounds of the form

$$\|\widehat{f}_1\|_{L^q(S_1, \mu_1)} \lesssim \|f_1\|_{L^p(\mathbf{R}^n)} \quad \text{and} \quad \|\widehat{f}_2\|_{L^q(S_2, \mu_2)} \lesssim \|f_2\|_{L^p(\mathbf{R}^m)}.$$

for all $f_1 \in \mathcal{S}(\mathbf{R}^n)$ and $f_2 \in \mathcal{S}(\mathbf{R}^m)$. Moreover, the operator norm of the restriction operator to S is the product of the operator norms of the restriction operators to S_1 and S_2 .

Proof. We calculate that for any pair f_1, f_2 ,

$$\|\widehat{f_1 \otimes f_2}\|_{L^q(S, \mu)} = \|\widehat{f_1}\|_{L^q(S_1, \mu_1)} \|\widehat{f_2}\|_{L^q(S_2, \mu_2)}.$$

and

$$\|f_1 \otimes f_2\|_{L^p(\mathbf{R}^{n+m})} = \|f_1\|_{L^p(\mathbf{R}^n)} \|f_2\|_{L^p(\mathbf{R}^m)}.$$

If a restriction estimate held on (S, μ) , it would then follow that

$$\|\widehat{f_1}\|_{L^q(S_1, \mu_1)} \lesssim \frac{\|f_2\|_{L^p(\mathbf{R}^d)}}{\|\widehat{f_2}\|_{L^q(S_2, \mu_2)}} \|f_1\|_{L^p(\mathbf{R}^n)}.$$

Choosing any nontrivial choice of f_2 gives a restriction estimate on (S_1, μ_1) . By symmetry, we also get a restriction estimate on (S_2, μ_2) .

Conversely, suppose we have a restriction estimate on (S_1, μ_1) and (S_2, μ_2) , and $f \in L^1(\mathbf{R}^{n+m}) \cap L^p(\mathbf{R}^{n+m})$. Without loss of generality, we may assume by Littlewood's principle that $q \geq p$. If we let \mathcal{F}_x and \mathcal{F}_y denote the Fourier transform in the first and second variables respectively, then by applying Minkowski's inequality and Fubini's theorem we find that

$$\begin{aligned} \|\widehat{f}\|_{L^q(S, \mu)} &= \left(\int_{\mathbf{R}^m} \int_{\mathbf{R}^n} |\mathcal{F}_x \mathcal{F}_y f(\xi, \eta)|^q d\mu_1(\xi) d\mu_2(\eta) \right)^{1/q} \\ &\lesssim \left(\int_{\mathbf{R}^m} \left(\int_{\mathbf{R}^n} |(\mathcal{F}_y f)(x, \eta)|^p dx \right)^{q/p} d\mu_2(\eta) \right)^{1/q} \\ &\leq \left(\int_{\mathbf{R}^n} \left(\int_{\mathbf{R}^m} |(\mathcal{F}_y f)(x, \eta)|^q d\mu_2(\eta) \right)^{p/q} dx_1 \right)^{1/p} \\ &\lesssim \left(\int_{\mathbf{R}^n} \int_{\mathbf{R}^m} |f(x, y)|^p dy dx \right)^{1/p} = \|f\|_{L^p(\mathbf{R}^d)}. \quad \square \end{aligned}$$

The main example of current research into restriction theory occurs when S is a smooth surface, and where $\mu = \psi\sigma$, where $\psi \in C_c^\infty(S)$. If $S \subset \mathbf{R}^d$ is the graph of some smooth function $\phi : U \rightarrow \mathbf{R}$, where U is an open subset of \mathbf{R}^{d-1} , then the restriction estimate is equivalent to an estimate of the form

$$\|\widehat{f}(\eta, \phi(\eta))\|_{L^q_\eta(U, \psi)} \lesssim \|f\|_{L^p_x},$$

and an extension estimate, equivalent to an estimate of the form

$$\left\| \int_U \psi(\eta) g(\eta) e^{2\pi i(\eta \cdot x + \phi(\eta)t)} d\eta \right\|_{L^p_x L^{p^*}_t} \lesssim \|g\|_{L^{q^*}(U, \psi)}.$$

Since one may always localize a restriction estimate to one of this form if one works with a compactly supported measure, such estimates are no more specific than a general restriction estimate to a compact section of a hypersurface. We shall find that in this setting there is a rich theory relating the existence of restriction maps to the curvature of the surface S . The complete resolution of the restriction problem in this setting is not solved, though much is known.

Restriction estimates often appear in the theory of partial differential equations. For instance, a tempered distribution u on \mathbf{R}^d which solves the

free Schrödinger equation $\partial_t u = (i/2\pi)\Delta_x u$ has space-time Fourier transform supported on the parabola defined by the equation $\tau + |\xi|^2 = 0$, and so there exists a tempered distribution v on \mathbf{R}^d such that

$$u(x, t) = \int v(\xi) e^{2\pi i(\xi \cdot x - |\xi|^2 t)} d\xi.$$

Similarly, a tempered distribution u on \mathbf{R}^n which solves the free wave equation $\partial_t^2 u = \Delta_x u$ has space-time Fourier transform support on the light cone defined by the equation $\tau^2 = |\xi|^2$, and thus there exists two tempered distributions v_1 and v_2 on \mathbf{R}^n such that

$$u(x, t) = \int v_1(\xi) e^{2\pi i(\xi \cdot x + |\xi|t)} d\xi + \int v_2(\xi) e^{2\pi i(\xi \cdot x - |\xi|t)} d\xi.$$

Thus understanding the regularity of solutions to these equations relates to understanding the extension problem for the parabola and light cone.

Example. *If S is a hyperplane, then no nontrivial estimates are possible for the restriction problem. This is because the only possible extension-type estimates of the form*

$$\left\| \int_{\mathbf{R}^d} \psi(\eta) g(\eta) e^{2\pi i(\eta \cdot x)} d\eta \right\|_{L_x^{p^*} L_t^{p^*}} \lesssim \|g\|_{L^{q^*}(U, \psi)}$$

must have $p = 1$, since the integral on the left hand side is independent of t . The general heuristic that has been developed in restriction theory is that ‘flatness’ is the main feature of the hyperplane preventing better estimates. For curved surfaces, one expects to get a much wider range of estimates.

Since flatness prevents getting good restriction estimates, we are lead to begin our study with the study of surfaces with non-vanishing curvature. The classical model examples of such results are the restriction estimates for the paraboloid and the sphere, where we have additional symmetries to use not present in all surfaces. There are several simple examples which limit the range of estimates one can get for hypersurfaces of nonvanishing curvature. No improvements to these limitations is known, and it is conjecture that outside of these limitations, one can always get an estimate. This is the *restriction conjecture*.

The first limitation follows from stationary phase. Let S be a hypersurface with nonvanishing curvature containing, without loss of generality, the origin. The standard theory of stationary phase shows that if $\psi \in C_c^\infty(\mathbf{R}^d)$ is supported on a small enough neighborhood of the origin, then for x lying in a cone around the normal vector to the surface at the origin, we have

$$|E_S \psi(x)| \gtrsim |x|^{-(d-1)/2}.$$

Thus $E_S \psi \in L^{p^*}(\mathbf{R}^d)$ if and only if $p^* > 2d/(d-1)$, i.e. if $p < 2d/(d+1)$. Thus we can only have a restriction estimate for $p < 2d/(d+1)$.

The second example is due to Knapp, and called the ‘Knapp example’ in the literature, which is really just studying the extension operator applied to a wave packet. It is obtained by exploiting the uncertainty principle. To construct the Knapp example, by translating and rotating (which by symmetry does not change any estimates in the Fourier transform) we may assume our surface S contains the origin and has normal vector pointing directly upwards at the origin. Consider a bump function $\eta \in C_c^\infty(S)$ supported on a small ball of radius r about the origin, with $\eta(x) = 1$ for $|x| \leq 1$, but with $\|\eta\|_{L^\infty(S)} \leq 1$. Then

$$\|\psi\|_{L^{q^*}(S, \mu)} \sim r^{(d-1)(1-1/q)}$$

On the other hand, $\eta \cdot \mu$ is support on a ‘cap’

$$\Theta_r = \{(\xi, \tau) : |\xi| \leq r, |\tau| \leq cr^2\}$$

for some constant $c > 0$ depending on the surface, but independent of r . The uncertainty principle heuristically implies that $E_S \eta$, which is the Fourier transform of $\eta \cdot \mu$, is locally constant on translations of the ‘tube’

$$\Theta_r^* = \{(x, t) : |x| \leq 1/r, |t| \leq C/r^2\}.$$

Since $\eta \cdot \mu$ has total mass $\sim r^{d-1}$, we conclude that $E_S \eta(x, t) \gtrsim r^{d-1}$ for $(x, t) \in \Theta_r^*$, and thus that

$$\|E_S \eta\|_{L^{p^*}(\mathbf{R}^d)} \gtrsim r^{d-1} |\Theta_r^*|^{1/p^*} = r^{(d+1)/p-2}.$$

To be more rigorous, since $E_S \eta(0) \gtrsim r^{d-1}$, it follows from (TODO: record Prop 5.5 of Wolff’s notes in these notes) that

$$\int |E_S \eta(x, t)| (1 + rx + r^2 t)^{-N} dx dt \gtrsim_N |\Theta_r^*| r^{d-1} \gtrsim r^{-2}$$

Hölder's inequality implies that if N is chosen larger than d , then

$$\int |E_S(\psi)(x, t)|(1 + rx + r^2t)^{-N} dx dx_d \lesssim r^{-(d+1)(1/p)} \|E_S\phi\|_{L^{p^*}(\mathbf{R}^d)},$$

and so we conclude that $\|E_S\phi\|_{L^{p^*}(\mathbf{R}^d)} \geq r^{(d+1)(1/p)-2}$. But this inequality implies that we must have $(d+1)/p - (d-1)/q \leq 2$ if we want an estimate of the form $\|E_S\eta\|_{L^{p^*}(\mathbf{R}^d)} \lesssim \|\eta\|_{L^{q^*}(S, \mu)}$ to hold.

Remark. TODO: If S is a surface passing through the origin whose derivatives vanish to order k , consider a variant of the Knapp example which shows how restriction fails in this domain.

The general belief in the field is that these examples are essentially the only barriers to restriction. Thus for any such pair $1 \leq p, q < \infty$ such that $p < 2d/(d+1)$ and $(d+1)/p - (d-1)/q \leq 2$, we have restriction (and thus extension) bounds from $L^{q^*}(S, \mu)$ to $L^{p^*}(\mathbf{R}^d)$, and thus a restriction bound from $L^p(\mathbf{R}^d)$ to $L^q(S, \mu)$. This is the *restriction conjecture*, which, to a large extent, still remains an open problem.

28.1 Stein-Tomas Theorem

The classical result for restriction estimates on surfaces with non-vanishing curvature is the *Stein-Tomas theorem*. If S is a hypersurface in \mathbf{R}^d with nonvanishing curvature, and $\mu = \psi\sigma$ for some $\psi \in C_c^\infty(\mathbf{R}^d)$, where σ is the surface measure on S , then for $p \leq 2(d+1)/(d+3)$,

$$\|\hat{f}\|_{L^2(S)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

This result is tight, as shown by the Knapp example. There are many proofs of the Stein-Tomas theorem. The simplest, a proof by Tomas (1975), uses the theory of stationary phase, but has the disadvantage that it does not include the endpoint case where $p = 2(d+1)/(d+3)$.

Theorem 28.2. *Let S be a hypersurface with nonvanishing curvature, and let $\mu = \psi d\sigma$, where σ is the surface measure on S , and $\psi \in C_c^\infty(S)$. Then if $p < 2(d+1)/(d+3)$, then for $f \in L^1(\mathbf{R}^d)$,*

$$\|R_S f\|_{L^2(S, \mu)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

Proof. We apply a TT^* argument, which can be used to study the boundedness of any operator whose codomain is a Hilbert space. For $f \in L^1(\mathbf{R}^d)$,

$$(E_S R_S f)(x) = \int_S \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\mu(\xi) = \mu(-D)f,$$

The TT^* method implies that the restriction estimate we wish to prove holds if and only if we have a bound of the form

$$\|E_S R_S f\|_{L^{p^*}(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

If k is the convolution kernel associated with $\mu(-D)$, then $E_S R_S f = k * f$, which is a much easier operator to understand than the individual restriction and extension operators. In particular, the fact that S has nonvanishing curvature means that

$$|k(x)| \sim \langle x \rangle^{-(d-1)/2}.$$

This implies that $k \in L^r(\mathbf{R}^d)$ for $r > 2d/(d-1)$, and lies in $L^{r,\infty}(\mathbf{R}^d)$ for $r = 2d/(d-1)$. Thus Young's inequality, and its weak variant, implies that

$$\|k * f\|_{L^{p^*}(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

for $p \leq 4d/(3d+1)$. To improve this bound to hold for $p < 2(d+1)/(d-1)$, we must exploit the oscillation of k , which is completely discarded by Young's inequality.

To argue this, we consider a dyadic decomposition, writing $k(x) = \sum_{j=0}^{\infty} k_j(x)$, where k_0 is smooth and compactly supported on a ball of radius $O(1)$, and for $j > 0$, $k_j(x) = \text{Dil}_{2^j} \psi \cdot k(x)$ for some $\psi \in C_c^\infty(\mathbf{R}^d)$. Since k_0 is smooth and compactly supported, we immediately obtain that

$$\|k_0 * f\|_{L^{p^*}(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}.$$

For $j > 0$, k_j has magnitude $O(2^{-j(d-1)/2})$ on this set, so we obtain from Young's inequality that

$$\|k_j * f\|_{L^\infty(\mathbf{R}^d)} \lesssim 2^{-j(d-1)/2} \|f\|_{L^1(\mathbf{R}^d)}.$$

We wish to obtain an (L^2, L^2) bound using orthogonality. We have

$$\hat{k}_j = 2^j (\hat{k} * \text{Dil}_{1/2^j} \hat{\psi}) = 2^{jd} (\mu * \text{Dil}_{1/2^j} \hat{\psi}) = 2^{jd} \int \hat{\psi}(2^j(\cdot - \xi)) d\mu(\xi).$$

Now μ is compactly supported on a hypersurface and has total mass $O(1)$ there. Since $\text{Dil}_{1/2^j} \hat{\psi}$ is supported in a ball of radius $1/2^j$, we should expect \hat{k}_j to distribute the mass of μ onto a $1/2^j$ thickening of a portion of a hypersurface, which has total mass $O(2^{-j(d-1)})$, so that the expect height of the convolution should be $O(2^{j(d-1)})$. Thus

$$\|\hat{k}_j\|_{L^\infty(\mathbf{R}^d)} \lesssim 2^{jd} \cdot 2^{-j(d-1)} = 2^j.$$

The rigorous argument is not too interesting (just use the fact that $\hat{\psi}$ is Schwartz) and left to the reader. Thus we conclude that

$$\|k_j * f\|_{L^2(\mathbf{R}^d)} \lesssim 2^j \|f\|_{L^2(\mathbf{R}^d)}.$$

For any $p < 2(d+1)/(d+3)$, interpolation therefore tells us that there is $\varepsilon > 0$ such that

$$\|k_j * f\|_{L^{p^*}(\mathbf{R}^d)} \lesssim 2^{-i\varepsilon} \|f\|_{L^p(\mathbf{R}^d)}.$$

Summing up gives the required bounds. \square

Remark. The only structure we used about S and μ here is that μ has a uniform, Frostman type bound, and that $\hat{\mu}$ has a geometric decay as frequencies become large. For such a pair one has a bound

$$\|R_S f\|_{L^2(S, \mu)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

provided that $1 \leq p < 2(d + \beta - \alpha)/(2d + \beta - 2\alpha)$, where $0 < \alpha < n$, $\beta > 0$, where $|\hat{\mu}(\xi)| \lesssim \langle \xi \rangle^{-\beta}$ and $|\mu(B_r(x))| \lesssim r^\alpha$.

The endpoint proof $p = 2(d+1)/(d+3)$ was first given by Stein (1975, unpublished). It involves the same idea, namely, to consider a dyadic decomposition, but to replace the inefficient use of the triangle inequality

$$\|k * f\|_{L^{p^*}(\mathbf{R}^d)} \leq \sum_{i=0}^{\infty} \|k_i * f\|_{L^{p^*}(\mathbf{R}^d)}$$

with a complex interpolation argument. Namely, we prove that for all $t \in \mathbf{R}$,

$$\left\| \sum_{j>0} 2^{[\frac{n-1}{2} + it]j} (f * k_j) \right\|_{L^\infty(\mathbf{R}^d)} \lesssim \|f\|_{L^1(\mathbf{R}^d)}$$

and that

$$\left\| \sum_{j>0} 2^{[-1+it]j} (f * k_j) \right\|_{L^2(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^d)},$$

so that we can apply analytic interpolation. From Young's inequality, the first inequality follows if we can show that

$$\left\| \sum_{j>0} 2^{[\frac{d-1}{2}+it]j} k_j \right\|_{L^\infty(\mathbf{R}^d)} \lesssim 1.$$

This just follows from the fact that all but consecutive kernels in the sequence $\{k_j\}$ have disjoint support (this is much more efficient than the triangle inequality), and the appropriate decay. To establish the second inequality, we must show that

$$\left\| \sum_{j>0} 2^{[(d-1)+it]j} (\mu * \text{Dil}_{1/2^j} \hat{\psi}) \right\|_{L^\infty(\mathbf{R}^d)}.$$

We must be slightly more efficient than we were before. TODO: See Tao's Notes.

Here is an argument that gives the endpoint for the special case of the sphere, relying on an alternate property of the sphere than curvature.

Lemma 28.3. *Let σ be the surface measure of the sphere $S = S^{n-1}$. Then $\sigma * \sigma$ is absolutely continuous with respect to the Lebesgue measure, supported on $|\xi| \leq 2$, and on this set,*

$$(\sigma * \sigma)(\xi) \lesssim \begin{cases} 1/|\xi| & : 0 < |\xi| \leq 1, \\ (2 - |\xi|)^{(n-3)/2} & : 1 \leq |\xi| \leq 2. \end{cases}$$

Proof. For each $\varepsilon > 0$, let $\sigma_\varepsilon = (1/\varepsilon)\mathbf{I}_{S_\varepsilon}$. Then σ_ε converges weakly to σ as $\varepsilon \rightarrow 0$, which implies that $(\sigma_\varepsilon * \sigma_\varepsilon)$ converges to $\sigma * \sigma$ weakly. Now $\sigma_\varepsilon * \sigma_\varepsilon$ is radial. Thus it suffices to bound $(\sigma_\varepsilon * \sigma_\varepsilon)(\xi)$ for $\xi = (r, 0)$, where $0 < r \leq 2$. Now

$$(\sigma_\varepsilon * \sigma_\varepsilon)(\xi) = |S_\varepsilon \cap (\xi + S_\varepsilon)|.$$

If $(\alpha, \beta) \in S_\varepsilon \cap (\xi + S_\varepsilon)$, where $\alpha \in \mathbf{R}$, $\beta \in \mathbf{R}^{n-1}$, then

$$1 - 2\varepsilon \leq \alpha^2 + \beta^2 \leq 1 + 3\varepsilon \quad \text{and} \quad 1 - 2\varepsilon \leq (\alpha - r)^2 + \beta^2 \leq 1 + 3\varepsilon.$$

Together, these inequalities imply that $\alpha = r/2 + O(\varepsilon/r)$ and thus $\beta^2 = 1 - r^2/4 + O(\varepsilon + \varepsilon^2/r)$, so $\beta = \sqrt{1 - r^2/4} + O(\varepsilon)$. ~~TODO~~ Thus α ranges over an interval of length $O(\varepsilon/r)$, and for each α , β can vary over a region of $n - 1$ dimensional volume $O(\varepsilon)$, so we conclude that \square

28.2 Bochner-Riesz Multipliers

One consequence of the Stein-Tomas theorem is a characterization of the boundedness for the Bochner-Riesz multipliers

$$m^\delta(\xi) = (1 - |\xi|^2)_+^\delta.$$

Herz showed a necessary condition for boundedness.

Theorem 28.4. *If $m^\delta \in M^p(\mathbf{R}^d)$, then*

$$\delta > n \left| \frac{1}{p} - \frac{1}{2} \right| - 1/2.$$

Proof. An oscillatory integral calculation shows that if $\widehat{K}^\delta = m^\delta$, then

$$K_\delta(x) = \frac{C_d \cos(2\pi|x|)}{|x|^{\frac{d+1}{2}+\delta}} + O(|x|^{-\frac{n+3}{2}+\delta}).$$

If χ is the indicator function of the unit ball, then Minkowski's inequality shows that for $1 \leq p \leq \infty$,

$$\|K^\delta\|_{L^p(\mathbf{R}^d)} \lesssim \|m^\delta(D)\chi\|_{L^p(\mathbf{R}^d)} \lesssim \|m^\delta\|_{M^p(\mathbf{R}^d)}.$$

Without loss of generality, we may assume that $p \geq 2$, in which case it suffices to show that one can only have a bound if $d(1/2 - 1/p) - 1/2 < \delta$. But K^δ lies in $L^p(\mathbf{R}^d)$ precisely for this range. \square

The Bochner Riesz conjecture is that this is *precisely* the range for which the operator is bounded.

Theorem 28.5. *Suppose that*

$$\|R$$

Proof. Assume $p \geq 2$. We break the kernel K^δ into dyadic space regions, i.e.

$$K^\delta = \sum_{j=0}^{\infty} K_j^\delta,$$

where K_0^δ is supported in the ball of radius 2, and

$$K_j^\delta(x) = \psi(x/2^j)K^\delta(x) = \psi_j(x)K^\delta(x).$$

Our proof will be complete if we can show that

$$\|K_j^\delta * f\|_{L^p(\mathbf{R}^d)} \lesssim 2^{[n(1/p-1/2)-1/2-\delta]j} \|f\|_{L^p(\mathbf{R}^d)}.$$

Since the support of $K_j^\delta * f$ lies on a 2^j thickening of the support of f , a decomposition argument implies that it suffices to show the result for any function f supported on a ball of radius 2^j at the origin. To make this look more like Tomas-Stein, we note that for such a function f , $K_j^\delta * f$ is supported on a ball of radius $O(2^j)$, and thus

$$\|K_j^\delta * f\|_{L^p(\mathbf{R}^d)} \lesssim 2^{dj(1/p-1/2)} \|K_j^\delta * f\|_{L^2(\mathbf{R}^d)}.$$

Thus it suffices to show that

$$\|K_j^\delta * f\|_{L^2(\mathbf{R}^d)} \lesssim 2^{-(1/2+\delta)j} \|f\|_{L^p(\mathbf{R}^d)}.$$

Plancherel shows that it suffices to prove that

$$\|\widehat{[\psi_j * m^\delta]} \cdot \widehat{f}(\xi)\|_{L_\xi^2} \lesssim 2^{-(1/2+\delta)j} \|f\|_{L_x^p}$$

Now ψ_j is supported away from the origin, and thus its Fourier transform is highly oscillatory. The multiplier m^δ is also proportionally smooth. Thus we should expect $\widehat{\psi_j * m^\delta}$ to have some decay by virtue of some cancellation. The ‘high frequency’ terms that make up m^δ are supported near the boundary of the sphere, since the function is smooth everywhere else. We will actually obtain that

$$|(\widehat{\psi_j * m^\delta})(\xi)| \lesssim_N 2^{-\delta j} \langle 2^j ||\xi| - 1| \rangle^{-N}.$$

Thus we get rapid decay outside of the annulus of thickness $1/2^j$. It thus suffices to show that if N is suitably large, then

$$\int \frac{|\widehat{f}(\xi)|^2}{\langle 2^j ||\xi| - 1| \rangle^N} d\xi \lesssim \|f\|_{L^p(\mathbf{R}^n)}.$$

The fact that f is compactly supported allows us to use the pointwise bound $|\widehat{f}(\xi)| \lesssim 2^{O(j)} \|f\|_{L^p(\mathbf{R}^d)}$, which gives the estimate for $||\xi| - 1| \geq 1/2$. The remaining integrand then follows by switching to polar coordinates, and then applying the restriction theorem.

Now let us justify that

$$|(\widehat{\psi_j} * m^\delta)(\xi)| \lesssim_N 2^{-\delta j} \langle 2^j ||\xi| - 1| \rangle^{-N}.$$

This motivates us to perform the decomposition

$$m^\delta(\xi) = m^\delta(\xi) \phi(2^j(|\xi| - 1)) + m^\delta(\xi) [1 - \phi(2^j(|\xi| - 1))] = m_1^\delta(\xi) + m_2^\delta(\xi),$$

where $\phi \in C_c^\infty[0, \infty)$ equals one in a neighborhood of the origin. The symbol $m_1^\delta(\xi)$ is now supported on the annulus of thickness $O(2^{-j})$ and radius 1. The height of the symbol is also $O(2^{-\delta j})$. Thus

$$|(\widehat{\psi_j} * m_1^\delta)(\xi)| \lesssim \int_{|r-1| \sim 2^{-j}} 2^{-\delta j} (|\widehat{\psi_j}| * \sigma_r) dr.$$

The fact that $\sigma_r * |\widehat{\psi_j}| \lesssim_N 2^j \langle 2^j ||x| - 1| \rangle^{-N}$ shows this contribution is acceptable. The remaining term is estimated via integration by parts (one may antidifferentiate ψ_j without introducing any problems since it vanishes near the origin). TODO. \square

28.3 Restriction on the Paraboloid

The main way we can extend local estimates to global estimates is to apply some kind of symmetry property, normally scaling. Let us use this technique to show that we can extend the result of the Tomas-Stein theorem to give the same bounds for restriction to the non-compact elliptic paraboloid

$$S = \{(\xi, \omega) \in \mathbf{R}^{d-1} \times \mathbf{R} : |\xi|^2 = \omega\}.$$

We work with extension estimates, setting, for a function $f : \mathbf{R}^{d-1} \rightarrow \mathbf{C}$,

$$Ef(x, t) = \int_{\mathbf{R}^{d-1}} e^{2\pi i(|\xi|^2 t + \xi \cdot x)} f(\xi) d\xi.$$

Our goal is to show bounds of the form $\|Ef\|_{L^p(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^{d-1})}$. This is possible, provided that $p = 2(d+1)/(d+3)$.

Theorem 28.6. *If $p = 2(d+1)/(d+3)$, then for any $f \in \mathcal{S}(\mathbf{R}^{d-1})$,*

$$\|Ef\|_{L^{p'}(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^{d-1})}.$$

Proof. The Tomas-Stein theorem gives the required bound for any $f \in C_c^\infty(\mathbf{R}^d)$, where the implicit constant depends on the support of f . For such f , define

$$Ef(x, t) = \int_{\mathbf{R}^{d-1}} e^{2\pi i(|\xi|^2 t + \xi \cdot x)} f(\xi) d\xi.$$

We note that $E(\text{Dil}_{\xi, a} f) = a^{d-1} \cdot \text{Dil}_{a^{-2}, t} \text{Dil}_{a^{-1}, x}(Ef)$, in light of the ‘parabolic symmetry’ of the extension operator. Given a function $f \in C_c^\infty(\mathbf{R}^d)$ supported on a ball of radius R at the origin, $\text{Dil}_{\xi, 1/R} f$ is supported on a ball of radius 1 at the origin, and so Tomas-Stein says that

$$\|E(\text{Dil}_{\xi, 1/R} f)\|_{L^{p'}(\mathbf{R}^d)} \lesssim \|\text{Dil}_{\xi, 1/R} f\|_{L^2(\mathbf{R}^{d-1})} = R^{-(d-1)/2} \cdot \|f\|_{L^2(\mathbf{R}^{d-1})},$$

where the implicit constant is independent of f . But we also know that

$$\begin{aligned} \|E(\text{Dil}_{\xi, 1/R} f)\|_{L^{p'}(\mathbf{R}^d)} &= R^{1-d} \|\text{Dil}_{R^2, t} \text{Dil}_{R, x}(Ef)\|_{L^{p'}(\mathbf{R}^d)} \\ &= R^{1-d} R^{2/p'} R^{(d-1)/p'} \|Ef\|_{L^{p'}(\mathbf{R}^d)}, \end{aligned}$$

and so we conclude that

$$\|Ef\|_{L^{p'}(\mathbf{R}^d)} \lesssim R^{(d-1)/2 - (d+1)/p'} \|f\|_{L^2(\mathbf{R}^{d-1})} = \|f\|_{L^2(\mathbf{R}^{d-1})}.$$

In other words, we have found a bound independent of the support of f . A simple approximation argument extends the bound to general $f \in \mathcal{S}(\mathbf{R}^d)$. \square

Remark. The same scaling symmetries show that we can only have an extension bound

$$\|Ef\|_{L^{p'}(\mathbf{R}^d)} \lesssim \|f\|_{L^{q'}(\mathbf{R}^{d-1})}$$

if $(d+1)/p' = (d-1)/q$. For $q = 2$, this shows the above result is tight.

It is a general heuristic that the extension theory for the paraboloid is essentially the same as the extension theory for a cap of a curve point. In particular, if we have a bound of the form

$$\|Ef\|_{L^{p'}(\mathbf{R}^d)} \lesssim \|f\|_{L^{q'}(S,\mu)},$$

where $(d+1)/p' = (d-1)/q$, and where μ is a smooth measure supported on a small cap around the neighbourhood of a point with non-vanishing curvature. One can then approximate the paraboloid by a rescaling of the cap, and thus obtain a bound of the form

$$\|Ef\|_{L^{p'}(\mathbf{R}^d)} \lesssim \|f\|_{L^{q'}(\mathbf{R}^{d-1})}$$

for any $f \in \mathcal{S}(\mathbf{R}^{d-1})$, where E is the extension function on the paraboloid
 TODO: Prove This. Thus, despite one theory being local and the other being global, the theories of these two extension operators are roughly equivalent. Every result obtained in one setting can essentially be translated to the other setting.

28.4 Restriction to the Cone

TODO: Conjectured Estimates

TODO: Lorentz Invariant

TODO: Application to the wave equation.

28.5 TODO: Hardy Littlewood Majorant Conjecture

Chapter 29

Almost Orthogonality

It is a standard result of Hilbert space theory that if $\{e_n\}$ are a family of pairwise orthogonal vectors in a Hilbert space H , then Parseval's inequality

$$\left\| \sum_n a_n e_n \right\|_H = \left(\sum |a_n|^2 \right)^{1/2}$$

holds. In many cases in analysis, we do not have *perfect* orthogonality, but we know that $\langle e_n, e_m \rangle$ is small for most pairs n and m . For general non-orthogonal vectors we do have the identity

$$\left\| \sum_n a_n e_n \right\|_H^2 = \sum_n a_n \overline{a_m} \langle e_n, e_m \rangle.$$

Here is a simple result which exploits this identity.

Lemma 29.1. *Suppose $\{e_n : n \in \mathbf{Z}^d\}$ is a family of vectors such that there is $\delta > 0$ with*

$$\langle e_n, e_m \rangle \leq \frac{C}{\langle n - m \rangle^{d+\delta}}.$$

Then

$$\left\| \sum_n a_n e_n \right\| \lesssim_\delta C^{1/2} \left(\sum |a_n|^2 \right)^{1/2}$$

Proof. Applying Cauchy-Schwartz and Young's convolution inequality, we

conclude that

$$\begin{aligned}
\left\| \sum_n a_n e_n \right\|_H^2 &\leq C \sum_n |a_n| \sum_m \frac{|a_m|}{\langle n-m \rangle^{d+\delta}} \\
&\leq C \left(\sum_n |a_n|^2 \right)^{1/2} \left(\sum_n \left(\sum_m \frac{|a_m|}{\langle n-m \rangle^{d+\delta}} \right)^2 \right)^{1/2} \\
&\lesssim_\delta C \left(\sum_n |a_n|^2 \right).
\end{aligned}$$

We then just take square roots on both sides of the formula. \square

Let us consider some examples of almost orthogonal systems.

Example. Consider $f \in L^2(\mathbf{R}^d)$ with compact support, and set $e_n = \text{Trans}_n f$. The $\langle e_n, e_m \rangle \neq 0$ only if $|n-m| \lesssim 1$, and then we have the trivial bound $\langle e_n, e_m \rangle \leq \|f\|_{L^2(\mathbf{R}^d)}^2$. Thus we conclude that

$$\left\| \sum a_n e_n \right\|_{L^2(\mathbf{R}^d)} \lesssim \|f\|_{L^2(\mathbf{R}^d)} \left(\sum |a_n|^2 \right)^{1/2}.$$

In this situation, another way to prove this inequality is to break up the sum into arithmetic subsequences which have a high enough gap so that the sums are over functions with disjoint support, in which case we have true orthogonality. Summing up the subsums then gives the result.

Example. Given $\phi \in C_c^\infty(\mathbf{R}^d)$, and consider $e_n = \text{Mod}_n f$. Then

$$\langle e_n, e_m \rangle = \int |\phi(x)|^2 e^{2\pi i(n-m) \cdot x} dx.$$

We can then apply the theorem of nonstationary phase, which gives

$$|\langle e_n, e_m \rangle| \lesssim_k \frac{1}{\langle n-m \rangle^k}$$

for any k . Thus this is an almost orthogonal system.

For almost orthogonal systems, we do *not* have a lower bound

$$\left\| \sum a_n e_n \right\|_{L^2(\mathbf{R}^d)} \gtrsim \left(\sum |a_n|^2 \right)^{1/2}.$$

Indeed, the decay does not preclude us from choosing the same two vectors for different indices, which give complete cancellation for appropriately chosen constants.

Example. Consider a family of vectors $\{e_n\}$ in $L^2(\mathbf{R}^d)$ such that for each n , the support of e_n intersects the support of $O(1)$ other vectors e_m . Do we have almost orthogonality here?

The most well used almost orthogonality result is the Cotler-Stein lemma, which enables us to bound sums of operators which are almost orthogonal (the bounded operators from a Hilbert space to itself form a Hilbert space). To begin with, let us treat the case where we are dealing with a family of operators which are pairwise orthogonal.

Lemma 29.2. Let $\{T_n\}$ be a sequence of operators on a Hilbert space H with $\sup_n \|T_n\| < \infty$ for each n , and such that $T_i^* T_j = T_i T_j^* = 0$ for any $i \neq j$. If

$$S_N = \sum_{n=1}^N T_n,$$

then S_N converges in the strong topology to an operator S with $\|S\| \leq 1$.

Proof. Since $T_i^* T_j = 0$ for $i \neq j$, it follows that the image of T_j is contained in the kernel of T_i^* , which is the orthogonal complement of the image of T_i . Thus the images $\{V_i\}$ of the operators $\{T_i\}$ are all orthogonal to one another. By symmetry, the images $\{W_i\}$ of the operators $\{T_i^*\}$ are all orthogonal to one another. Now W_i is the orthogonal complement of the kernel of T_i . Thus if $x = x_0 + \sum_i x_i$, with x_0 perpendicular to all W_i , and $x_i \in W_i$, then $\|T_i x\| = \|T_i x_i\|$. But this means that for any $x \in H$,

$$\begin{aligned} \|S_N x\|^2 &= \sum \|T_i x\|^2 \\ &= \sum \|T_i x_i\|^2 \\ &\leq \sum \|x_i\|^2 \\ &\leq \|x\|^2. \end{aligned}$$

Thus $\|S_N x\| \leq \|x\|$, and so $\|S_N\| \leq 1$. Now since $\sum_{i>N} \|x_i\|^2 \rightarrow 0$ as $i \rightarrow \infty$ for any $x \in H$, it follows that $\{S_N x\}$ is Cauchy, and so there exists a limit Sx . It is now simple to prove S is a bounded linear operator with $\|S\| \leq 1$. \square

Remark. The important point here is that not only are the outputs to all the operators mutually orthogonal, i.e. $T_i^* T_j = 0$, nor are the important inputs to each of the operators mutually orthogonal, i.e. $T_i T_j^* = 0$, but both points are actually true. If only one of these was true, we could only get a bound of the form $\|S_N\| \lesssim \sqrt{N}$.

Cotler-Stein gives an analogous result for almost orthogonal families of operators.

Theorem 29.3 (Cotler Stein). *Let H_1 and H_2 be Hilbert spaces, and consider a family of bounded operators $\{T_\alpha : H_1 \rightarrow H_2\}$. Let $a_{\alpha\beta} = \|T_\alpha T_\beta^*\|$ and let $b_{\alpha\beta} = \|T_\alpha^* T_\beta\|$. Then if*

$$A = \sup_\alpha \sum_\beta \sqrt{a_{\alpha\beta}} \quad \text{and} \quad B = \sup_\alpha \sum_\beta \sqrt{b_{\alpha\beta}}$$

are both finite, then $\sum T_\alpha$ converges unconditionally in the strong operator topology, and

$$\left\| \sum T_\alpha \right\| \leq \sqrt{AB}.$$

One can often extend almost orthogonality to obtain bounds in other L^p spaces via applying interpolation. Here is such a version I encountered in Heo, Nazarov, and Seeger's paper *Radial Fourier Multipliers in High Dimensions*.

Theorem 29.4. *Consider a family of functions $\{f_n : n \in \mathbf{Z}^d\}$ in $L^2(\mathbf{R}^d)$, together with a family of sidelength one cubes $\{Q_n\}$ in \mathbf{R}^d such that $\text{supp}(f_n) \subset Q_n$. Suppose*

$$|\langle f_n, f_m \rangle| \leq \frac{1}{\langle n - m \rangle^\beta}.$$

for some $\beta \in (0, d)$. Then for $p < 2d/(2d - \beta)$,

$$\left\| \sum a_n f_n \right\|_{L^p(\mathbf{R}^d)} \lesssim_{d, \beta, p} \left(\sum |a_n|^p \right)^{1/p}$$

Remark. If $\beta > d$, then Young's convolution inequality implies that

$$\begin{aligned}
\left\| \sum a_n f_n \right\|_{L^2(\mathbf{R}^d)} &= \left(\sum_{n,m} a_n \overline{a_m} \langle f_n, f_m \rangle \right)^{1/2} \\
&\leq \left(\sum_{n,m} \frac{a_n \overline{a_m}}{\langle n-m \rangle^\beta} \right)^{1/2} \\
&\leq \left(\sum_n |a_n|^2 \right)^{1/4} \left(\sum_n \left| \sum_m \frac{\overline{a_m}}{\langle n-m \rangle^\beta} \right|^2 \right)^{1/4} \\
&\leq \left(\sum_n |a_n|^2 \right)^{1/2} \left(\sum_n \frac{1}{\langle n \rangle^\beta} \right) \\
&\lesssim_{\beta,d} \left(\sum_n |a_n|^2 \right)^{1/2}
\end{aligned}$$

Thus one can view $\beta < d$ as a case where we have some, but not enough orthogonality to prove an L^2 orthogonality bound.

Proof. We view this result as proving the boundedness of the operator

$$a \mapsto \sum_n a_n f_n$$

from $l^p(\mathbf{Z}^d)$ to $L^p(\mathbf{R}^d)$. We shall prove that for $1 \leq p \leq 2d/(2d - \beta)$, a *restricted strong type inequality holds*, from which the general bound holds by interpolation. It suffices to show that for any finite set of indices $I \subset \mathbf{Z}^d$,

$$\left\| \sum_{n \in I} f_n \right\|_{L^p(\mathbf{R}^d)} \lesssim \#(I)^{1/p}.$$

Partition \mathbf{R}^d into an almost disjoint family of sidelength one cubes $\{R_\alpha\}$, define $I_\alpha = \{n \in I : Q_n \cap R_\alpha \neq \emptyset\}$, and set $F_\alpha = \sum_{n \in I_\alpha} f_n$. Now for each $x \in \mathbf{R}^d$, there are at most 3^d indices α such that $F_\alpha(x) \neq 0$. Thus

$$\left\| \sum_{n \in I} f_n \right\|_{L^p(\mathbf{R}^d)} = \left\| \sum_\alpha F_\alpha \right\|_{L^p(\mathbf{R}^d)} \leq 3^d \left(\sum_\alpha \|F_\alpha\|_{L^p(\mathbf{R}^d)}^p \right)^{1/p}.$$

Applying the almost-orthogonality of the functions $\{f_n\}$,

$$\begin{aligned}\|F_\alpha\|_{L^2(\mathbf{R}^d)}^2 &\leq \sum_{n,m \in I_\alpha} \frac{1}{\langle n-m \rangle^\beta} \\ &\leq \sum_{n \in I_\alpha} \sum_{|m| \lesssim \#(I_\alpha)^{1/d}} \frac{1}{\langle n-m \rangle^\beta} \\ &\lesssim \#(I_\alpha) \cdot \#(I_\alpha)^{1-\beta/d}\end{aligned}$$

Thus $\|F_\alpha\|_{L^2(\mathbf{R}^d)} \lesssim \#(I_\alpha)^{1-\beta/2d}$. Combined with the fact that F_α is supported on a sidelength $O(1)$ cube, we conclude that for $0 < p \leq 2$, $\|F_\alpha\|_{L^p(\mathbf{R}^d)} \lesssim_p \|F_\alpha\|_{L^2(\mathbf{R}^d)}$. But putting this together means that

$$\left(\sum_{\alpha} \|F_\alpha\|_{L^p(\mathbf{R}^d)}^p \right)^{1/p} = \left(\sum_{\alpha} \#(I_\alpha)^{p(1-\beta/2d)} \right)^{1/p}$$

Provided that $p(1 - \beta/2d) \leq 1$, i.e. $p \leq 2d/(2d - \beta)$, we have

$$\sum_{\alpha} \#(I_\alpha)^{p(1-\beta/2d)} \leq \sum \#(I_\alpha) = \#(I)$$

and so we conclude that

$$\left(\sum_{\alpha} \|F_\alpha\|_{L^p(\mathbf{R}^d)}^p \right)^{1/p} \lesssim \#(I)^{1/p},$$

which completes the proof of the restricted strong type bound. \square

Chapter 30

Weighted Estimates

30.1 Hardy-Littlewood Maximal Function

Let us consider a basic weighted estimate for the Hardy-Littlewood maximal function.

Theorem 30.1. *Suppose $w > 0$ is a measurable function. Then for any $f \in L^1_{loc}(\mathbf{R}^d)$, we have the weak type bound*

$$\int_{\mathbf{R}^d} \mathbf{I}(|Mf(x)| > \lambda) w(x) dx \lesssim_d \frac{1}{\lambda} \int_{\mathbf{R}^d} |f(x)| Mw(x) dx,$$

and for all $1 \leq p \leq \infty$, we have the strong type bound

$$\left(\int_{\mathbf{R}^d} |Mf(x)|^p w(x) dx \right)^{1/p} \lesssim_{d,p} \left(\int_{\mathbf{R}^d} |f(x)|^p Mw(x) dx \right)^{1/p}.$$

Proof. The result automatically follows for $p = \infty$, so by the Stein-Weiss interpolation theorem it suffices to obtain the weak-type bound. We work similarly to the standard Vitali-type approach. Renormalizing, to complete the proof it suffices to show that for any compact set K such that $|Mf(x)| > 1$ for all $x \in K$,

$$\int_K w(x) dx \lesssim_d \int_{\mathbf{R}^d} |f(x)| Mw(x) dx.$$

Find a disjoint family of balls B_1, \dots, B_N such that $3B_1, \dots, 3B_N$ covers K , and $\int_{B_i} |f(x)| dx \gtrsim_d 1$ for each i . Then

$$\int_K w(x) dx \leq \sum_{i=1}^N \int_{3B_i} w(x) dx$$

and so it suffices to show that

$$\int_{3B_i} w(x) dx \lesssim_d \int_{B_i} |f(x)| M w(x) dx.$$

But for $x \in B_i$, we have

$$M w(x) \gtrsim_d \frac{1}{|B_i|} \int_{3B_i} w(y) dy$$

from which the claim follows. \square

A simple corollary is a vector-valued generalization of the Hardy-Littlewood inequality.

Theorem 30.2. *If $1 < p, q < \infty$ and $\{f_n\}$ are any sequence of functions in $L^1_{loc}(\mathbf{R}^d)$, then*

$$\left\| \left(\sum_n |M f_n|^p \right)^{1/p} \right\|_{L^q(\mathbf{R}^d)} \lesssim_{d,p,q} \left\| \left(\sum_n |f_n|^p \right)^{1/p} \right\|_{L^q(\mathbf{R}^d)}$$

and

$$\left| \left\{ x : \left(\sum_n |M f_n(x)|^p \right)^{1/p} \geq \lambda \right\} \right| \lesssim_{d,p} \frac{1}{\lambda} \cdot \left\| \left(\sum_n |f_n(x)|^p \right)^{1/p} \right\|_{L^1(\mathbf{R}^d)}.$$

Proof. For $p = q$, the theorem follows from the standard Hardy-Littlewood maximal inequality. For $p < q$ we apply the equivalence between vector-valued bounds and weight bounds. To prove the remaining case, it suffices to prove the weak-type estimate for $p > 1$. By linearization, we may find radii $r_n(y)$ such that

$$|M f_n(y)| \leq \frac{1}{|B(y, r_n(y))|} \int_{\mathbf{R}^d} \psi_n(x, y) f_n(y) dy,$$

where $\psi_n(x, y)$ is a smooth bump function which equals one for $x \in B(y, r_n(y))$ and vanishes for $x \notin 2B(y, r_n(y))$. Thus it suffices to obtain a weak-type bound for the vector-valued operator

$$T(\{f_n\})(y) = \left\{ \int_{\mathbf{R}^d} \frac{1}{|B(y, r_n(y))|} \psi_n(x, y) f_n(x) dx \right\}.$$

This is a vector-valued kernel operator with kernel $K(x, y)$ the diagonal matrix with entry. TODO: SEE TAO. \square

Chapter 31

Bellman Function Methods

It is interesting to ask whether we can obtain bounds of the form

$$\|Mf\|_{L^p(\mathbf{R}^d)} \lesssim_{d,p} \|f\|_{L^p(\mathbf{R}^d)}$$

without employing any interpolation techniques. This is possible, though nontrivial. We begin with a Bellman function approach, which works best in the dyadic scheme, i.e. proving bounds on M_Δ .

The idea here is to perform an *induction on scales*, i.e. to induct on the complexity of the function f . For a fixed $f \in L^p(\mathbf{R}^d)$, our goal is to obtain bounds of the form

$$\left(\int |M_\Delta f(x)|^p dx \right)^{1/p} \lesssim \left(\int |f(x)|^p dx \right)^{1/p}$$

where the implicit constant is independent of p .

We begin by applying some monotone convergence arguments to simplify our analysis. For each $x \in \mathbf{R}^d$, $|M_\Delta f(x)| = \lim_{m \rightarrow -\infty} |M_{\geq m} f(x)|$, where $M_{\geq m}$ is the operator giving a maximal average over all dyadic cubes containing a point with sidelength exceeding 2^m , and the limit is monotone increasing. It follows that for any $f \in L^p(\mathbf{R}^d)$,

$$\|M_\Delta f\|_{L^p(\mathbf{R}^d)} = \lim_{m \rightarrow -\infty} \|M_{\geq m} f\|_{L^p(\mathbf{R}^d)}.$$

Thus if we can obtain a bound

$$\|M_{\geq m} f\|_{L^p(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

with a bound independent of m , we would obtain the required bound on M_Δ . But if we could obtain a bound

$$\|M_{\geq 0}f\|_{L^p(\mathbf{R}^d)} \lesssim \|f\|_{L^p(\mathbf{R}^d)}$$

for all $f \in L^p(\mathbf{R}^d)$, then a rescaling argument, using the fact that

$$M_{\geq m}f = \text{Dil}_{1/2^d} M_{\geq 0} \text{Dil}_{2^d} f$$

shows that we in fact have

$$\begin{aligned} \|M_{\geq m}f\|_{L^p(\mathbf{R}^d)} &= 2^{d/p} \|M_{\geq 0} \text{Dil}_{2^d} f\|_{L^p(\mathbf{R}^d)} \\ &\lesssim 2^{d/p} \|\text{Dil}_{2^d} f\|_{L^p(\mathbf{R}^d)} = \|f\|_{L^p(\mathbf{R}^d)}. \end{aligned}$$

Thus we need only concentrate on the operator $M_{\geq 0}$. Finally, we note we can *localize* our estimates. Given a function f supported on a dyadic cube Q with sidelength 2^n , and given $x \notin Q$, then there exists a smallest value $m_x > n$ such that x is contained in a dyadic cube with sidelength 2^{m_x} which also contains Q . It then follows that

$$(M_{\geq 0}f)(x) = \frac{\int_Q |f(y)| dy}{2^{dm_x}} = \frac{\|f\|_{L^1(Q)}}{2^{dm_x}}$$

For each $m > n$, if we set $E_m = \{x \in \mathbf{R}^d : m_x = m\}$, then E_m is contained in a dyadic cube of sidelength 2^m , so $|E_m| \leq 2^{dm}$. Thus we have

$$\begin{aligned} \|M_{\geq 0}f\|_{L^p(Q^c)} &= \left(\sum_{m=n+1}^{\infty} \|M_{\geq 0}f\|_{L^p(E_m)}^p \right)^{1/p} \\ &\leq \left(\sum_{m=n+1}^{\infty} \left(\|f\|_{L^1(Q)}^p / 2^{dpm} \right) 2^{dm} \right)^{1/p} \\ &\lesssim_{d,p} \|f\|_{L^1(Q)} 2^{dn(1/p-1)} = \|f\|_{L^1(Q)} |Q|^{1/p-1} \leq \|f\|_{L^p(Q)}. \end{aligned}$$

Thus, if we obtained the bound $\|M_{\geq 0}f\|_{L^p(Q)} \lesssim \|f\|_{L^p(Q)}$, then we would find

$$\|M_{\geq 0}f\|_{L^p(\mathbf{R}^d)} \leq \|M_{\geq 0}f\|_{L^p(Q)} + \|M_{\geq 0}f\|_{L^p(Q^c)} \lesssim \|f\|_{L^p(Q)}.$$

Thus if f is supported on a dyadic cube Q , it suffices to estimate $M_{\geq 0}f$ on the support of f . But by a final monotone convergence argument, it suffices to bound such functions, since given any n we can write $[-2^n, 2^n]$ as the almost disjoint union of 2^d sidelength 2^d dyadic cubes $Q_{n,1}, \dots, Q_{n,2^d}$. For any $f \in L^p(\mathbf{R}^d)$, we consider a pointwise limit $f = \lim_{n \rightarrow \infty} f_{n,1} + \dots + f_{n,2^d}$, where $f_{n,i}$ is equal to f restricted to $Q_{n,i}$, and the limit is monotone. We also have

$$M_{\geq 0}f = \lim_{n \rightarrow \infty} M_{\geq 0}f_{n,1} + \dots + M_{\geq 0}f_{n,2^d}.$$

where the limit is pointwise and monotone, so

$$\begin{aligned} \|M_{\geq 0}f\|_{L^p(\mathbf{R}^d)} &= \lim_{n \rightarrow \infty} \|M_{\geq 0}f_{n,1} + \dots + M_{\geq 0}f_{n,2^d}\|_{L^p(\mathbf{R}^d)} \\ &\lesssim \lim_{n \rightarrow \infty} \|f_{n,1}\|_{L^p(\mathbf{R}^d)} + \dots + \|f_{n,2^d}\|_{L^p(\mathbf{R}^d)} \lesssim 2^d \|f\|_{L^p(\mathbf{R}^d)}. \end{aligned}$$

Thus, after a technical reduction argument, we now show that we only have to establish a bound

$$\|M_{\geq 0}f\|_{L^p(Q)} \lesssim \|f\|_{L^p(Q)},$$

where $f \in L^p(Q)$, Q is a dyadic cube with sidelength ≥ 1 , and the implicit constant is independent of Q .

To carry out the inequality, we perform an *induction on scales*. For each $n \geq 0$, we let $C(n)$ denote the optimal constant such that for any function $f \in L^p(\mathbf{R}^d)$ supported on a dyadic cube Q of sidelength 2^n ,

$$\|M_{\geq 0}f\|_{L^p(Q)} \leq C(n) \cdot \|f\|_{L^p(Q)}.$$

If $n = 0$, the problem is trivial, since if Q is dyadic with sidelength 1 and $x \in Q$, then

$$M_{\geq 0}f = \int_Q |f(y)| \, dy$$

so $\|M_{\geq 0}f\|_{L^p(Q)} = \|f\|_{L^1(Q)}$, and $C(0) = 1$. Our goal is to show that $\sup_{n \geq 0} C(n) < \infty$. Given f supported on a cube Q with sidelength 2^n , the cube has 2^d children Q_1, \dots, Q_{2^d} with sidelength 2^{n-1} . If we decompose $f = f_1 + \dots + f_{2^d}$ onto these cubes, then by induction we know that

$$\|M_{\geq 0}f_i\|_{L^p(Q_i)} \leq C(n-1) \|f_i\|_{L^p(Q_i)}.$$

Now for $x \in Q_i$,

$$(M_{\geq 0}f)(x) = \max \left(M_{\geq 0}f_i(x), \int_Q |f(y)| dy \right).$$

Thus if $A = \int_Q |f(y)| dy$, then

$$\begin{aligned} \|M_{\geq 0}f\|_{L^p(Q)} &= \left(\|M_{\geq 0}f\|_{L^p(Q_1)}^p + \cdots + \|M_{\geq 0}f\|_{L^p(Q_{2^d})}^p \right)^{1/p} \\ &= \left(\|\max(M_{\geq 0}f_1, A)\|_{L^p(Q_1)}^p + \cdots + \|\max(M_{\geq 0}f_{2^d}, A)\|_{L^p(Q_{2^d})}^p \right)^{1/p} \end{aligned}$$

The bound $\max(M_{\geq 0}f_i, A) \leq M_{\geq 0}f_i + A$ gives

$$\|M_{\geq 0}f\|_{L^p(Q)} \leq C(n-1)\|f\|_{L^p(Q)} + 2^{d/p}|Q|^{1/p}A = (C(n-1) + 2^{d/p})\|f\|_{L^p(Q)}.$$

This gives $C(n) \leq C(n-1) + 2^{d/p}$, which is not enough to obtain a uniform bound. The idea here is to include more information in our induction hypothesis which gives control on $\max(M_{\geq 0}f_i, A)$. Since Q contains points not in Q_i , we need to treat A as an arbitrary quantity in our hypothesis.

To do this, we introduce *cost functions*. For each $A, B, D > 0$ and any integer $n \geq 0$, we let $V_n(A, B, D)$ be the optimal constant such that

$$\|\max(M_{\geq 0}f, A)^p\|_{L^p(Q)} \leq V_n(A, B, D)$$

For any function f supported on a dyadic cube Q with sidelength 2^n , with

$$\|f\|_{L^1(Q)} = B \quad \text{and} \quad \|f\|_{L^p(Q)} = D.$$

Our goal will be to show $V_n(A, B, D) \lesssim_p 2^{-dn/p}A + D$ which completes the proof. The role of B is subtle, but will soon become apparant. Of course, we have $\|f\|_{L^1(Q)} \leq 2^{dn(1-1/p)}\|f\|_{L^p(Q)}$, so we have $V_n(A, B, D) = -\infty$ unless $B \leq 2^{dn(1-1/p)}D$.

The recursive inequality gives an inequality for the values $V_n(A, B, D)$.
TODO: COMPLETE THIS PROOF.

Chapter 32

Maximal Averages Over Curves

32.1 Averages over a Parabola

Given any measurable function $f : \mathbf{R}^2 \rightarrow \mathbf{C}$ we can consider the maximal average

$$(Mf)(x, y) = \sup_{\varepsilon > 0} \frac{1}{2\varepsilon} \int_{-\varepsilon}^{\varepsilon} |f(x + t, y + t)| \, dt.$$

Thus Mf gives a maximal average over parabolas. Our goal is to show $\|Mf\|_{L^p(\mathbf{R}^d)} \lesssim_p \|f\|_{L^p(\mathbf{R}^d)}$ for $1 < p < \infty$.

It will be convenient to look at the operator

$$\tilde{M}f(x, y) = \sup_{\varepsilon > 0} \frac{1}{2\varepsilon} \int_{\varepsilon/2}^{\varepsilon} |f(x + t, y + t^2)| \, dt.$$

A dyadic decomposition shows that L^p bounds for \tilde{M} imply L^p bounds for M .

For each $k \in \mathbf{Z}$, let $\tilde{M}_k f(x, y) = 2^{-k} \int_{2^k}^{2^{k+1}} |f(x + t, y + t^2)| \, dt$. Rescaling shows that

$$\|\tilde{M}_k\|_{L^p(\mathbf{R}^2) \rightarrow L^p(\mathbf{R}^2)} = \|\tilde{M}_0\|_{L^p(\mathbf{R}^2) \rightarrow L^p(\mathbf{R}^2)}$$

so it suffices to focus on \tilde{M}_0 . The operator is translation invariant and therefore has a Fourier multiplier

$$\tilde{m}(\xi, \eta) = \int_1^2 e^{2\pi i(\xi t + \eta t^2)} \, dt.$$

Note that \tilde{m} is defined by an oscillatory integral with phase $\phi(t) = \xi t + \eta t^2$. We note that $\phi'(t) = \xi + 2\eta t$, so Van der Corput's lemma implies that for $|\xi| \geq 10|\eta|$,

$$|\tilde{m}(\xi, \eta)| \lesssim \frac{1}{|\xi|}.$$

Similarly, $\phi''(t) = 2\eta$, so we find

$$|\tilde{m}(\xi, \eta)| \lesssim \frac{1}{|\eta|^{1/2}}.$$

If $f \in L^2(\mathbf{R}^2)$ and \hat{f} is supported on the region

$$E_0 = \{(\xi, \eta) : |\eta| \geq 1 \text{ or } |\xi| \leq 1 \text{ and } |\eta| \geq 10\}$$

then $\|\tilde{m}\|_{L^\infty(E_0)} \lesssim 1$ and so

$$\|\tilde{M}_0 f\|_{L^2(\mathbf{R}^2)} = \|\tilde{m} \hat{f}\|_{L^2(\mathbf{R}^2)} \lesssim \|\hat{f}\|_{L^2(\mathbf{R}^2)} = \|f\|_{L^2(\mathbf{R}^2)}.$$

On the other hand, we can decompose $\mathbf{R}^2 - E_0$ into
suppose \hat{f} is supported on the region

$$E_1 = \{(\xi, \eta) : |\xi| \leq 1 \text{ and } |\eta| \leq 10\}.$$

Then the uncertainty principle implies that f is roughly constant on scales $|\Delta x| \leq 1$ and $|\Delta y| \leq 1/10$, which should imply good bounds for the maximal average. More precisely, \hat{f} is supported on the ellipsoid

$$\{(\xi, \eta) \in \mathbf{R}^2 : \xi^2/2 + \eta^2/20 \leq 1\}.$$

Thus the uncertainty principle implies that f is roughly constant on scales $|\Delta x|^2 \leq 1/2$ and $|\Delta y|^2 \leq 1/20$,

$$\phi(x) = \frac{1}{(1 + 2x^2 + 20y^2)^N}$$

Part IV

**Linear Partial Differential
Equations**

Here we study the techniques of the *constant coefficient* linear partial differential equations

$$Lf(x) = \sum a_\alpha D^\alpha f(x)$$

and the *non constant coefficient* partial differential equations

$$Lf(x) = \sum a_\alpha(x) D^\alpha f(x)$$

The central problems of the theory are *existence* and *regularity*: Given f , does a solution u to the equation $Lu = f$ exist, and if so, how does the regularity of f relate to the regularity of u . We often phrase the regularity problems in terms of function spaces. Let $\{X^s\}$ be a family of function spaces, indexed by a parameter $s \in \mathbf{R}$, measuring the *regularity* of functions in these spaces. Given a linear partial differential operator L , a regularity result might say that for some s and t , if $Lu = f$, where $f \in X^s$, then u in X^t . Such results often require one to use the special properties of the operator L ; general theories are rarely available, except for certain very special families of partial differential equations, especially the *elliptic partial differential equations*.

Chapter 33

Green's Functions

Let Ω be a bounded, open set on \mathbf{R}^n . Recall Green's second identity,

$$\int_{\Omega} (u \Delta v - v \Delta u) = \int_{\partial \Omega} u \frac{\partial v}{\partial \eta} - v \frac{\partial u}{\partial \eta}.$$

We introduce the quantity

$$u_x(y) = \frac{1}{|x - y|^{n-2}},$$

which is Newton's gravitational potential, or Coulomb's electrostatic potential given a point mass at the point x . A simple computation shows that $\Delta u_x(y) = 0$ for $y \neq x$, perhaps using the fact that if $u(x)$ is a radial function, with $u(x) = f(r)$ for $r = |x|$, then

$$\Delta u(x) = f''(r) + \frac{n-1}{r} f'(r).$$

Plugging this function into Green's identity, if $v \in C^2(\Omega)$ is a Harmonic function on a domain Ω , continuous on the closure of Ω , then for any x , applying Green's identity on the domain $\Omega_{x,\varepsilon}$, obtained from Ω by removing the ball B of radius ε centered at x , where ε is small enough so that the entire ball is contained in Ω , we find that

$$\begin{aligned} 0 &= \int_{\partial \Omega_{x,\varepsilon}} \left(u_x \frac{\partial v}{\partial \eta} - v \frac{\partial u_x}{\partial \eta} \right) \\ &= \int_{\partial \Omega} \left(u_x \frac{\partial v}{\partial \eta} - v \frac{\partial u_x}{\partial \eta} \right) - \int_{\partial B} u_x \frac{\partial v}{\partial \eta} - v \frac{\partial u_x}{\partial \eta}. \end{aligned}$$

Now for $y \in \partial B$, with $y = x + z$, $\eta(y) = \varepsilon^{-1}z$. And

$$\nabla u_x(y) = -(n-2) \frac{z}{|z|^n},$$

so

$$\int_{\partial B} u_x \frac{\partial v}{\partial \eta} - v \frac{\partial u_x}{\partial \eta} = \int_{|z|=1} \varepsilon \cdot \nabla v(x + \varepsilon z) \cdot z + (n-2)v(x + \varepsilon z).$$

Applying a Taylor series shows that as $\varepsilon \rightarrow 0$, this quantity converges to $(n-2)C_d v(x)$, where C_d is the surface area of S^{d-1} . Thus we conclude that

$$v(x) = \frac{-1}{C_d(n-2)} \int_{\partial \Omega} \left(u_x \frac{\partial v}{\partial \eta} - v \frac{\partial u_x}{\partial \eta} \right)$$

Thus v is determined by its boundary values, and boundary derivatives in the direction normal to Ω . To get rid of these normal boundary derivatives, we suppose we can compute a function h_x , which is harmonic in Ω , and equal to u_x on $\partial \Omega$. Then we can apply Green's second identity again, which yields

$$\begin{aligned} 0 &= \int_{\partial \Omega} v \frac{\partial h_x}{\partial \eta} - \frac{\partial v}{\partial \eta} G \\ &= \int_{\partial \Omega} v \frac{\partial h_x}{\partial \eta} - \frac{\partial v}{\partial \eta} u_x. \end{aligned}$$

Thus we conclude that

$$v(x) = \frac{-1}{C_d(n-2)} \int_{\partial \Omega} v \left(\frac{\partial h_x}{\partial \eta} - \frac{\partial u_x}{\partial \eta} \right).$$

The function

$$g_x(y) = \frac{-1}{C_d(n-2)} (h_x(y) - u_x(y))$$

is called the *Green's function* on the domain Ω . It solves the equation $\Delta g_x = \delta_x$ on Ω , while vanishing on the boundary. We have calculated this by virtue of the fact that for any harmonic function v on Ω ,

$$v(x) = \int_{\partial \Omega} \frac{\partial g_x}{\partial \eta} v \, dy.$$

Thus v is uniquely determined by its boundary values. What's more, we now have a natural operator which, given some sufficiently boundary conditions on $\partial \Omega$, allows us to find a unique solution to Laplace's equation on Ω .

Chapter 34

Elliptic Equations

Consider a partial differential operator $Lf(x) = \sum a_\alpha(x)D^\alpha f(x)$, where $a_\alpha \in C^\infty(\mathbf{R}^n)$. Associated with this operator is its symbol $a(x, \xi) = \sum a_\alpha(x)\xi^\alpha$, and if L has order m , its principle symbol is $a_m(x, \xi) = \sum_{|\alpha|=m} a_\alpha(x)\xi^\alpha$. We say L is *elliptic* near x_0 if $a_m(x_0, \xi) = 0$ only when $\xi = 0$. This is equivalent to assuming that there exists $M > 0$, and two constants c_1 and c_2 and a small neighborhood U of x_0 such that $c_1|\xi|^m \leq |a(x, \xi)| \leq c_2|\xi|^m$ for $|\xi| \geq M$.

Elliptic operators have a near complete theory, because they are *invertible* modulo smoothing operators. More precisely

Chapter 35

Dispersive Equations

A partial differential equation is *dispersive* if wave-like solutions to the partial differential equation with differing frequencies travel in different directions and speeds, so that linear combinations of such solutions tend to ‘spread out in space’ over time. This phenomenon presents itself in various *dispersive estimates* for such partial differential equations.

The simplest example where dispersive phenomena emerges is in the analysis of a constant-coefficient hyperbolic first order system of the form

$$\partial_t u = Lu$$

where $u : V \times \mathbf{R} \rightarrow \mathbf{R}$ for some finite dimensional Hilbert space V , and where $Lu = \sum c_\alpha D_x^\alpha u$ for some $c_\alpha \in \text{End}(V)$. In order to assume the differential equation is hyperbolic, and thus well-posed, we assume that L is formally skew-adjoint, i.e. that for all compactly supported, smooth functions $u : V \rightarrow \mathbf{R}$ and $v : V \rightarrow \mathbf{R}$,

$$\langle Lu, v \rangle = -\langle u, -Lv \rangle.$$

If we write $L = 2\pi i h(D_x)$ for some polynomial $h : \mathbf{R}^d \rightarrow \text{End}(V)$, then this is equivalent to the coefficients of h being self-adjoint. The polynomial h is called the *dispersion relation* of the equation.

Given such a PDE, we have *planar wave solutions* of the form

$$u(x, t) = e^{2\pi i(\xi \cdot x + \omega t)}$$

precisely when $\omega = h(\xi)$. Thus the dispersion relation relates the *wave number* ξ of a planar wave with the *angular frequency* ω of the planar waves

involved. In particular, looking at the phase $\xi \cdot x + \omega t$ and trying to solve in the x -variable, we see the wave will look like it is travelling through space at a *phase velocity*

$$v_p(\xi) = -\frac{\xi}{|\xi|} \frac{h(\xi)}{|\xi|}$$

In particular, the phase speed is $|h(\xi)|/|\xi|$.

More generally, we can consider superposition solutions using the Fourier inversion formula, i.e. writing

$$u(x, t) = \int \widehat{u_0}(\xi) e^{2\pi i[\xi \cdot x + h(\xi)t]} d\xi$$

Taking $u_0(\xi) = e^{2\pi i \xi \cdot x}$ gives the planar wave solutions above. A *wave packet* solution is obtained by taking u_0 to be supported on a small region of phase-space, i.e. localized to a particular position in space, and a particular frequency. One example localized near (x_0, ξ_0) would be of the form

$$u_0(x) = e^{2\pi i \xi_0 \cdot x} \phi\left(\frac{x - x_0}{\delta}\right),$$

where ϕ is a Schwartz function whose Fourier transform ψ is smooth and compactly supported. Fourier inversion tells us that

$$u(x, t) = \int \psi(\xi) e^{2\pi i[(\xi_0 + \xi) \cdot x + h(\xi_0 + \xi)t - \xi \cdot x_0]} d\xi.$$

The principle of nonstationary phase tells us that u has spatial concentration near $x = x_0 - \nabla h(\xi_0)t + O(t)$. Thus the phase envelope of u can be seen to be travelling starting at x_0 , and travelling at a *group velocity* $-\nabla h(\xi_0)$, though the slight differences in phase velocities for individual planar waves which give u as a superposition can lead to the phase envelope spreading apart in time.

Example. For $\omega_0 \in \mathbf{R}$, we can consider the phase rotation equation $\partial_t u = 2\pi i \omega_0 u$, which can be explicitly solved by writing

$$u(x, t) = e^{2\pi i \omega_0 t} u_0(x).$$

Thus solutions have the envelope defined by u_0 , and oscillate in time at the angular frequency ω_0 . In particular, we have planar-wave solutions of the form

$$e^{2\pi i(\xi \cdot x + \omega_0 t)}.$$

The phase velocity here is $-\omega_0/\xi$, i.e. planar waves with a high spatial frequency will appear to travel much more slowly in time. The group velocity, however, is 0, so a superposition of waves with similar spatial frequency will remain in the same position spatially.

Example. Consider the transport equation

$$\partial_t u = -v_0 \cdot \nabla_x u$$

for some $v_0 \in \mathbf{R}^d$. The solution to this equation is given by

$$u(x, t) = u_0(x - v_0 t).$$

The planar-wave solutions are of the form

$$e^{2\pi i(\xi \cdot x - (\xi \cdot v_0)t)}.$$

The dispersion relation is $h(\xi) = -v_0 \cdot \xi$. The phase velocity is $v_p(\xi) = \xi(v_0 \cdot \xi)/|\xi|^2$; if ξ is aligned in the direction v_0 , then the phase velocity will be v_0 , and if ξ is orthogonal to v_0 , the phase velocity will be zero, since the solution will be stationary in time. The group velocity is v_0 , i.e. a wave packet will always roughly move in the direction v_0 .

Example. In the free Schrödinger equation $i\partial_t u = -(\hbar/8\pi^2 m)\Delta u$, the dispersion relation is $h(\xi) = -(\hbar/2m)|\xi|^2$, and the phase velocity is $v_p(\xi) = (\hbar/2m)\xi$. The group velocity here is $v_g(\xi) = (\hbar/m)\xi$. Thus we see that the phase of a wave packet travels at half its group velocity. More interesting is the group velocity equation, which in physics is used as a derivation of De Broglie's law $mv_g = \hbar\xi$, i.e. that the frequency of a wave packet is tied to its velocity in time.

Example. The one-dimensional Airy equation is $\partial_t u = (2\pi)^{-3}\partial_x^3 u$. It has dispersion relation $h(\xi) = |\xi|^3$. The phase velocity is $v_p(\xi) = \xi|\xi|$, and the group velocity is $v_g(\xi) = 3\xi|\xi| = 3v_p(\xi)$.

For higher order hyperbolic equations $P(\partial_t, D_x)$, say of order m , the dispersion relation h is multi-valued, equal to the values $h(\xi)$ such that $P(ih(\xi), \xi) = 0$.

Example. Consider the wave equation $\square u = 0$, where \square is the d'Alembertian operator $(2\pi)^{-2}\Delta - \partial_t^2$. The resulting polynomial is $P(\tau, \xi) = -(\xi^2 + \tau^2)$. The dispersion relation here is thus given by the equation $(\xi^2 + (ih(\xi))^2) = 0$, i.e. we have $h(\xi) = \pm\xi$. The phase velocities are $v_p(\xi) = \pm\xi/|\xi|$, and the group velocities are also $v_g(\xi) = \pm\xi/|\xi|$.

Example. The Klein-Gordon equation $\square u = u$ describes the movement of electrons where relativity is relevant. It is given by the polynomial $P(\tau, \xi) = -(|\xi|^2 + \tau^2 + 1)$, and has dispersion relation $h(\xi) = \pm\sqrt{1 + |\xi|^2}$. For large ξ , we thus see waves here behave like the wave equation, though for small ξ , waves behave like the transport equation.

Our goal is to study *dispersive equations*, i.e. equations where wave packet solutions tend to spread apart over time. Since the velocity of a wave packet travelling at frequency ξ_0 is $\nabla h(\xi_0)$, wave packets near ξ_0 will travel at different frequencies if the Hessian $H(\xi_0)$ of h is invertible. If this is true, the equation is called *fully dispersive*, like the Schrödinger equation and Airy equation. In dimension one, for the wave equation, the Hessian is zero, so in this case the wave equation is very non-dispersive. But in d dimensions, we see the Hessian has rank $d - 1$, so we at least have ‘some dispersion’, but not in the radial direction.

35.1 Strichartz Estimates

Our goal is to obtain *space-time estimates* of the form $\|u\|_{L_t^q L_x^r}$ on solutions u to a dispersive partial differential equation. Let’s start with a dispersive equation of the form $\partial_t u = Lu$. let’s look at pointwise estimates. Let $T_t = e^{2\pi i t L}$.

$$T_t f(x) = \int e^{2\pi i [\xi \cdot x + h(\xi)t]} \hat{f}(\xi) d\xi.$$

This is an operator with Fourier multiplier $m_t(\xi) = e^{2\pi i h(\xi)t}$. To understand these operators, we apply Littlewood-Paley theory, writing

$$T_t = T_{t,0} + T_{t,1} + \dots$$

where $T_{t,l}$ is a Fourier multiplier with symbol $m_{t,l}(\xi) = \psi_l(\xi) e^{2\pi i h(\xi)t}$. Applying stationary phase, if $K_{t,l}$ is the inverse Fourier transform of $m_{t,l}$, then we have TODO

As $t \rightarrow \infty$, we can see Dispersion via the norm of the operators T_t .

Theorem 35.1. Suppose that T has a dispersion relation of the form $\|T_t\|_{L^q}$

Proof. Let T map functions on \mathbf{R}^d to functions on $\mathbf{R}^d \times \mathbf{R}$, given by

$$Tf(x, t) = e^{2\pi i t L} f(x) = T_t f(x).$$

The formal adjoint is T^* mapping functions on $\mathbf{R}^d \times \mathbf{R}$ to functions on \mathbf{R}^d by

$$T^*F(x) = \int e^{-2\pi i t L} F(x, t) dt = \int T_{-t} F(x, t) dt.$$

Thus

$$TT^*F(x, t) = \int e^{2\pi i(t-s)L} F(x, s) ds = \int T_{t-s} F_s(x) ds.$$

We have dispersive estimate

$$\|e^{2\pi i(t-s)L} f\|_{L_x^r} \lesssim |t-s|^{-(d/2-d/r)} \|f\|_{L_x^{r^*}}$$

for $1/r + 1/r^* = 1$. Thus

$$\begin{aligned} \|TT^*F\|_{L_t^q L_x^r} &\leq \left\| \int \|T_{t-s} F_s\|_{L_x^r} ds \right\|_{L_t^q} \\ &\lesssim \left\| \int \frac{\|F_s\|_{L_x^{r^*}}}{|t-s|^{d/2-d/r}} ds \right\|_{L_t^q}. \end{aligned}$$

Now applying Hardy-Littlewood-Sobolev gives the required bound, i.e. that

$$\|TT^*F\|_{L_t^q L_x^r} \lesssim \left\| \int \frac{\|F_s\|_{L_x^{r^*}}}{|t-s|^{d/2-d/r}} ds \right\|_{L_t^q} \lesssim \|F\|_{L_s^q L_x^r}.$$

The TT^* argument gives that

$$\|Tf\|_{L_t^q L_x^r} \lesssim \|f\|_{L_x^2}.$$

□

Part V

Geometry of Eigenfunctions

In these parts of the notes, we study *eigenfunctions* for the Laplacian on compact manifolds and bounded domains. This is the classical scenario under which harmonic analysis was first studied. For instance, in the study of the wave and heat equations

$$\left(\frac{\partial}{\partial t^2} - \Delta\right)u = 0 \quad \text{and} \quad \left(\frac{\partial}{\partial t} - \Delta\right)u = 0$$

on a vibrating membrane Ω fixed at the boundary, i.e. with $u(x, t) = 0$ for $x \in \partial\Omega$. If φ_λ solves the *Helmholtz equation* $\Delta\varphi_\lambda = -\lambda^2\varphi_\lambda$, vanishing on $\partial\Omega$, then a family of solutions to the wave equation is given by

$$u_\lambda(t, x) = \cos(\lambda t + \phi)\varphi_\lambda(x).$$

In this scenario, φ_λ are the *modes* of the wave equation, giving the profiles under which the functions u_λ oscillate. Studying how these eigenfunctions interact with one another allows us to understand more general solutions to the wave equation.

Another application of these eigenfunctions occurs in the study of the Schrödinger equation

$$\left(-\frac{\hbar^2}{2}\Delta + V\right)\psi = E\psi,$$

which describes the evolution of a wave function ψ under a potential V . The case $V = 0$ and $E = E_0$ a constant value (the free Schrödinger equation) is most clearly connected with the Helmholtz equation, i.e. with λ proportional to E_0/\hbar , and E_0 for which a solution exists are the *energy states* of the system. An intuition is that high energy states behave like the ‘classical versions’ of the physical systems that the Schrödinger equation models a quantum version of, i.e. to the classical dynamics of the Hamiltonian system given by the equation $H(x, \xi) = V(x) + |\xi|^2/2$.

There are various properties one can study of the eigenfunctions $\{\varphi_\lambda\}$. For instance, one could study their L^p norms, or other norms. One could also study the geometric properties of

The study of such eigenfunctions is closely related to *semiclassical analysis*, i.e. understanding the relation between classical and quantum mechanics as the scale under which a problem is being studied changes.

Another application occurs in the study of the Schrödinger equation

$$\left(-\frac{h^2}{2}\Delta + V\right)\psi = E\psi$$

where V stands for multiplication by a potential $V \in C^\infty(M)$, and h is Planck's constant, a small quantity. If $h = 1/\lambda$, $V = 0$, and $E = 1$, we obtain the wave equation.

Chapter 36

Self-Adjoint, Elliptic Operators on a Compact Manifold

Let M^d be a compact manifold. Given a scalar density $d\omega$ on a manifold M , we can consider $L^2(M)$ as a Hilbert space induced by the inner product

$$\langle u, v \rangle = \int u(x) \overline{v(x)} d\omega.$$

Thus given a scalar density we can consider the adjoint of a pseudodifferential operator on M by considering the operator as an unbounded operator on $L^2(M)$ with domain $H_c^k(M)$, and in particular, we can consider *self-adjoint unbounded* operators on M ; we are being somewhat lax with an overloaded set of terminology here, by a *self-adjoint operator* we mean an operator from $\mathcal{D}(M)$ to $\mathcal{D}(M)^*$, which is formally self adjoint on $\mathcal{D}(M)$, i.e. such that for all f, g in $\mathcal{D}(M)$,

$$\int_M T f \cdot \bar{g} d\omega = \int_M f \cdot \overline{T g} d\omega.$$

By a *self-adjoint unbounded operator*, we mean an operator from $\mathcal{D}(M)$ to $\mathcal{D}(M)^*$ which is essentially self-adjoint as an unbounded operator on $L^2(M)$, in the sense of the theory of self-adjointness. This is slightly stronger than being a self-adjoint operator, though we will soon see these properties are equivalent for an elliptic pseudodifferential operator. We now analyze the theory of self-adjoint, elliptic pseudodifferential operators on a compact manifold, in particular, proving the existence of the eigenfunction

expansions of such operators. We begin by showing that such operators are automatically self-adjoint in the sense of an unbounded operator.

Theorem 36.1. *Suppose T is a self-adjoint, elliptic pseudodifferential operator of order t on a compact manifold M . Then T is a closed, self-adjoint unbounded operator.*

Proof. Let us begin by showing T is closed. We begin by noting that T is a continuous operator from $\mathcal{D}(M)^*$ to itself. We desire to show

$$\{(u, v) \in H^t(M) \times L^2(M) : Tu = v\}$$

is closed in $L^2(M) \times L^2(M)$, and so T is a closed operator on $L^2(M)$ when viewed with domain $H^t(M)$. If $\{u_i\}$ is a sequence in $H^t(M)$ converging in the $L^2(M)$ norm to some $u \in L^2(M)$, and $\{Tu_i\}$ converges in the $L^2(M)$ norm to some $v \in L^2(M)$, then $\{u_i\}$ converges distributionally to u , and so $Tu = v$ by the distributional continuity of T . Applying elliptic regularity, since $v \in L^2(M)$, we conclude that $u \in H^t(M)$. Thus we have shown the set above is closed.

Next, we show that T is a self-adjoint unbounded operator. Because T is continuous from $H^t(M)$ to $L^2(M)$, it is simple to see via a density argument that on $L^2(M)$, we have

$$\langle Tu, v \rangle = \langle u, Tv \rangle.$$

Thus to show T is self-adjoint as an unbounded operator, it suffices to show that if $v \in L^2(M)$ induces an inequality of the form

$$|\langle Tu, v \rangle| \lesssim \|u\|_{L^2(M)}$$

for all $u \in H^t(M)$, then $v \in H^t(M)$. But this again follows by elliptic regularity. Indeed, the inequality then holds for $u \in C^\infty(M)$, and the distributional theory thus implies that for such u ,

$$\langle Tu, v \rangle = \langle u, Tv \rangle.$$

Thus we have an inequality of the form

$$|\langle u, Tv \rangle| \lesssim \|u\|_{L^2(M)}$$

for all $u \in C^\infty(M)$, and a density argument thus shows that $Tv \in L^2(M)$, so that $v \in H^t(M)$. \square

Now let T be a classical, self-adjoint pseudodifferential operator of order t . It follows from the calculus of pseudodifferential operators that its principal symbol is real-valued. If T is self-adjoint and also *elliptic*, and M is connected and has dimension bigger than one, it follows that the principal symbol is either positive or negative. By swapping T out with its negation, we might as well assume that the principal symbol is positive. We will begin by consider the case $t = 1$, though the general case will reduce to this question. Given that the principal symbol is positive, we might imagine that T is ‘positive-definite’ up to first order, an intuition which leads to the following lemma.

Lemma 36.2. *Let T be a classical self-adjoint elliptic pseudodifferential operator of order 1 on a manifold M . Then there exists a classical self-adjoint elliptic pseudodifferential operator S of order $1/2$ such that $T - S^*S$ is a smoothing operator.*

Proof. We proceed as in the proof that $\sqrt{-\Delta}$ is a classical pseudodifferential operator. Let S_0 be a classical self-adjoint pseudodifferential operator of order $1/2$ with principal symbol $b_0(x, \xi) = a(x, \xi)^{1/2}$, then $S_0^*S_0$ is positive-semidefinite, and $T - S_0^*S_0$ is a classical pseudodifferential operator of order zero. Let r_1 denote the principal symbol of this operator. More generally, let us suppose that $T - (S_0 + \cdots + S_n)^*(S_0 + \cdots + S_n)$ is a classical pseudodifferential operator of order $-n$, with principal symbol r_n . If we consider a classical pseudodifferential operator S_{n+1} of order $1/2 - (n + 1)$ with principal symbol $b_{n+1}(x, \xi) = r_n(x, \xi)/2b_0(x, \xi)$, then $T - (S_0 + \cdots + S_{n+1})^*(S_0 + \cdots + S_{n+1})$ is a classical pseudodifferential operator of order $-(n + 1)$. Thus if we choose a self-adjoint operator S of order $1/2$ such that $S \sim \sum_{i=0}^{\infty} S_i$, then $T - S^*S$ is smoothing. \square

Lemma 36.3. *Suppose T is a classical self-adjoint elliptic pseudodifferential operator of order 1 with a non-negative principal symbol on a manifold M . For any compact subset K of M , there exists $\lambda_0 > 0$ such that for $\lambda \geq \lambda_0$, and $u \in H^1(M)$ supported on K ,*

$$\langle (T + \lambda_0)u, u \rangle \sim \|u\|_{H^{1/2}(M)}^2.$$

In particular, if M is compact, one can do this analysis globally.

Proof. Without loss of generality, we may assume that the support of the kernel of T is compact. Choose a classical, self-adjoint, elliptic pseudodifferential operator S of order $1/2$ such that $T - S^*S$ is smoothing. Then for

$u \in H_c^1(M)$, there exists $\gamma_0 > 0$ such that

$$\left| \langle Tu, u \rangle - \|Su\|_{L^2(M)}^2 \right| = \left| \int_M (T - S^*S)u(x) \bar{u}(x) dx \right| \leq \gamma_0 \|u\|_{L^2(M)}^2.$$

Thus if $\gamma > \gamma_0$,

$$\langle (T + \gamma)u, u \rangle = \langle Tu, u \rangle + \gamma \|u\|_{L^2(M)}^2 \geq \|Su\|_{L^2(M)}^2 + (\gamma - \gamma_0) \|u\|_{L^2(M)}^2.$$

Thus we conclude that $T + \gamma$ is positive definite. Since S has principal symbol $a(x, \xi)^{1/2}$, it is elliptic, and so for $u \in H^{1/2}(M)$, if $\gamma > \gamma_0$.

$$\begin{aligned} \|u\|_{H^{1/2}(M)}^2 &\lesssim \|Su\|_{L^2(M)}^2 + \|u\|_{L^2(M)}^2 \\ &\lesssim |\langle Tu, u \rangle| + \|u\|_{L^2(M)}^2 \\ &= |\langle (T + \gamma)u, u \rangle - \langle \gamma u, u \rangle| + \|u\|_{L^2(M)}^2 \\ &\leq \langle (T + \gamma)u, u \rangle + \langle \gamma u, u \rangle + \|u\|_{L^2(M)}^2 \\ &\leq \langle (T + 2\gamma + 1)u, u \rangle. \end{aligned}$$

Thus if $\lambda_0 > 2\gamma_0 + 1$, then

$$\langle (T + \lambda_0)u, u \rangle \lesssim \|Su\|_{L^2(M)} + \|u\|_{L^2(M)} \lesssim \|u\|_{H^{1/2}(M)}. \quad \square$$

Theorem 36.4. *Suppose T is a classical, self-adjoint pseudodifferential operator of order one on a compact manifold M , with a non-negative principal symbol. Then for suitably large $\lambda_0 > 0$,*

- *For all $k > 0$, $(T + \lambda_0)^k$ is an isomorphism from $H^k(M)$ to $L^2(M)$.*
- *There exists an increasing family of real numbers $\{\lambda_i\}$ with $\lambda_i \rightarrow \infty$, and an orthogonal basis $\{e_i\}$ for $L^2(M)$ consisting solely of elements of $C^\infty(M)$, such that $Te_i = \lambda_i e_i$ for each $i > 0$. The functions $\{e_i\}$ satisfy*

$$\|e_i\|_{H^k(M)} \sim_k (\lambda_0 + \lambda_i)^k.$$

- *If $u \in H^k(M)$, then $\langle u, e_i \rangle \lesssim_k \|u\|_{H^k(M)} (\lambda_0 + \lambda_i)^{-k}$.*

Proof. The last result implies that $T + \lambda_0$ is an injective operator from $H^1(M)$ to $L^2(M)$ for sufficiently large $\lambda_0 > 0$. We have already seen that,

since M is compact, $T + \lambda_0$ is a closed, self-adjoint unbounded operator on $L^2(M)$. But the theory of unbounded operators thus implies that because $T + \lambda_0$ is injective, it has dense image in $L^2(M)$. Thus we can define its inverse $R = (T + \lambda_0)^{-1}$, which is a densely defined continuous operator in $L^2(M)$, and thus extends to a continuous operator on all of $L^2(M)$. Because $T + \lambda_0$ is positive-definite, i.e. $\langle (T + \lambda_0)u, u \rangle > 0$ for all non-zero $u \in H^1(M)$, density implies that $\langle Rv, v \rangle > 0$ for all $v \neq 0$. Thus R is positive definite. If $u \in H^1(M)$, and $(T + \lambda_0)u = v$, i.e. $u = Rv$, then ellipticity implies that

$$\|Rv\|_{H^1(M)} \lesssim \|v\|_{L^2(M)} + \|Rv\|_{L^2(M)} \lesssim \|v\|_{L^2(M)}.$$

Conversely, continuity of pseudodifferential operators implies that

$$\|v\|_{L^2(M)} = \|(T + \lambda_0)u\|_{L^2(M)} \lesssim \|u\|_{H^1(M)}.$$

Thus density implies that R is an *isomorphism* from $L^2(M)$ to $H^1(M)$. In particular we note that that $T + \lambda_0$ is an isomorphism from $H^1(M)$ to $L^2(M)$, i.e. it was surjective in the first place rather than merely having dense image. Moving on, The Rellich-Kondrachov theorem thus implies that R is a *compact*, positive definite operator on $L^2(M)$. Thus the spectral analysis of such operators implies that there exists an orthogonal basis $\{e_i\}$ and a family of positive, decreasing eigenvalues $\{\gamma_i\}$ such that if E_i is orthogonal projection onto the span of e_i , then $(T + \lambda_0)^{-1} = \sum_i \gamma_i^{-1} E_i$. If $\lambda_i = \gamma_i - \lambda_0$, then this means that $T = \sum_i \lambda_i E_i$.

Applying elliptic regularity, since $(T + \lambda_0)e_i = \gamma_i e_i$ lies in $L^2(M)$, we conclude that $e_i \in H^1(M)$. But iterating this argument gives that $e_i \in H^N(M)$ for any $N > 0$. Applying the Sobolev embedding theorem thus shows that $e_i \in C^\infty(M)$ for each i .

A simple generalization of the argument above (the spectral theory, ellipticity, and so on) shows that $(T + \lambda_0)^k$ is actually an isomorphism from $H^k(M)$ to $L^2(M)$ for each $k > 0$. We thus find that for any $u \in H^k(M)$,

$$\|u\|_{H^k(M)}^2 \sim_k \|(T + \lambda_0)^k u\|_{L^2(M)}^2 = \sum (\lambda_i + \lambda_0)^{2k} |\langle f, e_i \rangle|^2.$$

Thus we find that

$$|\langle u, e_i \rangle| \lesssim_k \|u\|_{H^k(M)} |\lambda_i + \lambda_0|^{-k} \lesssim_k \|u\|_{H^k(M)} |1 + \lambda_0|^{-k}. \quad \square$$

The following, known as *Hörmander's square root trick*, allows us to extend the theory above to elliptic, self-adjoint operators of arbitrary order.

Theorem 36.5. *Fix integers p and q with $q \neq 0$, and set $t = p/q$. Suppose T is a classical, positive-definite, elliptic pseudodifferential operator of order t on a compact manifold M . Then the operator $T^{q/p}$, defined using the spectral functional calculus, is a classical, positive-definite, elliptic pseudodifferential operator of order 1, with principal symbol $a(x, \xi)^{q/p}$.*

Proof. The case $t = -1$ is easily seen to be true, since the assumptions imply T is invertible, and its inverse is a pseudodifferential operator modulo smoothing. Thus, without loss of generality, we may assume that $q = 1$, and that $p > 0$. We can certainly construct an operator $S \in \Psi_{\text{cl}}^1(M)$ which is self-adjoint, with principal symbol $a(x, \xi)^{1/p}$, such that $S^p - T$ is a smoothing operator. Applying the theory above, we see that S has an eigenvalue decomposition, with eigenvectors lying in $C^\infty(M)$, and only finitely many eigenvalues being negative. Thus, modulo a smoothing operator, i.e. a finite sum of the projection operators $\{E_i\}$, we may actually replace S with a positive-definite operator. We claim that $S - T^{1/p}$ is also a smoothing operator, which would complete the proof since the principal symbol of S is equal to $a(x, \xi)^{q/p}$. Since S and T are both positive-definite, they are injective, and so by the spectral calculus of unbounded operators, T^{-1} and S^{-1} is a well-defined bounded, positive-definite operator on $L^2(M)$. If γ is a contour in the complex plane which winds around each element of $\sigma(T^{-1})$ exactly once, then we find using the holomorphic functional calculus that

$$T^{1/p} = \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} (z - T^{-1})^{-1} dz$$

and

$$S = \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} (z - S^{-p})^{-1} dz$$

Now S^{-p} is a pseudodifferential operator of order $-p$. Since $S^p - T$ is a smoothing operator, the composition calculus tells us that so is $T^{-1}(S^p - T)S^{-p} = T^{-1} - S^{-p}$. Write $S^{-p} = T^{-1} - R$. Then

$$\begin{aligned} S - T^{1/p} &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} ((z - T^{-1} + R)^{-1} - (z - T^{-1})^{-1}) \\ &= \frac{1}{2\pi i} \int_{\gamma} z^{-1/p} ((z - T^{-1} + R)^{-1} R (z - T^{-1})^{-1}). \end{aligned}$$

This is an integral over a compact curve over a continuous family of smoothing operators, which is therefore a smoothing operator. \square

Remark. As above, finding an operator S of order $t/2$ such that $T - S^*S$ is smoothing allows one to show that if T is a classical, elliptic, self-adjoint pseudodifferential operator of order t , then $T + \lambda_0$ is positive definite for suitably large $\lambda_0 > 0$, so that we may apply the theorem above to a wider situation.

Using the trick, we quickly extend the theory above to general operators of this form.

Theorem 36.6. *Suppose T is a classical, self-adjoint, elliptic pseudodifferential operator of order $t = p/q$ on a compact manifold M . Then for sufficiently large $\lambda_0 > 0$,*

- $(T + \lambda_0)^k$ is an isomorphism from $H^{tk}(M)$ to $L^2(M)$ for all $k > 0$.
- There is a basis $\{e_i\}$ for $L^2(M)$ consisting of elements of $C^\infty(M)$ and an increasing sequence of positive real numbers $\{\lambda_i\}$ with $\lambda_i \rightarrow \infty$, such that $Te_i = (\lambda_0 + \lambda_i)^t e_i$. The functions $\{e_i\}$ satisfy

$$\|e_i\|_{H^k(M)} \sim_k (\lambda_0 + |\lambda_i|)^k$$

- If $u \in H^k(M)$, then $|\langle u, e_i \rangle| \lesssim_k \|u\|_{H^k(M)} (\lambda_0 + |\lambda_i|)^{-k}$.

The special case of this result, where Δ is the Laplace-Beltrami operator on a compact Riemannian manifold, was first conjectured by Ohm and Raleigh, and proved by Hilbert in 1904.

Example. Let M be a compact Riemannian manifold. Then the Laplace-Beltrami operator Δ is a classical pseudodifferential operator of order two on M , and $-\Delta$ is formally positive-semidefinite. Thus we can see λ_0 above to be any positive number, which gives that there exists a discrete subset $\Lambda(M)$ of $[0, \infty)$, and a family of finite dimensional subspaces E_λ of $C^\infty(M)$ for each $\lambda \in \Lambda(M)$, such that $\Delta e_\lambda = -\lambda^2 e_\lambda$ for each $e_\lambda \in E_\lambda$. If $u \in H^k(\mathbf{T}^n)$ and $e_\lambda \in E_\lambda$, then

$$|\langle u, e_\lambda \rangle| \lesssim_{\varepsilon, k} \|e_\lambda\|_{L^2(M)} \|u\|_{H^k(M)} (\varepsilon + \lambda_i)^{-k}$$

On $\mathbf{T}^n = \mathbf{R}^n / \mathbf{Z}^n$, we can compute this expansion explicitly. The space E_λ consists of the span of any functions of the form $e_\xi(x) = e^{2\pi i \xi \cdot x}$, where $\xi \in \mathbf{Z}^n$,

and $\lambda = 2\pi|\xi|$. In this case the Fourier transform explicitly tells us the slightly stronger identity

$$|\langle u, e_\xi \rangle| = |\hat{u}(\xi)| \lesssim_k \min \left(\|u\|_{L^1(\mathbf{T}^n)}, \|D^k u\|_{L^1(\mathbf{T}^n)} |\xi|^{-k} \right).$$

Chapter 37

Distribution of Eigenvalues and the Half Wave Operator

Consider the setup we began our study of in the last section. We have a classical, self-adjoint, elliptic pseudodifferential operator T of order one on a compact manifold M with principal symbol $p(x, \xi) \geq 0$, and we have associated with T a sequence of orthonormal smooth functions $\{e_i\}$ in $C^\infty(M)$ which form a basis for $L^2(M)$, and an increasing sequence $\{\lambda_i\}$ tending to ∞ , such that $Te_i = \lambda_i e_i$. Our goal now is to further the study of the behaviour of the eigenfunctions $\{e_i\}$, and the distributions of the eigenvalues $\{\lambda_i\}$. A key object in the study of the distribution of the eigenvalues is the function $N(\lambda)$, which gives the number of eigenvalues of T less than or equal to λ . Our results already give a very basic estimate for N , which we will considerably sharpen over the course of this section.

Theorem 37.1. *The function N is tempered in λ . More precisely, for $\lambda > 0$,*

$$N(\lambda) \lesssim_\varepsilon \lambda^{n+\varepsilon}.$$

Proof. Define the operators S_λ , which are the projections in $L^2(M)$ onto the eigenspaces corresponding to eigenvalues with value $\leq \lambda$. Then S_λ clearly has kernel

$$K_\lambda(x, y) = \sum_{\lambda_i \leq \lambda} e_i(x) \overline{e_i(y)}.$$

This function is smooth, and

$$\int K_\lambda(x, x) dx = \sum_{\lambda_i \leq \lambda} |e_i(x)|^2 = N(\lambda).$$

Thus $N(\lambda)$ is the trace of the operator S_λ . Because $\|e_i\|_{H^k(M)} \lesssim_k (1 + |\lambda|)^k$ if $\lambda_i \leq \lambda$, we have the elementary estimate

$$\|S_\lambda f\|_{H^k(M)} \lesssim_k \lambda^k \|f\|_{L^2(M)}.$$

The Sobolev embedding theorem thus implies that if $\dim(M) = d$, and if $\varepsilon > 0$, then

$$\|S_\lambda f\|_{L^\infty(M)} \lesssim_\varepsilon \|S_\lambda f\|_{H^{d/2+\varepsilon}(M)} \lesssim \lambda^{d/2+\varepsilon} \|f\|_{L^2(M)}.$$

Thus we conclude that, for any $f \in C^\infty(M)$, and any $x \in M$,

$$\left| \int K_\lambda(x, y) f(y) dy \right| \lesssim_\varepsilon \lambda^{d/2+\varepsilon} \|f\|_{L^2(M)}.$$

By density, this implies that if $g_{\lambda, x}(y) = K_\lambda(x, y)$, then

$$\|g_{\lambda, x}\|_{L^2(M)} \lesssim_\varepsilon \lambda^{d/2+\varepsilon}.$$

But $g_{\lambda, x} \in \text{span}\{e_i : \lambda_i \leq \lambda\}$, so this means that

$$\|g_{\lambda, x}\|_{L^\infty(M)} = \|S_\lambda g_{\lambda, x}\|_{L^\infty(M)} \lesssim_\varepsilon \lambda^{d/2+\varepsilon} \|g_{\lambda, x}\|_{L^2(M)} \lesssim_\varepsilon \lambda^{d+2\varepsilon}.$$

But this means we conclude that $\|K_\lambda\|_{L^\infty(M \times M)} \lesssim_\varepsilon \lambda^{d+\varepsilon}$, and thus that

$$N(\lambda) = \int K_\lambda(x, x) dx \lesssim_\varepsilon \lambda^{d+\varepsilon}. \quad \square$$

Our next goal is now to establish the *Sharp Weyl formula*. If

$$c = \int_{\{(x, \xi) \in T^*M : a(x, \xi) \leq 1\}} d\xi dx,$$

then the result states that

$$N(\lambda) = c\lambda^n + O(\lambda^{n-1}).$$

We will continue to use the representation $N(\lambda) = \text{Tr}(S_\lambda)$. However, a key step here is to recognize that S_λ is a *function* of T , namely $S_\lambda = \chi_\lambda(T)$, where $\chi_\lambda(\xi) = \mathbf{I}(\xi \leq \lambda)$. The next trick is to use the Fourier multiplication

formula to obtain a useful representation of the structure of S_λ . For $f \in C^\infty(M)$, we can write

$$\begin{aligned} S_\lambda f &= \int_{-\infty}^{\infty} \chi_\lambda(\tau) \left(\sum_i \delta_{\lambda_i}(\tau) E_i f \right) d\tau \\ &= \int_{-\infty}^{\infty} \hat{\chi}_\lambda(t) \left(\sum_i e^{2\pi i t \lambda_i} E_i f \right) dt \\ &= \int_{-\infty}^{\infty} \hat{\chi}_\lambda(t) e^{2\pi i t T} dt \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{2\pi i t(T-\lambda)} f}{t + i0} dt. \end{aligned}$$

Thus we are lead to study the *half wave operator* $\partial_t - 2\pi i T$, since one sees that $(\partial_t - 2\pi i T)\{e^{2\pi i t T} f\} = 0$ for all $f \in L^2(M)$; it is easily verified for the eigenfunctions of T , and then the result follows by the distributional continuity of all operations involved.

We will understand the operators $e^{2\pi i t T}$ by constructing a (Lax) *parametrix* for the equation $\partial_t - 2\pi i T$ over small times, i.e. for $|t| \lesssim 1$. This equation is a pseudodifferential variant of a hyperbolic partial differential equation, so based on the parametrix construction in this scenario, we might hope to find a parametrix defined by an oscillatory integral of the form

$$Sf(x, t) = S(t)f(x) = \int s(t, x, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)} f(y) dy d\xi,$$

such that:

- $\Phi(t, x, y, \xi) = \phi(x, y, \xi) + tp(y, \xi)$, where ϕ is smooth away from $\xi = 0$, homogeneous of degree one, and $\phi(x, y, \xi) \approx (x - y) \cdot \xi$ in the sense that on the support of s ,

$$\partial_\xi^\beta \{\phi(x, y, \xi) - (x - y) \cdot \xi\} \lesssim_\beta |x - y|^2 |\xi|^{1-\beta}.$$

In particular, this implies that $\phi(x, y, \xi) = 0$ when $(x - y) \cdot \xi = 0$.

- s is a symbol of order zero, supported on $|x - y| \lesssim 1$ and on $|\xi| \geq 1$, in such a way that

$$|\nabla_\xi \phi(x, y, \xi)| \gtrsim |x - y|$$

and

$$|\nabla_x \phi(x, y, \xi)| \gtrsim |\xi|$$

for $(x, y) \in \text{supp}_x(s) \times \text{supp}_y(s)$.

If $(\partial_t - 2\pi i T) \circ S = 0$ is a smoothing operator on $(-\varepsilon, \varepsilon) \times M$ and $S(0)$ differs from the identity operator by a smoothing operator, then $S - e^{2\pi i t T}$ is smoothing for $|t| \leq \varepsilon$. This will therefore give us much more information about the behaviour of the propagators $e^{2\pi i t T}$ over small times.

To find a choice of ϕ and s which gives us this parametrix, let us start by determining what properties these functions should satisfy. Let us fix a coordinate system (x, U) , where $x(U)$ is a precompact subset of \mathbf{R}^n . Let us assume that in these coordinates, T has symbol $a(x, \xi)$. Then the kernel of $(\partial_t - 2\pi i T) \circ S$ in this coordinate system is

$$\int (\partial_t + 2\pi i T(x, D)) \left\{ s(t, \cdot, y, \xi) e^{2\pi i \Phi(t, \cdot, y, \xi)} \right\} d\xi.$$

If we set

$$s'(t, x, y, \xi) = e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t - 2\pi i T(x, D)) \left\{ s(t, \cdot, y, \xi) e^{2\pi i \Phi(t, \cdot, y, \xi)} \right\}$$

then the kernel is

$$\int s'(t, x, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)} d\xi.$$

Provided that s' is a symbol of order $-\infty$ for $0 < |t| \leq \varepsilon$, integration by parts shows that $(\partial_t + 2\pi i T) \circ S$ is smoothing, and so we will try to choose ϕ and s so as to obtain such a result.

In our discussion of pseudodifferential operators, we have already discussed an asymptotic formula for s' , namely, if

$$r_{x,y}(z) = \nabla_x \phi(x, z, \xi) \cdot (x - z) - \{\phi(x, y, \xi) - \phi(z, y, \xi)\}.$$

then for any $N > 0$, if $a \sim \sum_{k=-\infty}^1 a_k$, where a_k is homogeneous of degree

k , and if $\xi_\phi = \nabla_x \Phi(t, x, y, \xi) = \nabla_x \phi(x, y, \xi)$,

$$\begin{aligned}
& s'(t, x, y, \xi) \\
&= \underbrace{(p(y, \xi) - a(x, \xi_\phi)) \cdot s(t, x, y, \xi)}_{\text{symbols of order 1}} \\
&+ \underbrace{\partial_t s(t, x, y, \xi)}_{\text{symbol of order 0}} \\
&- \sum_{1 \leq |\beta| < N} \underbrace{\frac{2\pi i}{\beta! \cdot (2\pi i)^\beta} \cdot \partial_\xi^\beta a(x, \xi_\phi) \partial_z^\beta \{e^{2\pi i r_{x,y}(z)} s(t, z, y, \xi)\}}_{\text{symbols of order } 1 - \lfloor |\beta|/2 \rfloor} \Big|_{z=y} \\
&+ R_N(t, x, y, \xi).
\end{aligned}$$

where, because $|\nabla_x \Phi(t, x, y, \xi)| \gtrsim |\xi|$ on the support of s ,

$$\langle \xi \rangle^{t - \lfloor N/2 \rfloor} R_N \in L^\infty((-\varepsilon, \varepsilon) \times U \times U \times \mathbf{R}^d).$$

It is simple to establish estimates of the form

$$|\partial_x^\alpha \partial_y^\beta \partial_\xi^\lambda s'(t, x, y, \xi)| \lesssim \langle \xi \rangle^{N_{\alpha\beta\lambda}}.$$

Thus if we can justify that $|s'(t, x, y, \xi)| \lesssim_N \langle \xi \rangle^{-N}$ for all $N > 0$, then it will follow that s' is a symbol of order $-\infty$. We now determine the properties of the symbol s and the symbol ϕ which will give us these estimates.

To begin with, let us specify the function ϕ . In order to guarantee that s' is a symbol of order zero, the expansion above shows that $(p(y, \xi) - p(x, \xi_\phi)) \cdot s(t, x, y, \xi)$ must be a symbol of order zero. This will be true if we can pick ϕ such that, on the support of s , and for $|\xi| \gtrsim 1$,

$$p(x, \nabla_x \phi(x, y, \xi)) = p(y, \xi).$$

This is an example of an *Eikonal equation*, e.g. an equation of the form

$$q(z, \nabla_z \psi(z)) = 0$$

for some function $q(z, \zeta)$. In our case, $z = (x, y, \xi)$, so $\zeta = (\zeta_x, \zeta_y, \zeta_\xi)$, and so

$$q(z, \zeta) = p(x, \zeta_x) - p(y, \xi).$$

Let us make some further remarks we desire about our choice of function ϕ :

- We want ϕ to be homogeneous and smooth away from the origin. If we solve the equation for all $|\xi| = 1$, and then extend ϕ such that for $\lambda > 0$ and $|\xi| = 1$,

$$\phi(x, y, \lambda\xi) = \lambda\phi(x, y, \xi)\psi(\lambda),$$

where ψ is smooth, equal to one for $|\lambda| \geq 3/4$, and vanishing for $|\lambda| \leq 1/2$, then ϕ will satisfy the equation for all $|\xi| \gtrsim 1$. This means that

$$p(x, \nabla_x \phi(x, y, \xi)) - p(y, \xi)$$

is smooth and supported on $|\xi| \lesssim 1$, which implies it is a symbol of order $-\infty$, which suffices for our construction. Thus it suffices to solve the equation for $|\xi| = 1$.

- Since ϕ is smooth away from the origin and homogeneous, the equation

$$|\partial_\xi^\beta \{\phi(x, y, \xi) - (x - y) \cdot \xi\}| \lesssim_\beta |x - y|^2 \langle \xi \rangle^{1-\beta}$$

holds if, for $|\xi| = 1$, we have $\phi(x, y, \xi) = 0$ whenever $(x - y) \cdot \xi = 0$, and $\nabla_x \phi(x, y, \xi) = \xi$ whenever $x = y$. Thus we have some *initial conditions* for our Eikonal equation.

The second condition constitutes a type of initial condition for ϕ , since it specifies its behaviour on a hypersurface, a kind of Cauchy condition, and thus we should expect these are close to the conditions that give unique solutions to the equation. And the following Lemma indeed shows that there is a unique function ϕ , defined for $|x - y| \lesssim 1$ and $|\xi| = 1$ with these properties.

Lemma 37.2. *Let Z be a smooth manifold, and let $q(z, \zeta)$ be a real-valued, smooth function defined locally around a point $(z_0, \zeta_0) \in T^*Z$. Let S be a smooth hypersurface in Z passing through z_0 with conormal vector ζ_S at z_0 , such that*

$$\frac{\partial q}{\partial \zeta_S}(z_0, \zeta_0) = \lim_{t \rightarrow 0} \frac{q(z_0, \zeta_0 + t\zeta_S) - q(z_0, \zeta_0)}{t}$$

is nonzero. Suppose that ψ is any smooth function defined on S locally about z_0 , such that $d\psi(z_0)$ agrees with the action of ζ_0 on $T_{x_0}S$. Then there exists a unique smooth function ϕ defined in a neighborhood of z_0 , which agrees with ψ on S , satisfies the Eikonal equation $q(z, \nabla_z \phi(z)) = 0$, and has $\nabla_z \phi(z_0) = \zeta_0$.

Proof. TODO: See Sogge, Theorem 4.1.1. □

In our case,

$$Z = \{(x, y, \xi) : |\xi| = 1\}.$$

We have $z_0 = (x_0, x_0, \xi_0)$, $\zeta_0 = (\xi, \xi, 0)$, and

$$S = \{(x, y, \xi) : |\xi| = 1 \text{ and } (x - y) \cdot \xi = 0\}.$$

The conormal vector ξ_S of S at z_0 is a multiple of $(\xi_0, -\xi_0, 0)$, and so by homogeneity,

$$\frac{\partial q}{\partial \xi_S} = \lim_{t \rightarrow 0} \frac{p(x_0, (1+t)\xi_0) - p(x_0, \xi_0)}{t} = p(x_0, \xi_0),$$

which is nonvanishing because T is elliptic. If we define ψ equal to zero on S , then $d\psi = 0$, which agrees with the action of ζ_0 on S . Thus the theorem applies local uniqueness and existence to solutions to the Eikonal equation, and by compactness of Z we can patch such solutions together to find a solution defined for all $|x - y| \lesssim 1$.

We therefore conclude that there exists a unique choice of ϕ such that, if s has small enough support, $s'(t, x, y, \xi)$ is a symbol of order zero. Next, let us see what constraints are forced on us in order to ensure that $S(0)$ differs from the identity by a smoothing operator. The kernel of U is precisely

$$\int s(0, x, y, \xi) e^{2\pi i \phi(x, y, \xi)} d\xi.$$

We now show that this operator is actually a *pseudodifferential operator* of order zero, and determine its symbol up to first order.

To do this, we write $\phi_\alpha(x, y, \xi) = (1 - \alpha)\phi(x, y, \xi) + \alpha(x - y) \cdot \xi$. Let U_α be the operator with kernel

$$\int s(0, x, y, \xi) e^{2\pi i \phi_\alpha(x, y, \xi)} d\xi.$$

Assume the support of s is close enough to the diagonal such that

$$|\nabla_\xi \phi_\alpha(x, y, \xi)| \gtrsim |x - y|$$

on the support of s . Then $\partial_\alpha^n U_t$ has kernel

$$\int (2\pi i)^n (\phi_1 - \phi_0)^n s(0, x, y, \xi) e^{2\pi i \phi_\alpha(x, y, \xi)} d\xi.$$

This is an oscillatory integral defined by a symbol of order n . However, when $t = 1$, the fact that $\phi(x, y, \xi) \approx (x - y) \cdot \xi$, together with the formula for converting pseudodifferential operators with compound symbols into standard Kohn-Nirenberg type symbols shows that $\partial_\alpha^n U_1$ is actually a pseudodifferential operator of order $-n$. Integration by parts, similarly, shows that $\partial_\alpha^n U_t$ is defined by an oscillator integral against a symbol of order $-n$. But this means that if we define a pseudodifferential operator by the asymptotic formula

$$V \sim \sum \frac{(-1)^n}{n!} \partial_\alpha^n U_1,$$

then $U - V$ is smoothing. Indeed, for any n , by Taylor's formula we have

$$U = \sum_{k=0}^{n-1} \frac{(-1)^k}{k!} \partial_\alpha^k U_1 + \frac{(-1)^n}{n!} \int_0^1 \alpha^{n-1} \partial_\alpha^n U_\alpha d\alpha$$

The integral here is an oscillatory integral defined against a symbol of order $-n$, and thus taking $n \rightarrow \infty$ verifies the claim.

It is an important remark that reversing this argument shows that *any* pseudodifferential operator can be written in the form above for the particular choice of ϕ we have given. This is a special case of the *equivalence of phase functions* theorem. This in particular guarantees that we can choose a symbol $I(x, y, \xi)$ of order zero such that $U - 1$ is smoothing if and only if $s(0, x, y, \xi) - I(x, y, \xi)$ is a symbol of order $-\infty$. The symbol I can be chosen to be vanishing for $|x - y| \gtrsim 1$, since the difference will be a smoothing pseudodifferential operator.

Next, the quantity

$$\begin{aligned} & \partial_t s(t, x, y, \xi) \\ & + \sum_{k=1}^d \partial_\xi^k a(x, \xi_\phi) \partial_x^k s(t, x, y, \xi) \\ & + \left(a_0(x, \xi_\phi) + \frac{1}{2\pi} \sum_{|\beta|=2} \partial_\xi^\beta p(x, \xi_\phi) \partial_x^\beta \phi(x, y, \xi) \right) s(t, x, y, \xi). \end{aligned}$$

must be a symbol of order -1 . But because the coefficients of this equation are smooth, and all derivatives are bounded, it follows from the general

theory of transport equations that there exists a unique smooth, function s_0 defined for $|t| \leq \varepsilon$, which is a symbol of order zero, such that $s_0(0, x, y, \xi) = I(x, y, \xi)$, s_0 vanishes for $|x - y| \gtrsim 1$, and satisfies the transport equation

$$\begin{aligned} & \partial_t s_0(t, x, y, \xi) \\ & + \sum_{k=1}^d \partial_\xi^k a(x, \xi_\phi) \partial_x^k s_0(t, x, y, \xi) \\ & + \left(a_0(x, \xi_\phi) + \frac{1}{2\pi} \sum_{|\beta|=2} \partial_\xi^\beta p(x, \xi_\phi) \partial_x^\beta \phi(x, y, \xi) \right) s_0(t, x, y, \xi) = 0. \end{aligned}$$

We have thus justified that the quantity

$$R_0(t, x, y, \xi) = e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t - 2\pi i T)(s_0(t, \cdot, y, \xi) e^{2\pi i \Phi(t, x, y, \xi)})$$

is a symbol of order -1 . Now we come to a quirk of this parametrix, which does not occur in the study of hyperbolic partial differential equations. Since the operator $P(x, D)$ is only *pseudolocal* rather than completely local, the remainder term R_0 is *not* necessarily supported on a neighborhood of the origin. To fix this, we now successively define the terms $\{s_k\}$ for $k < 0$, which are symbols of order $-k$, such that $s_k(0, x, y, \xi) = 0$, and

TODO : SPECIFY REQUIRED EQUATION.

Again, solutions exist for small time periods. And this implies that $e^{-2\pi i \Phi(t, x, y, \xi)} (\partial_t + 2\pi i T)((s_0 + \dots + s_{-k}) e^{2\pi i \Phi(t, x, y, \xi)})$ is a symbol of order $-k$ (TODO: Is It), and we can continue the calculation to complete the argument.

Let us now use this equation to prove the sharp Weyl formula, which immediately is given by the following, more precise estimate.

Theorem 37.3. *Let M be a compact manifold, and let T be a classical, self-adjoint, elliptic pseudodifferential operator of order one on M , with principal symbol $p(x, \xi)$. If S_λ is the projection operator onto the span of eigenfunctions with eigenvalue $\leq \lambda$, and K_λ is the kernel of the operator S_λ , then*

$$K_\lambda(x, x) = c(x) \lambda^n + O(\lambda^{n-1}),$$

where

$$c(x) = \int_{p(x, \xi) \leq 1} d\xi.$$

Proof. Recall that we have

$$S_\lambda f(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{2\pi i t(T-\lambda)} f}{t + i0} dt.$$

This integral only has a singularity when $t = 0$, so it makes sense to decompose the integral into two parts using an even function $\rho \in C_c^\infty(\mathbf{R})$ supported on $|t| \leq \varepsilon/2$ and equal to one for $|t| \leq \varepsilon/4$, i.e. writing

$$S_\lambda f = S_{\lambda, \text{low}} f + S_{\lambda, \text{high}} f,$$

where

$$S_{\lambda, \text{low}} f = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\rho(t)}{t + i0} e^{2\pi i t(T-\lambda)} f dt$$

and

$$S_{\lambda, \text{high}} f = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1 - \rho(t)}{t} e^{2\pi i t(T-\lambda)} f dt.$$

To understand $S_{\lambda, \text{low}} f$, we use the parametrix for the wave equation. If we switch to coordinates around a particular point, then we find that the kernel of $S_{\lambda, \text{low}}$ is equal to $\tilde{K}_\lambda + R_\lambda$, where \tilde{K}_λ is supported on a neighborhood of the diagonal, and locally in coordinates we can write

$$\tilde{K}_\lambda(x, y) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t + i0} s(t, x, y, \xi) e^{2\pi i (\phi(x, y, \xi) + t(p(y, \xi) - \lambda))} d\xi dt.$$

and where

$$R_\lambda(x, y) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\rho(t)}{t + i0} A(t, x, y) e^{2\pi i t \lambda} dt,$$

where A is smooth for $|t| \leq \varepsilon$ and for $x, y \in M$. But since $\rho \cdot A$ is smooth, with compact support in the t variable taking inverse Fourier transforms in the t variable shows that

$$|R_\lambda(x, y)| = \left| \int_{-\infty}^{\lambda} \mathcal{F}_t^{-1}\{\rho A\}(\lambda - \tau, x, y) d\tau \right| \lesssim 1,$$

By Taylor's formula, we can write $s(t, x, x, \xi) = s(0, x, x, \xi) + t \cdot r(t, x, \xi)$,

where r is a symbol of order zero. But if N is sufficiently large, we have

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t+i0} (t \cdot r(t, x, \xi)) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \rho(t) r(t, x, \xi) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt \\
&= \frac{1}{2\pi} \int_{\mathbf{R}^n} \mathcal{F}_t^{-1} \{\rho \cdot r\} (p(x, \xi) - \lambda, x, \xi) d\xi \\
&= \int O_N (1 + |p(x, \xi) - \lambda|)^{-N} d\xi = O(\lambda^{n-1}).
\end{aligned}$$

We also have $s(0, x, x, \xi) = 1 + s_{-1}(x, \xi)$, where s_{-1} is a symbol of order -1 . Now if $\tilde{\chi}_\lambda = \chi_\lambda * \mathcal{F}_t^{-1} \{\rho\}$, then

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t+i0} s_{-1}(x, \xi) e^{2\pi i t(p(x, \xi) - \lambda)} d\xi dt = \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) s_{-1}(x, \xi) d\xi.$$

Then $\tilde{\chi}_\lambda(p(x, \xi)) \sim 1$ for $|\xi| \lesssim \lambda$, and for $|\xi| \gtrsim \lambda$,

$$\tilde{\chi}_\lambda(p(x, \xi)) \lesssim_N \langle |\xi| - \lambda \rangle^{-N}$$

This implies that

$$\left| \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) s_{-1}(x, \xi) d\xi \right| \lesssim \int_0^{\lambda+O(1)} r^{n-2} + \int_{\lambda+O(1)}^{\infty} r^{n-1} (r - \lambda)^{-N} \lesssim \lambda^{n-1}.$$

Thus we have

$$\begin{aligned}
\tilde{K}_\lambda(x, x) &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} \frac{\rho(t)}{t+i0} e^{2\pi i t(p(x, \xi) - \lambda)} + O(\lambda^{n-1}) \\
&= \int_{\mathbf{R}^n} \tilde{\chi}_\lambda(p(x, \xi)) d\xi + O(\lambda^{n-1}) \\
&= \int_{\mathbf{R}^n} \tilde{\chi}(p(x, \xi) - \lambda) d\xi + O(\lambda^{n-1}) \\
&= c(x) \lambda^n + \int_{\mathbf{R}^n} (\tilde{\chi} - \chi)(p(x, \xi) - \lambda) d\xi + O(\lambda^{n-1}).
\end{aligned}$$

Now $\tilde{\chi} - \chi = \chi * u$, where u is the inverse Fourier transform of $\rho(t) - 1$. But then

$$(\chi * u)(\tau) = \int_{-\infty}^0 \hat{\rho}(\tau - \omega) d\omega - \mathbf{I}(\tau \geq 0),$$

from which we verify that

$$|(\chi * u)(\tau)| \lesssim_N (1 + |\tau|)^{-N}.$$

But we can use this to again show that

$$\begin{aligned} \left| \int_{\mathbf{R}^n} (\tilde{\chi} - \chi)(p(x, \xi) - \lambda) d\xi \right| &\lesssim \int_0^{\lambda+O(1)} r^{n-1} |\xi|^{-N} dr \\ &\quad + \int_{\lambda-O(1)}^{\lambda+O(1)} (\lambda - r)^{-N} r^{n-1} dr \\ &\quad + \int_{\lambda+O(1)}^{\infty} r^{n-1} (r - \lambda)^{-N} dr \\ &\lesssim \lambda^{n-1}. \end{aligned}$$

Thus we have established the result for the kernel $K_{\lambda, \text{low}}$ of the operator $S_{\lambda, \text{low}}$. The proof of this theorem would be finished if we were able to justify that

$$|K_\lambda(x, x) - K_{\lambda, \text{low}}(x, x)| \lesssim \lambda^{n-1}.$$

Define the function

$$\begin{aligned} g(\lambda, x) &= K_\lambda(x, x) - K_{\lambda, \text{low}}(x, x) \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} (1 - \rho(t)) \{e^{-2\pi i t(T-\lambda)}(x, x)\} dt. \end{aligned}$$

Then the Fourier transform in the λ variable is

$$\hat{g}(t, x) = \frac{1}{2\pi i} (1 - \rho(t)) e^{-2\pi i t T}.$$

Thus the support of \hat{g} is supported away from $|t| \leq \varepsilon/2$. A Lemma of Tauberian type following this proof shows that it suffices to show that for $\lambda > 0$, and $0 < \tau \leq 1$,

$$|g(\lambda + \tau, x) - g(\lambda, x)| \lesssim (1 + \lambda)^{n-1}$$

But this follows if we can show that

$$|K_{\lambda+\tau}(x, x) - K_\lambda(x, x)| \lesssim (1 + \lambda)^{n-1}$$

and that

$$|K_{\lambda+\tau, \text{low}}(x, x) - K_{\lambda, \text{low}}(x, x)| \lesssim (1 + \lambda)^{n-1}.$$

These inequalities are equivalent to the (L^1, L^2) *discrete restriction theorem* for eigenfunctions of the Laplacian, which we will analyze in the next section. Indeed, if we let $\chi_{\lambda, \tau}$ be the operator with kernel $K_{\lambda+\tau} - K_{\lambda}$, then the discrete restriction theorem tells us that

$$\|\chi_{\lambda, \tau} f\|_{L^2(M)} \lesssim (1 + \lambda)^{(n-1)/2} \|f\|_{L^1(M)}$$

which by Schur's test, is equivalent to us having

$$\sup_{x \in M} \int_M |K_{\lambda+\tau}(x, y) - K_{\lambda}(x, y)|^2 dy \lesssim (1 + \lambda)^{n-1}$$

But the left hand side, by orthogonality, is equal to

$$\sup_{x \in M} \sum_j \mathbf{I}(\lambda \leq \lambda_j \leq \lambda + \tau) |e_j(x)|^2 = \sup_{x \in M} |K_{\lambda+\tau}(x, x) - K_{\lambda}(x, x)|,$$

which gives the required inequality. The analysis of $K_{\lambda, \text{low}}$ follows because we can write

$$K_{\lambda+\tau, \text{low}}(x, y) - K_{\lambda, \text{low}}(x, y) = \frac{1}{2\pi i} \int \frac{\rho(t)}{t + i0} (e^{-2\pi i \tau t} - 1) e^{2\pi i (T - \lambda)t} dt.$$

The $e^{-2\pi i \tau t} - 1$ term annihilates the singularity at the origin, and one can now argue as above to show this quantity is $O(\lambda^{n-1})$. \square

Lemma 37.4. *Let $g(\lambda)$ be a piecewise continuous function on \mathbf{R} , such that for $\lambda > 0$, and $0 < \tau \leq 1$,*

$$|g(\lambda + \tau) - g(\lambda)| \lesssim (1 + \lambda)^a.$$

If $\widehat{g}(t)$ vanishes for $|t| \leq 1$, then

$$|g(\lambda)| \lesssim (1 + \lambda)^a.$$

Proof. If

$$G(\lambda) = \int_{\lambda}^{\lambda+1} g(\tau) d\tau,$$

then G is absolutely continuous, and for almost all λ , G is differentiable with

$$|G'(\lambda)| = |g(\lambda + 1) - g(\lambda)| \lesssim (1 + \lambda)^a.$$

The Fourier transform of G also vanishes for $|t| \leq 1$. But

$$|g(\lambda)| \leq |G(\lambda)| + O((1 + \lambda)^a)$$

so it suffices to prove the estimates for G . If $\eta(t) = 1$ for $|t| > 1$, and $\eta(t) = 0$ for $|t| \leq 1/2$, and if ψ has Fourier transform $\eta(t)/2\pi it$, then ψ is bounded and rapidly decreasing at infinity. But $G' * \psi = G$, which gives

$$|G(\lambda)| \lesssim (1 + \lambda)^a \int |\psi(s)|(1 + |s|)^a ds \lesssim (1 + \lambda)^a. \quad \square$$

Chapter 38

Nodal Sets

Let M^d be a compact manifold, and consider a classical, self-adjoint, elliptic pseudodifferential operator P with positive principal symbol. Then we have seen that one can decompose $L^2(M)$ into orthogonal eigenspaces $\oplus_\lambda E_\lambda$ of the Laplace-Beltrami operator on M , where λ ranges over a discrete subset of real numbers bounded from below, and E_λ is a finite dimensional subspace of $C^\infty(M)$ such that $PE_\lambda = \lambda E_\lambda$. The *node* of a real-valued eigenfunction e_λ is the zero set of e_λ , denoted N_λ or $N(e_\lambda)$. We are interested in the asymptotic behaviour of N_λ as $\lambda \rightarrow \infty$. In particular, one might ask questions such as:

- How does the number of connected components of $M - N_\lambda$, the *nodal count* of the eigenfunction, grow as $\lambda \rightarrow \infty$.
- What is the volume of N_λ . S.T. Yau's nodal size conjecture is that if $P = \sqrt{-\Delta}$, where Δ is the Laplace Beltrami operator, then

$$H^{n-1}(N_\lambda) \sim_M \lambda,$$

where the implicit constant depends solely on the manifold M .

- One cannot 'hear the shape of a drum', i.e. there exists non-isomorphic compact Riemannian manifolds which are *isospectral*, i.e. whose Laplace-Beltrami operators have the same spectrum. However Smilansky has posed the question of whether one can 'count the shape of a drum', i.e. does the nodal count of N_λ for all λ determine the Riemannian manifold.

The analysis often breaks up into complex-variable techniques, which assume M is an analytic Riemannian manifold, and real-variable techniques, which depend only on the fact that M is a smooth manifold. For concreteness, we will focus on the case where $P = \sqrt{-\Delta}$, which has been the scenario with the most focused study.

Example. The only compact Riemannian manifold of dimension one, up to scaling, is $\mathbf{T} = \mathbf{R}/\mathbf{Z}$. In this case, λ ranges over the non-negative integers, E_0 is one dimensional, the span of the constant function, and more generally, E_n is two dimensional, spanned by $e^{2\pi nix}$ and $e^{-2\pi nix}$. A general element of E_n can be written as

$$e_n = A \cos(2\pi nx + \phi)$$

for some constants A and ϕ , and we see the nodal set $N(e_n)$ consists of $2n$ equidistributed points on the torus, and thus the nodal count is $NC(e_n) = 2n$.

More generally, we can consider a flat torus $T = \mathbf{R}^d/\Gamma$, where Γ is a lattice with some basis $\{z_1, \dots, z_d\}$. We define the dual lattice Γ^* to be the space of all vectors w such that $w \cdot z \in \mathbf{Z}$ for any $z \in \Gamma$. If $w \in \Gamma^*$, then

$$e_w(x) = e^{2\pi i w \cdot x}$$

is seen to be an eigenfunction of the Laplacian on T with eigenvalue $2\pi|w|$. These form an orthogonal basis for $L^2(T)$, and therefore span the space of all eigenvalues. Due to the multiplicity of the eigenspaces, it seems very difficult to picture a general nodal set. But if we consider an eigenfunction of the form

$$e(x) = Ae^{2\pi i w \cdot x} + Be^{-2\pi i w \cdot x} = C \cos(2\pi w \cdot x + \phi)$$

then we see that the nodal set here is a union of $2v(w)$ hyperplanes, where $v(w)$ is the minimum positive value of $w \cdot \gamma$, where γ ranges over elements of Γ . If we consider the basis of Γ^* given by $\{w_1, \dots, w_d\}$, where $w_i \cdot z_j = \delta_{i,j}$, and $w = \sum m_i w_i$, then $v(w)$ is the greatest common divisor of the integers $\{m_1, \dots, m_n\}$.

Example. The eigenfunctions of the sphere S^d consists of the spherical harmonics. The eigenvalues λ range over numbers of the form $n^{1/2}(n+d-1)^{1/2}$ for some $n \geq 0$, and for such λ , the eigenspace E_λ can be identified with the space of all harmonic polynomials in $d+1$ variables, homogeneous of degree n . The nodal sets of a general harmonic polynomial, however, is much more difficult to analyze than the torus, since the eigenspaces E_λ have high multiplicity for large λ .

Knowledge of general nodal sets is very restricted. We know that the $d - 1$ dimensional Hausdorff measure of a nodal set is finite, as well as the $d - 2$ dimensional Hausdorff measure of the singularities of the nodal set, i.e. where e_λ and ∇e_λ vanishes.

38.1 Nodal Sets for Open Subsets of \mathbf{R}^d

Before we discuss the general setting, let us gain some intuition by studying the basic properties of the nodal sets of eigenfunctions of the Laplacian on bounded, open subsets of \mathbf{R}^d . We fix some open, bounded subset Ω of \mathbf{R}^d , and consider $C^2(\Omega)$, real-valued functions $e_\lambda : \Omega \rightarrow \mathbf{R}^d$ such that $\Delta e_\lambda = -\lambda^2 \phi$. In this case, we can reduce the study of e_λ to a harmonic function on $\Omega \times \mathbf{R}$, by setting $\phi(x, t) = e_\lambda(x)e^{\lambda t}$. Then

$$\Delta_{x,t}\phi = \Delta_x\phi + \Delta_t\phi = -\lambda^2\phi + \lambda^2\phi = 0.$$

Since $e^{\lambda t}$ never vanishes, we have $N(u) = N(\phi) \times \mathbf{R}$. Thus in this situation we can shift our analysis from the understanding of general eigenfunctions to the understanding of harmonic functions. In particular, we see that any eigenfunction lies in $C^\infty(\Omega)$.

Theorem 38.1. *Let B be the unit ball in \mathbf{R}^d . If $e_\lambda \in C^\infty(B)$, and $\Delta e_\lambda = -\lambda^2 e_\lambda$, then $B/2$ is contained in a $c\lambda^{-1}$ neighborhood of N_λ .*

Proof. We prove the result by contradiction. Suppose that $B/2$ was not contained in an $r < 1/4$ neighborhood of N_λ . Then e_λ does not change sign in some other ball B' with center lying in $B/2$ and with radius r , and we may assume without loss of generality that e_λ is positive here. Consider the harmonic function

$$\phi : B \times \mathbf{R} \rightarrow \mathbf{R}.$$

associated with e_λ by setting $\phi(x, t) = e_\lambda(x)e^{\lambda t}$. Harnack's inequality tells us that there exists a constant C such that

$$e^{\lambda r/2} \sup_{x \in B'} e_\lambda(x) = \sup_{|t| \leq r/2} \sup_{x \in B'} \phi(x, t) \leq C \inf_{|t| \leq r/2} \inf_{x \in B'} \phi(x, t) = Ce^{-\lambda r/2} \inf_{x \in B'} e_\lambda(x).$$

Thus

$$e^{\lambda r} \sup_{x \in B'} e_\lambda(x) \leq C \inf_{x \in B'} e_\lambda(x) \leq C \sup_{x \in B'} e_\lambda(x).$$

Thus, dividing out, we conclude that $e^{\lambda r} \leq C$, which implies that if $c = 2\log(C)$, then $r < c/\lambda$. \square

The solution to the Cauchy uniqueness problem for harmonic functions tells us that non-zero harmonic functions can only vanish of order up to one on hyperplanes. This also remains true for eigenfunctions of the Laplacian by this extension trick.

38.2 Nodal Sets on the Sphere

To gain some intuition, let us restrict our analysis even further, to the study of *homogeneous harmonic polynomials* of degree k \mathbf{H}_k on \mathbf{R}^{d+1} , which is equivalent to study eigenfunctions of the Laplacian on the sphere S^d . The space \mathbf{H}_k then has dimension

$$\binom{d+k}{d} - \binom{d+k-2}{d}$$

General nodal sets are impossible to calculate, but the concreteness here should make a stronger analysis more possible.

Example. The space \mathbf{H}_k on \mathbf{R}^2 has dimension 2 for all k . The space therefore consists of functions expressible in polar coordinates in the form

$$(r, \theta) \mapsto r^k \sin(k\theta + \phi).$$

Alternatively, we can get these results from the real and imaginary parts of the complex homogeneous polynomial $(x, y) \mapsto (x + iy)^k$. The nodal sets of these polynomials in \mathbf{R}^2 are precisely k lines through the origin, spaced out by a fixed angle. The nodal set on S^1 therefore has zero dimensional Hausdorff measure equal to k .

Example. The space \mathbf{H}_k on \mathbf{R}^3 has dimension $2k+1$. One can find a basis of \mathbf{H}_k using Legendre functions, but the nodal sets become hard to picture. But there are analogues here to the case of \mathbf{R}^2 , e.g. the real and imaginary parts of the functions $(x + iy)^k$, $(x + iz)^k$, and $(y + iz)^k$, whose nodal sets are k planes with a common intersection line, and are spaced out at equal angles radially outward from this perpendicular line. Thus the one dimensional Hausdorff measure of this nodal set on S^2 is equal to $2k\pi$. One can perform a similar analysis in \mathbf{R}^n ,

with the functions $(x + iy)^k$ being formed from k hyperplanes in \mathbf{R}^{d+1} with a common d dimensional plane lying at their intersection, and thus the nodal set on S^d has measure kc_d where c_d is the volume of an $d - 1$ dimensional great circle on S^d .

To estimate the size of a general nodal set on a sphere, we utilize an integral geometric formula. Namely, if E is a smooth, n dimensional hypersurface in \mathbf{R}^{n+1} , and we define

$$A_i = \int_{\mathbf{R}^n} H^0(E \cap H_{i,y}) dy,$$

where $H_{i,y}$ is the hyperplane given by $\{x \in \mathbf{R}^{n+1} : (x_1, \dots, \hat{x}_j, \dots, x_{n+1}) = y\}$ for $y \in \mathbf{R}^n$, then

$$\left(\sum_{i=1}^{n+1} A_i^2 \right)^{1/2} \leq H^{n-1}(E) \leq \sum_{i=1}^{n+1} A_i.$$

This leads to a general bound.

Theorem 38.2. *If $f \in \mathbf{H}^k$, then $N(f) \subset S^d$ satisfies*

$$H^{d-1}(N(f)) \lesssim_d k.$$

Proof. Observe that the number of points in $N(f) = E \cap H_{i,y}$ is equal to the number of zeroes of the degree $\leq k$ polynomial $f(y_1, \dots, t, \dots, y_n)$. Since f is harmonic, and thus does not vanish to order greater than one on any hyperplane, for almost every y , we conclude that $H^0(N(f) \cap H_{i,y}) \leq k$, and applying the formula above immediately yields the result. \square

Remark. A lower bound here is immediately seen more difficult, since a general single variable degree $\leq k$ polynomial can have any number of zeroes on the real line, so this technique cannot work so well to get lower bounds unless we use analyticity in some way, and this probably depends on the underlying manifold we are working with to have some analytic structure. But one can obtain this result for the sphere, precisely because it has this analytic structure. TODO: See Han's notes on Nodal Sets.

38.3 Brownian Motion and Nodal Sets

Eigenfunctions e_λ to the Laplace-Beltrami operator on a compact manifold M^d behave well under the heat equation, i.e. if $e^{\Delta t}$ are the propagators for the heat equation $\partial_t = \Delta$ on M , and $e_\lambda u = -\lambda^2 u$, then

$$(e^{t\Delta} e_\lambda)(x) = e^{-\lambda^2 t} e_\lambda(x).$$

The heat equation is mathematically describing the distribution of a large number of particles, each diffusing through a medium in which they are subject to random molecular bombardments. The theory of *diffusions* in probability give us an alternate viewpoint through which to model this situation, so it makes sense this theory will bring light upon the theory of eigenfunctions.

38.3.1 Probabilistic Tools

To define diffusions, let us introduce the required machinery. We work over a fixed probability space Ω . A *continuous stochastic process* valued in some space M is then a Borel-measurable function $X : \Omega \rightarrow C([0, \infty), M)$, which we also view as a family of M valued random variables $\{X_t : t \in [0, \infty)\}$. Intuitively, a stochastic process X is a random path formed by a particle travelling through space. Given a fixed process on M and $x \in M$, we let $\mathbf{E}^x[\cdot] = \mathbf{E}[\cdot | X_0 = x]$ denote the conditional expectation given that the process starts at the position x .

The most basic diffusion process is *Brownian motion*, and is used to generate the more general family. A one dimensional Brownian motion B is a continuous stochastic process such that for any interval $I = [t, s]$, the random variables $\Delta_I B = B_s - B_t$ are mean zero, variance $s - t$ Gaussians, and for any disjoint family of closed intervals $\{I_1, \dots, I_n\}$ in $[0, \infty)$, the random variables $\{\Delta_{I_1} B, \dots, \Delta_{I_n} B\}$ are independent of another. An n dimensional Brownian motion on \mathbf{R}^d is precisely a stochastic process whose coordinates are independent Brownian motions. An *Itô diffusion* with *drift coefficient* $b(x)$ (valued in \mathbf{R}^d) and *diffusion coefficient* $A(x)$ (a $d \times d$ matrix) is a continuous random process X satisfying a *stochastic differential equation* of the form $dX = b(X)dt + A(X)dB$. The formal definition of this stochastic differential equation is quite difficult, but intuitively, there exists a Brownian

motion B such that

$$X_{t+\delta} \approx X_t + b(X_t)\delta + A(X_t)[B_{t+\delta} - B_t]$$

which the approximation becomes equality in a technical sense as $\delta \rightarrow 0$, i.e. the error $E_{t,\delta}$ between the two random variables is a random variable with mean $o(\delta)$ and third moment $O(\delta^3)$. As one might expect, one can define Itô diffusions on a compact Riemannian manifold M , given a vector field b on M operating as the drift coefficient, and a section of linear transformations on TM .

An important fact about the diffusions above is that they are *Markov*, i.e. if one observes the values of the Brownian motion $\{X_t\}$, up to a certain time T , the only information one observes which tells us about the evolution of the process past the time T is the position the diffusion arrives to at time T , i.e. the random variable X_T . More formally, a probabilist would write this by stating that for $S > T$, and for a function f on M ,

$$\mathbf{E}[f(X_S)|\Sigma_T] = \mathbf{E}[f(X_S)|X_T],$$

where the conditional expectation is taken with respect to a σ algebra Σ_T , which models the information one sees when observing the process up to time T . One can also let the times T and S be random *stopping times*, so a diffusion satisfies the *strong Markov property*.

Now we connect diffusions to the study of second order partial differential equations. For any $f \in C^2(M)$, if we define a differential operator L on M , the *generator of the diffusion*, by setting

$$Lf(x) = b(x) \cdot \nabla f(x) + \frac{A(x)A(x)^T \cdot Hf(x)}{2},$$

Then the derivative of the function $t \mapsto \mathbf{E}^x[f(X_t)]$ at $t = 0$ is precisely $Lf(x)$. Dynkin's formula follows by formally applying the fundamental theorem of calculus to this derivative, i.e. that

$$\mathbf{E}^x[f(X_T)] = \mathbf{E}^x[f(X_0)] + \mathbf{E}^x \left[\int_0^T Lf(X_s) ds \right] = f(x) + \mathbf{E}^x \left[\int_0^T Lf(X_s) ds \right],$$

and we can allow T to be a stopping time as well, provided that $\mathbf{E}^x[T] < \infty$. The *Feynman-Kac formula* reverses this process; if we define

$$u(x, t) = \mathbf{E}^x[f(X_t)]$$

then u is the solution to the partial differential equation $\partial_t u = Lu$, with initial conditions f . Thus one can convert problems about second order parabolic PDEs to problems about diffusions, and vice versa. In particular, we see that we can obtain a natural definition of Brownian motion on a manifold by letting B be a diffusion with generator given by the Laplace Beltrami operator.

TODO: Showing that it takes R^2/n time on average to leave a ball of radius R .

Next, we move onto the study of boundary value problems for semielliptic PDEs L via diffusions. Let us start with a bounded region D of \mathbf{R}^d . Consider a diffusion process X with generator L , and let $T_D = \inf\{t > 0 : X_t \notin D\}$. Then for sufficiently regular inputs, the Dynkin formula tells us that the unique solution to the problem $Lv = -h$ on D with boundary conditions ϕ on ∂D is given by

$$v(x) = \mathbf{E}^x[\phi(X_{T_D})] + \mathbf{E}^x \left[\int_0^{T_D} h(X_t) dt \right].$$

We can also solve the heat equation $\partial_t u = Lu$ with absorbing boundary conditions $u(x, t) = 0$ for $x \in \partial D$, and initial condition $u(x, 0) = f(x)$ by setting

$$u(x, t) = \mathbf{E}^x[f(X_t)\chi_t],$$

where $\chi_t = 1$ if $t < T_D$, and $\chi_t = 0$ otherwise (we kill paths that reach the boundary). Using linearity, we can now solve the problem $\partial_t u - Lu = h$ with a general boundary condition $u(x, t) = \phi(x)$ on ∂D and with initial conditions $u(x, 0) = f(x)$, namely, we have

$$u(x, t) = \mathbf{E}^x[\phi(X_{T_D})] + \mathbf{E}^x \left[\int_0^{T_D} h(X_t) dt \right] + \mathbf{E}^x[f(X_t)\chi_t].$$

Note that if D is bounded, and $f \in L^\infty(D)$, then dominated convergence implies that for each $x \in D$, $u(x, t) \rightarrow v(x)$ as $t \rightarrow \infty$, i.e. so the solution to Dirichlet's problem is really the long term distribution of the process.

38.4 Nodal Sets Via Brownian Motion

To exploit this relation to control nodal sets, we let Ω be a nodal domain of e_λ , i.e. a connected component of M . The Feynman-Kac formula tells us

that solutions to the equation $\partial_t = \Delta$ on an open set $\Omega \subset M$, which are held equal to one on $\partial\Omega$ and vanish initially on the interior can be understood in terms of the probability that Brownian motion will hit $\partial\Omega$ in a given number of units of time.

We will get information about the nodal set by constructing diffusion processes which behave differently near the boundary via the Feynman-Kac formula,

TODO: WHY IS NEUMANN VS DIRICHLET CONNECTED TO THE NODAL SET OF AN EIGENFUNCTION? NEUMANN CONDITIONS DON'T WORK, BECAUSE THE ASSOCIATED BROWNIAN MOTION MUST BE REFLECTED ABOUT THE SINGULAR SET OF e_λ , WHICH CAUSES PROBLEMS.

Instead of imposing boundary conditions, we instead construct a stochastic process which looks like a diffusion for small times, but converges back to initial data at large times. This leads to a bound of the form

$$H^{d-1}(N_\lambda) \gtrsim_M \lambda^{-\frac{n-3}{4}}$$

A related problem is to show that nodal domains can be ‘concentrated’ near flat surfaces. Call a smooth, embedded $d - 1$ dimensional hypersurface Σ in M *admissible up to distance r* if the geodesic flow outward from Σ is injective up to a distance r , i.e. any point in the r -thickening of Σ has a unique closest point on Σ . Roughly speaking, this implies that Σ is ‘locally flat at a scale r ’.

Theorem 38.3. *For any manifold M , there exists a constant $c > 0$ such that if Σ is admissible up to a distance $\lambda^{-1/2}$, then no nodal domain N_λ can be contained in a $c\lambda^{-1/2}$ neighborhood of Σ .*

The proof will show one can generalize this result to Σ formed from a finite union of embedded manifolds admissible up to a distance $\lambda^{-1/2}$, provided these manifolds are sufficiently transversal to one another. This leads to a new proof of a classic result due to Hayman.

Corollary 38.4. *There exists $c \geq 1/900$ such that if $\Omega \subset \mathbf{R}^2$ is simply connected with inradius ρ (ρ is the largest radius of a ball contained in Ω), $\lambda_1(\Omega) \geq c/\rho^2$.*

TODO: DO WE EXPECT DENSITY OF THE NODAL SET FOR LARGE λ , i.e. $\lambda^{-1/2}$ density.

(Citation: Green, Brown, and Probability) (Citation: Stochastic Differential Equations) (Citation: Tao Brownian Motion)

Part VI

Harmonic Analysis and Fractals

Chapter 39

Riemann's Theory of Trigonometric Series

Using the techniques of measure theory, we can actually prove that the Fourier series is essentially the unique way of representing a function on any part of its domain as a trigonometric series.

Lemma 39.1. *For any sequence u_n and set E of finite measure,*

$$\lim_{n \rightarrow \infty} \int_E \cos^2(nx + u_n) dx = |E|/2$$

Proof. We have

$$\cos^2(nx + u_n) = \frac{1 + \cos(2nx + 2u_n)}{2} = \frac{1}{2} + \frac{\cos(2nx)\cos(2u_n) - \sin(2nx)\sin(2u_n)}{2}$$

Since $\cos(2u_n)$ and $\sin(2u_n)$ are bounded, we have $\int \chi_E(x) \cos(2nx)$ and $\int \chi_E(x) \sin(2nx) \rightarrow 0$ as $n \rightarrow \infty$, and the same is true for the latter component of the sum since $\cos(2u_n)$ and $\sin(2u_n)$ are bounded, we conclude that

$$\int_E \cos^2(nx + u_n) = \int \chi_E(x) \cos^2(nx + u_n) = |E|/2$$

completing the proof. \square

Theorem 39.2 (Cantor-Lebesgue Theorem). *If, for some pair of sequences a_0, a_1, \dots and b_0, b_1, \dots are chosen such that*

$$\sum_{n=0}^{\infty} a_n \cos(2\pi nx) + b_n \sin(2\pi nx)$$

converges on a set of positive measure in $[0, 1]$, then $a_n, b_n \rightarrow 0$.

Proof. Let E be the set of points upon which the trigonometric series converges. We write $a_n \cos(2\pi nx) + b_n \sin(2\pi nx) = r_n \cos(nx + c_n)$. The result of the theorem is then precisely that $r_n \rightarrow 0$. If this is not true, then we must have $\cos(nx + c_n) \rightarrow 0$ for every $x \in E$. In particular, the dominated convergence theorem implies that

$$\lim_{n \rightarrow \infty} \int_E \cos(nx + c_n)^2 dx = 0$$

Yet we know this tends to $|E|/2$ as $n \rightarrow \infty$, which is a contradiction. \square

TODO: EXPAND ON THIS FACT.

Part VII

Abstract Harmonic Analysis

The main property of spaces where Fourier analysis applies is symmetry – for a function \mathbf{R} , we can translate and negate. On \mathbf{R}^n we have not only translational symmetry but also rotational symmetry. It turns out that we can apply Fourier analysis to any ‘space with symmetry’. That is, functions on an Abelian group. We shall begin with the study of finite abelian groups, where convergence questions disappear, and with it much of the analytical questions involved in the theory. We then proceed to generalize to a study of infinite abelian groups with topological structure.

Chapter 40

Topological Groups

In abstract harmonic analysis, the main subject matter is the **topological group**, a group G equipped with a topology which makes the operation of multiplication and inversion continuous. In the mid 20th century, it was realized that basic Fourier analysis could be generalized to a large class of groups. The nicest generalization occurs over those groups which are locally compact.

Example. *There are a few groups we should keep in mind for intuition in the general topological group.*

- *The classical groups \mathbf{R}^n and \mathbf{T}^n , from which Fourier analysis originated.*
- *The group μ of roots of unity, rational numbers \mathbf{Q} , and cyclic groups \mathbf{Z}_n , which occur in number theory.*
- *The matrix subgroups of the general linear group $GL(n)$, which occur in representation theory.*
- *The product \mathbf{T}^ω of Torii, occurring in the study of Dirichlet series.*
- *The product \mathbf{Z}_2^ω , which occurs in probability theory, and other contexts.*
- *The field of p -adic numbers \mathbf{Q}_p , which are the completion of \mathbf{Q} with respect to the absolute value $|p^{-m}q|_p = p^m$.*

40.1 Basic Results

The topological structure of a topological group naturally possesses large amounts of symmetry, simplifying the spatial structure. For any topological group, the maps

$$x \mapsto gx \quad x \mapsto xg \quad x \mapsto x^{-1}$$

are homeomorphisms. Thus if U is a neighbourhood of x , then gU is a neighbourhood of gx , Ug a neighbourhood of xg , and U^{-1} a neighbourhood of x^{-1} , and as we vary U through all neighbourhoods of x , we obtain all neighbourhoods of the other points. Understanding the topological structure at any point reduces to studying the neighbourhoods of the identity element of the group.

In topological group theory it is even more important than in basic group theory to discuss set multiplication. If U and V are subsets of a group, then we define

$$U^{-1} = \{x^{-1} : x \in U\} \quad UV = \{xy : x \in U, y \in V\}$$

We let $V^2 = VV$, $V^3 = VVV$, and so on.

Theorem 40.1. *Let U and V be subsets of a topological group.*

- (i) *If U is open, then UV is open.*
- (ii) *If U is compact, and V closed, then UV is closed.*
- (iii) *If U and V are connected, UV is connected.*
- (iv) *If U and V are compact, then UV is compact.*

Proof. To see that (i) holds, we see that

$$UV = \bigcup_{x \in V} Ux$$

and each Ux is open. To see (ii), suppose $u_i v_i \rightarrow x$. Since U is compact, there is a subnet u_{i_k} converging to y . Then $y \in U$, and we find

$$v_{i_k} = u_{i_k}^{-1}(u_{i_k} v_{i_k}) \rightarrow y^{-1}x$$

Thus $y^{-1}x \in V$, and so $x = yy^{-1}x \in UV$. (iii) follows immediately from the continuity of multiplication, and the fact that $U \times V$ is connected, and (iv) follows from similar reasoning. \square

Example. If U is merely closed, then (ii) need not hold. For instance, in \mathbf{R} , take $U = \alpha\mathbf{Z}$, and $V = \mathbf{Z}$, where α is an irrational number. Then $U + V = \alpha\mathbf{Z} + \mathbf{Z}$ is dense in \mathbf{R} , and is hence not closed.

There are useful ways we can construct neighbourhoods under the group operations, which we list below.

Lemma 40.2. *Let U be a neighbourhood of the identity. Then*

- (1) *There is an open V such that $V^2 \subset U$.*
- (2) *There is an open V such that $V^{-1} \subset U$.*
- (3) *For any $x \in U$, there is an open V such that $xV \subset U$.*
- (4) *For any x , there is an open V such that $xVx^{-1} \subset U$.*

Proof. (1) follows simply from the continuity of multiplication, and (2) from the continuity of inversion. (3) is verified because $x^{-1}U$ is a neighbourhood of the origin, so if $V = x^{-1}U$, then $xV = U \subset U$. Finally (4) follows in a manner analogously to (3) because $x^{-1}Ux$ contains the origin. \square

If \mathcal{U} is an open basis at the origin, then it is only a slight generalization to show that for any of the above situations, we can always select $V \in \mathcal{U}$. Conversely, suppose that \mathcal{V} is a family of subsets of a (not yet topological) group G containing e such that (1), (2), (3), and (4) hold. Then the family $\mathcal{V}' = \{xV : V \in \mathcal{V}, x \in G\}$ forms a subbasis for a topology on G which forms a topological group. If \mathcal{V} also has the base property, then \mathcal{V}' is a basis.

Theorem 40.3. *If K and C are disjoint, K is compact, and C is closed, then there is a neighbourhood V of the origin for which KV and CV is disjoint. If G is locally compact, then we can select V such that KV is precompact.*

Proof. For each $x \in K$, C^c is an open neighbourhood containing x , so by applying the last lemma recursively we find that there is a symmetric neighbourhood V_x such that $xV_x^4 \subset C^c$. Since K is compact, finitely many of the xV_x cover K . If we then let V be the open set obtained by intersecting the finite subfamily of the V_x , then KV is disjoint from CV . \square

Taking K to be a point, we find that any open neighbourhood of a point contains a closed neighbourhood. Provided points are closed, we can set C to be a point as well.

Corollary 40.4. *Every Kolmogorov topological group is Hausdorff.*

Theorem 40.5. *For any set $A \subset G$,*

$$\overline{A} = \bigcap_V AV$$

Where V ranges over the set of neighbourhoods of the origin.

Proof. If $x \notin \overline{A}$, then the last theorem guarantees that there is V for which $\overline{A}V$ and Ax are disjoint. We conclude $\bigcap_V AV \subset \overline{A}$. Conversely, any neighbourhood contains a closed neighbourhood, so that $\overline{A} \subset AV$ for a fixed V , and hence $\overline{A} \subset \bigcap_V AV$. \square

Theorem 40.6. *Every open subgroup of G is closed.*

Proof. Let H be an open subgroup of G . Then $\overline{H} = \bigcap_V HV$. If W is a neighbourhood of the origin contained in H , then we find $\overline{H} \subset HW \subset H$, so H is closed. \square

We see that open subgroups of a group therefore correspond to connected components of the group, so that connected groups have no proper open subgroups. This also tells us that a locally compact group is σ -compact on each of its components, for if V is a pre-compact neighbourhood of the origin, then V^2, V^3, \dots are all precompact, and $\bigcup_{k=1}^{\infty} V^k$ is an open subgroup of G , which therefore contains the component of e , and is σ -compact. Since the topology of a topological group is homogenous, we can conclude that all components of the group are σ compact.

40.2 Quotient Groups

If G is a topological group, and H is a subgroup, then G/H can be given a topological structure in the obvious way. The quotient map is open, because VH is open in G for any open set V , and if H is normal, G/H is also a topological group, because multiplication is just induced from the quotient map of $G \times G$ to $G/H \times G/H$, and inversion from G to G/H . We should think the quotient structure is pleasant, but if no conditions on H are given, then G/H can have pathological structure. One particular example is the quotient \mathbf{T}/μ_{∞} of the torus modulo the roots of unity, where the quotient is lumpy.

Theorem 40.7. *If H is closed, G/H is Hausdorff.*

Proof. If $x \neq y \in G/H$, then xHy^{-1} is a closed set in G , not containing e , so we may conclude there is a neighbourhood V for which V and $VxHy^{-1}$ are disjoint, so VyH and VxH are disjoint. This implies that the open sets $V(xH)$ and $V(yH)$ are disjoint in G/H . \square

Theorem 40.8. *If G is locally compact, G/H is also.*

Proof. If $\{U_i\}$ is a basis of precompact neighbourhoods at the origin, then U_iH is a family of precompact neighbourhoods of the origin in G/H , and is in fact a basis, for if V is any neighbourhood of the origin, there is $U_i \subset \pi^{-1}(V)$, and so $U_iH \subset V$. \square

If G is a non-Hausdorff group, then $\overline{\{e\}} \neq \{e\}$, and $G/\overline{\{e\}}$ is Hausdorff. Thus we can get away with assuming all our topological groups are Hausdorff, because a slight modification in the algebraic structure of the topological group gives us this property.

40.3 Uniform Continuity

An advantage of the real line \mathbf{R} is that continuity can be explained in a *uniform sense*, because we can transport any topological questions about a certain point x to questions about topological structure near the origin via the map $g \mapsto x^{-1}g$. We can then define a uniformly continuous function $f : \mathbf{R} \rightarrow \mathbf{R}$ to be a function possessing, for every $\varepsilon > 0$, a $\delta > 0$ such that if $|y| < \delta$, $|f(x+y) - f(x)| < \varepsilon$. Instead of having to specify a δ for every point on the domain, the δ works uniformly everywhere. The group structure is all we need to talk about these questions.

We say a function $f : G \rightarrow H$ between topological groups is (left) uniformly continuous if, for any open neighbourhood U of the origin in H , there is a neighbourhood V of the origin in G such that for each x , $f(xV) \subset f(x)U$. Right continuity requires $f(Vx) \subset Uf(x)$. The requirement of distinguishing between left and right uniformity is important when we study non-commutative groups, for there are certainly left uniform maps which are not right uniform in these groups. If $f : G \rightarrow \mathbf{C}$, then left uniform continuity is equivalent to the fact that $\|L_x f - f\|_\infty \rightarrow 0$ as $x \rightarrow 1$, where $(L_x f)(y) = f(xy)$. Right uniform continuity requires $\|R_x f - f\|_\infty \rightarrow 0$,

where $(R_x f)(y) = f(yx)$. R_x is a homomorphism, but L_x is what is called an antihomomorphism.

Example. Let G be any Hausdorff non-commutative topological group, with sequences x_i and y_i for which $x_i y_i \rightarrow e$, $y_i x_i \rightarrow z \neq e$. Then the uniform structures on G are not equivalent.

It is hopeless to express uniform continuity in terms of a new topology on G , because the topology only gives a local description of continuity, which prevents us from describing things uniformly across the whole group. However, we can express uniform continuity in terms of a new topology on $G \times G$. If $U \subset G$ is an open neighbourhood of the origin, let

$$L_U = \{(x, y) : yx^{-1} \in U\} \quad R_U = \{(x, y) : x^{-1}y \in U\}$$

The family of all L_U (resp. R_U) is known as the left (right) uniform structure on G , denoted $LU(G)$ and $RU(G)$. Fix a map $f : G \rightarrow H$, and consider the map

$$g(x, y) = (f(x), f(y))$$

from G^2 to H^2 . Then f is left (right) uniformly continuous if and only if g is continuous with respect to $LU(G)$ and $LU(H)$ ($RU(G)$ and $RU(H)$). $LU(G)$ and $RU(G)$ are weaker than the product topologies on G and H , which reflects the fact that uniform continuity is a strong condition than normal continuity. We can also consider uniform maps with respect to $LU(G)$ and $RU(H)$, and so on and so forth. We can also consider uniform continuity on functions defined on an open subset of a group.

Example. Here are a few examples of easily verified continuous maps.

- If the identity map on G is left-right uniformly continuous, then $LU(G) = RU(G)$, and so uniform continuity is invariant of the uniform structure chosen.
- Translation maps $x \mapsto axb$, for $a, b \in G$, are left and right uniform.
- Inversion is uniformly continuous.

Theorem 40.9. All continuous maps on compact subsets of topological groups are uniformly continuous.

Proof. Let K be a compact subset of a group G , and let $f : K \rightarrow H$ be a continuous map into a topological group. We claim that f is then uniformly continuous. Fix an open neighbourhood V of the origin, and let V' be a symmetric neighbourhood such that $V'^2 \subset V$. For any x , there is U_x such that

$$f(x)^{-1}f(xU_x) \subset V'$$

Choose U'_x such that $U'^2_x \subset U_x$. The xU'_x cover K , so there is a finite sub-cover corresponding to sets $U'_{x_1}, \dots, U'_{x_n}$. Let $U = U'_{x_1} \cap \dots \cap U'_{x_n}$. Fix $y \in G$, and suppose $y \in x_k U'_{x_k}$. Then

$$\begin{aligned} f(y)^{-1}f(yU) &= f(y)^{-1}f(x_k)f(x_k)^{-1}f(yU) \\ &\subset f(y)^{-1}f(x_k)f(x_k)^{-1}f(x_k U x_k) \\ &\subset f(y)^{-1}f(x_k)V' \\ &\subset V'^2 \subset V \end{aligned}$$

So that f is left uniformly continuous. Right uniform continuity is proven in the exact same way. \square

Corollary 40.10. *All maps with compact support are uniformly continuous.*

Corollary 40.11. *Uniform continuity on compact groups is invariant of the uniform structure chosen.*

40.4 Ordered Groups

In this section we describe a general class of groups which contain both interesting and pathological examples. Let G be a group with an ordering $<$ preserved by the group operations, so that $a < b$ implies both $ag < bg$ and $ga < gb$. We now prove that the order topology gives G the structure of a normal topological group (the normality follows because of general properties of order topologies).

First note, that $a < b$ implies $a^{-1} < b^{-1}$. This results from a simple algebraic trick, because

$$a^{-1} = a^{-1}bb^{-1} > a^{-1}ab^{-1} = b^{-1}$$

This implies that the inverse image of an interval (a, b) under inversion is (b^{-1}, a^{-1}) , hence inversion is continuous.

Now let $e < b < a$. We claim that there is then $e < c$ such that $c^2 < a$. This follows because if $b^2 \geq a$, then $b \geq ab^{-1}$ and so

$$(ab^{-1})^2 = ab^{-1}ab^{-1} \leq ab^{-1}b = a$$

Now suppose $a < e < b$. If $\inf\{y : y > e\} = x > e$, then $(x^{-1}, x) = \{e\}$, and the topology on G is discrete, hence the continuity of operations is obvious. Otherwise, we may always find c such that $c^2 < b$, $a < c^{-2}$, and then if $c^{-1} < g, h < c$, then

$$a < c^{-2} < gh < c^2 < b$$

so multiplication is continuous at every pair (x, x^{-1}) . In the general case, if $a < gh < b$, then $g^{-1}ah^{-1} < e < g^{-1}bh^{-1}$, so there is c such that if $c^{-1} < g', h' < c$, then $g^{-1}ah^{-1} < g'h' < g^{-1}bh^{-1}$, so $a < gg'h'h < b$. The set of gg' , where $c^{-1} < g' < c$, is really just the set of $gc^{-1} < x < gc$, and the set of $h'h$ is really just the set of $c^{-1}h < x < ch$. Thus multiplication is continuous everywhere.

Example (Dieudonne). *For any well ordered set S , the dictionary ordering on \mathbf{R}^S induces a linear ordering inducing a topological group structure on the set of maps from S to \mathbf{R} .*

Let us study Dieudonne's topological group in more detail. If S is a finite set, or more generally possesses a maximal element w , then the topology on \mathbf{R}^S can be defined such that $f_i \rightarrow f$ if eventually $f_i(s) = f(s)$ for all $s < w$ simultaneously, and $f_i(w) \rightarrow f(w)$. Thus \mathbf{R}^S is isomorphic (topologically) to a discrete union of a certain number of copies of \mathbf{R} , one for each tuple in $S - \{w\}$.

If S has a countable cofinal subset $\{s_i\}$, the topology is no longer so simple, but \mathbf{R}^S is still first countable, because the sets

$$U_i = \{f : (\forall w < s_i : f(w) = 0)\}$$

provide a countable neighbourhood basis of the origin.

The strangest properties of \mathbf{R}^S occur when S has no countable cofinal set. Suppose that $f_i \rightarrow f$. We claim that it follows that $f_i = f$ eventually. To prove by contradiction, we assume without loss of generality (by thinning the sequence) that no f_i is equal to f . For each f_i , find the largest $w_i \in S$ such that for $s < w_i$, $f_i(s) = f(s)$ (since S is well ordered, the set of

elements for which $f_i(s) \neq f(s)$ has a minimal element). Then the w_i form a countable cofinal set, because if $v \in S$ is arbitrary, the f_i eventually satisfy $f_i(s) = f(s)$ for $s < v$, hence the corresponding w_i is greater than v_i . Hence, if $f_i \rightarrow f$ in \mathbf{R}^S , where S does not have a countable cofinal subset, then eventually $f_i = f$. We conclude all countable sets in \mathbf{R}^S are closed, and this proof easily generalises to show that if S does not have a cofinal set of cardinality \mathfrak{a} , then every set of cardinality $\leq \mathfrak{a}$ is closed.

The simple corollary to this proof is that compact subsets are finite. Let $X = f_1, f_2, \dots$ be a denumerable, compact set. Since all subsets of X are compact, we may assume $f_1 < f_2 < \dots$ (or $f_1 > f_2 > \dots$, which does not change the proof in any interesting way). There is certainly $g \in \mathbf{R}^S$ such that $g < f_1$, and then the sets $(g, f_2), (f_1, f_3), (f_2, f_4), \dots$ form an open cover of X with no finite subcover, hence X cannot be compact. We conclude that the only compact subsets of \mathbf{R}^S are finite.

Furthermore, the class of open sets is closed under countable intersections. Consider a series of functions

$$f_1 \leq f_2 \leq \dots < h < \dots \leq g_2 \leq g_1$$

Suppose that $f_i \leq k < h < k' \leq g_j$. Then the intersection of the (f_i, g_i) contains an interval (k, k') around h , so that the intersection is open near h . The only other possibility is that $f_i \rightarrow h$ or $g_i \rightarrow h$, which can only occur if $f_i = h$ or $g_i = h$ eventually, in which case we cannot have $f_i < h, h < g_i$. We conclude the intersection of countably many intervals is open, because we can always adjust any intersection to an intersection of this form without changing the resulting intersecting set (except if the set is empty, in which case the claim is trivial). The general case results from noting that any open set in an ordered group is a union of intervals.

40.5 Topological Groups arising from Normal subgroups

Let G be a group, and \mathcal{N} a family of normal subgroups closed under intersection. If we interpret \mathcal{N} as a neighbourhood base at the origin, the resulting topology gives G the structure of a totally disconnected topological group, which is Hausdorff if and only if $\bigcap \mathcal{N} = \{e\}$. First note that $g_i \rightarrow g$ if g_i is eventually in gN , for every $N \in \mathcal{N}$, which implies

$g_i^{-1} \in Ng^{-1} = g^{-1}N$, hence inversion is continuous. Furthermore, if h_i is eventually in hN , then $g_i h_i \in gNhN = ghN$, so multiplication is continuous. Finally note that $N^c = \bigcup_{g \neq e} gN$ is open, so that every open set is closed.

Example. Consider $\mathcal{N} = \{\mathbf{Z}, 2\mathbf{Z}, 3\mathbf{Z}, \dots\}$. Then \mathcal{N} induces a Hausdorff topology on \mathbf{Z} , such that $g_i \rightarrow g$, if and only if g_i is eventually in $g + n\mathbf{Z}$ for all n . In this topology, the series $1, 2, 3, \dots$ converges to zero!

This example gives us a novel proof, due to Furstenburg, that there are infinitely many primes. Suppose that there were only finitely many, $\{p_1, p_2, \dots, p_n\}$. By the fundamental theorem of arithmetic,

$$\{-1, 1\} = (\mathbf{Z}p_1)^c \cap \dots \cap (\mathbf{Z}p_n)^c$$

and is therefore an open set. But this is clearly not the case as open sets must contain infinite sequences.

Chapter 41

The Haar Measure

One of the reasons that we isolate locally compact groups to study is that they possess an incredibly useful object allowing us to understand functions on the group, and thus the group itself. A **left (right) Haar measure** for a group G is a Radon measure μ for which $\mu(xE) = \mu(E)$ for any $x \in G$ and measurable E ($\mu(Ex) = \mu(E)$ for all x and E). For commutative groups, all left Haar measures are right Haar measures, but in non-commutative groups this need not hold. However, if μ is a right Haar measure, then $\nu(E) = \mu(E^{-1})$ is a left Haar measure, so there is no loss of generality in focusing our study on left Haar measures.

Example. *The example of a Haar measure that everyone knows is the Lebesgue measure on \mathbf{R} (or \mathbf{R}^n). It commutes with translations because it is the measure induced by the linear functional corresponding to Riemann integration on $C_c^+(\mathbf{R}^n)$. A similar theory of Darboux integration can be applied to linearly ordered groups, leading to the construction of a Haar measure on such a group.*

Example. *If G is a Lie group, consider a 2-tensor $g_e \in T_e^2(G)$ inducing an inner product at the origin. Then the diffeomorphism $f : a \mapsto b^{-1}a$ allows us to consider $g_b = f^* \lambda \in T_b^2(G)$, and this is easily verified to be an inner product, hence we have a Riemannian metric. The associated Riemannian volume element can be integrated, producing a Haar measure on G .*

Example. *If G and H have Haar measures μ and ν , then $G \times H$ has a Haar measure $\mu \times \nu$, so that the class of topological groups with Haar measures is closed under the product operation. We can even allow infinite products, provided that the groups involved are compact, and the Haar measures are normalized*

to probability measures. This gives us measures on F_2^ω and \mathbf{T}^ω , which models the probability of an infinite sequence of coin flips.

Example. dx/x is a Haar measure for the multiplicative group of positive real numbers, since

$$\int_a^b \frac{1}{x} = \log(b) - \log(a) = \log(cb) - \log(ca) = \int_{ca}^{cb} \frac{1}{x}$$

If we take the multiplicative group of all non-negative real numbers, the Haar measure becomes $dx/|x|$.

Example. $dxdy/(x^2 + y^2)$ is a Haar measure for the multiplicative group of complex numbers, since we have a basis of ‘arcs’ around the origin, and by a change of variables to polar coordinates, we verify the integral is changed by multiplication. Another way to obtain this measure is by noticing that \mathbf{C}^\times is topologically isomorphic to the product of the circle group and the multiplicative group of real numbers, and hence the measure obtained should be the product of these measures. Since

$$\frac{dxdy}{x^2 + y^2} = \frac{drd\theta}{r}$$

We see that this is just the product of the Haar measure on \mathbf{R}^+ , dr/r , and the Haar measure on \mathbf{T} , $d\theta$.

Example. The space $M_n(\mathbf{R})$ of all n by n real matrices under addition has a Haar measure dM , which is essentially the Lebesgue measure on \mathbf{R}^{n^2} . If we consider the measure on $GL_n(\mathbf{R})$, defined by

$$\frac{dM}{\det(M)^n}$$

To see this, note the determinant of the map $M \mapsto NM$ on $M_n(\mathbf{R})$ is $\det(N)^n$, because we can view $M_n(\mathbf{R})$ as the product of \mathbf{R}^n n times, multiplication operates on the space componentwise, and the volume of the image of the unit parallelliped in each \mathbf{R}^n is $\det(N)$. Since the multiplicative group of complex numbers $z = x + iy$ can be identified with the group of matrices of the form

$$\begin{pmatrix} x & -y \\ y & x \end{pmatrix}$$

and the measure on $\mathbf{C} - \{0\}$ then takes the form $dM/\det(M)$. More generally, if G is an open subset of \mathbf{R}^n , and left multiplication acts affinely, $xy = A(x)y + b(x)$, then $dx/|\det(A(x))|$ is a left Haar measure on G , where dx is Lebesgue measure.

It turns out that there is a Haar measure on any locally compact group, and what's more, it is unique up to scaling. The construction of the measure involves constructing a positive linear functional $\phi : C_c(G) \rightarrow \mathbf{R}$ such that $\phi(L_x f) = \phi(f)$ for all x . The Riesz representation theorem then guarantees the existence of a Radon measure μ which represents this linear functional, and one then immediately verifies that this measure is a Haar measure.

Theorem 41.1. *Every locally compact group G has a Haar measure.*

Proof. The idea of the proof is fairly simple. If μ was a Haar measure, $f \in C_c^+(G)$ was fixed, and $\phi \in C_c^+(G)$ was a function supported on a small set, and behaving like a step function, then we could approximate f well by translates of ϕ ,

$$f(x) \approx \sum c_i(L_{x_i}\phi)$$

Hence

$$\int f(x)d\mu \approx \sum c_i \int L_{x_i}\phi = \sum c_i \int \phi$$

If $\int \phi = 1$, then we could approximate $\int f(x)d\mu$ as literal sums of coefficients c_i . Since μ is outer regular, and ϕ is supported on neighbourhoods, one can show $\int f(x)d\mu$ is the infimum of $\sum c_i$, over all choices of $c_i > 0$ and $\int \phi \geq 1$, for which $f \leq \sum c_i L_{x_i}\phi$. Without the integral, we cannot measure the size of the functions ϕ , so we have to normalize by a different factor. We define $(f : \phi)$ to be the infimum of the sums $\sum c_i$, where $f \leq \sum c_i L_{x_i}\phi$ for some $x_i \in G$. We would then have

$$\int f d\mu \leq (f : \phi) \int \phi d\mu$$

If k is fixed with $\int k = 1$, then we would have

$$\int f d\mu \leq (f : \phi)(\phi : k)$$

We cannot change k if we wish to provide a limiting result in ϕ , so we notice that $(f : g)(g : h) \leq (f : h)$, which allows us to write

$$\int f d\mu \leq \frac{(f : \phi)}{(k : \phi)}$$

Taking the support of ϕ to be smaller and smaller, this value should approximate the integral perfectly accurately.

Define the linear functional

$$I_\phi(f) = \frac{(f : \phi)}{(k : \phi)}$$

Then I_ϕ is a sublinear, monotone, function with a functional bound

$$(k : f)^{-1} \leq I_\phi(f) \leq (f : k)$$

Which effectively says that, regardless of how badly we choose ϕ , the approximation factor $(f : \phi)$ is normalized by the approximation factor $(k : \phi)$ so that the integral is bounded. Now we need only prove that I_ϕ approximates a linear functional well enough that we can perform a limiting process to obtain a Haar integral. If $\varepsilon > 0$, and $g \in C_c^+(G)$ with $g = 1$ on $\text{supp}(f_1 + f_2)$, then the functions

$$h = f_1 + f_2 + \varepsilon g$$

$$h_1 = f_1/h \quad h_2 = f_2/h$$

are in $C_0^+(G)$, if we define $h_i(x) = 0$ if $f_i(x) = 0$. This implies that there is a neighbourhood V of e such that if $x \in V$, and y is arbitrary, then

$$|h_1(xy) - h_1(y)| \leq \varepsilon \quad |h_2(xy) - h_2(y)| < \varepsilon$$

If $\text{supp}(\phi) \subset V$, and $h \leq \sum c_i L_{x_i} \phi$, then

$$f_j(x) = h(x)h_j(x) \leq \sum c_i \phi(x_i x) h_j(x) \leq \sum c_i \phi(x_i x) [h_j(x_i^{-1}) + \varepsilon]$$

since we may assume that $x_i x \in \text{supp}(\phi) \subset V$. Then, because $h_1 + h_2 \leq 1$,

$$(f_1 : \phi) + (f_2 : \phi) \leq \sum c_j [h_1(x_j^{-1}) + \varepsilon] + \sum c_j [h_2(x_j^{-1}) + \varepsilon] \leq \sum c_j [1 + 2\varepsilon]$$

Now we find, by taking infimums, that

$$I_\phi(f_1) + I_\phi(f_2) \leq I_\phi(h)(1 + 2\varepsilon) \leq [I_\phi(f_1 + f_2) + \varepsilon I_\phi(g)][1 + 2\varepsilon]$$

Since g is fixed, and we have a bound $I_\phi(g) \leq (g : k)$, we may always find a neighbourhood V (dependant on f_1, f_2) for any $\varepsilon > 0$ such that

$$I_\phi(f_1) + I_\phi(f_2) \leq I_\phi(f_1 + f_2) + \varepsilon$$

if $\text{supp}(\phi) \subset V$.

Now we have estimates on how well I_ϕ approximates a linear function, so we can apply a limiting process. Consider the product

$$X = \prod_{f \in C_0^+(G)} [(k : f)^{-1}, (k : f_0)]$$

a compact space, by Tychonoff's theorem, consisting of $F : C_c^+(G) \rightarrow \mathbf{R}$ such that $(k : f)^{-1} \leq F(f) \leq (f : k)$. For each neighbourhood V of the identity, let $K(V)$ be the closure of the set of I_ϕ such that $\text{supp}(\phi) \subset V$. Then the set of all $K(V)$ has the finite intersection property, so we conclude there is some $I : C_c^+(G) \rightarrow \mathbf{R}$ contained in $\bigcap K(V)$. This means that every neighbourhood of I contains I_ϕ with $\text{supp}(\phi) \subset V$, for all ϕ . This means that if $f_1, f_2 \in C_c^+(G)$, $\varepsilon > 0$, and V is arbitrary, there is ϕ with $\text{supp}(\phi) \subset V$, and

$$|I(f_1) - I_\phi(f_1)| < \varepsilon \quad |I(f_2) - I_\phi(f_2)| < \varepsilon$$

$$|I(f_1 + f_2) - I_\phi(f_1 + f_2)| < \varepsilon$$

this implies that if V is chosen small enough, then

$$|I(f_1 + f_2) - (I(f_1) + I(f_2))| \leq 2\varepsilon + |I_\phi(f_1 + f_2) - (I_\phi(f_1) + I_\phi(f_2))| < 3\varepsilon$$

Taking $\varepsilon \rightarrow 0$, we conclude I is linear. Similar limiting arguments show that I is homogenous of degree 1, and commutes with all left translations. We conclude the extension of I to a linear functional on $C_0(G)$ is well defined, and the Radon measure obtained by the Riesz representation theorem is a Haar measure. \square

We shall prove that the Haar measure is unique, but first we show an incredibly useful regularity property.

Proposition 41.2. *If U is open, and μ is a Haar measure, then $\mu(U) > 0$. It follows that if f is in $C_c^+(G)$, then $\int f d\mu > 0$.*

Proof. If $\mu(U) = 0$, then for any $x_1, \dots, x_n \in G$,

$$\mu\left(\bigcup_{i=1}^n x_i U\right) \leq \sum_{i=1}^n \mu(x_i U) = 0$$

If K is compact, then K can be covered by finitely many translates of U , so $\mu(K) = 0$. But then $\mu = 0$ by regularity, a contradiction. \square

Theorem 41.3. *Haar measures are unique up to a multiplicative constant.*

Proof. Let μ and ν be Haar measures. Fix a compact neighbourhood V of the identity. If $f, g \in C_c^+(G)$, consider the compact sets

$$A = \text{supp}(f)V \cup V\text{supp}(f) \quad B = \text{supp}(g)V \cup V\text{supp}(g)$$

Then the functions $F_y(x) = f(xy) - f(yx)$ and $G_y(x) = g(xy) - g(yx)$ are supported on A and B . There is a neighbourhood $W \subset V$ of the identity such that $\|F_y\|_\infty, \|G_y\|_\infty < \varepsilon$ if $y \in W$. Now find $h \in C_c^+(G)$ with $h(x) = h(x^{-1})$ and $\text{supp}(h) \subset W$ (take $h(x) = k(x)k(x^{-1})$ for some function $k \in C_c^+(G)$ with $\text{supp}(k) \subset W$, and $k = 1$ on a symmetric neighbourhood of the origin). Then

$$\begin{aligned} \left(\int h d\mu\right) \left(\int f d\lambda\right) &= \int h(y)f(x) d\mu(y) d\lambda(x) \\ &= \int h(y)f(yx) d\mu(y) d\lambda(x) \end{aligned}$$

and

$$\begin{aligned} \left(\int h d\lambda\right) \left(\int f d\mu\right) &= \int h(x)f(y) d\mu(y) d\lambda(x) \\ &= \int h(y^{-1}x)f(y) d\mu(y) d\lambda(x) \\ &= \int h(x^{-1}y)f(y) d\mu(y) d\lambda(x) \\ &= \int h(y)f(xy) d\mu(y) d\lambda(x) \end{aligned}$$

Hence, applying Fubini's theorem,

$$\begin{aligned} \left| \int h d\mu \int f d\lambda - \int h d\lambda \int f d\mu \right| &\leq \int h(y) |F_y(x)| d\mu(y) d\lambda(x) \\ &\leq \varepsilon \lambda(A) \int h d\mu \end{aligned}$$

In the same way, we find this is also true when f is swapped with g , and A with B . Dividing these inequalities by $\int h d\mu \int f d\mu$, we find

$$\left| \frac{\int f d\lambda}{\int f d\mu} - \frac{\int h d\lambda}{\int h d\mu} \right| \leq \frac{\varepsilon \lambda(A)}{\int f d\mu}$$

and this inequality holds with f swapped out with g , A with B . We then combine these inequalities to conclude

$$\left| \frac{\int f d\lambda}{\int f d\mu} - \frac{\int g d\lambda}{\int g d\mu} \right| \leq \varepsilon \left[\frac{\lambda(A)}{\int f d\mu} + \frac{\lambda(B)}{\int g d\mu} \right]$$

Taking ε to zero, we find $\lambda(A), \lambda(B)$ remain bounded, and hence

$$\frac{\int f d\lambda}{\int f d\mu} = \frac{\int g d\lambda}{\int g d\mu}$$

Thus there is a constant $c > 0$ such that $\int f d\lambda = c \int f d\mu$ for any function $f \in C_c^+(G)$, and we conclude that $\lambda = c\mu$. \square

The theorem can also be proven by looking at the translation invariant properties of the derivative $f = d\mu/d\nu$, where $\nu = \mu + \lambda$ (We assume our group is σ compact for now). Consider the function $g(x) = f(yx)$. Then

$$\int_A g(x) d\nu = \int_{yA} f(x) d\nu = \mu(yA) = \mu(A)$$

so g is derivative, and thus $f = g$ almost everywhere. Our interpretation is that for a fixed y , $f(yx) = f(x)$ almost everywhere with respect to ν . Then (applying a discrete version of Fubini's theorem), we find that for almost all x with respect to ν , $f(yx) = f(x)$ holds for almost all y . But this implies that there exists an x for which $f(yx) = f(x)$ holds almost everywhere. Thus for any measurable A ,

$$\mu(A) = \int_A f(y) d\nu(y) = f(x) \nu(A) = f(x) \mu(A) + f(x) \nu(A)$$

Now $(1 - f(x))\mu(A) = f(x)\nu(A)$ for all A , implying (since $\mu, \nu \neq 0$), that $f(x) \neq 0, 1$, and so

$$\frac{1 - f(x)}{f(x)}\mu(A) = \nu(A)$$

for all A . This shows the uniqueness property for all σ compact groups. If G is an arbitrary group with two measures μ and ν , then there is c such that $\mu = c\nu$ on every component of G , and thus on the union of countably many components. If A intersects uncountably many components, then either $\mu(A) = \nu(A) = \infty$, or the intersection of A on each set has positive measure on only countably many components, and in either case we have $\mu(A) = \nu(A)$.

41.1 Fubini, Radon Nikodym, and Duality

Before we continue, we briefly mention that integration theory is particularly nice over locally compact groups, even if we do not have σ finiteness. This essentially follows because the component of the identity in G is σ compact (take a compact neighbourhood and its iterated multiples), hence all components in G are σ compact. The three theorems that break down outside of the σ compact domain are Fubini's theorem, the Radon Nikodym theory, and the duality between $L^1(X)$ and $L^\infty(X)$. We show here that all three hold if X is a locally compact topological group.

First, suppose that $f \in L^1(G \times G)$. Then the essential support of f is contained within countably many components of $G \times G$ (which are simply products of components in G). Thus f is supported on a σ compact subset of $G \times G$ (as a locally compact topological group, each component of $G \times G$ is σ compact), and we may apply Fubini's theorem on the countably many components (the countable union of σ compact sets is σ compact). The functions in $L^p(G)$, for $1 \leq p < \infty$, also vanish outside of a σ compact subset (for if $f \in L^p(G)$, $|f|^p \in L^1(G)$ and thus vanishes outside of a σ compact set). What's more, all finite sums and products of functions from these sets (in either variable) vanish outside of σ compact subsets, so we almost never need to explicitly check the conditions for satisfying Fubini's theorem, and from now on we apply it wantonly.

Now suppose μ and ν are both Radon measures, with $\nu \ll \mu$, and ν is σ -finite. By inner regularity, the support of ν is a σ compact set E . By inner regularity, μ restricted to E is σ finite, and so we may find a Radon

Nikodym derivative on E . This derivative can be extended to all of G because ν vanishes on G .

Finally, we note that $L^\infty(X) = L^1(X)^*$ can be made to hold if X is not σ finite, but locally compact and Hausdorff, provided we are integrating with respect to a Radon measure μ , and we modify $L^\infty(G)$ slightly. Call a set $E \subset X$ **locally Borel** if $E \cap F$ is Borel whenever F is Borel and $\mu(F) < \infty$. A locally Borel set is **locally null** if $\mu(E \cap F) = 0$ whenever $\mu(F) < \infty$ and F is Borel. We say a property holds **locally almost everywhere** if it is true except on a locally null set. $f : X \rightarrow \mathbf{C}$ is **locally measurable** if $f^{-1}(U)$ is locally Borel for every borel set $U \subset \mathbf{C}$. We now define $L^\infty(X)$ to be the space of all functions bounded except on a locally null set, modulo functions that are locally zero. That is, we define a norm

$$\|f\|_\infty = \inf\{c : |f(x)| \leq c \text{ locally almost everywhere}\}$$

and then $L^\infty(X)$ consists of the functions that have finite norm. It then follows that if $f \in L^\infty(X)$ and $g \in L^1(X)$, then fg vanishes outside of a σ -finite set Y , so $fg \in L^1(X)$, and if we let $Y_1 \subset Y_2 \subset \dots \rightarrow Y$ be an increasing subsequence such that $\mu(Y_i) < \infty$, then $|f(x)| \leq \|f\|_\infty$ almost everywhere for $x \in Y_i$, and so by the monotone convergence theorem

$$\int |fg| d\mu = \lim_{Y_i \rightarrow \infty} \int_{Y_i} |fg| d\mu \leq \|f\|_\infty \int_{Y_i} |g| d\mu \leq \|f\|_\infty \|g\|_1$$

Thus the map $g \mapsto \int fg d\mu$ is a well defined, continuous linear functional with norm $\|f\|_\infty$. That $L^1(X)^* = L^\infty(X)$ follows from the decomposability of the Carathéodory extension of μ , a fact we leave to the general measure theorists.

41.2 Unimodularity

We have thus defined a left invariant measure, but make sure to note that such a function is not right invariant. We call a group whose left Haar measure is also right invariant **unimodular**. Obviously all abelian groups are unimodular.

Given a fixed y , the measure $\mu_y(A) = \mu(Ay)$ is a new Haar measure on the space, hence there is a constant $\Delta(y) > 0$ depending only on y such that $\mu(Ay) = \Delta(y)\mu(A)$ for all measurable A . Since $\mu(Axy) = \Delta(y)\mu(Ay) =$

$\Delta(x)\Delta(y)\mu(A)$, we find that $\Delta(xy) = \Delta(x)\Delta(y)$, so Δ is a homomorphism from G to the multiplicative group of real numbers. For any $f \in L^1(\mu)$, we have

$$\int f(xy)d\mu(x) = \Delta(y^{-1}) \int f(x)d\mu(x)$$

If $y_i \rightarrow e$, and $f \in C_c(G)$, then $\|R_{y_i}f - f\|_\infty \rightarrow 0$, so

$$\Delta(y_i^{-1}) \int f(x)d\mu = \int f(xy_i)d\mu \rightarrow \int f(x)d\mu$$

Hence $\Delta(y_i^{-1}) \rightarrow 1$. This implies Δ , known as the *modular function*, is a continuous homomorphism from G to the real numbers. Note that Δ is trivial if and only if G is unimodular.

Theorem 41.4. *Any compact group is unimodular.*

Proof. $\Delta : G \rightarrow \mathbf{R}^*$ is a continuous homomorphism, hence $\Delta(G)$ is compact. But the only compact subgroup of \mathbf{R} is trivial, hence Δ is trivial. \square

Let G^c be the smallest closed subgroup of G containing the commutators $[x, y] = xyx^{-1}y^{-1}$. It is verified to be a normal subgroup of G by simple group theory.

Theorem 41.5. *If G/G^c is compact, then G is unimodular.*

Proof. Δ factors through G/G^c since it is abelian. But if Δ is trivial on G/G^c , it must also be trivial on G . \square

Corollary 41.6. *If G is a connected, semi-simple Lie group, then G is unimodular.*

Proof. The Lie algebra \mathfrak{g} of G is a direct sum of simple Lie algebras, and so $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$. But $[\mathfrak{g}, \mathfrak{g}]$ is the Lie algebra of $[G, G]$, so $G = [G, G]$ since G is connected. \square

More generally, if G is a connected Lie group, and Ad denotes the adjoint representation of G on \mathfrak{g} , then $\Delta(x) = \det(\text{Ad}(x^{-1}))$. TODO

The modular function relates right multiplication to left multiplication in the group. In particular, if $d\mu$ is a Left Haar measure, then $\Delta^{-1}d\mu$ is a right Haar measure. Hence any right Haar measure is a constant multiple

of $\Delta^{-1}d\mu$. Hence the measure $\nu(A) = \mu(A^{-1})$ has a value c such that for any function f ,

$$\int \frac{f(x)}{\Delta(x)} d\mu(x) = c \int f(x) d\nu(x) = c \int f(x^{-1}) d\mu$$

If $c \neq 1$, pick a symmetric neighbourhood U such that for $x \in U$, $|\Delta(x) - 1| \leq \varepsilon|c - 1|$. Then if $f > 0$

$$|c - 1|\mu(U) = |c\mu(U^{-1}) - \mu(U)| = \left| \int_U [\Delta(x^{-1}) - 1] d\mu(x) \right| \leq \varepsilon\mu(U)|c - 1|$$

A contradiction if $\varepsilon < 1$. Thus we have

$$\int f(x^{-1}) d\mu(x) = \int \frac{f(x)}{\Delta(x)} d\mu(x)$$

A useful integration trick. When Δ is unbounded, then it follows that $L^p(\mu)$ and $L^p(\nu)$ do not consist of the same functions. There are two ways of mapping the sets isomorphically onto one another – the map $f(x) \mapsto f(x^{-1})$, and the map $f(x) \mapsto \Delta(x)^{1/p} f(x)$.

From now on, we assume a left invariant Haar measure is fixed over an entire group. Since a Haar measure is uniquely determined up to a constant, this is no loss of generality, and we might as well denote our integration factors $d\mu(x)$ and $d\mu(y)$ as dx and dy , where it is assumed that this integration is over the Lebesgue measure.

41.3 Convolution

If G is a topological group, then $C(G)$ does not contain enough algebraic structure to identify G – for instance, if G is a discrete group, then $C(G)$ is defined solely by the cardinality of G . The algebras we wish to study over G is the space $M(G)$ of all complex valued Radon measures over G and the space $L^1(G)$ of integrable functions with respect to the Haar measure, because here we can place a Banach algebra structure with an involution. We note that $L^1(G)$ can be isometrically identified as the space of all measures $\mu \in M(G)$ which are absolutely continuous with respect to the Haar measure. Given $\mu, \nu \in M(G)$, we define the convolution measure

$$\int \phi d(\mu * \nu) = \int \phi(xy) d\mu(x) d\nu(y)$$

The measure is well defined, for if $\phi \in C_c^+(X)$ is supported on a compact set K , then

$$\begin{aligned} \left| \int \phi(xy) d\mu(x) d\nu(y) \right| &\leq \int_G \int_G \phi(xy) d|\mu|(x) d|\nu|(y) \\ &\leq \|\mu\| \|\nu\| \|\phi\|_\infty \end{aligned}$$

This defines an operation on $M(G)$ which is associative, since, by applying the associativity of G and Fubini's theorem.

$$\begin{aligned} \int \phi d((\mu * \nu) * \lambda) &= \int \int \phi(xz) d(\mu * \nu)(x) d\lambda(z) \\ &= \int \int \int \phi((xy)z) d\mu(x) d\nu(y) d\lambda(z) \\ &= \int \int \int \phi(x(yz)) d\mu(x) d\nu(y) d\lambda(z) \\ &= \int \int \phi(xz) d\mu(x) d(\nu * \lambda)(z) \\ &= \int \phi d(\mu * (\nu * \lambda)) \end{aligned}$$

Thus we begin to see how the structure of G gives us structure on $M(G)$. Another example is that convolution is commutative if and only if G is commutative. We have the estimate $\|\mu * \nu\| \leq \|\mu\| \|\nu\|$, because of the bound we placed on the integrals above. $M(G)$ is therefore an involutive Banach algebra, which has a unit, the dirac delta measure at the identity.

As a remark, we note that involutive Banach algebras have nowhere as near a nice of a theory than that of C^* algebras. $M(G)$ cannot be renormed to be a C^* algebra, since every weakly convergent Cauchy sequence converges, which is impossible in a C^* algebra, except in the finite dimensional case.

A **discrete measure** on G is a measure in $M(G)$ which vanishes outside a countable set of points, and the set of all such measures is denoted $M_d(G)$. A **continuous measure** on G is a measure μ such that $\mu(\{x\}) = 0$ for all $x \in G$. We then have a decomposition $M(G) = M_d(G) \oplus M_c(G)$, for if μ is any measure, then $\mu(\{x\}) \neq 0$ for at most countably many points x , for

$$\|\mu\| \geq \sum_{x \in G} |\mu|(x)$$

This gives rise to a discrete measure ν , and $\mu - \nu$ is continuous. If we had another decomposition, $\mu = \psi + \phi$, then $\mu(\{x\}) = \psi(\{x\}) = \nu(\{x\})$, so $\psi = \nu$ by discreteness, and we then conclude $\phi = \mu - \nu$. $M_c(G)$ is actually a closed subspace of $M(G)$, since if $\mu_i \rightarrow \mu$, and $\mu_i \in M_c(G)$, and $\|\mu_i - \mu\| < \varepsilon$, then for any $x \in G$,

$$\varepsilon > \|\mu - \mu_i\| \geq |(\mu_i - \mu)(\{x\})| = |\mu(\{x\})|$$

Letting $\varepsilon \rightarrow 0$ shows continuity.

The convolution on $M(G)$ gives rise to a convolution on $L^1(G)$, where

$$(f * g)(x) = \int f(y)g(y^{-1}x)dy$$

which satisfies $\|f * g\|_1 \leq \|f\|_1 \|g\|_1$. This is induced by the identification of f with $f(x)dx$, because then

$$\begin{aligned} \int \phi(f(x)dx * g(x)dx) &= \int \int \phi(yx)f(y)g(x)dydx \\ &= \int \phi(y) \left(\int f(y)g(y^{-1}x)dx \right) dy \end{aligned}$$

Hence $f d\mu * g d\mu = (f * g)d\mu$. What's more,

$$\|f\|_1 = \|f d\mu\|$$

If $\nu \in M(G)$, then we can still define $\nu * f \in L^1(G)$

$$(\nu * f)(x) = \int f(y^{-1}x)d\mu(y)$$

which holds since

$$\int \phi d(\nu * f\mu) = \int \phi(yx)f(x)d\nu(y)d\mu(x) = \int \phi(x)f(y^{-1}x)d\nu(y)d\mu(x)$$

If G is unimodular, then we also find

$$\int \phi d(f\mu * \nu) = \int \phi(yx)f(y)d\mu(y)d\nu(x) = \int \phi(x)f(y)d\mu(y)d\nu(y^{-1}x)$$

So we let $f * \mu(x) = \int f(y)d\mu(y^{-1}x)$.

Theorem 41.7. $L^1(G)$ and $M_c(G)$ are closed ideals in $M(G)$, and $M_d(G)$ is a closed subalgebra.

Proof. If $\mu_i \rightarrow \mu$, and each μ_i is discrete, the μ is discrete, because there is a countable set K such that all μ_i are equal to zero outside of K , so μ must also vanish outside of K (here we have used the fact that $M(G)$ is a Banach space, so that we need only consider sequences). Thus $M_d(G)$ is closed, and is easily verified to be subalgebra, essentially because $\delta_x * \delta_y = \delta_{xy}$. If $\mu_i \rightarrow \mu$, then $\mu_i(\{x\}) \rightarrow \mu(\{x\})$, so that $M_c(G)$ is closed in $M(G)$. If ν is an arbitrary measure, and μ is continuous, then

$$(\mu * \nu)(\{x\}) = \int_G \mu(\{y\}) d\nu(y^{-1}x) = 0$$

$$(\nu * \mu)(\{x\}) = \int_G \mu(\{y\}) d\nu(xy^{-1}) = 0$$

so $M_c(G)$ is an ideal. Finally, we verify $L^1(G)$ is closed, because it is complete, and if $\nu \in M(G)$ is arbitrary, and if U has null Haar measure, then

$$(f dx * \nu)(U) = \int \chi_U(xy) f(x) dx d\nu(y) = \int_G \int_{y^{-1}U} f(x) dx d\nu(y) = 0$$

$$(\nu * f dx)(U) = \int \chi_U(xy) d\nu(x) f(y) dy = \int_G \int_{Ux^{-1}} f(y) dy d\nu(x) = 0$$

So $L^1(G)$ is a two-sided ideal. □

If we wish to integrate by right multiplication instead of left multiplication, we find by the substitution $y \mapsto xy$ that

$$\begin{aligned} (f * g)(x) &= \int f(y) g(y^{-1}x) dy \\ &= \int \int f(xy) g(y^{-1}) dy \\ &= \int \int \frac{f(xy^{-1}) g(y)}{\Delta(y)} dy \end{aligned}$$

Observe that

$$f * g = \int f(y) L_{y^{-1}} g dy$$

which can be interpreted as a vector valued integral, since for $\phi \in L^\infty(\mu)$,

$$\int (f * g)(x) \phi(x) dx = \int f(y) g(y^{-1}x) \phi(x) dx dy$$

so we can see convolution as a generalized ‘averaging’ of translate of g with respect to the values of f . If G is commutative, this is the same as the averaging of translates of f , but not in the noncommutative case. It then easily follows from operator computations $L_z(f * g) = (L_z f) * g$, and $R_z(f * g) = f * (R_z g)$, or from the fact that

$$(f * g)(zx) = \int f(y) g(y^{-1}zx) dy = \int f(zy) g(y^{-1}x) dy = [(L_z f) * g](x)$$

$$(f * g)(xz) = \int f(y) g(y^{-1}xz) dy = [f * (R_z g)](x)$$

Convolution can also be applied to the other L^p spaces, but we have to be a bit more careful with our integration.

Theorem 41.8. *If $f \in L^1(G)$ and $g \in L^p(G)$, then $f * g$ is defined for almost all x , $f * g \in L^p(G)$, and $\|f * g\|_p \leq \|f\|_1 \|g\|_p$. If G is unimodular, then the same results hold for $g * f$, or if G is not unimodular and f has compact support.*

Proof. We use Minkowski’s inequality to find

$$\begin{aligned} \|f * g\|_p &= \left(\int \left| \int f(y) |g(y^{-1}x)| dy \right|^p dx \right)^{1/p} \\ &\leq \int |f(y)| \left(\int |g(y^{-1}x)|^p dx \right)^{1/p} dy \\ &= \|f\|_1 \|g\|_p \end{aligned}$$

If G is unimodular, then

$$\|g * f\|_p = \left(\int \left| \int g(xy^{-1}) f(y) dy \right|^p dx \right)^{1/p}$$

and we may apply the same trick as used before.

If f has compact support K , then $1/\Delta$ is bounded above by $M > 0$ on K and

$$\begin{aligned}\|g * f\|_p &= \left(\int \left| \int \frac{g(xy^{-1})f(y)}{\Delta(y)} dy \right|^p dx \right)^{1/p} \\ &\leq \int \left(\int \left| \frac{g(xy^{-1})f(y)}{\Delta(y)} \right|^p dx \right)^{1/p} dy \\ &= \|g\|_p \int_K \frac{|f(y)|}{\Delta(y)} d\mu(y) \\ &\leq M \|g\|_p \|f\|_1\end{aligned}$$

which shows that $g * f$ is defined almost everywhere. \square

Theorem 41.9. *If G is unimodular, $f \in L^p(G)$, $g \in L^q(G)$, and $p = q^*$, then $f * g \in C_0(G)$ and $\|f * g\|_\infty \leq \|f\|_p \|g\|_q$.*

Proof. First, note that

$$\begin{aligned}|(f * g)(x)| &\leq \int |f(y)| |g(y^{-1}x)| dy \\ &\leq \|f\|_p \left(\int |g(y^{-1}x)|^q dy \right)^{1/q} \\ &= \|f\|_p \|g\|_q\end{aligned}$$

For each x and y , applying Hölder's inequality, we find

$$\begin{aligned}|(f * g)(x) - (f * g)(y)| &\leq \int |f(z)| |g(z^{-1}x) - g(z^{-1}y)| dz \\ &\leq \|f\|_p \left(\int |g(z^{-1}x) - g(z^{-1}y)|^q dz \right)^{1/q} \\ &= \|f\|_p \left(\int |g(z) - g(zx^{-1}y)|^q dz \right)^{1/q} \\ &= \|f\|_p \|g - R_{x^{-1}y}g\|_q\end{aligned}$$

Thus to prove continuity (and in fact uniform continuity), we need only prove that $\|g - R_x g\|_q \rightarrow 0$ for $q \neq \infty$ as $x \rightarrow \infty$ or $x \rightarrow 0$. This is the content of the next lemma. \square

We now show that the map $x \mapsto L_x$ is a continuous operation from G to the weak $*$ topology on the L_p spaces, for $p \neq \infty$. It is easily verified that translation is not continuous on L_∞ , by taking a suitable bumpy function.

Theorem 41.10. *If $p \neq \infty$, then $\|g - R_x g\|_p \rightarrow 0$ and $\|g - L_x g\|_p \rightarrow 0$ as $x \rightarrow 0$.*

Proof. If $g \in C_c(G)$, then one verifies the theorem by using left and right uniform continuity. In general, we let $g_i \in C_c(G)$ be a sequence of functions converging to g in the L_p norm, and we then find

$$\|g - L_x g\|_p \leq \|g - g_i\|_p + \|g_i - L_x g_i\|_p + \|L_x(g_i - g)\|_p = 2\|g - g_i\|_p + \|g_i - L_x g_i\|_p$$

Taking i large enough, x small enough, we find $\|g - L_x g\|_p \rightarrow 0$. The only problem for right translation is the appearance of the modular function

$$\|R_x(g - g_i)\|_p = \frac{\|g - g_i\|_p}{\Delta(x)^{1/p}}$$

If we assume our x values range only over a compact neighbourhood K of the origin, we find that $\Delta(x)$ is bounded below, and hence $\|R_x(g - g_i)\|_p \rightarrow 0$, which effectively removes the problems in the proof. \square

Since the map is linear, we have verified that the map $x \mapsto L_x f$ is uniformly continuous in L^p for each $f \in L^p$. In the case where $p = \infty$, the same theorem cannot hold, but we have even better conditions that do not even require unimodularity.

Theorem 41.11. *If $f \in L^1(G)$ and $g \in L^\infty(G)$, then $f * g$ is left uniformly continuous, and $g * f$ is right uniformly continuous.*

Proof. We have

$$\|L_z(f * g) - (f * g)\|_\infty = \|(L_z f - f) * g\|_\infty \leq \|L_z f - f\|_1 \|g\|_\infty$$

$$\|R_z(g * f) - (g * f)\|_\infty = \|g * (R_z f - f)\|_\infty \leq \|g\|_\infty \|R_z f - f\|_1$$

and both integrals converge to zero as $z \rightarrow 1$. \square

The passage from $M(G)$ to $L^1(G)$ removes an identity from the Banach algebra in question (except if G is discrete), but there is always a way to approximate an identity.

Theorem 41.12. For each neighbourhood U of the origin, pick a function $f_U \in (L^1)^+(G)$, with $\int \phi_U = 1$, $\text{supp}(f_U) \subset U$. Then if g is any function in $L^p(G)$,

$$\|f_U * g - g\|_p \rightarrow 0$$

where we assume g is left uniformly continuous if $p = \infty$, and if f_U is viewed as a net with neighbourhoods ordered by inclusion. If in addition $f_U(x) = f_U(x^{-1})$, then $\|g * f_U - g\|_p \rightarrow 0$, where g is right uniformly continuous for $p = \infty$.

Proof. Let us first prove the theorem for $p \neq \infty$. If $g \in C_c(G)$ is supported on a compact K , and if U is small enough that $|g(y^{-1}x) - g(x)| < \varepsilon$ for $y \in U$, then because $\int_U f_U(y) dy = 1$, and by applying Minkowski's inequality, we find

$$\begin{aligned} \|f_U * g - g\|_p &= \left(\int \left| \int f_U(y) [g(y^{-1}x) - g(x)] dy \right|^p dx \right)^{1/p} \\ &\leq \int f_U(y) \left(\int |g(y^{-1}x) - g(x)|^p dx \right)^{1/p} dy \\ &\leq 2\mu(K)\varepsilon \int f_U(y) dy \leq 2\mu(K)\varepsilon \end{aligned}$$

Results are then found for all of L^p by taking limits. If g is left uniformly continuous, then we may find U such that $|g(y^{-1}x) - g(x)| < \varepsilon$ for $y \in U$ then

$$|(f_U * g - g)(x)| = \left| \int f_U(y) [g(y^{-1}x) - g(x)] dy \right| \leq \varepsilon$$

For right convolution, we find that for $g \in C_c(G)$, where $|g(xy) - g(x)| < \varepsilon$

for $y \in U$, then

$$\begin{aligned}
\|g * f_U - g\|_p &= \left(\int \left| \int g(y) f_U(y^{-1}x) - g(x) dy \right|^p dx \right)^{1/p} \\
&= \left(\int \left| \int [g(xy) - g(x)] f_U(y) dy \right|^p dx \right)^{1/p} \\
&\leq \int \left(\int |g(xy) - g(x)|^p dx \right)^{1/p} f_U(y) dy \\
&\leq \mu(K) \varepsilon \int f_U(y) (1 + \Delta(y)) dy \\
&= \mu(K) \varepsilon + \mu(K) \varepsilon \int f_U(y) \Delta(y) dy
\end{aligned}$$

We may always choose U small enough that $\Delta(y) < \varepsilon$ for $y \in U$, so we obtain a complete estimate $\mu(K)(\varepsilon + \varepsilon^2)$. If g is right uniformly continuous, then choosing U for which $|g(xy) - g(x)| < \varepsilon$, then

$$|(g * f_U - g)(x)| = \left| \int [g(xy) - g(x)] f_U(y) dy \right| \leq \varepsilon$$

We will always assume from hereon out that the approximate identities in $L^1(G)$ are of this form. \square

We have already obtained enough information to characterize the closed ideals of $L^1(G)$.

Theorem 41.13. *If V is a closed subspace of $L^1(G)$, then V is a left ideal if and only if it is closed under left translations, and a right ideal if and only if it is closed under right translations.*

Proof. If V is a closed left ideal, and f_U is an approximate identity at the origin, then for any g ,

$$\|(L_z f_U) * g - L_z g\|_1 = \|L_z(f_U * g - g)\|_1 = \|f_U * g - g\| \rightarrow 0$$

so $L_z g \in V$. Conversely, if V is closed under left translations, $g \in L^1(G)$, and $f \in V$, then

$$g * f = \int g(y) L_{y^{-1}} f dy$$

which is in the closed linear space of the translates of f . Right translation is verified very similarly. \square

41.4 The Riesz Thorin Theorem

We finalize our basic discussion by looking at convolutions of functions in $L^p * L^q$. Certainly $L^p * L^1 \subset L^p$, and $L^p * L^q \subset L^\infty$ for $q = p^*$. To prove general results, we require a foundational interpolation result.

Theorem 41.14. *For any $0 < \theta < 1$, and $0 < p, q \leq \infty$. If we define*

$$1/r_\theta = (1 - \theta)/p + \theta/q$$

to be the inverse interpolation of the two numbers. Then

$$\|f\|_{r_\theta} \leq \|f\|_p^{1-\theta} \|f\|_q^\theta$$

Proof. We apply Hölder's inequality to find

$$\|f\|_{r_\theta} \leq \|f\|_{p/(1-\theta)} \|f\|_{q/\theta} = \left(\int |f|^{p/(1-\theta)} \right)^{(1-\theta)/p} \left(\int |f|^{q/\theta} \right)^{\theta/q}$$

so it suffices to prove $\|f\|_{p/(1-\theta)} \leq \|f\|_p^{1-\theta}$, $\|f\|_{q/\theta} \leq \|f\|_q^\theta$.

The map $x \mapsto x^p$ is concave for $0 < p < 1$, so we may apply Jensen's inequality in reverse to conclude

$$\left(\int |f|^{p/(1-\theta)} \right)^{(1-\theta)/p} \leq \left(\int |f|^p \right)^{1/p}$$

□

The Riesz Thorin interpolation theorem then implies $L^p * L^q \subset L^r$, for $p^{-1} + q^{-1} = 1 + r^{-1}$. However, these estimates only guarantee $L^1(G)$ is closed under convolution. If G is compact, then $L_p(G)$ is closed under convolution for all p (TODO). The L_p conjecture says that this is true if and only if G is compact. This was only resolved in 1990.

41.5 Homogenous Spaces and Haar Measures

The natural way for a locally compact topological group G to act on a locally compact Hausdorff space X is via a representation of G in the homeomorphisms of X . We assume the action is transitive on X . The standard

example are the action of G on G/H , where H is a closed subspace. These are effectively all examples, because if we fix $x \in X$, then the map $y \mapsto yx$ induces a continuous bijection from G/H to X , where H is the set of all y for which $yx = x$. If G is a σ compact space, then this map is a homeomorphism.

Theorem 41.15. *If a σ compact topological group G has a transitive topological action on X , and $x \in X$, then the continuous bijection from G/G_x to X is a homeomorphism.*

Proof. It suffices to show that the map $\phi : G \rightarrow X$ is open, and we need only verify this for the neighbourhood basis of compact neighbourhoods V of the origin by properties of the action. G is covered by countably many translates y_1V, y_2V, \dots , and since each $\phi(y_kV) = y_k\phi(V)$ is closed (compactness), we conclude that $y_k\phi(V)$ has non-empty interior for some y_k , and hence $\phi(V)$ has a non-empty interior point $\phi(y_0)$. But then for any $y \in V$, y is in the interior of $\phi(yVy_0^{-1}) \subset \phi(VVy_0^{-1})$, so if we fix a compact U , and find V with $V^3 \subset U$, we have shown $\phi(U)$ is open in X . \square

We shall say a space X is homogenous if it is homeomorphic to G/H for some group action of G over X . The H depends on our choice of basepoint x , but only up to conjugation, for if we switch to a new basepoint y , and c maps x to y , then $ax = x$ holds if and only if $cac^{-1}y = y$. The question here is to determine whether we have a G -invariant measure on X . This is certainly not always possible. If we had a measure on \mathbf{R} invariant under the affine maps $ax + b$, then it would be equal to the Haar measure by uniqueness, but the Haar measure is not invariant under dilation $x \mapsto ax$.

Let G and H have left Haar measures μ and ν respectively, denote the projection of G onto G/H as $\pi : G \rightarrow G/H$, and let Δ_G and Δ_H be the respective modular functions. Define a map $P : C_c(G) \rightarrow C_c(G/H)$ by

$$(Pf)(Hx) = \int_H f(xy) d\nu(y) = \int_H$$

this is well defined by the invariance properties of ν . Pf is obviously continuous, and $\text{supp}(Pf) \subset \pi(\text{supp}(f))$. Moreover, if $\phi \in C(G/H)$ we have

$$P((\phi \circ \pi) \cdot f)(Hx) = \phi(xH) \int_H f(xy) d\nu(y)$$

so $P((\phi \circ \pi) \cdot f) = \phi P(f)$.

Lemma 41.16. *If E is a compact subset of G/H , there is a compact $K \subset G$ with $\pi(K) = E$.*

Proof. Let V be a compact neighbourhood of the origin, and cover E by finitely many translates of $\pi(V)$. We conclude that $\pi^{-1}(E)$ is covered by finitely many of the translates, and taking the intersections of these translates with $\pi^{-1}(E)$ gives us the desired K . \square

Lemma 41.17. *A compact $F \subset G/H$ gives rise to a function $f \geq 0$ in $C_c(G)$ such that $Pf = 1$ on E .*

Proof. Let E be a compact neighbourhood containing F , and if $\pi(K) = E$, there is a function $g \in C_c(G)$ with $g > 0$ on K , and $\phi \in C_c(G/H)$ is supported on E and $\phi(x) = 1$ for $x \in F$, let

$$f = \frac{\phi \circ \pi}{Pg \circ \pi} g$$

Hence

$$Pf = \frac{\phi}{Pg} Pg = \phi$$

\square

Lemma 41.18. *If $\phi \in C_c(G/H)$, there is $f \in C_c(G)$ with $Pf = \phi$, and $\pi(\text{supp } f) = \text{supp }(\phi)$, and also $f \geq 0$ if $\phi \geq 0$.*

Proof. There exists $g \geq 0$ in $C_c(G/H)$ with $Pg = 1$ on $\text{supp }(\phi)$, and then $f = (\phi \circ \pi)g$ satisfies the properties of the theorem. \square

We can now provide conditions on the existence of a measure on G/H .

Theorem 41.19. *There is a G invariant measure ψ on G/H if and only if $\Delta_G = \Delta_H$ when restricted to H . In this case, the measure is unique up to a common factor, and if the factor is chosen, we have*

$$\int_G f d\mu = \int_{G/H} Pf d\psi = \int_{G/H} \int_H f(xy) d\nu(y) d\psi(xH)$$

Proof. Suppose ψ existed. Then $f \mapsto \int Pf d\psi$ is a non-zero left invariant positive linear functional on G/H , so $\int Pf d\psi = c \int f d\mu$ for some $c > 0$.

Since $P(C_c(G)) = C_c(G/H)$, we find that ψ is determined up to a constant factor. We then compute, for $y \in H$,

$$\begin{aligned}\Delta_G(y) \int f(x) d\mu(x) &= \int f(xy^{-1}) d\mu(x) \\ &= \int_{G/H} \int_H f(xzy^{-1}) d\nu(z) d\psi(xH) \\ &= \Delta_H(y) \int_{G/H} \int_H f(xz) d\nu(z) d\psi(xH) \\ &= \Delta_H(y) \int f(x) d\mu(x)\end{aligned}$$

Hence $\Delta_G = \Delta_H$. Conversely, suppose $\Delta_G = \Delta_H$. First, we claim if $f \in C_c(G)$ and $Pf = 0$, then $\int f d\mu = 0$. Indeed if $P\phi = 1$ on $\pi(\text{supp} f)$ then

$$0 = Pf(xH) = \int_H f(xy) d\nu(y) = \Delta_G(y^{-1}) \int_H f(xy^{-1}) d\nu(y)$$

so

$$\begin{aligned}0 &= \int_G \int_H \Delta_G(y^{-1}) \phi(x) f(xy^{-1}) d\nu(y) d\mu(x) \\ &= \int_H \int_G \phi(xy) f(x) d\mu(x) d\nu(y) \\ &= \int_G P\phi(xH) f(x) d\mu(x) \\ &= \int_G f(x) d\mu(x)\end{aligned}$$

This implies that if $Pf = Pg$, then $\int_G f = \int_G g$. Thus the map $Pf \mapsto \int_G f$ is a well defined G invariant positive linear functional on $C_c(G/H)$, and we obtain a Radon measure from the Riesz representation theorem. \square

If H is compact, then Δ_G and Δ_H are both continuous homomorphisms from H to \mathbf{R}^+ , so Δ_G and Δ_H are both trivial, and we conclude a G invariant measure exists on G/H .

Chapter 42

The Character Space

Let G be a locally compact group. A character on G is a *continuous* homomorphism from G to \mathbf{T} . The space of all characters of a group will be denoted $\Gamma(G)$.

Example. *Determining the characters of \mathbf{T} involves much of classical Fourier analysis. Let $f : \mathbf{T} \rightarrow \mathbf{T}$ be an arbitrary continuous character. For each $w \in \mathbf{T}$, consider the function $g(z) = f(zw) = f(z)f(w)$. We know the Fourier series acts nicely under translation, telling us that*

$$\hat{g}(n) = w^n \hat{f}(n)$$

Conversely, since $g(z) = f(z)f(w)$,

$$\hat{g}(n) = f(w) \hat{f}(n)$$

Thus $(w^n - f(w))\hat{f}(n) = 0$ for all $w \in \mathbf{T}$, $n \in \mathbf{Z}$. Fixing n , we either have $f(w) = w^n$ for all w , or $\hat{f}(n) = 0$. This implies that if $f \neq 0$, then f is just a power map for some $n \in \mathbf{Z}$.

Example. *The characters of \mathbf{R} are of the form $t \mapsto e(t\xi)$, for $\xi \in \mathbf{R}$. To see this, let $e : \mathbf{R} \rightarrow \mathbf{T}$ be an arbitrary character. Define*

$$F(x) = \int_0^x e(t) dt$$

Then $F'(x) = e(x)$. Since $e(0) = 1$, for suitably small δ we have

$$F(\delta) = \int_0^\delta e(t) dt = c > 0$$

and then it follows that

$$F(x + \delta) - F(x) = \int_x^{x+\delta} e(t)dt = \int_0^\delta e(x+t)dt = ce(x)$$

As a function of x , F is differentiable, and by the fundamental theorem of calculus,

$$\frac{dF(x + \delta) - F(x)}{dt} = F'(x + \delta) - F'(x) = e(x + \delta) - e(x)$$

This implies the right side of the above equation is differentiable, and so

$$ce'(x) = e(x + \delta) - e(x) = e(x)[e(\delta) - 1]$$

Implying $e'(x) = Ae(x)$ for some $A \in \mathbf{C}$, so $e(x) = e^{Ax}$. We require that $e(x) \in \mathbf{T}$ for all x , so $A = \xi i$ for some $\xi \in \mathbf{R}$.

Example. Consider the group \mathbf{R}^+ of positive real numbers under multiplication. The map $x \mapsto \log x$ is an isomorphism from \mathbf{R}^+ and \mathbf{R} , so that every character on \mathbf{R}^+ is of the form $e(s \log x) = x^{is}$, for some $s \in \mathbf{R}$. The character group is then \mathbf{R} , since $x^{is}x^{is'} = x^{i(s+s')}$.

There is a connection between characters on G and characters on $L^1(G)$ that is invaluable to the generalization of Fourier analysis to arbitrary groups.

Theorem 42.1. For any character $\phi : G \rightarrow \mathbf{C}$, the map

$$\varphi(f) = \int \frac{f(x)}{\phi(x)} dx$$

is a non-zero character on the convolution algebra $L^1(G)$, and all characters arise this way.

Proof. The induced map is certainly linear, and

$$\begin{aligned} \varphi(f * g) &= \int \int \frac{f(y)g(y^{-1}x)}{\phi(x)} dy dx \\ &= \int \int \frac{f(y)g(x)}{\phi(y)\phi(x)} dy dx \\ &= \int \frac{f(y)}{\phi(y)} dy \int \frac{g(x)}{\phi(x)} dx \end{aligned}$$

Since ϕ is continuous, there is a compact subset K of G where $\phi > \varepsilon$ for some $\varepsilon > 0$, and we may then choose a positive f supported on K in such a way that $\varphi(f)$ is non-zero.

The converse results from applying the duality theory of the L^p spaces. Any character on $L^1(G)$ is a linear functional, hence is of the form

$$f \mapsto \int f(x)\phi(x)dx$$

for some $\phi \in L^\infty(G)$. Now

$$\begin{aligned} \int \int f(y)g(x)\phi(yx)dydx &= \int \int f(y)g(y^{-1}x)\phi(x)dydx \\ &= \int f(x)\phi(x)dx \int g(y)\phi(y)dy \\ &= \int f(x)g(y)\phi(x)\phi(y)dx dy \end{aligned}$$

Since this holds for all functions f and g in $L^1(G)$, we must have $\phi(yx) = \phi(x)\phi(y)$ almost everywhere. Also

$$\begin{aligned} \int \varphi(f)g(y)\phi(y)dy &= \varphi(f * g) \\ &= \int \int g(y)f(y^{-1}x)\phi(x)dydx \\ &= \int \int (L_{y^{-1}}f)(x)g(y)\phi(x)dydx \\ &= \int \varphi(L_{y^{-1}}f)g(y)dy \end{aligned}$$

which implies $\varphi(f)\phi(y) = \varphi(L_{y^{-1}}f)$ almost everywhere. Since the map $\varphi(L_{y^{-1}}f)/\varphi(f)$ is a uniformly continuous function of y , ϕ is continuous almost everywhere, and we might as well assume ϕ is continuous. We then conclude $\phi(xy) = \phi(x)\phi(y)$. Since $\|\phi\|_\infty = 1$ (this is the norm of any character operator on $L^1(G)$), we find ϕ maps into \mathbb{T} , for if $\|\phi(x)\| < 1$ for any particular x , $\|\phi(x^{-1})\| > 1$. \square

Thus there is a one-to-one correspondence with $\Gamma(G)$ and $\Gamma(L^1(G))$, which implies a connection with the Gelfand theory and the character

theory of locally compact groups. This also gives us a locally compact topological structure on $\Gamma(G)$, induced by the Gelfand representation on $\Gamma(L^1(G))$. A sequence $\phi_i \rightarrow \phi$ if and only if

$$\int \frac{f(x)}{\phi_i(x)} dx \rightarrow \int \frac{f(x)}{\phi(x)} dx$$

for all functions $f \in L^1(G)$. This actually makes the map

$$(f, \phi) \mapsto \int \frac{f(x)}{\phi(x)} dx$$

a jointly continuous map, because as we verified in the proof above,

$$\widehat{f}(\phi)\phi(y) = \widehat{L_y f}(\phi)$$

And the map $y \mapsto L_y f$ is a continuous map into $L^1(G)$. If $K \subset G$ and $C \subset \Gamma(G)$ are compact, this allows us to find open sets in G and $\Gamma(G)$ of the form

$$\{\gamma : \|1 - \gamma(x)\| < \varepsilon \text{ for all } x \in K\} \quad \{x : \|1 - \gamma(x)\| < \varepsilon \text{ for all } \gamma \in C\}$$

And these sets actually form a base for the topology on $\Gamma(G)$.

Theorem 42.2. *If G is discrete, $\Gamma(G)$ is compact, and if G is compact, $\Gamma(G)$ is discrete.*

Proof. If G is discrete, then $L^1(G)$ contains an identity, so $\Gamma(G) = \Gamma(L^1(G))$ is compact. Conversely, if G is compact, then it contains the constant 1 function, and

$$\widehat{1}(\phi) = \int \frac{dx}{\phi(x)}$$

And

$$\frac{1}{\phi(y)} \widehat{1}(\phi) = \int \frac{dx}{\phi(yx)} = \int \frac{dx}{\phi(x)} = \widehat{1}(\phi)$$

So either $\phi(y) = 1$ for all y , and it is then verified by calculation that $\widehat{1}(\phi) = 1$, or $\widehat{1}(\phi) = 0$. Since $\widehat{1}$ is continuous, the trivial character must be an open set by itself, and hence $\Gamma(G)$ is discrete. \square

Given a function $f \in L^1(G)$, we may take the Gelfand transform, obtaining a function on $C_0(\Gamma(L^1(G)))$. The identification then gives us a function on $C_0(\Gamma(G))$, if we give $\Gamma(G)$ the topology induced by the correspondence (which also makes $\Gamma(G)$ into a topological group). The formula is

$$\widehat{f}(\phi) = \phi(f) = \int \frac{f(x)}{\phi(x)}$$

This gives us the classical correspondence between $L^1(\mathbf{T})$ and $C_0(\mathbf{Z})$, and $L^1(\mathbf{R})$ and $C_0(\mathbf{R})$, which is just the Fourier transform. Thus we see the Gelfand representation as a natural generalization of the Fourier transform. We shall also denote the Fourier transform by \mathcal{F} , especially when we try and understand its properties as an operator. Gelfand's theory (and some basic computation) tells us instantly that

- $\widehat{f * g} = \widehat{f} \widehat{g}$ (The transform is a homomorphism).
- \mathcal{F} is norm decreasing and therefore continuous: $\|\widehat{f}\|_\infty \leq \|f\|_1$.
- If G is unimodular, and $\gamma \in \Gamma(G)$, then $(f * \gamma)(x) = \gamma(x) \widehat{f}(\gamma)$.

Whenever we integrate a function with respect to the Haar measure, there is a natural generalization of the concept to the space of all measures on G . Thus, for $\mu \in M(G)$, we define

$$\widehat{\mu}(\phi) = \int \frac{d\mu}{\phi(x)}$$

which we call the **Fourier-Stieltjes transform** on G . It is essentially an extension of the Gelfand representation on $L^1(G)$ to $M(G)$. Each $\widehat{\mu}$ is a bounded, uniformly continuous function on $\Gamma(G)$, because the transform is still contracting, i.e.

$$\left| \int \frac{d\mu(x)}{\phi(x)} \right| \leq \|\mu\|$$

It is uniformly continuous, because

$$(L_\nu \widehat{\mu} - \widehat{\mu})(\phi) = \int \frac{1 - \nu(x)}{\nu(x)\phi(x)} d\mu(x)$$

The regularity of μ implies that there is a compact set K such that $|\mu|(K^c) < \varepsilon$. If $v_i \rightarrow 0$, then eventually we must have $|v_i(x) - 1| < \varepsilon$ for all $x \in K$, and then

$$|(L_\nu \hat{\mu} - \hat{\mu})(\phi)| \leq 2|\mu|(K^c) + \varepsilon\|\mu\| \leq \varepsilon(2 + \|\mu\|)$$

Which implies uniform continuity.

Let us consider why it is natural to generalize operators on $L^1(G)$ to $M(G)$. The first reason is due to the intuition of physicists; most of classical Fourier analysis emerged from physical considerations, and it is in this field that $L^1(G)$ is often confused with $M(G)$. Take, for instance, the determination of the electric charge at a point in space. To determine this experimentally, we take the ratio of the charge over some region in space to the volume of the region, and then we limit the size of the region to zero. This is the historical way to obtain the density of a measure with respect to the Lebesgue measure, so that the function we obtain can be integrated to find the charge over a region. However, it is more natural to avoid taking limits, and to just think of charge as an element of $M(\mathbf{R}^3)$. If we consider a finite number of discrete charges, then we obtain a discrete measure, whose density with respect to the Lebesgue measure does not exist. This doesn't prevent physicists from trying, so they think of the density obtained as shooting off to infinity at points. Essentially, we obtain the Dirac Delta function as a 'generalized function'. This is fine for intuition, but things seem to get less intuitive when we consider the charge on a subsurface of \mathbf{R}^3 , where the 'density' is 'dirac'-esque near the function, where as measure theoretically we just obtain a density with respect to the two-dimensional Hausdorff measure on the surface. Thus, when physicists discuss quantities as functions, they are really thinking of measures, and trying to take densities, where really they may not exist.

There is a more austere explanation, which results from the fact that, with respect to integration, $L^1(G)$ is essentially equivalent to $M(G)$. Notice that if $\mu_i \rightarrow \mu$ in the weak-* topology, then $\hat{\mu}_i \rightarrow \hat{\mu}$ pointwise, because

$$\int \frac{d\mu_i(x)}{\phi(x)} \rightarrow \int \frac{d\mu(x)}{\phi(x)}$$

(This makes sense, because weak-* convergence is essentially pointwise convergence in $M(G)$). Thus the Fourier-Stieltjes transform is continuous with respect to these topologies. It is the unique continuous extension of the Fourier transform, because

Theorem 42.3. $L^1(G)$ is weak-* dense in $M(G)$.

Proof. First, note that the Dirac delta function can be weak-* approximated by elements of $L^1(G)$, since we have an approximate identity in the space.

First, note that if $\mu_i \rightarrow \mu$, then $\mu_i * \nu \rightarrow \mu * \nu$, because

$$\int f d(\mu_i * \nu) = \int \int f(xy) d\mu_i(x) d\nu(y)$$

The functions $y \mapsto \int f(xy) d\mu_i(x)$ converge pointwise to $\int f(xy) d\mu(y)$. Since

$$\left| \int f(xy) d\mu_i(x) \right| \leq \|f\|_1 \|\mu_i\|$$

If i is taken large enough that \square

If $\phi_\alpha \rightarrow \phi$, in the sense that $\phi_\alpha(x) \rightarrow \phi(x)$ for all $x \in G$, then, because $\|\phi_\alpha(x)\| = 1$ for all x , we can apply the dominated convergence theorem on any compact subset K of G to conclude

$$\int_K \frac{d\mu(x)}{\phi_\alpha(x)} \rightarrow \int_K \frac{d\mu(x)}{\phi(x)}$$

It is immediately verified to be a map into $L^1(\Gamma(G))$, because

$$\int \left| \int \frac{d\mu(x)}{\phi(x)} \right| d\phi \leq \int \|\mu\|$$

The formula above immediately suggests a generalization to a transform on $M(G)$. For $\nu \in M(G)$, we define

$$\mathcal{F}(\nu)(\phi) = \int \frac{d\nu}{\phi}$$

If $\mathcal{G} : L^1(G) \rightarrow C_0(\Gamma(G))$ is the Gelfand transform, then the transform induces a map $\mathcal{G}^* : M(\Gamma(G)) \rightarrow L^\infty(G)$.

The duality in classical Fourier analysis is shown through the inversion formulas. That is, we have inversion functions

$$\mathcal{F}^{-1}(\{a_k\}) = \sum a_k e_k(t) \quad \mathcal{F}^{-1}(f)(x) = \int f(t) e(xt)$$

which reverses the fourier transform on \mathbf{T} and \mathbf{R} respectively, on a certain subclass of L^1 . One of the challenges of Harmonic analysis is trying to find where this holds for the general class of measurable functions.

The first problem is to determine surjectivity. We denote by $A(G)$ the space of all continuous functions which can be represented as the fourier transform of some function in $L^1(G)$. It is to even determine $A(\mathbf{T})$, the most basic example. $A(G)$ always separates the points of $\Gamma(G)$, by Gelfand theory, and if G is unimodular, then it is closed under conjugation. If we let $g(x) = \overline{f(x^{-1})}$, we find

$$\mathcal{F}(g)(\phi) = \int \frac{g(x)}{\phi(x)} dx = \overline{\int \frac{f(x^{-1})}{\phi(x^{-1})} dx} = \int \frac{f(x)}{\phi(x)} dx = \overline{\mathcal{F}(f)(\phi)}$$

so that by the Stone Weirstrass theorem $A(G)$ is dense in $C_0(\Gamma(L^1(G)))$.

Chapter 43

Banach Algebra Techniques

In the mid 20th century, it was realized that much of the analytic information about a topological group can be captured in various C^* algebras related to the group. For instance, consider the Gelfand space of $L^1(\mathbf{Z})$ is \mathbf{T} , which represents the fact that one can represent functions over \mathbf{T} as sequences of numbers. Similarly, we find the characters of $L^1(\mathbf{R})$ are the maps $f \mapsto \hat{f}(x)$, so that the Gelfand space of \mathbf{R} is \mathbf{R} , and the Gelfand transform is the Fourier transform on this space. For a general G , we may hope to find a generalized Fourier transform by understanding the Gelfand transform on $L^1(G)$. We can also generalize results by extending our understanding to the class $M(G)$ of regular, Borel measures on G .

Chapter 44

Vector Spaces

If \mathbf{K} is a closed, multiplicative subgroup of the complex numbers, then \mathbf{K} is also a locally compact abelian group, and we can therefore understand \mathbf{K} by looking at its dual group \mathbf{K}^* . The map $\langle x, y \rangle = xy$ is bilinear, in the set that it is a homomorphism in the variable y for each fixed x , and a homomorphism in the variable x for each y .

If \mathbf{K} is a subfield of the complex numbers, then \mathbf{K} is also an abelian group under addition, and we can consider the dual group \mathbf{K}^* . The inner product $\langle x, y \rangle = xy$ gives a continuous bilinear map $\mathbf{K} \times \mathbf{K} \rightarrow \mathbf{C}$, and therefore we can define $x^* \in \mathbf{K}^*$ by $x^*(y) = \langle x, y \rangle$. If $x^*(y) = xy = 0$ for all y , then in particular $x^*(1) = x$, so $x = 0$. This means that the homomorphism $\mathbf{K} \rightarrow \mathbf{K}^*$ is injective.

Chapter 45

Interpolation of Besov and Sobolev spaces

An important class of operators arise as singular integrals, that is, they arise as convolution operators T given by $T(f) = f * K$, where K is an appropriate distribution. Taking Fourier transforms, these operators can also be defined by $\widehat{T(f)} = \widehat{f}\widehat{K}$. The function \widehat{K} is known as a **Fourier multiplier**, because it operates by multiplication on the frequencies of the function f . We say \widehat{K} is a **Fourier multiplier on $L^p(\mathbf{R}^n)$** if T is a bounded map from $S(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$, under the L^p norms. Such maps clearly extend uniquely to maps from $L^p(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$, and so we can think of T as operating by convolution on the space of L^p functions. We will denote the space of all Fourier multipliers on L^p by M_p . We define the L^p norm on these distributions K , denoted $\|K\|_p$, to be the operator norm of the associated operator T .

Example. Consider the space M_∞ . If K is a distribution in M_∞ , then $\|K\|_\infty < \infty$, and since convolution commutes with translations, in the sense that $f_h * K = (f * K)_h$, then

$$\|K\|_\infty = \sup_{f \in L^\infty(\mathbf{R}^n)} \frac{|(f * K)(0)|}{\|f\|_\infty}$$

But then the map $f \mapsto (f * K)(0)$ is a bounded operator on the space of bounded continuous functions, and so the Riesz representation says there is a bounded Radon measure μ such that

$$(f * K)(0) = \int f(-y) d\mu(y)$$

But now we know

$$(f * K)(x) = (f_{-x} * K)(0) = \int f(x - y) d\mu(y) = (f * \mu)(x)$$

Thus M_∞ is really just the space of all bounded Radon measures, and

$$\|K\|_\infty = \sup_{f \in L^\infty(\mathbf{R}^n)} \frac{|\int f(y) d\mu(y)|}{\|f\|_\infty} = \|\mu\|_1$$

so M_∞ even has the same norm as the space of all bounded Radon measures. Note that it becomes a Banach algebra under convolution of distributions, since the convolution of two bounded Radon measures is a bounded Radon measure.

Theorem 45.1. For any $1 \leq p \leq \infty$, and $q = p^*$, then $M_p = M_q$.

Proof. Let $f \in L^p$, and $g \in L^q$, then Hölder's inequality gives

$$|(K * f * g)(0)| \leq \|K * f\|_p \|g\|_q \leq \|K\|_p \|f\|_p \|g\|_p$$

Thus $K * g \in L_q$, and that $K \in M_q$ with $\|K\|_q \leq \|K\|_p$. By symmetry, we find $\|K\|_p = \|K\|_q$. \square

Example. Consider M_2 . If K is a distribution with $\|f * K\|_2 \leq A\|f\|_2$, then Parseval's inequality implies that

$$\|\hat{f}\hat{K}\|_2 = \|f * K\|_2 \leq A\|f\|_2 = A\|\hat{f}\|_2$$

so for each \hat{f} , TODO: PROVE THAT THIS IS REALLY JUST THE SPACE $L^\infty(\mathbf{R}^n)$, with the supremum norm. Note that this is also a Banach algebra under pointwise multiplication.

Using the Riesz-Thorin interpolation theorem, we find that if $1/p = (1-\theta)/p_0 + \theta/p_1$, then $\|K\|_p \leq \|K\|_{p_0}^{1-\theta} \|K\|_{p_1}^\theta$, when K lies in the three spaces. In particular, $\|K\|_p$ is a decreasing function of p for $1 \leq p \leq 2$, so we find $M_1 \subset M_p \subset M_q \subset M_2$ for $1 \leq p < q \leq 2$. In particular, all Fourier multipliers can be viewed as Fourier multipliers with respect to bounded, measurable functions on L^∞ . Riesz interpolation shows that each M_p is a Banach algebra under multiplication in the frequency domain, or convolution in the spatial domain.

Theorem 45.2. *Let $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a surjective affine transformation. Then the endomorphism T^* on $M_p(\mathbf{R}^n)$ defined by $(T^*f)(\xi) = f(T(\xi))$ is an isometry, and if T is a bijection, so too is T^* .*

Proof. TODO □

The next theorem is the main tool to prove results about Sobolev and Besov space. Note that it assumes $1 < p < \infty$, and cannot be applied for $p = 1$ or $p = \infty$. The proof relies on two lemmas, the first of which is used frequently later, and the second is used universally in modern harmonic analysis.

Lemma 45.3. *There exists a Schwartz function φ on \mathbf{R}^n which is supported on the annulus*

$$\{\xi : 1/2 \leq |\xi| \leq 2\}$$

is positive for $1/2 < |\xi| < 2$, and satisfies

$$\sum_{k=-\infty}^{\infty} \varphi(2^{-k}\xi) = 1$$

for all $\xi \neq 0$.

Lemma 45.4 (Calderon-Zygmund Decomposition). *Let $f \in L^1(\mathbf{R}^n)$, and $\sigma > 0$. Then there are pairwise almost disjoint cubes I_1, I_2, \dots with edges parallel to the coordinate axis and*

$$\sigma < \frac{1}{|I_n|} \int_{I_n} |f(x)| dx \leq 2^n \sigma$$

and with $|f(x)| \leq \sigma$ for almost all x outside these cubes.

Theorem 45.5 (The Mihlin Multiplier Theorem). *Let m be a bounded function on \mathbf{R}^n which is smooth except possibly at the origin, such that*

$$\sup_{\substack{\xi \in \mathbf{R}^n \\ |\alpha| \leq L}} |\xi|^{|\alpha|} |(D^\alpha m)(x)| < \infty$$

Then m is an L^p Fourier multiplier for $1 < p < \infty$.

45.1 Besov Spaces

Recall the Schwarz function φ used to prove the Mihlin multiplier theorem. We now define functions φ_k such that

$$\widehat{\varphi_n}(\xi) = \varphi(2^{-n}\xi) \quad \widehat{\psi}(\xi) = 1 - \sum_{n=1}^{\infty} \varphi(2^{-n}\xi)$$

Thus φ_n essentially covers the annulus $2^{n-1} \leq |\xi| \leq 2^{n+1}$, and the function ψ covers the remaining low frequency parts covered in the frequency ball of radius 2. We have

$$\varphi_n(\xi) = \widetilde{\varphi_{2^{-n}}}(\xi) = 2^{dn} \check{\varphi}(2^n \xi)$$

Given $s \in \mathbf{R}$, and $1 \leq p, q \leq \infty$, we write

$$\|f\|_{pq}^s = \|\psi * f\|_p + \left(\sum_{n=1}^{\infty} (2^{sn} \|\varphi_n * f\|_p)^q \right)^{1/q}$$

The convolution $\varphi_n * f$ essentially captures the portion of f whose frequencies lie in the annulus $2^{n-1} \leq |\xi| \leq 2^{n+1}$

45.2 Proof of The Projection Result

As with Marstrand's projection theorem, we require an energy integral variant. Rather than considering the Riesz kernel on \mathbf{R}^n , we consider the kernel on balls

$$K_\alpha(x) = \frac{\chi_{B(0,R)}(x)}{|x|^\alpha}$$

where R is a fixed radius. If $\alpha < \beta$, and μ is measure supported on a β dimensional subset of \mathbf{R}^n , then $\mu * K_\alpha \in L^\infty(\mathbf{R}^d)$ because μ cancels out the singular part of K_α . Assuming $\beta < d$, we conclude $\mu * K_\alpha \in L^1(\mathbf{R}^d)$. Applying interpolation (TODO: Which interpolation), we conclude that $\nu * K_\rho$

Chapter 46

The Cap Set Problem

The cap set problem comes out of additive combinatorics, whose goal is to understand additive structure in some abelian group, typically the integers. For instance, we can think of a set A as being roughly closed under addition if $|A + A| = O(|A|)$. Over rings, we can study the interplay between additive and multiplicative structure. For instance, one conjecture of Erdős and Szemerédi says that if A is a finite subset of real numbers, then $\max(|A + A|, |A \cdot A|) \gtrsim |A|^{1+c}$ for some positive $c \in (0, 1)$. The best known c so far is $c \sim 1/3$, though it is conjectured that we can take c arbitrarily close to 1. This can be seen as a discrete version of the results of Bourgain and Edgar-Miller on the Hausdorff dimensions of Borel subrings.

Theorem 46.1 (Van Der Waerden - 1927). *For any positive integers r and k , there is N such that if the integers in $[1, N]$ are given an r coloring, then there is a monochromatic k term arithmetic progression.*

The coloring itself is not so important, more just the partitioning. We just pigeonhole, using the density of k term arithmetic progressions. This problem suggests the Ramsey type problem of determining the largest set A of the integers $[1, N]$ which does not contain k term arithmetic progressions. Behrend's theorem says we can choose A to be on the order of $N \exp(-c\sqrt{\log N})$.

Theorem 46.2 (Roth - 1956). *If A is a set of integers in $[1, N]$ which is free of three term arithmetic progressions, then $|A| = O(N/\log \log N)$.*

Szemerédi proved that if A is free of k term arithmetic progressions, $|A| = o(N)$. If Erdős Turan, if $\sum_{x \in X} 1/x$ diverges, then X contains arbitrarily long arithmetic progressions. For now, we'll restrict our attention to three term arithmetic progressions. Heath and Brown showed that three term arithmetic progressions are $O(N/(\log N)^c)$ for some constant c . In 2016, the best known bound was given by Bloom, given $O(N(\log \log N)^4/\log N)$.

One way we can simplify our problem is to note that avoiding three term arithmetic progressions is a local issue, so we can embed $[1, N]$ in $\mathbf{Z}/M\mathbf{Z}$ for suitably large M , and we lose none of the problems we had over the integers. A heuristic is that it is easier to solve these kind of problems in \mathbf{F}_p^n , where p is small and n is large, which should behave like $\{1, \dots, p^n\}$. This leads naturally to the cap set problem.

Theorem 46.3 (Cap Set Problem). *What is the largest subset of \mathbf{F}_3^n containing no three term arithmetic progressions?*

We look at \mathbf{F}_3 because it is the smallest case where three term arithmetic progressions become important.

Theorem 46.4 (Mészáros - 1995). *Let $A \subset \mathbf{F}_3^n$ be a cap set. Then $|A| = O(3^n/n)$. This is analogous to a $N/\log N$ case over the integers, giving evidence that the finite field case is easier.*

In 2012, Bateman and Katz showed $|A| = O(3^n/n^{1+\varepsilon})$ for some $c > 0$. This was a difficult proof. In 2016, there was a more significant breakthrough, which gave an easy proof using the polynomial method of an exponentially small bound of c^n , where $c < 4$, over $\mathbf{Z}/4\mathbf{Z}$, and a week later Ellenberg-Gijswijt used this argument in the \mathbf{F}_3 case to prove that if A is a capset in \mathbf{F}_3 , then $|A| = O(c^n)$, for $c = 2.7551\dots$.

The idea of the polynomial method is to take combinatorial information about some set, encode it as some algebraic structural information, and then apply the theory of polynomials to encode this algebraic information and use it to limit and enable certain properties to occur.

If V is the space of polynomials of degree d vanishing on a set A , then we know $\dim V \geq \dim \mathcal{P}_d - |A|$. This gives a lower bound on the size of A , whereas we want an upper bound. To get an upper bound, we take $|A|^c$ instead, which shows

$$\dim V \geq \dim \mathcal{P}_d - |A| - 3^n$$

whichs gives $|A| \leq 3^n + \dim V - \dim \mathcal{P}_d$. Now using linear algebra, we can find a polynomial P vanishing on A^c with support of cardinality greater than or equal to $\dim V$, hence

$$|A| \leq 3^n - \dim \mathcal{P}_d + \max |\text{supp}(P)|$$

It follows that A is a cap set if and only if $x + y = 2z$, or $x + y + z = 0$ holds if and only if $x = y = z$. This is an algebraic property which says directly that A has no nontrivial three term arithmetic progressions. Thus for any $a_1, \dots, a_m \in A$, $P(-a_i - a_j) = 0$ when $i \neq j$. Equivalently, this means $P(-a_i - a_j) \neq 0$ when $i = j$. This suggests we consider the $|A|$ by $|A|$ matrix M with $M_{ij} = P(-a_i - a_j)$. This is a diagonal matrix, with $M_{ii} = P(a_i)$. Thus the rank of this matrix is the dimension of the support of P , so it suffices to upper bound the rank of M . The key observation, where we now explicitly employ the fact that P is a polynomial, is that $P(-x - y)$ is a polynomial in $2n$ variables $x, y \in \mathbb{F}_3^n$,

Part VIII

Decoupling

Decoupling Theory is an in depth study of how ‘interference patterns’ can show up when combined waves with frequency supports in disjoint regions of space. The geometry of these regions effects how much constructive interference can happen. Of course decoupling theory is essential to studying many dispersive partial differential equations, but also has surprising applications in number theory as well, as well as other areas of harmonic analysis, such as restriction theory. The theory of decoupling was initiated by Wolff in the early 2000s, and Laba-Wolff, Laba-Pramanik, Pramanik-Seeger, and Garrigos and Seeger in the 2000s. But the theory was brought to the forefront of harmonic analysis by Bourgain, Demeter, and Guth in the mid 2010s.

Chapter 47

The General Framework

In any norm space X , given $x_1, \dots, x_N \in X$, one can apply the Cauchy-Schwartz inequality to obtain the estimate

$$\|x_1 + \dots + x_N\|_X \leq \|x_1\|_X + \dots + \|x_N\|_X.$$

Such a result is often sharp for general x_1, \dots, x_N . For instance, when $X = L^1(\mathbf{R}^d)$, and the x_1, \dots, x_N are functions with disjoint supports, but with equal L^1 norm. Iff $\|x_i\|_X \sim 1$ for each i , then we conclude that

$$\|x_1 + \dots + x_N\|_X \lesssim N.$$

However, if the x_1, \dots, x_n are ‘uncorrelated’, then one can often expect this result to be substantially improved. For instance, if X is a Hilbert space, and if x_1, \dots, x_N are pairwise orthogonal, Bessel’s inequality allows us to conclude that

$$\|x_1 + \dots + x_N\|_X \leq \left(\|x_1\|_X^2 + \dots + \|x_N\|_X^2 \right)^{1/2}.$$

In this case, if $\|x_i\|_X \sim 1$, then

$$\|x_1 + \dots + x_N\|_X \lesssim N^{1/2}.$$

Thus we obtain a significant ‘square root cancellation’ in N . For instance, in $L^2(\mathbf{R}^d)$, this occurs if x_1, \dots, x_N have disjoint supports, or perhaps more interestingly in the context of Fourier analysis, if their Fourier transforms have disjoint supports.

The most basic case of square root cancellation occurs if we replace the property of orthogonality with *almost orthogonality*. If x_1, \dots, x_N are a family of vectors in a Hilbert space H , then

$$\|x_1 + \dots + x_N\|_H^2 = \sum \|x_i\|_H^2 + 2 \sum_{1 \leq i < j \leq N} \langle x_i, x_j \rangle.$$

Thus, in particular, if each x_i is orthogonal to all but at most M of the other vectors in this family, then we conclude that

$$\|x_1 + \dots + x_N\|_H^2 \leq \sum_i \|x_i\|_H^2 + 2M \sum_i \|x_i\|_H^2 \leq (2M + 1) \sum \|x_i\|_H^2.$$

Thus, provided that M is much smaller than N , we also obtain square root cancellation in this setting.

We are interested in determining what causes ‘square root cancellation’ in more general norm spaces than just Hilbert. The theory of *almost orthogonality* studies this phenomena in Hilbert spaces, but we are interested in studying when this phenomenon in other norm spaces. Informally, we say x_1, \dots, x_N satisfies a *decoupling inequality* in a norm space X if for all $\varepsilon > 0$, we have

$$\|x_1 + \dots + x_N\|_X \lesssim_\varepsilon N^\varepsilon (\|x_1\|_X^2 + \dots + \|x_N\|_X^2)^{1/2}.$$

Thus decoupling theory is the study of when certain values can be correlated with one another, in general norm spaces. Of particular importance in harmonic analysis is the determination of the properties the Fourier transform of a family of functions must have to enable us to obtain decoupling phenomena.

Remark. We are primarily interested in studying decoupling in $L^p(\Omega)$. However, in this case we should only expect decoupling to occur when $p \geq 2$. Functions in L^p spaces are ‘most orthogonal’ when their supports are disjoint. If a family of functions $\{f_i\}$ have disjoint supports, then we actually have

$$\|f_1 + \dots + f_N\|_{L^p(\Omega)} = \left(\|f_1\|_{L^p(\Omega)}^p + \dots + \|f_N\|_{L^p(\Omega)}^p \right)^{1/p}.$$

For $p < 2$, we have a sharp inequality

$$\left(\|f_1\|_{L^p(\Omega)}^p + \dots + \|f_N\|_{L^p(\Omega)}^p \right)^{1/p} \leq N^{1/p-1/2} \left(\|f_1\|_{L^2(\Omega)}^2 + \dots + \|f_N\|_{L^2(\Omega)}^2 \right)^{1/2},$$

so we do not have square root cancelation in this setting.

The most basic case where we can obtain decoupling in $L^p(\Omega)$ for $p > 2$ is when p is an even integer. As a basic example, say a family of functions $f_1, \dots, f_N \in L^4(\Omega)$ are *biorthogonal* if $\{f_i f_j : i < j\}$ forms an orthogonal family in $L^2(\Omega)$.

Theorem 47.1. *If f_1, \dots, f_N are a biorthogonal family, then*

$$\|f_1 + \dots + f_N\|_{L^4(\Omega)} \lesssim \left(\|f_1\|_{L^4(\Omega)}^2 + \dots + \|f_N\|_{L^4(\Omega)}^2 \right)^{1/2}.$$

Proof. We write

$$\begin{aligned} \|f_1 + \dots + f_N\|_{L^4(\Omega)}^2 &= \|(f_1 + \dots + f_N)^2\|_{L^2(\Omega)} \\ &= \left\| \sum_{1 \leq i, j \leq N} f_i f_j \right\|_{L^2(\Omega)} \\ &\lesssim \sum_{i=1}^N \|f_i^2\|_{L^2(\Omega)} + \left\| \sum_{1 \leq i < j \leq N} f_i f_j \right\|_{L^2(\Omega)} \end{aligned}$$

Applying Bessel's inequality, we conclude that

$$\begin{aligned} \left\| \sum_{1 \leq i < j \leq N} f_i f_j \right\|_{L^2(\Omega)} &= \left(\sum_{1 \leq i < j \leq N} \|f_i f_j\|_{L^2(\Omega)}^2 \right)^{1/2} \\ &= \left\| \sum_{i=1}^N |f_i|^2 \right\|_{L^2(\Omega)} \lesssim \sum_{i=1}^N \|f_i^2\|_{L^2(\Omega)}. \end{aligned}$$

Combining these calculations, noticing that $\|f_i^2\|_{L^2(\Omega)} = \|f_i\|_{L^4(\Omega)}^2$, and taking in square roots completes the claim. \square

Remark. The calculation above shows a decoupling inequality still holds if the family $f_i f_j$ is almost biorthogonal, e.g. if each of the elements of the family $\{f_i f_j\}$ are orthogonal to all but $O_\varepsilon(N^\varepsilon)$ of the other elements of the family.

Remark. Similarly, if $f_1, \dots, f_N \in L^6(\Omega)$ are chosen to be *triorthogonal*, in the sense that the family of functions $\{f_i f_j f_k\}$ are orthogonal to one another, one can obtain a decoupling inequality in the L^6 norm.

We will be most interested in studying families of functions f_1, \dots, f_N with disjoint Fourier supports in $L^p(\mathbf{R}^d)$, where $p \geq 2$. It is then certainly true that

$$\|f_1 + \dots + f_N\|_{L^2(\mathbf{R}^d)} \leq \left(\|f_1\|_{L^2(\mathbf{R}^d)}^2 + \dots + \|f_N\|_{L^2(\mathbf{R}^d)}^2 \right)^{1/2}.$$

However, for $p > 2$ having disjoint Fourier supports does not immediately imply decoupling, because constructive interference can still occur in the L^p norm in a way not detected in the L^2 norm, and we require additional features of the family in order to guarantee that this constructive interference does not occur.

Theorem 47.2. *If $f_1, \dots, f_N \in L^p(\mathbf{R}^d)$, have disjoint Fourier support, and $2 \leq p \leq \infty$, then*

$$\|f_1 + \dots + f_N\|_{L^p(\mathbf{R}^d)} \leq N^{1/2-1/p} \left(\|f_1\|_{L^p(\mathbf{R}^d)}^2 + \dots + \|f_N\|_{L^p(\mathbf{R}^d)}^2 \right)^{1/2}.$$

Proof. For $p = \infty$, we have the trivial inequality

$$\begin{aligned} \|f_1 + \dots + f_N\|_{L^\infty(\mathbf{R}^d)} &\leq \|f_1\|_{L^\infty(\mathbf{R}^d)} + \dots + \|f_N\|_{L^\infty(\mathbf{R}^d)} \\ &\leq N^{1/2} \left(\|f_1\|_{L^\infty(\mathbf{R}^d)}^2 + \dots + \|f_N\|_{L^\infty(\mathbf{R}^d)}^2 \right)^{1/2}. \end{aligned}$$

By a density argument (applying a cutoff in frequency space), we may assume without loss of generality that the functions $\{f_i\}$ are Schwartz. But then by orthogonality of the Fourier transform implies the functions themselves are orthogonal, so we have

$$\|f_1 + \dots + f_N\|_{L^2(\mathbf{R}^d)} \leq \left(\|f_1\|_{L^2(\mathbf{R}^d)}^2 + \dots + \|f_N\|_{L^2(\mathbf{R}^d)}^2 \right)^{1/2}.$$

We can then interpolate with the case $p = \infty$. □

In general, this result is optimal, as the next result shows one can have significant constructive interference for $p > 2$ if our functions are spaced about on a periodic family of frequencies.

Example. Fix $\phi \in \mathcal{S}(\mathbf{R}^d)$ with $\phi(0) = 1$, and with Fourier support in $[0, 1]$. For each $k \in \{1, \dots, N\}$, set $f_k = e^{2\pi i k x} \phi(x)$. Then f_k has Fourier support in $[2k, 2k + 1]$, and $f_k(0) = 1$. The uncertainty principle thus implies that the

functions f_k are all roughly constant at a scale $O(1/N)$, which implies that in a ball of radius $O(1/N)$ around the origin, we have $f_1(x) + \dots + f_n(x) \approx 1$. Thus we conclude that

$$\|f_1 + \dots + f_N\|_{L^p(\mathbf{R})} \gtrsim N^{1-1/p}.$$

On the other hand, we have

$$\left(\|f_1\|_{L^p(\mathbf{R})}^2 + \dots + \|f_N\|_{L^p(\mathbf{R})}^2 \right)^{1/2} \lesssim N^{1/2},$$

Thus

$$\|f_1 + \dots + f_N\|_{L^p(\mathbf{R})} \gtrsim N^{1/2-1/p} \left(\|f_1\|_{L^p(\mathbf{R})}^2 + \dots + \|f_N\|_{L^p(\mathbf{R})}^2 \right)^{1/2},$$

which shows our result is tight up to constants.

In the face of this result, we are interested in knowing, for a given family \mathcal{S} of disjoint sets in \mathbf{R}^d , what the smallest constant $\text{Dec}(\mathcal{S}, p)$ is for which it is true that if f_1, \dots, f_N have Fourier support on distinct regions $S_1, \dots, S_N \in \mathcal{S}$, we have

$$\|f_1 + \dots + f_N\|_{L^p(\mathbf{R}^d)} \leq \text{Dec}(\mathcal{S}, p) \left(\|f_1\|_{L^p(\mathbf{R}^d)}^2 + \dots + \|f_N\|_{L^p(\mathbf{R}^d)}^2 \right)^{1/2}.$$

Pure orthogonality gives $\text{Dec}(\mathcal{S}, 2) = 1$. The triangle inequality gives $\text{Dec}(\mathcal{S}, \infty) \leq \#(\mathcal{S})^{1/2}$, and this is actually sharp: we have $\text{Dec}(\mathcal{S}, \infty)$

If each element of \mathcal{S} contains a ball of radius δ , then by summing up modulated bump functions on these balls, we can find functions such that $\text{Dec}(\mathcal{S}, \infty) \gtrsim \delta^d \#(\mathcal{S})^{1/2}$. Thus non-trivial decoupling in the framework we are considering is impossible in $L^\infty(\mathbf{R}^d)$, i.e. $\text{Dec}(\mathcal{S}, \infty) \sim_\delta \#(\mathcal{S})^{1/2}$. In between, we can interpolate to get $\text{Dec}(\mathcal{S}, p) \leq \#(\mathcal{S})^{1/2-1/p}$.

Such a result depends significantly on the geometric structure of the regions in \mathcal{S} . The techniques we will use (e.g. induction on scales) imply the need for the ' ε loss' given by the N^ε factor above. Below is a positive result for a particular family \mathcal{S} , easily proved using the biorthogonality arguments established above.

Theorem 47.3. *Suppose \mathcal{S} is a family of sets in \mathbf{R}^d such that for any four sets $S_1, S_2, S_3, S_4 \in \mathcal{S}$ with $\{S_1, S_2\} \neq \{S_3, S_4\}$, the sets $S_1 + S_2$ are disjoint from $S_3 + S_4$. Then if distinct sets $S_1, \dots, S_N \in \mathcal{S}$ are selected from \mathcal{S} , and f_1, \dots, f_N*

are a family of Schwartz functions in \mathbf{R}^d such that f_i has Fourier support in S_i for each i , we find''

$$\|f_1 + \cdots + f_N\|_{L^4(\Omega)} \lesssim \left(\|f_1\|_{L^4(\Omega)}^2 + \cdots + \|f_N\|_{L^4(\Omega)}^2 \right)^{1/2}.$$

Remark. We say a set of integers $A \subset \{1, \dots, N\}$ is a *Sidon set* if there does not exist a nontrivial solution to the equation $a_1 + a_2 = a_3 + a_4$. If A is Sidon, then $\mathcal{S} = \{[2k, 2k+1] : k \in A\}$ satisfies the constraints of the result above, and so we obtain that if $\{f_k : k \in A\}$ are a family of Schwartz functions such that f_k has Fourier support in $[2k, 2k+1]$, then

$$\left\| \sum_{k \in A} f_k \right\|_{L^4(\mathbf{R})} \lesssim \left(\sum_{k \in A} \|f_k\|_{L^4(\mathbf{R})}^2 \right)^{1/2}.$$

On the other hand, a variant of the example above shows that for any Sidon set A , there is a family of functions $\{f_k : k \in A\}$ with f_k having Fourier support on $[2k, 2k+1]$, and with

$$\left\| \sum_{k \in A} f_k \right\|_{L^4(\mathbf{R})} \gtrsim \frac{\#(A)^{1/2}}{N^{1/4}} \left(\sum_{k \in A} \|f_k\|_{L^4(\mathbf{R})}^2 \right)^{1/2}.$$

Combining this inequality with the decoupling inequality, we obtain an interesting number theory result: any Sidon set A , must satisfy the bound $\#(A) \lesssim N^{1/2}$. More generally, we can extend this result to show that any set $A \subset \{0, \dots, N-1\}$ having no nontrivial solutions to the equation

$$x_1 + \cdots + x_m = y_1 + \cdots + y_m$$

should satisfy $\#(A) \lesssim N^{1/m}$.

Another example family of sets \mathcal{S} where Decoupling occurs occurs in Littlewood-Paley theory.

Theorem 47.4. *Let \mathcal{S} be the collection of all boxes in \mathbf{R}^d of the form*

$$I_k = I_{\pm k_1} \times \cdots \times I_{\pm k_d}$$

such that $I_{+k_1} = [2^{k_1}, 2^{k_1+1}]$ and $I_{-k_1} = [-2^{k_1+1}, -2^{k_1}]$. Littlewood-Paley theory implies that if $S_1, \dots, S_N \in \mathcal{S}$ and f_1, \dots, f_N are Schwartz functions with f_i having Fourier support on S_i for each i , then for each $2 \leq p < \infty$,

$$\|f_1 + \cdots + f_N\|_{L^p(\mathbf{R}^d)} \sim_{p,d} \|f_i\|_{L^p(\mathbf{R}^d) l_N^2} \lesssim \|f_i\|_{l_N^2 L^p(\mathbf{R}^d)}.$$

This is precisely a decoupling inequality. More generally, whenever we have a reverse square function estimate

$$\|f_1 + \cdots + f_N\|_{L^p(\mathbf{R}^d)} \lesssim \|f_i\|_{L^p_x L^2_N},$$

we get a decoupling inequality by changing norms.

We say a set $\omega \subset \mathbf{R}^d$ is an *almost rectangular box* with sidelengths L_1, \dots, L_d if there is a rectangular box R with sidelengths L_1, \dots, L_d centered at the origin and $\theta \in \mathbf{R}^d$ such that $\theta + C^{-1}R \subset \omega \subset \theta + C \cdot R$ for some universal constant C . As a special case, we have *almost cubes* as well. For each $R \geq 1$, partition $[-1, 1]^{d-1}$ into $\sim R^{(d-1)/2}$ almost rectangular boxes of sidelength $R^{-1/2}$. If we consider the paraboloid $\mathbf{P}^{n-1} = \{\xi, |\xi|^2\}$, then we obtain a partition of $N(\mathbf{P}^{n-1} \cap [0, 1]^d, R^{-1})$ by a family $\Theta(1/R)$ of $R^{-1/2} \times R^{-1}$ almost rectangular boxes, by setting

$$\Theta(1/R) = \{(\omega \times \mathbf{R}) \cap N(\mathbf{P}^{n-1} \cap [0, 1]^d)\}$$

It is conjectured that for $p = 2n/(n-1)$, and for Schwartz functions f_1, \dots, f_N with Fourier support on distinct elements of $\Theta(1/R)$, we have

$$\|f_1 + \cdots + f_N\|_{L^p(\mathbf{R}^d)} \lesssim_R \|f_i\|_{L^p(\mathbf{R}^d) l_N^2},$$

This reverse square function estimates is strong. In particular, it implies the restriction conjecture. But it also implies the decoupling estimate

$$\|f_1 + \cdots + f_N\|_{L^p(\mathbf{R}^d)} \lesssim_\varepsilon R^\varepsilon \left(\|f_1\|_{L^p(\mathbf{R}^d)} + \cdots + \|f_N\|_{L^p(\mathbf{R}^d)} \right)^{1/2}$$

. However, unlike the restriction conjecture, this problem is closed: we have a proof of this decoupling estimate not only when $p = 2d/(d-1)$, but even when $2 \leq p \leq 2(d+1)/(d-1)$.

47.1 Localized Estimates

Suppose f_1, \dots, f_N are Schwartz functions in \mathbf{R}^d with disjoint Fourier supports, and $\Omega \subset \mathbf{R}^d$. A natural question to ask is when one should expect

$$\|f_1 + \cdots + f_N\|_{L^2(\Omega)}^2 \lesssim \|f_1\|_{L^2(\Omega)}^2 + \cdots + \|f_N\|_{L^2(\Omega)}^2.$$

If we consider the bump function counterexample constructed from earlier, and let $\Omega = \{x \in \mathbf{R} : |x| \lesssim 1/N\}$, then $\|f_1 + \cdots + f_N\|_{L^2(\Omega)} \gtrsim N$, whereas $\|f_k\|_{L^2(\Omega)}^2 \lesssim 1/N$ so $\|f_1\|_{L^2(\Omega)}^2 + \cdots + \|f_N\|_{L^2(\Omega)}^2 \lesssim 1$, which means such a result cannot be obtained. However, we shall find that such a result holds if Ω is large enough, depending on the supports of f_1, \dots, f_N , and if we allow weighted estimates.

Let us begin with the case in one dimension. Given an interval I with centre x_0 , and length R , we consider the weight function

$$w_I(x) = \left(1 + \frac{|x - x_0|}{R}\right)^{-M}$$

It is a useful heuristic that if f has Fourier support in I , then f is ‘locally constant’ on intervals of length $1/|I|$.

In \mathbf{R}^d , given a ball B with centre x_0 and radius R , we consider the weight function

$$w_B(x) = \left(1 + \frac{|x - x_0|}{R}\right)^{-M},$$

where M is a large integer. Then

$$\int w_B(x) dx$$

TODO FINISH THIS

47.2 Vinogradov Systems

One long standing number theoretic conjecture that has been particularly amenable to the use of decoupling techniques is *Vinogradov’s Conjecture*. Let $J_{s,k}(N)$ denote the number of solutions to the *system* of equations

$$x_1^i + \cdots + x_s^i = y_1^i + \cdots + y_s^i$$

for $1 \leq i \leq k$, where $x_1, \dots, x_s, y_1, \dots, y_s \in \{1, \dots, N\}$. Our primary interest is determining the asymptotic growth in this quantity as $N \rightarrow \infty$. Setting $y_i = x_i$ gives a family of N^s solutions. Thus $J_{s,k}(N) \geq N^s$. On the other hand, dyadic pigeonholing implies that there exists some family of integer tuples $I \subset [1, N] \times \cdots \times [1, N^k]$ and some integer M with $N^s = M\#(I)$

such that for each tuple $(r_1, \dots, r_k) \in I$, there are $\Omega(M)$ tuples of values (x_1, \dots, x_s) such that for each $1 \leq i \leq k$,

$$x_1^i + \dots + x_s^i = r_i.$$

For any i , $x_1^i + x_s^i$ takes on values between 1 and N^i . Since $1 \leq \#(I) \leq N^{k(k+1)/2}$, we have $N^{s-k(k+1)/2} \lesssim M \lesssim N^s$, and so

$$J_{s,k}(N) \gtrsim \#(I)M^2 \gtrsim N^{2s-k(k+1)/2},$$

a bound that is much tighter if the number of variables is large, i.e. $s \geq k(k+1)/2$. Vinogradov's mean value conjecture is that

$$J_{s,k}(N) \lesssim_{s,k,\varepsilon} N^\varepsilon (N^{2s-k(k+1)/2} + N^s),$$

for any $\varepsilon > 0$. One result of decoupling is a proof of this conjecture.

Remark. Another approach to the Vinogradov mean value conjecture is to use more number theoretic techniques, for instance, efficient congruencing. There has even been some feedback from this alternate approach in the field of decoupling, e.g. using efficient congruencing to obtain decoupling inequalities (see Guo-Li-Yung-Zorin Kranich, 2021).

Similar types of number theoretic problems had been studied using harmonic analysis techniques. Classically, Hardy and Littlewood studied the quantity $\text{HL}_{k,s}(N)$, which counts the number of solutions to the *single equation*

$$x_1^k + \dots + x_s^k = y_1^k + \dots + y_s^k$$

with $x_1, \dots, x_s, y_1, \dots, y_s \in \{1, \dots, N\}$. To study this quantity, Hardy and Littlewood introduced the function

$$f_{k,N}(x) = \sum_{n=1}^N e^{2\pi i n^k x}.$$

Orthogonality then implies that

$$\text{HL}_{k,s}(N) = \|h_{k,N}\|_{L^{2s}[0,1]}.$$

Bounding the nature of the function $h_{k,N}$ then involves Hardy and Littlewood's circle method. On the other hand, for Vinogradov's conjecture we are lead to study a higher dimensional function

$$j_{k,N}(x) = \sum_{n=1}^N e^{2\pi i (nx_1 + n^2 x_2 + \dots + n^k x_k)},$$

in which case

$$J_{k,s}(N) = \|j_{k,N}\|_{L^{2s}[0,1]^k}.$$

Though in higher dimensions, Vinogradov initially expected his result to be *easier* than Hardy and Littlewood's problem, because of the *rescalable* nature of the problem, which we will get to later. Indeed, decoupling has lead to a solution of Vinogradov's conjecture for almost all parameters, but a complete solution to the conjectured bound

$$\text{HL}_{k,s}(N) \lesssim_\varepsilon N^\varepsilon (N^s + N^{2s-k})$$

has not yet been resolved.

To do our analysis, it will be helpful to rescale our quantities. Thus we let

$$f_{k,N}(x) = \sum_{n=1}^N e^{2\pi i((n/N)x_1 + \dots + (n/N)^k x_k)}.$$

The function $f_{k,N}$ is then N -periodic, and so Vinogradov's conjecture will follow from showing that

$$\|f_{k,N}\|_{L^{2s}[0,N^k]^k} = N^{k^2/2s} \|j_{k,N}\|_{L^{2s}[0,1]^k}$$

satisfies a bound

$$\|f_{k,N}\|_{L^{2s}[0,N^k]^k} \lesssim_\varepsilon N^{k^2/2s+\varepsilon} (N^s + N^{2s-k}).$$

This is what we will address using the theory of decoupling.

From the perspective of decoupling, it will be more handy to study the quantities

$$\tilde{j}_{k,N}(x) = \sum_{n=1}^N e^{2\pi i(nx_1 + \dots + n^k x_k)} \phi(x),$$

and

$$\tilde{f}_{k,N}(x) = \sum_{n=1}^N e^{2\pi i((n/N)x_1 + \dots + (n/N)^k x_k)} \phi(x_1/N, \dots, x_k/N^k),$$

where $\phi \in \mathcal{S}(\mathbf{R}^d)$ has Fourier support in $|\xi| \leq 1/10$ and $\phi(x) \geq 1$ in a neighborhood of the origin. Bounding the L^{2s} norm of $\tilde{j}_{k,N}$ and $\tilde{f}_{k,N}$ is essentially

equivalent to bounding $j_{k,N}$ since we can upper bound $j_{k,N}$ and $f_{k,N}$ pointwise by $O(1)$ sums of this form on the domain upon which we are taking the L^{2s} norm. Now if we let

$$f_i = e^{2\pi i((n/N)x_1 + \dots + (n/N)^k x_k)} \phi(x_1/N, \dots, x_k/N^k),$$

then \hat{f}_i is supported on a rectangle with sidelengths $O(N) \times \dots \times O(N^k)$ centered at the point $(n/N, \dots, (n/N)^k)$ of the moment curve. The l^2 decoupling inequality for the moment curve thus implies that

$$\|f\|_{L^{2s}(\mathbf{R}^d)} \lesssim_\varepsilon N^{1/2+\varepsilon},$$

which proves Vinogradov's mean value result for $s \geq k^3/(2k^2 - 1)$, i.e. for $s \gtrsim k$, which, up to constants, is the current state of the art for the mean-value theorem.