

Large Sets Avoiding Rough Patterns

Jacob Denson* Malabika Pramanik* Joshua Zahl*

March 21, 2019

Abstract

The pattern avoidance problem seeks to construct a set $X \subset \mathbf{R}^d$ with large dimension that avoids a prescribed pattern such as three term arithmetic progressions, or more general patterns such as avoiding points x_1, \dots, x_n such that $f(x_1, \dots, x_n) = 0$ (three term arithmetic progressions are specified by the pattern $x_1 - 2x_2 + x_3 = 0$). Previous work on the subject has considered patterns described by polynomials, or by functions f satisfying certain regularity conditions. We consider the case of ‘rough patterns’. There are several problems that fit into the framework of rough pattern avoidance. As a first application, if $Y \subset \mathbf{R}^d$ is a set with Minkowski dimension α , we construct a set X with Hausdorff dimension $1 - \alpha$ such that $X + X$ is disjoint from Y . As a second application, given a set Y of dimension close to one, we can construct a subset $X \subset Y$ of dimension $1/2$ that avoids isosceles triangles. [**TODO:** I’d like to replace this with a rectifiability statement]

A major question in modern geometric measure theory is whether sets with sufficiently large Hausdorff dimension necessarily contain copies of certain patterns. For example, Theorem 6.8 of [11] shows that each set $X \subset \mathbf{R}^d$ with Hausdorff dimension exceeding one must contain three colinear points. On the other hand, the main result of [9] constructs a set $X \subset \mathbf{R}^2$ with full Hausdorff dimension such that no four points in X form the vertices of a parallelogram. At the time of this writing, many fundamental questions about the relation between the geometric patterns of a set and its fractional dimension remain open. The threshold dimension above which patterns are guaranteed can vary for different patterns, or not even exist at all. In this paper, we provide lower bounds for the threshold by solving the pattern avoidance problem; constructing sets with large dimension avoiding patterns.

One natural way to approach the pattern avoidance problem is to find general methods for constructing pattern avoiding sets by exploiting a particular geometric feature of the pattern. In [10], Máthé shows that, given a pattern specified by a countable union of rational coefficient bounded degree polynomials f_1, f_2, \dots in nd variables, one can construct a set $X \subset \mathbf{R}^d$ such that for every collection of distinct points x_1, \dots, x_n , and each index k , $f_k(x_1, \dots, x_n) \neq 0$. In [5], Fraser and the second author consider a related problem where the polynomials f_k are replaced with full rank C^1 functions.

In this paper, we also find general methods for constructing pattern avoid sets. But rather than avoiding the zeroes of a function, we fix $Z \subset \mathbf{R}^{dn}$, and construct sets X such

*University of British Columbia, Vancouver BC, {denson, malabika, jzahl}@math.ubc.ca.

that for any distinct $x_1, \dots, x_n \in X$, $(x_1, \dots, x_n) \notin Z$. We then say X avoids the pattern specified by Z . For instance, if $Z = \{(x_1, x_2, x_3) \in (\mathbf{R}^d)^3 : (x_1, x_2, x_3) \text{ are colinear}\}$, then $X \subset \mathbf{R}^d$ avoids the pattern specified by Z precisely when X does not contain three colinear points. Our problem generalizes the problem statements considered in [10] and [5] if we set $Z = \bigcup_{k=1}^{\infty} f_k^{-1}(0)$. From this perspective, Máthé constructs sets avoiding a set Z formed from a countable union of algebraic varieties, while Fraser and the second author construct sets avoiding a set Z formed from the countable union of C^1 manifolds.

The advantage of the formulation of pattern avoidance we consider is that it is now natural to consider ‘rough’ sets $Z \subset \mathbf{R}^{dn}$ which are not naturally specified as the zero set of a function. In particular, in this paper we consider Z formed from the countable union of sets, each with lower Minkowski dimension bounded above by some α . In contrast with previous work, no further assumptions are made on Z . The dimension of the avoiding set X we will eventually construct depends only on the codimension $nd - \alpha$ of the set Z , and the number of variables n .

Theorem 1. *Let $Z \subset \mathbf{R}^{nd}$ be a countable union of compact sets, each with lower Minkowski dimension at most α , with $d \leq \alpha < dn$. Then there exists a set $X \subset [0, 1]^d$ with Hausdorff dimension at least $(nd - \alpha)/(n - 1)$ such that whenever $x_1, \dots, x_n \in X$ are distinct, then $(x_1, \dots, x_n) \notin Z$.*

Remark. *When $\alpha < d$, the pattern avoidance problem is trivial, since $X = [0, 1]^d - \pi(Z)$ is full dimensional and solves the pattern avoidance problem, where $\pi(x_1, \dots, x_n) = x_1$ is a projection map from \mathbf{R}^{dn} to \mathbf{R}^d . The case $\alpha = dn$ is trivial as well, since we can set $X = \emptyset$.*

When Z is a countable union of smooth manifolds, Theorem 1 generalizes Theorem 1.1 and 1.2 from [5]. Section 6 is devoted to a comparison of our methods with [5], as well as other general pattern avoidance methods. But our result is more interesting when it can be applied to truly ‘rough’ sets. One surprising application to our method is obtained by considering a ‘rough’ set Y in addition to a set Z , and finding a set $X \subset Y$ of large dimension avoiding Z . Previous methods in the literature fundamentally exploit the ability to select points on the entirety of Euclidean space, and so it remains unlikely that we can obtain any result of this form using their methods, unless Y is suitably flat, i.e. a smooth surface. To contrast this, our result even applies for certain sets Y which are of Cantor type. We discuss applications of our result in Section 5.

Theorem 1 is proved using a Cantor-type construction, a common theme in the surrounding literature, described explicitly in Section 3. For a sequence of decreasing lengths $\{l_n\}$ with $l_n \rightarrow 0$, the construction specifies a selection mechanism for a nested family of sets $X_n \rightarrow X$, with X_n a union of sidelength l_n cubes and avoiding Z at scales close to l_n . Our *key* contribution to this process is using the probabilistic method to guarantee the existence of efficient selections at each scale, described in Section 2. This selection also assigns mass uniformly on intermediate scales, akin to [5], ensuring that the selection procedure at a single scale is the sole reason for the Hausdorff dimension we calculate in Section 4. Furthermore, the randomization allows us to avoid the complicated queueing techniques in [8] and [5], which has the additional benefit that we can confront the entire behaviour of Z at scales close to l_n simultaneously during the n th step of the construction.

1 Frequently Used Notation and Terminology

Our argument heavily depends upon discretizing sets into unions of cubes. Throughout our argument, we use the following notation:

- (A) A *dyadic length* is a length l equal to 2^{-k} for some non-negative integer k .
- (B) Given a length $l > 0$, we let \mathcal{B}_l^d denote the family of all half open cubes in \mathbf{R}^d with sidelength l and corners on the lattice $(l \cdot \mathbf{Z})^d$. That is,

$$\mathcal{B}_l^d = \{[a_1, a_1 + l] \times \cdots \times [a_d, a_d + l] : a_k \in l \cdot \mathbf{Z}\}.$$

If $E \subset \mathbf{R}^d$, $\mathcal{B}_l^d(E)$ is the family of cubes in \mathcal{B}_l^d intersecting E , i.e.

$$\mathcal{B}_l^d(E) = \{I \in \mathcal{B}_l^d : I \cap E = \emptyset\}.$$

- (C) The *lower* and *upper Minkowski dimension* of a compact set $Z \subset \mathbf{R}^d$ are defined as

$$\underline{\dim}_{\mathbf{M}}(Z) = \liminf_{l \rightarrow 0} \frac{\log(\#\mathcal{B}_l^d(Z))}{\log(1/l)} \quad \text{and} \quad \overline{\dim}_{\mathbf{M}}(Z) = \limsup_{l \rightarrow 0} \frac{\log(\#\mathcal{B}_l^d(Z))}{\log(1/l)}.$$

If $\underline{\dim}_{\mathbf{M}}(Z) = \overline{\dim}_{\mathbf{M}}(Z)$, then we define $\dim_{\mathbf{M}}(Z)$ equal to the common value.

- (D) If $0 \leq \alpha$ and $\delta > 0$, we define the dyadic Hausdorff content of a set $E \subset \mathbf{R}^d$ as

$$H_\delta^\alpha(E) = \inf \left\{ \sum_{k=1}^m l_k^\alpha : E \subset \bigcup_{k=1}^m I_k \text{ and } I_k \in \mathcal{B}_{l_k}^d, l_k \leq \delta \text{ for all } k \right\}.$$

The α -dimensional dyadic Hausdorff measure H^α on \mathbf{R}^d is $H^\alpha(E) = \lim_{\delta \rightarrow 0} H_\delta^\alpha(E)$, and the *Hausdorff dimension* of a set E is $\dim_{\mathbf{H}}(E) = \inf\{\alpha \geq 0 : H^\alpha(E) = 0\}$.

- (E) Given $I \in \mathcal{B}_l^{dn}$, we can decompose I as $I_1 \times \cdots \times I_n$ for unique cubes $I_1, \dots, I_n \in \mathcal{B}_l^d$. We say I is *strongly non-diagonal* if the cubes I_1, \dots, I_n are distinct. Strongly non-diagonal cubes will play an important role in Section 2, when we solve a discrete version of Theorem 1.

- (F) Adopting the terminology of [7], we say a collection of sets $\{U_k\}$ is a *strong cover* of a set E if $E \subset \limsup U_k$, which means every element of E is contained in infinitely many of the sets U_k . This idea will be useful in Section 3.

- (G) A *Frostman measure* of dimension α is a non-zero compactly supported probability measure μ on \mathbf{R}^d such that for every cube I of sidelength l , $\mu(I) \lesssim l^\alpha$. Note that a measure μ satisfies this inequality for every cube I if and only if it satisfies the inequality for cubes whose sidelengths are dyadic lengths. *Frostman's lemma* says that

$$\dim_{\mathbf{H}}(E) = \sup \left\{ \alpha : \begin{array}{l} \text{there is an } \alpha \text{ dimensional Frostman} \\ \text{measure supported on } E \end{array} \right\}.$$

2 Avoidance at Discrete Scales

In this section we describe a method for avoiding Z at a single scale. We apply this technique in Section 3 at many scales to construct a set X avoiding Z at all scales. This single scale avoidance technique is the core part of our construction, and the efficiency with which we can avoid Z at a single scale has direct consequences on the Hausdorff dimension of the set X obtained in Theorem 1.

At a single scale, we solve a discretized version of the problem, where all sets are unions of cubes at two dyadic lengths $l \geq s$. In the discrete setting, Z is replaced by a discretization version of itself, i.e. a union of cubes in \mathcal{B}_s^{dn} , denoted by Z_s . Given a set E , which is a union of cubes in \mathcal{B}_l^d , our goal is to construct a union of cubes in \mathcal{B}_s^d , denoted F , such that $F \subset E$ and F^n is disjoint from the strongly non-diagonal cubes of Z_s (see Definition (E)).

In order to ensure the final set X obtained in Theorem 1 has large Hausdorff dimension regardless of the rapid decay of scales used in the construction of X , it is crucial that F is spread uniformly over E . We achieve this by decomposing E into sub-cubes in \mathcal{B}_r^d for some intermediate scale $r \in [s, l]$, and distributing F evenly among as many of these intermediate sub-cubes as possible. Assuming a mild regularity condition on the volume of Z_s , this is possible.

Lemma 1. *Fix two dyadic lengths $l \geq s$. Let E be a nonempty union of cubes in \mathcal{B}_l^d , and let Z_s be a union of cubes in \mathcal{B}_s^{dn} such that $|Z_s| \leq l^{dn}/2$. Then there exists a dyadic length $r \in [s, l]$ such that*

$$A_l |Z_s|^{1/d(n-1)} \leq r \leq \max(s, 2A_l |Z_s|^{1/d(n-1)}) \quad \text{where} \quad A_l = (2^{1/d}/l)^{1/(n-1)}. \quad (2.1)$$

For this scale, there exists a set $F \subset E$, which is a nonempty union of cubes in \mathcal{B}_s^d , satisfying the following three properties:

- (A) Avoidance: For any distinct $J_1, \dots, J_n \in \mathcal{B}_s^d(F)$, $J_1 \times \dots \times J_n \notin \mathcal{B}_s^{dn}(Z_s)$.
- (B) Non-Concentration: For any $I \in \mathcal{B}_r^d(E)$, there is at most one $J \in \mathcal{B}_s^d(F)$ with $J \subset I$.
- (C) Large Size: For any $I \in \mathcal{B}_l^d(E)$, $\# \mathcal{B}_s^d(F \cap I) \geq \# \mathcal{B}_r^d(I)/2 = (l/r)^d/2$.

In other words, F avoids strongly non-diagonal cubes in Z_s , and contains a single sidelength s portion of more than half of the sidelength r cubes contained in any sidelength l cube in E .

Proof. Let r be the smallest dyadic length larger than s and $A_l |Z_s|^{1/d(n-1)}$, so that (2.1) is satisfied. The assumption that $|Z_s| \leq l^{dn}/2$ implies

$$A_l |Z_s|^{1/d(n-1)} \leq A_l l^{n/(n-1)} / 2^{1/d(n-1)} = l.$$

Thus we have guaranteed that $r \in [s, l]$. For each $I \in \mathcal{B}_r^d(E)$, let J_I be a random element of $\mathcal{B}_s^d(I)$ chosen uniformly at random, independantly from the other random variables $J_{I'}$ with $I \neq I'$. Define a random set

$$U = \bigcup \{J_I : I \in \mathcal{B}_r^d(E)\},$$

and for each U , set

$$\mathcal{K}(U) = \{K \in \mathcal{B}_s^{dn}(Z_s) : K \in U^n, K \text{ strongly non-diagonal}\}$$

Then set

$$F_U = U - \{\pi(K) : K \in \mathcal{K}(U), K \text{ is strongly diagonal}\}, \quad (2.2)$$

where $\pi : \mathbf{R}^{dn} \rightarrow \mathbf{R}^d$ maps $K_1 \times \cdots \times K_n$ to K_1 for each $K_i \in \mathbf{R}^d$. Given any strongly non-diagonal cube $J_1 \times \cdots \times J_n \in \mathcal{B}_s^{dn}(Z_s)$, either $J_1 \times \cdots \times J_n \notin \mathcal{B}_s^{dn}(U^n)$, or $J_1 \times \cdots \times J_n \in \mathcal{B}_s^{dn}(U^n)$. If the former occurs then $J_1 \times \cdots \times J_n \notin \mathcal{B}_s^{dn}(F_U^n)$ since $F_U \subset U$, while if the latter occurs then $K \in \mathcal{K}(U)$, so $J_1 \notin \mathcal{B}_s^d(F_U)$. In either case, $J_1 \times \cdots \times J_n \notin \mathcal{B}_s^{dn}(F_U^n)$, so F_U satisfies Property (A). By construction, U contains at most one subcube $J \in \mathcal{B}_s^{dn}$ for each $I \in \mathcal{B}_l^{dn}(E)$. Since $F_U \subset U$, F_U satisfies Property (B). These properties are satisfied for any instance of U , but it is not true that Property (C) holds for each F_U . The remainder of this proof is devoted to showing this property holds for F_U with non-zero probability, guaranteeing the existence of the required set F satisfying the properties of the lemma.

For each cube $J \in \mathcal{B}_s^d(E)$, there is a unique ‘parent’ cube $I \in \mathcal{B}_r^d(E)$ such that $J \subset I$. Since I contains $(r/s)^d$ elements of $\mathcal{B}_s^d(E)$, and J_I is chosen uniformly at random from $\mathcal{B}_s^d(I)$,

$$\mathbf{P}(J \subset U) = \mathbf{P}(J_I = J) = (s/r)^d.$$

The cubes J_I are chosen independantly, so if J_1, \dots, J_k are distinct cubes in $\mathcal{B}_s^d(E)$, then the last calculation combined with Property (B) shows that

$$\mathbf{P}(J_1, \dots, J_k \in U) = \begin{cases} (s/r)^{dk} & : \text{if } J_1, \dots, J_k \text{ have distinct parents} \\ 0 & : \text{otherwise} \end{cases}$$

Let $K = J_1 \times \cdots \times J_n \in \mathcal{B}_s^{dn}(Z_s)$ be a strongly non-diagonal cube. Then cubes J_1, \dots, J_n are distinct, so the calculation we just performed implies

$$\mathbf{P}(K \subset U^n) = \mathbf{P}(J_1, \dots, J_k \in U) \leq (s/r)^{dn}.$$

Together with linearity of expectation, and (2.1), if K ranges over the strongly non-diagonal cubes of $\mathcal{B}_s^{dn}(Z_s)$, we find

$$\begin{aligned} \mathbf{E}(\#\mathcal{K}(U)) &= \sum_K \mathbf{P}(K \subset U^n) \leq \#\mathcal{B}_s^{dn}(Z_s) \cdot (s/r)^{dn} = |Z_s| r^{-dn} \\ &= [|Z_s| r^{-d(n-1)}] r^{-d} \leq [|Z_s| (A_l |Z_s|^{1/d(n-1)})^{-d(n-1)}] r^{-d} = [l^d/2] r^{-d} = (l/r)^d/2. \end{aligned}$$

Thus there exists at least one (non-random) set U_0 such that

$$\#\mathcal{K}(U_0) \leq \mathbf{E}(\#\mathcal{K}(U)) \leq (l/r)^d/2. \quad (2.3)$$

This means that the resultant set F_{U_0} is obtained by removing at most $(l/r)^d/2$ cubes in \mathcal{B}_s^d from U_0 . Since U_0 contains $(l/r)^d$ cubes in $\mathcal{B}_s^d(I)$ for each $I \in \mathcal{B}_l^d(E)$, F_{U_0} contains at least $(l/r)^d/2$ cubes in $\mathcal{B}_s^d(I)$ for each $I \in \mathcal{B}_l^d(E)$, so F_{U_0} satisfies Property (C). Setting $F = F_{U_0}$ completes the proof. \square

Remark. While the existence of the set F in Lemma 1 was obtained by probabilistic techniques, we emphasize that it's existence is a purely deterministic statement. One can find a candidate F constructively by checking all of the finitely many possible choice of U to find one particular choice U_0 which satisfies (2.3), and then defining F by (2.2). Thus the set we obtain in Theorem 1 exists by purely constructive means.

Our inability to select almost every cube in Lemma 1 means that repeated applications of the result will lead to a loss in Hausdorff dimension. In fact, in the worst case, applying the lemma causes us to lose as much Hausdorff dimension as is permitted by Theorem 1. Note that the value r specified in Theorem 1 is always bounded below by $|Z_s|$. We will later see that if Z is the countable union of sets with lower Minkowski dimension α , the scale s discretization Z_s satisfies $|Z_s| \leq s^{dn-\alpha-\varepsilon}$, for some small positive ε converging to zero as $s \rightarrow 0$. But we can certainly find sets Z with lower Minkowski dimension α satisfying $|Z_s| \geq s^{dn-\alpha}$ for all s . In this situation, (2.1) shows

$$r \geq A_l |Z_s|^{1/d(n-1)} \geq A_l s^{(dn-\alpha)/d(n-1)} \geq s^{(dn-\alpha)/d(n-1)} \quad (2.4)$$

If we combine this inequality with Property (B) of Lemma 1, we conclude

$$\frac{\log \# \mathcal{B}_s^d(F)}{\log(1/s)} \leq \frac{\log \# \mathcal{B}_r^d(E)}{\log(1/s)} = \frac{\log |E| r^{-d}}{\log(1/s)} = \frac{d \log(1/r) - \log |E|^{-1}}{\log(1/s)} \leq \frac{dn - \alpha}{n - 1}.$$

Given a set $F = \bigcap F_k$, where $\{F_k\}$ are infinitely many sets obtained from an application of Lemma 1 at a sequence of scales $\{s_k\}$ with $s_k \rightarrow 0$ as $k \rightarrow \infty$, then we can show

$$\dim_{\mathbf{H}}(F) \leq \lim_{s_k \rightarrow 0} \frac{\log \# \mathcal{B}_{s_k}^d(F_k)}{\log(1/s_k)} \leq \frac{dn - \alpha}{n - 1}.$$

This is why we can only lower the Hausdorff dimension of the set X obtained in Theorem 1 by $(dn - \alpha)/(n - 1)$. Furthermore, we see that we must be very careful to ensure applications of the discrete lemma are the only place in our proof where dimension is lost.

Remark. *Lemma 1 is the core method in our avoidance technique. The remaining argument is fairly modular. If, for a special case of Z , one can improve the result of Lemma 1 so that r is chosen on the order of $s^{\beta/d}$, then the remaining parts of our paper can be applied near verbatim to yield a set X with Hausdorff dimension β , as in Theorem 1. The last paragraph shows that when $|Z_s| \geq s^{dn-\alpha}$, which is certainly possible given the hypothesis of Theorem 1, the length r is chosen on the order of $s^{(dn-\alpha)/d(n-1)}$, as we saw in (2.4), which is why we obtain a Hausdorff dimension $(dn - \alpha)/(n - 1)$ set.*

3 Fractal Discretization

In this section we will construct the set X by applying Lemma 1 at many scales. Since Z is a countable union of compact sets with Minkowski dimension at most α , there exists a strong cover (see Definition (F)) of Z by cubes restricted to a sequence of dyadic lengths $\{l_k\}$. We will select this strong cover so that the scales l_k converge to 0 very quickly.

Lemma 2. *Let $Z \subset \mathbf{R}^{dn}$ be a countable union of compact sets, each with lower Minkowski dimension at most α . Let $\{\varepsilon_k\}$ be a sequence of positive numbers and let $\{f_k\}$ be a sequence of functions such that $f_k: (0, \infty) \rightarrow (0, \infty)$. Then there exists a sequence of dyadic lengths $\{l_k\}$ and compact sets $\{Z_k\}$ such that*

(A) *For each index $k \geq 2$, $l_k \leq f_{k-1}(l_{k-1})$.*

(B) For each index k , Z_k is a union of cubes in $\mathcal{B}_{l_k}^{dn}$.

(C) Z is strongly covered by the sets $\{Z_k\}$.

(D) For each index k , $\#\mathcal{B}_{l_k}^{dn}(Z_k) \leq (1/l_k)^{\alpha+\varepsilon_k}$.

Proof. Let Z be the union of sets $\{Y_k\}$ with $\underline{\dim}_{\mathbf{M}}(Y_k) \leq \alpha$ for any k . Let m_1, m_2, \dots be a sequence of integers that repeats each integer infinitely often. For each positive integer k , since $\underline{\dim}_{\mathbf{M}}(Y_{m_k}) \leq \alpha$, definition (C) implies that there exists arbitrarily small lengths l which satisfy $\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq 1/l^{\alpha+(\varepsilon_k/2)}$. Replacing l with a dyadic length at most twice the size of l , there are infinitely many *dyadic* scales l with

$$\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq \frac{1}{(l/2)^{\alpha+\varepsilon_k}} \leq \frac{2^{dn}}{l^{\alpha+(\varepsilon_k/2)}} = \frac{(2^{dn}l^{\varepsilon_k/2})}{l^{\alpha+\varepsilon_k}}.$$

In particular, we may select a dyadic length l such that $2^{dn}l^{\varepsilon_k/2} \leq 1$, and, if $k \geq 2$, also satisfying $l \leq f_{k-1}(l_{k-1})$. The first constraint together with the last calculation implies $\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq 1/l^{\alpha+\varepsilon_k}$. We then set $l_k = l$, and define Z_k to be the union of all cubes in $\mathcal{B}_{l_k}^{dn}(Y_{m_k})$. \square

We now construct X by avoiding the various discretizations of Z at each scale. The aim is to find a nested decreasing family of discretized sets $\{X_k\}$ with $X = \bigcap X_k$. One condition guaranteeing that X avoids Z is that X_k^n is disjoint from *strongly non-diagonal* cubes in Z_k .

Lemma 3. *Let $Z \subset \mathbf{R}^{dn}$ and let $\{l_k\}$ be a sequence of lengths converging to zero. For each index k , let Z_k be a union of cubes in $\mathcal{B}_{l_k}^{dn}$, and suppose the sets $\{Z_k\}$ strongly cover Z . For each index k , let X_k be a union of cubes in $\mathcal{B}_{l_k}^d$. Suppose that for any k , X_k^n avoids strongly non-diagonal cubes in Z_k . If $X = \bigcap X_k$, then $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$.*

Proof. Let $z \in Z$ be a point with distinct coordinates z_1, \dots, z_n . Set

$$\Delta = \{(w_1, \dots, w_n) \in \mathbf{R}^{dn} : \text{there exists } i \neq j \text{ such that } w_i = w_j\}.$$

Then $d(\Delta, z) > 0$, where d is the Hausdorff distance between Δ and z . Since $\{Z_k\}$ strongly covers Z , there is a subsequence $\{k_m\}$ such that $z \in Z_{k_m}$ for any index m . For suitably large m , the sidelength l_k cube I in Z_{k_m} containing z is disjoint from Δ . But this means I is strongly non-diagonal, and so $z \notin X_{k_m}^n$. In particular, z is not an element of X^n . \square

We are now ready to construct the set X in Theorem 1. Let $l_0 = 1$ and $X_0 = [0, 1)^d$. For each $k \geq 1$, define $\varepsilon_k = c_0/k$, where we view $c_0 = (dn - \alpha)/4$ as a irrelevant constant small enough that $dn - \alpha - 2\varepsilon_k > 0$ for any k . We set

$$f_k(x) = \min \left(x^{k^2}, (x^{dn}/2)^{1/(dn-\alpha-\varepsilon_{k+1})}, (1/2A_x)^{d(n-1)/\varepsilon_{k+1}} \right).$$

Apply Lemma 2 to Z with this choice of $\{\varepsilon_k\}$ and $\{f_k\}$; let $\{l_k\}$ be the resulting sequence of dyadic lengths and let $\{Z_k\}$ be the resulting strong cover of Z . Observe that the definition of f_k implies that for any index k ,

$$l_{k+1} \leq l_k^{k^2}, \quad l_{k+1}^{dn-\alpha-\varepsilon_{k+1}} \leq l_k^{dn}/2, \quad \text{and} \quad 2A_{l_k} l_{k+1}^{\varepsilon_{k+1}/d(n-1)} \leq 1. \quad (3.1)$$

For each index $k \geq 1$, define $l = l_k$ and $s = l_{k+1}$. Observe that X_k is a non-empty union of cubes in \mathcal{B}_l^d ; that Z_{k+1} is a union of cubes in \mathcal{B}_s^{dn} , and that (3.1) implies

$$|Z_{k+1}| \leq l_{k+1}^{dn-\alpha-\varepsilon_{k+1}} \leq l_k^{dn}/2 = l^{dn}/2.$$

Setting $Z_s = Z_{k+1}$, we are therefore justified in applying Lemma 1. This produces a dyadic length r , which we denote by r_{k+1} . Assuming $\alpha \geq d$, by (2.1) and (3.1), we find

$$\begin{aligned} r_{k+1} &\leq \max(l_{k+1}, 2A_{l_k}|Z_{k+1}|^{1/d(n-1)}) \leq \max\left(l_{k+1}, 2A_{l_k}l_{k+1}^{(dn-\alpha-\varepsilon_{k+1})/d(n-1)}\right) \\ &\leq \max\left(l_{k+1}, l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)}\right) = l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)}. \end{aligned} \quad (3.2)$$

For this choice of r , we obtain a set $F \subset X_k$ which is a union of cubes in $\mathcal{B}_s^d(E)$ satisfying Properties (A), (B), and (C) from the lemma, and we define $X_{k+1} = F$. Property (A) implies X_{k+1} avoids strongly non-diagonal cubes in Z_{k+1} , so if we define $X = \bigcap X_k$, then Lemma 3 implies $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$.

Lemma 4.

$$\overline{\dim}_{\mathbf{M}}(X) \geq \frac{dn - \alpha}{n - 1}$$

Proof. Consider X as the limit of the sequence $\{X_k\}$. Since every cube in $\mathcal{B}_{l_{k+1}}^d(X_{k+1})$ intersects X , $\mathcal{B}_{l_{k+1}}^d(X_{k+1}) = \mathcal{B}_{l_{k+1}}^d(X)$. And so by Property (C) of Lemma 1, (3.1), and (3.2), we conclude

$$\begin{aligned} \frac{\log(\#\mathcal{B}_{l_{k+1}}^d(X))}{\log(1/l_{k+1})} &= \frac{\log(\#\mathcal{B}_{l_{k+1}}^d(X_{k+1}))}{\log(1/l_{k+1})} \geq \frac{\log((l_k/r_{k+1})^d \cdot \#\mathcal{B}_{l_k}^d(X_k))}{\log(1/l_{k+1})} \\ &= \frac{d \log(1/r_{k+1})}{\log(1/l_{k+1})} - \frac{d \log(1/l_k)}{\log(1/l_{k+1})} \\ &\geq \frac{d \log\left(1/l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)}\right)}{\log(1/l_{k+1})} - \frac{d \log(1/l_k)}{\log(1/l_k^{k^2})} \\ &= \frac{dn - \alpha - 2\varepsilon_{k+1}}{n - 1} - \frac{d}{k^2} \\ &= \frac{dn - \alpha}{n - 1} - o(1) \end{aligned}$$

Taking $k \rightarrow \infty$, we conclude $\overline{\dim}_{\mathbf{M}}(X) \geq (dn - \alpha)/(n - 1)$. □

4 Dimension Bounds

To complete the proof of Theorem 1, we must show that $\dim_{\mathbf{H}}(X) \geq \beta$, where

$$\beta = \frac{dn - \alpha}{n - 1}.$$

We begin with a rough outline of our proof strategy. Recall that from the previous section, we have a decreasing sequence of lengths $\{l_k\}$. The most convenient way to examine the

dimension of X at various scales is to use Frostman's lemma (see Definition (G)). We construct a probability measure μ supported on X such that for all $\varepsilon > 0$, for all dyadic lengths l , and for all $I \in \mathcal{B}_l^d$, $\mu(I) \lesssim_\varepsilon l^{\beta-\varepsilon}$. We begin by proving the bound $\mu(I) \lesssim l_k^{\beta-O(1/k)}$ when $I \in \mathcal{B}_{l_k}^d$, which we view as saying X looks like a set with dimension $\beta - O(1/k)$ at the lengths $\{l_k\}$. To obtain the complete dimension bound, it then suffices to interpolate to get an acceptable bound at all intermediate scales. In this construction, as in [5], the rapid decay of the lengths $\{l_k\}$ forced on us in the construction means interpolation poses a significant difficulty. We avoid this difficulty because of the uniform way that we have selected cubes in consecutive scales. This will imply that between the scales l_k and r_{k+1} , the mass of μ distributes with similar properties to the full dimensional Lebesgue measure, which makes interpolation easy.

We now define the measure μ , by first defining it as a premeasure on $\bigcup_{i=1}^\infty \mathcal{B}_{l_i}^d[0,1]^d$. We initially set $\mu([0,1]^d) = 1$. Given $I \in \mathcal{B}_{l_k}^d$, we find a parent cube $I' \in \mathcal{B}_{l_{k-1}}^d$ with $I \subset I'$. If $I \subset X_k$, then we set

$$\mu(I) = \frac{\mu(I')}{\#\mathcal{B}_{l_k}^d(X_k \cap I')}. \quad (4.1)$$

Otherwise, if $I \not\subset X_k$, we set $\mu(I) = 0$. Notice

$$\mu(I') = \sum_{I \in \mathcal{B}_{l_k}^d(X_k \cap I')} \frac{\mu(I')}{\#\mathcal{B}_{l_k}^d(X_k \cap I')} = \sum_{I \in \mathcal{B}_{l_k}^d(I')} \mu(I).$$

Thus mass is maintained at each stage of the construction, and then Proposition 1.7 of [4] implies μ is a premeasure, and thus extends to a measure on the entire Borel sigma algebra. The remainder of this section is devoted to showing that μ is a Frostman measure of dimension $\beta - \varepsilon$ for any $\varepsilon > 0$.

Lemma 5. *If $I \in \mathcal{B}_{l_k}^d$, then*

$$\mu(I) \leq 2^k \left[\frac{r_k \cdots r_1}{l_{k-1} \cdots l_1} \right]^d. \quad (4.2)$$

Proof. We prove the theorem inductively on k . For $k = 0$, the theorem is obvious, because $\mu(I) \leq 1$. For the purposes of induction, let $I \in \mathcal{B}_{l_k}^d$, together with a parent cube $I' \in \mathcal{B}_{l_{k-1}}^d$ with $I \subset I'$. If $\mu(I) > 0$, $I \subset X_k$, so $I' \subset X_{k-1}$. Because X_k was obtained from X_{k-1} via an application of Lemma 1, Property (C) of that lemma states that $\#\mathcal{B}_{l_k}^d(X_k \cap I) \geq (l_{k-1}/r_k)^d/2$, so together with the inductive hypothesis and (4.1), we conclude

$$\mu(I) = \frac{\mu(I')}{\#\mathcal{B}_{l_k}^d(X_k \cap I)} \leq \frac{\mu(I')}{(l_{k-1}/r_k)^d/2} \leq \frac{2^{k-1} \cdot \left[\frac{r_{k-1} \cdots r_1}{l_{k-2} \cdots l_1} \right]^d}{(l_{k-1}/r_k)^d/2} = 2^k \left[\frac{r_k \cdots r_1}{l_{k-1} \cdots l_1} \right]^d. \quad \square$$

Treating all parameters in (4.2) which depend on indices smaller than k as essentially constant, and using (3.2), we 'conclude' that

$$\mu(I) \lesssim r_k^d \lesssim l_k^{\beta-2\varepsilon_k/(n-1)} = l_k^{\beta-O(1/k)}.$$

The bounds in (3.1) imply l_k decays very rapidly, which enables us to ignore quantities depending on previous indices, and obtain a true inequality.

Corollary 1. *There exists $c > 0$ such that for all $I \in \mathcal{B}_{l_k}^d$, $\mu(I) \lesssim l_k^{\beta-c/k}$.*

Proof. Given $\varepsilon > 0$, Lemma 5, Equation (3.2), the inequality $l_k \leq l_{k-1}^{(k-1)^2}$ from (3.1), and the fact that there exists a constant c_0 such that $\varepsilon_{k+1} = c_0/k$, we find

$$\begin{aligned} \mu(I) &\leq 2^k \left[\frac{r_k \cdots r_1}{l_{k-1} \cdots l_1} \right]^d \leq \left(\frac{2^k}{l_{k-1}^d \cdots l_1^d} \right) l_k^{\beta-\varepsilon_k/(n-1)} = \left(\frac{2^k l_k^{2d/k}}{l_{k-1}^{d(k-1)}} \right) l_k^{\beta-\varepsilon_k/(n-1)-2d/k} \\ &\leq \left(2^k l_{k-1}^{(2d/k)(k-1)^2-d(k-1)} \right) l_k^{\beta-(c_0/(n-1)+2d)/k} = o \left(l_k^{\beta-(c_0/(n-1)+2d)/k} \right), \end{aligned}$$

so we can then set $c = c_0/(n-1) + 2d$. \square

Corollary 3 gives a clean expression of the β dimensional behaviour of μ at discrete scales. To obtain a Frostman measure bound at *all* scales, we need to apply a covering argument. This is where the uniform mass assignment technique comes into play. Because μ behaves like a full dimensional set between the scales l_k and r_{k+1} , we won't be penalized for making the gap between l_k and r_{k+1} arbitrarily large. This is essential to our argument, because l_k decays faster than 2^{-k^m} for any $m > 0$.

Lemma 6. *If l is dyadic and $I \in \mathcal{B}_l^d$, then $\mu(I) \lesssim_k l^{\beta-c/k}$ for each integer k .*

Proof. We begin by assuming $l \leq l_k$. To bound $\mu(I)$, we apply a covering argument, which breaks into cases depending on the size of l in proportion to the scales l_k and r_k :

- If $r_{k+1} \leq l \leq l_k$, we can cover I by $(l/r_{k+1})^d$ cubes in $\mathcal{B}_{r_{k+1}}^d$. Because of Property (B) and (C) of Lemma 1, we know that the mass of each cube in $\mathcal{B}_{r_{k+1}}^d$ is bounded by at most $2(r_{k+1}/l_{k+1})^d$ times the mass of a cube in $\mathcal{B}_{l_k}^d$. Thus

$$\mu(I) \lesssim (l/r_{k+1})^d (2(r_{k+1}/l_k)^d) l_k^{\beta-c/k} \leq 2l^d/l_k^{d-\beta+c/k} \leq 2l^{\beta-c/k},$$

where we used the fact that $d - \beta + c/k \geq 0$, so $l_k^{d-\beta+c/k} \geq l^{d-\beta+c/k}$.

- If $l_{k+1} \leq l \leq r_{k+1}$, we can cover I by a single cube in $\mathcal{B}_{r_{k+1}}^d$. Because of Property (B) of Lemma 1, each cube in $\mathcal{B}_{r_{k+1}}^d$ contains at most one cube of $\mathcal{B}_{l_{k+1}}^d(X_{k+1})$, so

$$\mu(I) \lesssim l_{k+1}^{\beta-c/k} \leq l^{\beta-c/k}.$$

- If $l \leq l_{k+1}$, there certainly exists m such that $l_{m+1} \leq l \leq l_m$, and one of the previous cases yields that $\mu(I) \lesssim l^{\beta-c/m} \leq l^{\beta-c/k}$.

If $l \geq l_k$, then $\mu(I) \leq 1 \lesssim_k l_k^{\beta-c/k} \leq l^{\beta-c/k}$, so $\mu(I) \lesssim_k l^{\beta-c/k}$ for arbitrary dyadic l . \square

Applying Frostman's lemma to Lemma 6 gives $\dim_{\mathbf{H}}(X) \geq \beta - c/k$ for each $k > 0$. Taking $k \rightarrow \infty$ proves the needed dimension bound. Since we have already shown in Section 3 that $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$, this concludes the proof of Theorem 1.

5 Applications

As discussed in the introduction, Theorem 1 generalizes Theorems 1.1 and 1.2 from [5]. In this section, we present two applications of Theorem 1 in settings where previous methods cannot obtain any results.

Theorem 2 (Sum-sets avoiding specified sets). *Let $Y \subset \mathbf{R}^d$ be a countable union of sets of Minkowski dimension at most α . Then there exists a set $X \subset \mathbf{R}^d$ with Hausdorff dimension at least $1 - \alpha$ such that $X + X$ is disjoint from Y .*

Proof. Define $Z = Z_1 \cup Z_2$, where

$$Z_1 = \{(x, y) : x + y \in Y\} \quad \text{and} \quad Z_2 = \{(x, y) : y \in Y/2\}.$$

Since Y is a countable union of sets of Minkowski dimension at most α , Z is a countable union of sets with lower Minkowski dimension at most $1 + \alpha$. Applying Theorem 1 with $d = 1$, $n = 2$, giving a set $X \subset \mathbf{R}^d$ with Hausdorff dimension $1 - \alpha$ avoiding Z . Since X avoids Z_1 , whenever $x, y \in X$ are distinct, $x + y \notin Y$. Since X avoids Z_2 , $X \cap (Y/2) = \emptyset$, and thus for any $x \in X$, $x + x \notin Y$. Thus $X + X$ is disjoint from Y . \square

Remark. *One weakness of our result is that as the number of variables n increases, the dimension of X tends to zero. If we try and make the n -fold sum $X + \dots + X$ disjoint from Y , current techniques only yield a set of dimension $(1 - \alpha)/(n - 1)$. We have ideas on how to improve our main result when Z is ‘flat’, in addition to being low dimension, which will enable us to remove the dependence of $\dim_{\mathbf{H}}(X)$ on n . In particular, we expect to be able to construct a set X of dimension $1 - \alpha$, such that X is disjoint from Y , and X is closed under addition, and multiplication by rational numbers. In particular, given a \mathbf{Q} subspace V of \mathbf{R}^d with dimension α , we can always find a ‘complementary’ \mathbf{Q} vector space W with complementary fractional dimension $d - \alpha$ such that $V \cap W = (0)$.*

One of the most interesting uses of our method is to construct subsets of fractals avoiding patterns. In [5], Fraser and the second author show that if γ is a C^2 curve with non-vanishing curvature, then there exists a set $E \subset \gamma$ of Hausdorff dimension $1/2$ that does not contain isocles triangles. Our method can extend this result from the case of curves to more general sets, which gives an example of the flexibility of our method. For simplicity, we stick to an analysis of planar sets.

Theorem 3 (Restricted sets avoiding isocles triangles). *Let $Y \subset \mathbf{R}^2$ and let $\pi : \mathbf{R}^2 \rightarrow \mathbf{R}$ be an orthogonal projection such that $\pi(Y)$ has non-empty interior. Let d be an arbitrary metric on \mathbf{R}^2 . Suppose that*

$$Z_0 = \{(y_1, y_2, y_3) \in Y^3 : d(y_1, y_2) = d(y_1, y_3)\}$$

is the countable union of sets with lower Minkowski dimension at most α , for $\varepsilon \geq 0$. Then there exists a set $X \subset Y$ with dimension at least $(3 - \alpha)/2$ so that no triple of points $(x_1, x_2, x_3) \in X^3$ form the vertices of an isocles triangle.

Proof. Without loss of generality, by translation and rescaling, we may assume $\pi(Y)$ contains $[0, 1)$. Form the set

$$Z = \pi(Z_0) = \{(\pi(y_1), \pi(y_2), \pi(y_3)) : y \in Z_0\}$$

Then Z is the projection of an α dimensional set, and therefore has dimension at most α . Applying Theorem 1 with $d = 1$ and $n = 3$, we construct a set $X_0 \subset [0, 1)$ with Hausdorff dimension at least $(3 - \alpha)/2$ such that for any distinct $x_1, x_2, x_3 \in X_0$, $(x_1, x_2, x_3) \notin Z$. Thus if we form a set X by picking, from each $x \in X_0$, a single element of $\pi^{-1}(x)$, then X avoids isosceles triangles, and has Hausdorff dimension at least as large as X_0 . \square

To see that Theorem 3 indeed generalizes the result of Fraser and the second author, observe that if d is the Euclidean metric, then for every pair of points $x, y \in \mathbf{R}^2$, the set

$$\{z \in \mathbf{R}^2 : d(x, z) = d(y, z)\}$$

is the perpendicular bisector B_{xy} of x and y . If γ is a compact portion of a smooth curve with non-vanishing curvature, then the number of points in $\gamma \cap B_{xy}$ is bounded independently of x and y . Thus the set Z_0 in the statement of Theorem 3 has Minkowski dimension at most 2, and we can find a set with Hausdorff dimension $(3 - 2)/2 = 1/2$ on the curve avoiding isosceles triangles.

Results about slice of measures, such as those detailed in Chapter 6 of [11], show that for any one dimensional set Y , for almost every line L , $L \cap Y$ consists of a finite collection of points. This suggests that if Y is any set with fractional dimension one, then Z_0 has dimension at most 2. This implies that we can find a subset of Y with dimension $1/2$ avoiding curves. We are unsure if this is true for every set with dimension one, but we provide two examples suggesting this is true for a generic set. The first result shows that for any $\varepsilon > 0$, there is an infinite family of Cantor-type sets with dimension $1 + \varepsilon$ such that Z_0 has dimension at most $2 + \varepsilon$. The second shows that for any rectifiable curve, Z_0 has dimension at most 2. Thus Theorem 3 can be applied in settings where Y is incredibly ‘rough’, i.e. totally disconnected.

To study the first example, we consider a probabilistic model for a Cantor-type set which almost surely has the required properties. This model is obtained by considering a nested decreasing family of discretized random sets $\{C_k\}$, with each C_k a union of sidelength $1/2^k$ squares. First, we fix $p \in [0, 1]$. Then we set $C_0 = [0, 1]^2$. To construct C_{k+1} , we split each sidelength $1/2^k$ cube in C_k into four sidelength $1/2^{k+1}$ squares, and keep each square in C_{k+1} with probability p . To study this model, we employ some results about tail bounds and asymptotics for branching processes.

Lemma 7. *If $p > 1/4$, then with non-zero probability, $\dim_{\mathbf{M}}(C) = 2 - \log_2(1/p)$.*

Proof. Let $p > 1/4$. For each k , let Z_k denote the number of sidelength $1/2^k$ cubes in $[0, 1]^2$. Then the sequence $\{Z_k\}$ is a branching process, where each cube can produce between zero and four subcubes, with each of these four cubes kept with probability p . Thus $\mathbf{E}[Z_{k+1}|Z_k] = (4p)Z_k$. Since $4p > 1$, A simple calculation, summarized in Theorem 8.1 of [6], shows that the process $W_k = Z_k/(4p)^k$ is an L^2 bounded martingale, and so there exists a random variable W such that $W_k \rightarrow W$ almost surely, and $\mathbf{E}(W|W_k) = W_k$ for all k . Whenever W

is non-zero,

$$\dim_{\mathbf{M}}(C) = \lim_{k \rightarrow \infty} \frac{\log Z_k}{k \log 2} = \lim_{k \rightarrow \infty} \frac{\log(W_k/W) + \log(W(4p)^k)}{k \log 2} = 2 - \log_2(1/p).$$

Since $\mathbf{E}(W) = \mathbf{E}(\mathbf{E}(W|W_0)) = \mathbf{E}(W_0) = 1$, W is non-zero with positive probability. \square

Similar asymptotics for branching processes show that the set Z_0 associated with C as in Theorem 3 almost surely has Minkowski dimension $3 - \alpha$. We do this first by proving a supplementary result.

Lemma 8. *Let $\{Z_k\}$ be a supercritical branching process with extinction probability q . If we set $\tilde{M} = q + M$, then there exists small positive constants λ and ε , depending only on the offspring law of $\{Z_k\}$, such that $\mathbf{P}(Z_k \geq k\tilde{M}^k) \lesssim \exp(-\lambda k^{1+\varepsilon})$.*

Proof. Let q_0, q_1, \dots, q_M denote the offspring law for the branching process, so $q = q_0$. Then there exists a grid of i.i.d discrete random variables X_{ij} with $\mathbf{P}(X_{ij} = k) = q_k$ such that

$$Z_{k+1} = \sum_{j=1}^{Z_k} X_{ij}.$$

Now consider the branching process $\{\tilde{Z}_k\}$ defined by setting $\tilde{Z}_0 = 1$, and

$$\tilde{Z}_{k+1} = \sum_{j=1}^{\tilde{Z}_k} \max(X_{ij}, 1).$$

We find $Z_k \leq \tilde{Z}_k$, and if $\tilde{M} = q + M$, then

$$\mathbf{E}(\tilde{Z}_{k+1}|\tilde{Z}_k) = \left(q + \sum_{k=1}^N kq_k \right) \tilde{Z}_k = (q + M)\tilde{Z}_k = \tilde{M}\tilde{Z}_k.$$

Most importantly for our purposes, $\{\tilde{Z}_k\}$ has zero chance of extinction. Theorem 5 of [1] implies that for such a supercritical branching process, there exists $\lambda > 0$ depending only on the offspring distribution of \tilde{Z} , such that if $\tilde{W}_k = \tilde{Z}_k/\tilde{M}^k$, and $\tilde{W} = \lim \tilde{W}_k$, then

$$\mathbf{P}(|\tilde{W} - \tilde{W}_k| \geq t) \lesssim \exp\left(-\lambda t^{2/3} \tilde{M}^{k/3}\right)$$

In particular, this means

$$\begin{aligned} \mathbf{P}\left(Z_k \geq (1 + \tilde{W})\tilde{M}^k\right) &\leq \mathbf{P}(\tilde{Z}_k \geq (1 + \tilde{W})\tilde{M}^k) \\ &\leq \mathbf{P}\left(|\tilde{W}\tilde{M}^k - \tilde{Z}_k| \geq \tilde{M}^k\right) \\ &= \mathbf{P}\left(|\tilde{W} - \tilde{W}_k| \geq 1\right) \lesssim \exp\left(-\lambda \tilde{M}^{k/3}\right), \end{aligned}$$

Theorem 2 of [3] implies that as $t \rightarrow \infty$, there are constants C and $\varepsilon > 0$ depending only on the offspring distribution of \tilde{Z} such that

$$-\log \mathbf{P}(\tilde{W} \geq t) \geq (C + o(1))t^{1+\varepsilon}$$

In particular, there exists a small constant λ such that

$$\mathbf{P}(1 + \tilde{W} \geq k) \leq \exp(-\lambda k^{1+\varepsilon})$$

Applying a union bound gives

$$\begin{aligned} \mathbf{P}(Z_k \geq k\tilde{M}^k) &\leq \mathbf{P}(Z_k \geq (1 + \tilde{W})\tilde{M}^k) + \mathbf{P}(1 + \tilde{W} \geq k) \\ &\lesssim \exp(-\lambda \tilde{M}^{k/3}) + \exp(-\lambda k^{1+\varepsilon}) \lesssim \exp(-\lambda k^{1+\varepsilon}). \end{aligned} \quad \square$$

For a set E , let $E_\delta = \{x : d(x, E) \leq \delta\}$ denote the δ thickened version of E .

Lemma 9. *There exists a constant B such that for any k , we can find lines $L_{k,1}, \dots, L_{k,M}$ with $M \leq B \cdot 8^{kN}$ such that for any line L , there exists i such that $[0, 1]^2 \cap L_{1/2^{kN+1}} \subset (L_{k,i})_{2^{-kN}}$.*

Proof. For each $(x, \theta) \in [0, 1]^2 \times [0, 1]$, let

$$L^{\theta,x} = \{x + te^{2\pi i\theta} : t \in \mathbf{R}\}$$

Note that

$$L^{\theta,x} \cap [0, 1]^2 \subset \{x + te^{2\pi i\theta} : t \in [-2, 2]\}$$

Any line intersecting $[0, 1]^2$ is equal to $L^{\theta,x}$ for some θ and some x . For any k , we consider the set of $8^5 \cdot 8^{kN}$ points $E = (\mathbf{Z}/2^{kN+5})^3 \cap [0, 1]^3$. For any (x_1, θ_1) and (x_2, θ_2) , and $t \in [-2, 2]$, we calculate that

$$|(x_1 + te^{2\pi i\theta_1}) - (x_2 + te^{2\pi i\theta_2})| \leq |x_1 - x_2| + 2\pi t|\theta_1 - \theta_2| \leq |x_1 - x_2| + 4\pi|\theta_1 - \theta_2|$$

Thus $L^{\theta_1,x_1} \cap [0, 1]^2$ is contained in the $|x_1 - x_2| + 4\pi|\theta_1 - \theta_2|$ thickening of L^{θ_2,x_2} . For any line $L^{\theta,x}$, there exists $(\theta_0, x_0) \in E$ such that $|\theta - \theta_0| \leq 1/2^{kN+2}$ and $|x - x_0| \leq \sqrt{2}/2^{kN+5}$, and so $L^{\theta,x} \cap [0, 1]^2$ is contained in the $\sqrt{2}/2^{kN+5} + 4\pi/2^{kN+5} \leq 14/2^{kN+5} \leq 1/2^{kN+1}$, and thus $L_{1/2^{kN+1}}^{\theta,x} \cap [0, 1]^2 \subset L_{1/2^{kN}}$. Thus we may set $B = 8^5$, completing the proof. \square

Lemma 10. *Fix N . If $p > 1/2$, then almost surely, there exists a value k_0 such that if $k \geq k_0$, and L is any line, then*

$$\# \mathcal{B}_{2^{-kN}}^2(L_{2^{-kN}}) \leq k \cdot 10^k (2p)^{kN}$$

Proof. Fix N , and for any index k let $\delta_k = 1/2^{Nk}$. Given a line L , let I be a sidelength δ_k square intersecting the δ_k thickened line L_{δ_k} . Then L passes from one edge of I to another edge, and so if we split I into 4^N sidelength δ_{k+1} squares, then $L_{\delta_{k+1}}$ intersects at most $5 \cdot 2^N$ of these boxes. Conditioned on I being contained in C_{kN} , each of these subboxes occurs in $C_{(k+1)N}$ with probability p^N . We let $Z_k(L)$ denote the number of sidelength $1/2^{Nk}$ boxes

in C_{Nk} intersecting L_{δ_k} . We now find $\tilde{Z}_k(L) \geq Z_k(L)$ by adding *exactly* $5 \cdot 2^N$ potential subboxes of each box at each subsequent stage, then $\tilde{Z}_k(L)$ is a branching process, whose offspring distribution is independent of L . Since each subbox is added with probability p^N , the extinction probability of \tilde{Z}_k is $(1-p^N)^{5 \cdot 2^N}$. Also, $\mathbf{E}(\tilde{Z}_{k+1}(L)|\tilde{Z}_k(L)) = 5 \cdot (2p)^N \tilde{Z}_k$. Setting $M = 5(2p)^N$ and $q = (1-p^N)^{5 \cdot 2^N}$, Lemma 8 shows that there exists positive constants λ and ε , independent of L , such that if $\tilde{M} = M + q$,

$$\mathbf{P}(\tilde{Z}_k(L) \geq k\tilde{M}^k) \lesssim \exp(-\lambda k^{1+\varepsilon}).$$

Elementary bounds on the logarithm show that

$$5 \cdot 2^N \log(1-p^N) \leq \frac{-5(2p)^N}{2-p^N} \leq -5(2p)^N$$

so if $p > 1/2$,

$$q = (1-p^N)^{5 \cdot 2^N} \leq e^{-5(2p)^N} \leq 1 \leq 5(2p)^N$$

and this implies $\tilde{M} \leq 10(2p)^N$, so we have shown

$$\mathbf{P}(\tilde{Z}_k(L) \geq k \cdot 10^k (2p)^{kN}) \lesssim \exp(-\lambda k^{1+\varepsilon}).$$

This gives an upper bound on $\tilde{Z}_k(L)$ up to a superexponentially decaying term in k .

For each k , Lemma 9 shows we can find lines $L_{k,1}, \dots, L_{k,M}$, with $M \leq B \cdot 8^{kN}$ such that for any line L , there exists i such that $[0, 1]^2 \cap L_{\delta_k/2} \subset (L_{k,i})_{\delta_k}$. Applying a union bound, we find that

$$\mathbf{P}(\text{there is } i \text{ such that } Z_k(L_{k,i}) \geq k \cdot 10^k (2p)^{kN}) \lesssim B \cdot 8^{kN} \exp(-ck^\lambda)$$

But since $\lambda > 1$, we therefore find

$$\sum_{k=1}^{\infty} \mathbf{P}(\text{there is } i \text{ such that } Z_k(L_{k,i}) \geq k \cdot 10^k (2p)^{kN}) < \infty$$

Applying the Borel-Cantelli lemma, we conclude that almost surely, it is eventually true for sufficiently large k that $Z_k(L_{k,i}) \leq k \cdot 10^k (2p)^{kN}$ for all i . In particular, the number of cubes intersecting $L_{\delta_k/2}$ for any line L is upper bounded by $k \cdot 10^k (2p)^{kN}$. \square

Lemma 11. *There exists a constant A such that if $I, J \in \mathcal{B}_{1/2^{kN}}^2[0, 1]^2$, with $d(I, J) \geq A/2^{kN}$, then*

$$\bigcup \{L_{xy} : x \in I, y \in J\} \subset L_{1/2^{kN+1}}$$

where we recall that L_{xy} is the bisector of the points x and y .

Proof. Denote the bottom left vertices of the boxes I and J by $(N_I, M_I) \cdot 1/2^{kN}$ and $(N_J, M_J) \cdot 1/2^{kN}$, with $0 \leq N_I, M_I, N_J, M_J \leq 2^{kN}$. Let $\Delta_N = |N_I - N_J|$ and $\Delta_M = |M_I - M_J|$. We may assume for simplicity that $N_I \leq N_J$ and $M_I \leq M_J$. Since $d(I, J) \geq A/2^{kN}$,

$$(\Delta_N^2 + \Delta_M^2)^{1/2} \geq A$$

If we let θ_1 and θ_2 denote the minimal and maximal angle between the lines connecting pairs of points in I and J and the horizontal line, then

$$\begin{aligned}\sin(\theta_2 - \theta_1) &= \sin(\theta_2) \cos(\theta_1) - \sin(\theta_1) \cos(\theta_2) \\ &= \frac{(\Delta_M + 1)(\Delta_N + 1)}{\Delta_N^2 + \Delta_M^2} - \frac{\Delta_N \Delta_M}{\Delta_N^2 + \Delta_M^2} \\ &\leq \frac{\Delta_N + \Delta_M + 1}{\Delta_N^2 + \Delta_M^2} \leq 1/A\end{aligned}$$

Thus $\theta_2 - \theta_1 \leq 2/A$. Thus all the bisectors are contained in a

$$\sqrt{2}/2^{kN} + 4\pi(2/A) \leq 1/2^{kN}$$

□

Theorem 4. *If $p > 1/2$, then almost surely, the set Z_0 associated with C has lower Minkowski dimension at most $5 - 3\log_2(1/p)$.*

Proof. Almost surely, Lemma 7 shows that for sufficiently large k we can cover C_{kN} by $O(2^{\alpha kN})$ boxes $I_1, \dots, I_M \in \mathcal{B}_{2^{-kN}}^2$, with $\alpha = 2 - \log_2(1/p)$. Lemma 10 shows that if k is sufficiently large, then the $1/2^{kN+1}$ thickened line L intersects at most $k \cdot 10^k (2p)^{kN}$ squares in $\mathcal{B}_{2^{-kN}}^2$. There exists a constant A such that for any i and j , if $d(I_i, I_j) > A2^{-kN}$, then there exists a line L_{ij} such that as x ranges over all points in I_i , and y over all points in I_j ,

$$\bigcup_{x,y} B_{xy} \cap [0, 1]^2 \subset (L_{ij})_{\delta_k/2}$$

This means that $Z_0 \cap (I_i \times I_j \times [0, 1]^2) \subset I_i \times I_j \times (L_{ij})_{\delta_k/2}$, and the last paragraph implies that $(L_{ij})_{\delta_k/2}$ is covered by $k \cdot 10^k (2p)^{kN}$ cubes in $\mathcal{B}_{l_k}^2[0, 1]^2$. On the other hand, if $d(I_i, I_j) < A\delta_k$, we can apply the obvious bound that $Z_0 \cap (I_i \times I_j \times [0, 1]^2) \subset I_i \times I_j \times C_{kN}$, and C_{kN} is coverable by $O(2^{\alpha kN})$ boxes. Thus we conclude that almost surely, for sufficiently large k ,

$$\begin{aligned}\# \mathcal{B}_{\delta_k}^6(Z_0) &= \sum_{i,j} \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times [0, 1]^2)) \\ &\leq \sum_i \left(\sum_{d(I_i, I_j) \leq A\delta_k} \# \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times (L_{ij})_{\delta_k/2})) \right) \\ &\quad + \left(\sum_{d(I_i, I_j) > A\delta_k} \# \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times C_{kN})) \right) \\ &\leq \sum_i \left(\sum_{d(I_i, I_j) \leq A\delta_k} \# \mathcal{B}_{\delta_k}^2((L_{ij})_{\delta_k/2}) \right) + \left(\sum_{d(I_i, I_j) > A\delta_k} \# \mathcal{B}_{\delta_k}^2(C_{kN}) \right) \\ &\lesssim \sum_i 2^{\alpha kN} \cdot k 10^k (2p)^{kN} + 2^{\alpha kN} \lesssim 2^{2\alpha kN} k 10^k (2p)^{kN}\end{aligned}$$

Thus we conclude that almost surely,

$$\begin{aligned}
\dim_{\mathbf{M}}(Z_0) &= \lim_{k \rightarrow \infty} \frac{\log(\#\mathcal{B}_{\delta_k}^6(Z_0))}{\log(1/\delta_k)} \\
&\leq \lim_{k \rightarrow \infty} \frac{\log(2^{2\alpha k N} k 10^k (2p)^{kN}) + O(1)}{\log(2^{kN})} \\
&= \lim_{k \rightarrow \infty} \frac{2\alpha k N \log(2) + k \log 10 + kN \log(2p)}{kN \log(2)} \\
&= 2\alpha + 3/N + 1 - \log_2(1/p) \\
&= 5 - 3 \log_2(1/p) + 3/N
\end{aligned}$$

Taking $N \rightarrow \infty$, we obtain the required result. \square

TODO: does a projection of C have non-empty interior almost surely?

Corollary 2. *If $p = 1/2^{1-\varepsilon}$, then conditioned on C having Minkowski dimension $1 + \varepsilon$, we can find a subset X of C avoiding isocoles triangles with $\dim_{\mathbf{H}}(X) \geq 1/2 - (3/2)\varepsilon$.*

6 Relation to Literature, and Future Work

Our result is part of a growing body of work finding general methods to find sets avoiding patterns. The main focus of this section is comparing our method to the two other major results in the literature. [5] constructs sets with dimension $k/(n-1)$ avoiding the zero sets of rank k C^1 functions. In [10], sets of dimension d/l are constructed avoiding a degree l algebraic hypersurface specified by a polynomial with rational coefficients.

We can view our result as a robust version of Pramanik and Fraser's result. Indeed, if we try and avoid the zero set of a C^1 rank k function, then we are really avoiding a dimension $dn - k$ dimensional manifold. Our method gives a dimension

$$\frac{dn - (dn - k)}{n - 1} = \frac{k}{n - 1}$$

set, which is exactly the result obtained in [5].

That our result generalizes [5] should be expected because the technical skeleton of our construction is heavily modeled after their construction technique. Their result also reduces the problem to a discrete avoidance problem. But they *deterministically* select a particular side length S cube in every side length R cube. For arbitrary Z , this selection procedure can easily be exploited for a particularly nasty Z , so their method must rely on smoothness in order to ensure some cubes are selected at each stage. Our discrete avoidance technique was motivated by other combinatorial optimization problems, where adding a random quantity prevents inefficient selections from being made in expectation. This allows us to rely purely on combinatorial calculations, rather than employing smoothness, and greatly increases the applicability of the sets Z we can apply our method to. Furthermore, it shows that the underlying problem is robust to changes in dimension; slightly 'thickening' Z only slightly perturbs the dimension of X .

One useful technique in [5], and its predecessor [8], is the use of a Cantor set construction ‘with memory’; a queue in the construction algorithm for their sets allows storage of particular discrete versions of the problem to be stored, and then retrieved at a much later stage of the construction process. This enables them to ‘separate’ variables in the discrete version of the problem, i.e. instead of forming a single set F from a set E , they form n sets F_1, \dots, F_n from disjoint sets E_1, \dots, E_n . The fact that our result is more general, yet does not rely on this technique is an interesting anomaly. An obvious advantage is that the description of the technique is much more simple. But an additional advantage is that we can attack ‘one scale’ of the problem at a time, rather than having to rely on stored memory from a vast number of steps before the current one. We believe that we can exploit the single scale approach to the problem to generalize our theorem to a much wider family of ‘dimension α ’ sets Z , which we plan to discuss in a later paper.

As a generalization of the result in [5], our result has the same issues when compared to the result of [10]. When the parameter n is large, the dimension of our result suffers greatly, as with the n fold sum application in the last section. Furthermore, our result can’t even beat trivial results if Z is almost full dimensional, as the next example shows.

Example. Consider an α dimensional set of angles Y , and try and find $X \subset \mathbf{R}^2$ such that the angle formed from any collection of three points in X avoids Y . If we form the set

$$Z = \left\{ (x, y, z) : \text{There is } \theta \in Y \text{ such that } \frac{(x - y) \cdot (x - z)}{|x - y||x - z|} = \cos \theta \right\}$$

Then we can find X avoiding Z . But one calculates that Z has dimension $3d + \alpha - 1$, which means X has dimension $(1 - \alpha)/2$. Provided the set of angles does not contain π , the trivial example of a straight line beats our result.

Nonetheless, we still believe our method is a useful inspiration for new techniques in the ‘high dimensional’ setting. Most prior literature studies sets Z only implicitly as zero sets of some regular function f . The features of the function f imply geometric features of Z , which are exploited by these results. But some geometric features are not obvious from the functional perspective; in particular, the fractional dimension of the zero set of f is not an obvious property to study. We believe obtaining methods by looking at the explicit geometric structure of Z should lead to new techniques in the field, and we already have several ideas in mind when Z has geometric structure in addition to a dimension bound, which we plan to publish in a later paper.

We can compare our randomized selection technique to a discrete phenomenon that has been recently noticed, for instance in [2]. There, certain combinatorial problems can be rephrased as abstract problems on hypergraphs, and one can then generalize the solutions of these problems using some strategy to improve the result where the hypergraph is sparse. Our result is a continuous analogue of this phenomenon, where sparsity is represented by the dimension of the set Z we are trying to avoid. One can even view Lemma 1 as a solution to a problem about independent sets in hypergraphs. In particular, we can form a hypergraph by taking the cubes $\mathcal{B}_s^d(E)$ as vertices, and adding an edge (I_1, \dots, I_n) between n distinct cubes $I_k \in \mathcal{B}_s^d(E)$ if $I_1 \times \dots \times I_n$ intersects Z_s . An independent set of cubes in this hypergraph corresponds precisely to a set F with F^n disjoint except on a discretization of the diagonal.

And so Lemma 1 really just finds a ‘uniformly chosen’ independent set in a sparse graph. Thus we really just applied the discrete phenomenon at many scales to obtain a continuous version of the phenomenon.

References

- [1] Krishna Athreya, *Large deviation rates for branching processes—i. single type case*, Ann. Appl. Probab. **4** (1994), 779–790.
- [2] József Balogh, Robert Morris, and Wojciech Samotij, *Independent sets in hypergraphs*, J. Amer. Math. Soc. **28** (2015), 669–709.
- [3] J.D. Biggins and N.H Bingham, *Large deviations in the supercritical branching process*, Adv. in Appl. Probab. **25**, 757–772.
- [4] Kenneth Falconer, *Fractal geometry: Mathematical foundations and applications*, Wiley, 2003.
- [5] Robert Fraser and Malabika Pramanik, *Large sets avoiding patterns*, Anal. PDE **11** (2018), 1083–1111.
- [6] Theodore Harris, *The theory of branching processes*, Springer-Verlag, 1963.
- [7] Nets Hawk Katz and Terrence Tao, *Some connections between Falconer’s distance set conjecture, and sets of Furstenburg type*, New York J. Math. **7** (2001), 149–187.
- [8] Tamás Keleti, *A 1-dimensional subset of the reals that intersects each of its translates in at most a single point*, Real Anal. Exchange **24** (1999), 843–845.
- [9] Péter Maga, *Full dimensional sets without given patterns*, Real Anal. Exchange **36** (2010), 79–90.
- [10] András Máthé, *Sets of large dimension not containing polynomial configurations*, Adv. Math. **316** (2017), 691–709.
- [11] Pertti Mattila, *Fourier analysis and Hausdorff dimension*, Cambridge University Press, 2015.