

Analysis SEP Problems & Solutions

Max Bacharach and Jacob Denson

July-August 2022

Contents

1	Basic Analysis Notes	3
2	Day 1: Basic Analysis	7
3	Day 2: Basic Analysis	17
4	Day 3: Basic Analysis	29
5	Measure Theory Notes	33
6	Day 4: Measure Theory	35
7	Functional Analysis Notes	39
8	Day 5: Functional Analysis	42
9	Day 6: Functional Analysis	49
10	Baire Category Notes	54
11	Day 7: Baire Category	56
12	Distribution Theory Notes	59
13	Day 8: Intro to Distribution	61
14	Fourier Analysis / Tempered Distribution Notes	72
15	Day 9: Fourier Analysis + Distribution Theory	74
16	Day 10: Bonus Questions	79
17	Day 11: Bonus Questions	87
18	Day 12: Bonus Questions	96
19	Day 14: Analysis Qualifying Exam 2022	103
20	Day 13: Interchanging limits and integrals + Arzela-Ascoli Theorem	108
	20.1 Limits and Integrals (DCT and friends)	108
	20.2 Arzela-Ascoli	110

21 Day 14: Misc. Topics	115
22 Day 15: Spring 2021 Final Qualifying Exam	121
23 Questions that need solutions	126

1 Basic Analysis Notes

Let's begin by reviewing the fundamental techniques to analysis. These include techniques from calculus, sequences and series, and integration techniques. In this section we give a brief summary of the techniques that are applied to the problems in this sheet, which we feel are representative of the majority of basic analysis questions on the exam.

- To show that a sequence $\{c_n\}$ converges to c as $n \rightarrow \infty$, it suffices to show that

$$\limsup_{n \rightarrow \infty} c_n \leq c$$

and

$$\liminf_{n \rightarrow \infty} c_n \geq c.$$

- The *ε of room* technique. To show that an inequality $a \leq b$ is true, it suffices to show that $a \leq b + \varepsilon$ is true for any $\varepsilon > 0$. This technique is often applied to show that two quantities are equal to one another, i.e. to show that $a = b$, it suffices to show that $a \leq b + \varepsilon$ and $b \leq a + \varepsilon$ for any $\varepsilon > 0$.
- A *dyadic decomposition* is often useful for rough estimates, i.e. breaking up regions of integration / regions of summation into powers of two, and applying crude estimates on these regions.
- The *Stone-Weirstrass theorem* tells us that in \mathbb{R}^n , the family of multi-variate polynomials on \mathbb{R}^n form a dense subspace of $C(K)$, for any compact set $K \subset \mathbb{R}^n$, where $C(K)$ is the Banach space given by the L^∞ norm. It follows that this class is also dense in most other function spaces encountered in analysis, e.g. the spaces $L^p(\mathbb{R}^n)$ for $1 \leq p < \infty$.
- Inequalities involving sine are often useful on the qualifying exam, for instance, the inequality

$$\sin x \geq x/2,$$

which holds for $x \in [0, \pi/2]$, and the inequality

$$x - x^2/2 \leq \sin x \leq x$$

holds for all $x \in \mathbb{R}$.

- The *mean value theorem*, which implies that for $a < b$, and for $f \in C^1[a, b]$, there exists $c \in [a, b]$ such that

$$f(b) - f(a) = f'(c)(b - a).$$

More generally, in higher dimensions, we have *Taylor's formula*, which allows us to write a function $f \in C^{k+1}(\mathbb{R}^n)$ as

$$f(x) = \sum_{|\alpha| \leq k} \frac{D^\alpha f(x_0)}{\alpha!} (x - x_0)^\alpha + \sum_{|\alpha| = k+1} R_\alpha(x) (x - x_0)^\alpha,$$

where

$$R_\alpha(x) = \frac{|\alpha|}{\alpha!} \int_0^1 (1-t)^{|\alpha|-1} (D^\alpha f)(x_0 + t(x - x_0)) dt = o(|x - x_0|^{k+1}).$$

- The Cauchy-Schwartz inequality implies that

$$|\sum a_n b_n| \leq \left(\sum |a_n|^2 \right)^{1/2} \left(\sum |b_n|^2 \right)^{1/2}$$

and more generally, we have *Hölder's inequality*

$$|\sum a_n b_n| \leq \left(\sum |a_n|^p \right)^{1/p} \left(\sum |b_n|^q \right)^{1/q}$$

where $1 \leq p, q \leq \infty$ and $1/p + 1/q = 1$. We also have similar inequalities for integrals.

- The *Arzela-Ascoli theorem*, which says that a set K of functions in $C[0, 1]$ is compact in the L^∞ norm if K is *uniformly bounded*, i.e. there is $M > 0$ such that $\|f\|_{L^\infty} \leq M$ for all $f \in K$, and *uniformly equicontinuous*, i.e. for any $\varepsilon > 0$, there is $\delta > 0$ such that if $|x - y| \leq \delta$ then for any $f \in K$, $|f(x) - f(y)| \leq \varepsilon$.

Sequences, Series, and Integrals

There are many questions on analysis qualifying exams asking to determine whether a given sequence

$$\sum_{n=1}^{\infty} a_n$$

converges. Depending on the sequence, one of various techniques may apply:

- If the sequence $\{a_n\}$ is a sequence of positive integer values, then the convergence of sequence is equivalent to the convergence of the sum

$$\sum_{k=1}^{\infty} \#(n : a_n = k) \cdot k.$$

Similarly, if the sequence takes values in a countable set $S \subset (0, \infty)$, then the convergence of the sequence is equivalent to the convergence of the sum

$$\sum_{s \in S} \#(n : a_n = s) \cdot s.$$

- If the sequence $\{a_n\}$ is positive and non-increasing, then one can apply *Cauchy's condensation theorem*, which says that the convergence of the series is equivalent to convergence of the sequence

$$\sum_{k=1}^{\infty} 2^k \cdot a_{2^k}.$$

This is often useful if the sequence $\{a_n\}$ grows somewhat logarithmically, since logarithms will be turned into linear terms in the sequence $\{a_{2^k}\}$.

- One can often convert sums into integrals, and vice versa. If the sequence $\{a_n\}$ is positive and non-increasing, and $a_n = a(n)$ for some function $a : [1, \infty) \rightarrow [0, \infty)$ then the convergence of the sum is equivalent to the convergence of the integral

$$\int_1^{\infty} a(x) dx.$$

One can often use a variant of this technique if the sum is not necessarily positive, or not necessarily increasing if one employs other tricks. For instance, provided that one can justify that

$$\sum_{k=1}^{\infty} \left(\sup_{0 \leq x \leq 1} |a(k+x) - a(k)| \right) < \infty,$$

then the convergence of the sum is also equivalent to convergence of the integral.

- If a function f is smooth (the derivative of f is well behaved), and a function g is oscillating very fast, then one can often understand an integral via an integral by parts, e.g. writing

$$\int_a^b f(x)g(x) dx = f(b)G(b) - \int_a^b f'(x)G(x) dx$$

where $G(x) = \int_a^x g(x) dx$ is likely small since g is oscillating fast. A similar phenomenon in the theory of series is a *summation by parts*, e.g. writing

$$\sum_{k=1}^n a_k b_k = a_n B_n + \sum_{k=1}^{n-1} (a_k - a_{k+1}) B_k,$$

which might help understand the sum if the discrete derivative $a_k - a_{k+1}$ is well behaved, and the sequence $\{b_k\}$ is oscillating fast.

- The limit comparison test, to determine if a series converges absolutely. If $\{a_n\}$ and $\{b_n\}$ are sequences of non-negative numbers, and

$$0 < \lim_{n \rightarrow \infty} \frac{a_n}{b_n} < \infty,$$

then $\sum a_n$ converges if and only if $\sum b_n$ converges.

- If a sequence $\{a_n\}$ *converges absolutely*, then any rearrangement of $\{a_n\}$ converges, and converges to the same value, i.e. for any bijection $\pi : \{1, 2, \dots\} \rightarrow \{1, 2, \dots\}$,

$$\sum a_n = \sum a_{\pi(n)}.$$

Day 1: Warm Up Question

1. (Fall 2016) For $n \geq 2$ an integer, define

$$F(n) = \max \{k \in \mathbb{Z} : 2^k/k \leq n\}.$$

Does the infinite series

$$\sum_{n=2}^{\infty} 2^{-F(n)}$$

converge or diverge?

Solution: The series diverges.

One solution follows by applying the intuitive identity

$$\sum_{n=2}^{\infty} 2^{-F(n)} = \sum_{k=1}^{\infty} 2^{-k} \cdot \#\{n \geq 1 : F(n) = k\},$$

which can also be interpreted as viewing the left hand side as a kind of discrete Lebesgue integral of a non-negative step function. But $F(n) = k$ holds if and only if

$$2^k/k \leq n < 2^{k+1}/(k+1),$$

and there are at most $2^{k+1}/(k+1) - 2^k/k \sim 2^k/k$ integers between $2^k/k$ and $2^{k+1}/(k+1)$, which implies that

$$\begin{aligned} \sum_{k=1}^{\infty} \#\{n \geq 1 : F(n) = k\} &\geq \sum_{k=1}^{\infty} 2^{-k} \cdot \left(\frac{2^{k+1}}{k+1} - \frac{2^k}{k} \right) \\ &\sim \sum_{k=1}^{\infty} 2^{-k} \cdot (2^k/k) \\ &= \sum_{k=1}^{\infty} 1/k = \infty. \end{aligned}$$

Thus we have proved the series diverges.

Another method to prove this result is to use the *Cauchy condensation theorem*, which states that for any non-increasing sequence of non-negative real numbers $\{a(n)\}$, the series

$$\sum_n a(n)$$

converges if and only if the sum

$$\sum_n 2^n a(2^n)$$

converges. The advantage of the theorem is that in many series, especially those involving *logarithms*, the values $\{a(2^n)\}$ will be easier to study than the sequence $\{a(n)\}$. Though $F(n)$ is not explicitly a logarithm, it behaves very similarly to a logarithm, i.e. we notice that there exists $C > 0$ such that

$$\log_2(n) \leq F(n) \leq \log_2(n) + \log_2 \log_2(n) + C.$$

This means that

$$\sum_n 2^{-F(n)} = \sum_n 2^n 2^{-F(2^n)} \geq \sum_n 2^n 2^{-n - \log_2 \log_2(n) - C} \gtrsim \sum_n \frac{1}{\log_2(n)},$$

and another application of the Cauchy condensation theorem shows that this series diverges.

2 Day 1: Basic Analysis

2. (Fall 2017) Let $\{a_n\}$ be a sequence of complex numbers and let

$$c_n = n^{-5} \sum_{k=1}^n k^4 a_k.$$

- (a) Prove or Disprove: If $\lim_{n \rightarrow \infty} a_n = a$ exists, then $\lim_{n \rightarrow \infty} c_n = c$ exists.

Solution: The result is *true*, so let us prove it. Roughly speaking, the value of c_n is the weighted average of the sequence $\{a_1, \dots, a_n\}$, heavily weighted in the favor of the later elements of the sequence. Thus it is intuitive that our proof should decompose the sum defining c_n into earlier terms, which must be shown to be negligible, and later terms, which are more well behaved because they are close to the value a . Indeed, fix $\varepsilon > 0$. Then there is k_0 such that $|a_k - a| < \varepsilon$ for $k > k_0$. But this means that

$$\begin{aligned} c_n &= n^{-5} \sum_{k=1}^n k^4 a_k \\ &= n^{-5} \sum_{k=1}^{k_0} k^4 a_k + \sum_{k=k_0+1}^n k^4 a_k \\ &= O_{k_0}(n^{-5}) + n^{-5} \sum_{k=k_0+1}^n k^4 a_k \\ &= O_{k_0}(n^{-5}) + n^{-5} \sum_{k=k_0+1}^n k^4 a + n^{-5} \sum_{k=k_0+1}^n k^4 (a_k - a) \\ &= O_{k_0}(n^{-5}) + a \cdot \left(\frac{1}{n^5} \sum_{k=k_0+1}^n k^4 \right) + O(\varepsilon n^{-5} \sum_{k=k_0+1}^n k^4) \\ &= O_{k_0}(n^{-5}) + a(1/5 + O_{k_0}(1/n)) + O(\varepsilon). \end{aligned}$$

But this calculation implies that there exists a positive integer n_0 and a constant $C > 0$ such that for $n \geq n_0$,

$$|c_n - a/5| \leq C\varepsilon.$$

Since ε was arbitrary, we have shown that $c_n \rightarrow a/5$.

A key step in the calculation of the limit above was that

$$\frac{1}{n^5} \sum_{k=k_0+1}^n k^4 = 1/5 + O_{k_0}(1/n).$$

One can prove this using the technique of *replacing sums with integrals*, which we are often more used to manipulating. Observe that

$$\frac{1}{n^5} \sum_{k=k_0}^n k^4 = \frac{1}{n^5} \sum_{k=k_0}^n \int_k^{k+1} k^4 dx \leq \frac{1}{n^5} \int_{k_0}^{n+1} x^4 dx = \frac{(n+1)^5 - k_0^5}{5n^5} = \frac{1}{5} + O_{k_0}(1/n).$$

A similar argument shows the opposite inequality, namely, that

$$\frac{1}{n^5} \sum_{k=k_0}^n k^4 \geq \frac{1}{5} + O_{k_0}(1/n),$$

which gives this result.

One can perform a similar calculation using *limsup* and *liminf* arguments, which yields a slightly different method and avoids some of the more quantitative calculations as above. It is intuitive that for large n we should have

$$c_n \approx \frac{a}{n^5} \sum_{k=1}^n k^4.$$

Indeed, we will show that

$$\limsup_{n \rightarrow \infty} \left| c_n - \frac{a}{n^5} \sum_{k=1}^n k^4 \right| = 0.$$

To do this, we utilize a ε of room argument, i.e. showing that for any $\varepsilon > 0$,

$$\limsup_{n \rightarrow \infty} \left| c_n - \frac{a}{n^5} \sum_{k=1}^n k^4 \right| \leq \varepsilon.$$

One can pick k_0 such that for $k \geq k_0$, $|a_k - a| \leq \varepsilon$. Thus

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left| c_n - \frac{a}{n^5} \sum_{k=1}^n k^4 \right| &= \limsup_{n \rightarrow \infty} \left| \frac{1}{n^5} \sum_{k=1}^n k^4 (a_k - a) \right| \\ &= \limsup_{n \rightarrow \infty} \left| \frac{1}{n^5} \sum_{k=k_0}^n k^4 (a_k - a) \right| \\ &= \limsup_{n \rightarrow \infty} \frac{\varepsilon}{n^5} \sum_{k=k_0}^n k^4 \\ &= \varepsilon \cdot (1/5) \leq \varepsilon. \end{aligned}$$

But now we find that

$$\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \frac{a}{n^5} \sum_{k=1}^n k^4 = a \lim_{n \rightarrow \infty} (1/5 + O(1/n)) = a/5.$$

- (b) Prove or Disprove: If $\lim_{n \rightarrow \infty} c_n = c$ exists, then $\lim_{n \rightarrow \infty} a_n = a$ exists.

Solution: This is false, as we might expect since averages like those defining the sequence $\{c_n\}$ are often much more well behaved than the original sequence $\{a_n\}$ upon which they are defined. In particular, if the sequence $\{a_n\}$ has a lot of *oscillation*, then we might expect cancellation in the average to yield a much smaller sequence $\{c_n\}$. Thus we are lead to consider the sequence $a_n = (-1)^n$ to obtain a counterexample. The sequence $\{a_n\}$ clearly does not converge. But we claim that the sequence $\{c_n\}$ *does* converge. We have

$$c_n = \frac{1}{n^5} \sum_{k=1}^n (-1)^k k^4 = \frac{S_n}{n^5},$$

where $S_n = \sum_{k=1}^n (-1)^k k^4$. We will prove by induction that for any $n \geq 1$, $|S_n| \leq n^4$, and $\text{Sign}(S_n) = (-1)^n$. The case $n = 1$ is obvious. Assuming the result for n , we have

$$S_{n+1} = S_n + (-1)^{n+1} (n+1)^4.$$

Now S_n and $(-1)^{n+1} (n+1)^4$ have opposite signs, $|S_n| \leq n^4$, and $(n+1)^4 > n^4$, so we conclude

that $S_n + (-1)^{n+1}(n+1)^4$ has sign $(-1)^{n+1}$, and

$$|S_{n+1}| \leq (n+1)^4.$$

But this implies that $c_n = S_n/n^5 = O(1/n)$, so $c_n \rightarrow 0$.

Another counterexample follows if we let $\{a_n\}$ be a *sparse sequence*, i.e. most elements are zero, in which case the averaged sequence $\{c_n\}$ will be small. Let

$$a_k = \begin{cases} 1 & : k = 2^l \text{ for some } l = 0, 1, 2, \dots, \\ 0 & : \text{otherwise.} \end{cases}$$

Then we find that for any $n \geq 1$, and any $m < 2^n$, $c_{2^n+m} < c_{2^n}$, and

$$c_{2^n} = \frac{1}{2^{5n}} \sum_{k=0}^n 2^{4k} = \frac{1}{2^{5n}} \frac{2^{4n} - 1}{2^4 - 1} \leq 2^{-n} \rightarrow 0.$$

Thus $\{c_n\} \rightarrow 0$.

3. (Fall 2018) For $c_k \in \mathbb{R}$, say that $\prod c_k$ converges if $\lim_{K \rightarrow \infty} \prod_{k=1}^K c_k = C$ exists with $C \neq 0, \infty$.

(a) Prove that if $0 < a_k < 1$ for all k , or if $-1 < a_k < 0$, for all k , then $\prod(1 + a_k)$ converges if and only if $\sum_k a_k$ converges.

Solution: Let

$$P_n = \prod_{k=1}^n (1 + a_k) \quad \text{and} \quad S_n = \sum_{k=1}^n a_k.$$

Then we have

$$P_n = \exp \left(\sum_{k=1}^n \log(1 + a_k) \right) = \exp(Q_n).$$

Thus $\{P_n\}$ converges to a nonzero value if and only if $\{Q_n\}$ converges to a finite value, and it suffices to show that the convergence of $\{Q_n\}$ is equivalent to the convergence of $\{S_n\}$. To do this, we apply the *limit comparison test* for series, which states that for any two non-negative sequences $\{a_n\}$ and $\{b_n\}$, if $\lim_{n \rightarrow \infty} (a_n/b_n) = c$, where $0 < c < \infty$, then either both of the series $\sum a_n$ and $\sum b_n$ converge, or they both diverge. We employ the inequality

$$\frac{x}{1+x} \leq \log(1+x) \leq x,$$

which holds for any $x \geq 1$. Thus

$$\frac{\log(1+a_k)}{a_k} \leq \frac{a_k}{a_k} = 1$$

and

$$\frac{\log(1+a_k)}{a_k} \geq \frac{1}{1+a_k} \geq 1-a_k.$$

Thus if $\{a_k\}$ converges to zero, then the convergence of $\{Q_n\}$ and $\{S_n\}$ are equivalent. But if $\{S_n\}$ converges, it is obvious that $\{a_k\}$ converges to zero, and if $\{Q_n\}$ converges, then $\log(1+a_k) \rightarrow 0$, which implies $a_k \rightarrow 0$. Thus we see that the convergence of $\{Q_n\}$ and $\{S_n\}$ are always equivalent.

Here is another approach, avoiding the limit comparison test, but at the cost of requiring some more quantitative calculations:

Case 1: Assume $a_k \in (0, 1)$ for all k .

First assume $\lim_{n \rightarrow \infty} S_n = L < \infty$. We wish to show that $\lim_{n \rightarrow \infty} P_n$ converges. Since $\{P_n\}$ is an increasing sequence, it suffices to show that $\{P_n\}$ is bounded above. Since $e^x \geq 1 + x$ for all $x \in \mathbb{R}$, we have

$$P_n \leq \prod_{k=1}^n e^{a_k} = e^{S_n} \leq e^L$$

where we have used the fact that S_n is increasing and convergent, so that $S_n \leq L$.

Next, assume that $\lim_{n \rightarrow \infty} P_n = M$ where $M \neq 0, \infty$. We wish to show that $\lim_{n \rightarrow \infty} S_n$ exists. Since $\{S_n\}$ is increasing, it again suffices to show that $\{S_n\}$ is bounded above. This will follow from an argument showing $S_n \leq P_n - 1$, since we then have $S_n \leq M - 1$ for all n . We prove this using induction. The case $n = 1$ is trivial since $S_1 = P_1 - 1$. For $n > 1$, observe that

$$\begin{aligned} P_n &= (1 + a_n)P_{n-1} \\ &\geq (1 + a_n)(1 + S_{n-1}) && \text{by induction hypothesis} \\ &= 1 + S_n + a_n S_{n-1} \\ &\geq 1 + S_n \end{aligned}$$

This proves the claim. Alternatively, using the inequality

$$\log(1 + x) \geq \frac{x}{1 + x},$$

which holds for all $x > -1$, we conclude that

$$P_n = \exp \left(\sum_{k=1}^n \log(1 + a_k) \right) \geq \exp \left(\sum_{k=1}^n \frac{a_k}{1 + a_k} \right) \geq \exp \left(\frac{1}{2} \sum_{k=1}^n a_k \right) = e^{S_n/2},$$

which implies $\{S_n\}$ is bounded from above if $\{P_n\}$ is bounded from above.

Case 2: Assume that $a_k \in (-1, 0)$ for all k .

Suppose $\lim_{n \rightarrow \infty} P_n = L$, with $L \neq 0$. We wish to show that S_n converges as well. Since $\{S_n\}$ is now decreasing, it suffices to show that $\{S_n\}$ is bounded from below. But this follows because $S_n \geq \log P_n$, so that $S_n \geq \log L$. Indeed, using the inequality $\log(1 + x) \leq x$, which holds for $x > -1$, we have

$$P_n = \exp \left(\sum_{k=1}^n \log(1 + a_k) \right) \leq e^{S_n}.$$

Taking logarithms on both sides verifies the inequality.

Next, assume that $\{S_n\}$ converges. We will show that $\{P_n\}$ converges. It suffices to show that $\{P_n\}$ is bounded from below, since it is decreasing. But this follows from the inequality

$$\log(1 + x) \geq \frac{x}{1 + x},$$

which holds for $x > -1$, so that

$$P_n = \exp \left(\sum_{k=1}^n \log(1 + a_k) \right) \geq \exp \left(\sum_{k=1}^n \frac{a_k}{1 + a_k} \right) \geq \exp \left(\sum_{k=1}^n a_k(1 - a_k) \right),$$

The fact that $\{S_k\}$ tends to zero implies that $\{a_k\}$ tends to zero, so that there is $c > 0$ such that $1 - a_k \geq c$, and thus

$$P_n \geq \exp(c \cdot S_n).$$

Thus $\{P_n\}$ is lower bounded.

- (b) However, prove that $\prod_{k>1} \left(1 + \frac{(-1)^k}{\sqrt{k}}\right)$ diverges.

Solution: Proof of Part (b): Let $P_n = \prod_{k=2}^n \left(1 + \frac{(-1)^k}{\sqrt{k}}\right)$. Our goal is to show the sequence $\{P_n\}$ diverges. Though one cannot apply the result above directly to this product, since the terms are not all positive or negative, we can apply the result to a subsequence of the products $\{P_n\}$ above by noting that *consecutive products* are negative. Thus

$$\begin{aligned} P_{2n-1} &= \prod_{k=2}^{2n-1} \left(1 + \frac{(-1)^k}{\sqrt{k}}\right) \\ &= \prod_{k=1}^n \left(1 - \frac{1}{\sqrt{2k+1}}\right) \left(1 + \frac{1}{\sqrt{2k}}\right) \\ &= \prod_{k=1}^n \left(1 - \frac{1 - (\sqrt{2k+1} - \sqrt{2k})}{\sqrt{2k}\sqrt{2k+1}}\right). \end{aligned}$$

Using the fact that

$$\sqrt{x} - \sqrt{y} = (\sqrt{x} - \sqrt{y}) \frac{\sqrt{x} + \sqrt{y}}{\sqrt{x} + \sqrt{y}} = \frac{x - y}{\sqrt{x} + \sqrt{y}},$$

we find that

$$0 < \frac{1 - (\sqrt{2k+1} - \sqrt{2k})}{\sqrt{2k}\sqrt{2k+1}} \leq \frac{1 - 1/(2k+1)}{2k} = \frac{1}{2k+1} < 1.$$

Thus the previous result implies $\{P_{2n-1}\}$ converges to zero. Since

$$\sum_{k=1}^n \frac{1 - (\sqrt{2k+1} - \sqrt{2k})}{\sqrt{2k}\sqrt{2k+1}} \sim \sum_{k=1}^n \frac{1}{k}$$

diverges, it follows that the product diverges.

4. (Fall 2019) Let f be a continuous function on \mathbb{R} satisfying

$$|f(x)| \leq \frac{1}{1+x^2}.$$

Define a function F on \mathbb{R} by

$$F(x) = \sum_{n=-\infty}^{\infty} f(x+n).$$

- (a) Prove that F is continuous and periodic with period 1.

Solution: For each $x \in \mathbb{R}$, the sum

$$\sum_{n=-\infty}^{\infty} f(x+n)$$

converges *absolutely*. This justifies that the limit of the infinite series remains the same if we rearrange the series. Thus

$$F(x+m) = \sum_{n=-\infty}^{\infty} f(x+n+m) = \sum_{n=-\infty}^{\infty} f(x+n) = F(x).$$

Thus F is periodic. Since the finite sum of continuous functions is continuous, each of the partial sums

$$F_K(x) = \sum_{n=-K}^K f(x+n) = \sum_{n=-K}^K f_n(x)$$

is continuous. Fix $N > 0$ and define $I_N = [-N, N]$. That F is continuous will follow if we can show that $\{F_N\}$ converges to F uniformly on I_N for each N , since the uniform limit of continuous functions is continuous. Thus we find F is continuous on I_N for each N , and taking $N \rightarrow \infty$ gives that F is continuous everywhere.

So let us show that $\{F_K\}$ converges to F uniformly in $L^\infty(I_N)$ for each N . If $|n| \geq 2N$, if $x \in I_N$, then

$$|x+n| \geq |n| - |x| \geq n/2.$$

Thus

$$|f_n(x)| \leq \frac{1}{1+(n/2)^2} \leq \frac{1}{1+n^2/4} \lesssim 1/n^2.$$

Thus for $K_2 \geq K_1 \geq T \geq 2N$, and $x \in I_N$,

$$\|F_{K_2} - F_{K_1}\|_{L^\infty(I_N)} = \left\| \sum_{K_1 < |n| \leq K_2} f_n \right\|_{L^\infty(I_N)} \lesssim \sum_{K_1 < |n| \leq K_2} 1/n^2 \lesssim 1/K_1 \leq 1/T.$$

Taking $T \rightarrow \infty$ shows that the sequence $\{F_K\}$ is Cauchy in $L^\infty(I_N)$, and thus converges uniformly to F .

(b) Prove that if G is continuous and periodic with period one, then

$$\int_0^1 F(x)G(x) dx = \int_{-\infty}^{\infty} f(x)G(x) dx.$$

Solution: Since F is the locally uniform limit of the functions $\{F_N\}$ defined above, we can interchange integrals and limits, writing

$$\begin{aligned} \int_0^1 F(x)G(x) dx &= \lim_{N \rightarrow \infty} \int_0^1 F_N(x)G(x) dx \\ &= \lim_{N \rightarrow \infty} \sum_{n=-N}^N \int_0^1 f_n(x)G(x) dx \\ &= \lim_{N \rightarrow \infty} \int_{-N}^{N+1} f(x)G(x) dx \\ &= \int_{-\infty}^{\infty} f(x)G(x) dx, \end{aligned}$$

where the last result follows because G is continuous and periodic, and thus bounded, and so

$$|f(x)G(x)| \lesssim |f(x)| \lesssim \frac{1}{1+x^2}.$$

5. (Fall 2015) Let a_1, a_2, \dots be a sequence of positive real numbers and assume that

$$\lim_{n \rightarrow \infty} \frac{a_1 + \dots + a_n}{n} = 1.$$

- (a) Show that $\lim_{n \rightarrow \infty} a_n n^{-1} = 0$.

Solution: We can extract the value a_n/n from the sequence $\{S_n/n\}$, since

$$\frac{a_n}{n} = \frac{S_n}{n} - \frac{n-1}{n} \frac{S_{n-1}}{n-1}.$$

Thus

$$\lim_{n \rightarrow \infty} \frac{a_n}{n} = \lim_{n \rightarrow \infty} \frac{S_n}{n} - \lim_{n \rightarrow \infty} \frac{n-1}{n} \frac{S_{n-1}}{n-1} = 1 - 1 = 0.$$

- (b) If $b_n = \max(a_1, \dots, a_n)$, show that $\lim_{n \rightarrow \infty} b_n n^{-1} = 0$.

Solution: We have

$$b_n/n = \max(1/n \cdot a_1/1, 2/n \cdot a_2/2, \dots, (n/n) \cdot (a_n/n)).$$

Fix $\varepsilon > 0$. There exists $N > 0$ such that for $n \geq N$, $|a_n/n| \leq \varepsilon$. Thus if $A = \max(a_1/1, \dots, a_N/N)$, we conclude that for $M \geq N$,

$$b_M/M \leq \max(A(N/M), \varepsilon).$$

Thus if $M \geq AN/\varepsilon$, $b_M/M \leq \varepsilon$. Taking $\varepsilon \rightarrow 0$ completes the proof.

- (c) Show that

$$\lim_{n \rightarrow \infty} \frac{a_1^\beta + \dots + a_n^\beta}{n^\beta} = \begin{cases} 0 & : \beta > 1 \\ \infty & : \beta < 1. \end{cases}$$

Solution: For $\beta > 1$, we have

$$\begin{aligned} \frac{a_1^\beta + \dots + a_n^\beta}{n^\beta} &= \frac{a_1 a_1^{\beta-1} + \dots + a_n a_n^{\beta-1}}{n^\beta} \\ &\leq \frac{a_1 + \dots + a_n}{n} \max(a_1/n, \dots, a_n/n)^{\beta-1} \\ &\leq (S_n/n) \cdot (b_n/n)^{\beta-1}. \end{aligned}$$

Thus

$$\lim_{n \rightarrow \infty} \frac{a_1^\beta + \dots + a_n^\beta}{n^\beta} = \lim_{n \rightarrow \infty} (S_n/n) \cdot \left(\lim_{n \rightarrow \infty} b_n/n \right)^{\beta-1} = 1 \cdot 0 = 0.$$

To prove the result for $\beta < 1$, let us begin with an intuitive argument. For $\beta < 1$, if $x \ll 1$, then $x^\beta \gg x$. Because $b_n/n \rightarrow 0$, for suitably large n we have $a_k/n \ll 1$ for $1 \leq k \leq n$. But this means that $(a_k/n)^\beta \gg a_k/n$, which implies

$$\frac{a_1^\beta + \dots + a_n^\beta}{n^\beta} \gg \frac{a_1 + \dots + a_n}{n} \approx 1.$$

More precisely, if $\delta > 0$, then there exists n_0 such that for $n \geq n_0$, and $1 \leq k \leq n$,

$$a_k/n \leq \delta \quad \text{and} \quad S_n/n \geq 1/2.$$

But this implies that

$$(a_k/n)^\beta = \frac{(a_k/n)}{(a_k/n)^{1-\beta}} \geq \frac{(a_k/n)}{\delta^{1-\beta}}.$$

Thus

$$\frac{a_1^\beta + \cdots + a_n^\beta}{n^\beta} = (a_1/n)^\beta + \cdots + (a_n/n)^\beta \geq \frac{1}{\delta^{1-\beta}} \frac{a_1 + \cdots + a_n}{n} \geq (1/2) \frac{1}{\delta^{1-\beta}}.$$

Taking $\delta \rightarrow 0$ completes the proof.

Another solution is to use an *interpolation argument*. We have

$$\left(\frac{a_1^\beta + \cdots + a_n^\beta}{n^\beta} \right) = \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^\beta}^\beta.$$

It thus suffices to show that for $\beta > 1$,

$$\limsup_{n \rightarrow \infty} \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^\beta} = 0,$$

and for $\beta < 1$,

$$\liminf_{n \rightarrow \infty} \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^\beta} = \infty,$$

For each $1 \leq \beta \leq \infty$, let

$$A_n(\beta) = \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^\beta}.$$

Then

$$\limsup_{n \rightarrow \infty} A_n(1) = \limsup_{n \rightarrow \infty} \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^1} = 1$$

and

$$\limsup_{n \rightarrow \infty} A_n(\infty) = \limsup_{n \rightarrow \infty} \left\| \frac{a_1 e_1 + \cdots + a_n e_n}{n} \right\|_{l^\infty} = \max_{1 \leq i \leq n} (a_i/n) = 0.$$

Hölder's inequality implies that

$$A_n(\beta) \leq A_n(1)^{1/\beta} A_n(\infty)^{1-1/\beta}.$$

Thus

$$\limsup_{n \rightarrow \infty} A_n(\beta) \leq (\limsup_n A_n(1))^{1/\beta} (\limsup_n A_n(\infty))^{1-1/\beta} = 1^{1/\beta} \cdot 0^{1-1/\beta} = 0.$$

This is the interpolation argument (upper bounding a quantity for two parameters β_0 and β_1 , and then using those results to upper bound a quantity for a parameter β lying in between β_0 and β_1). For $0 < \beta < 1$, another application of Hölder's inequality shows that if

$$\lambda = \frac{1/\beta - 1}{1/\beta - 1/2}$$

then $0 < \lambda < 1$, and

$$A_n(1) \leq A_n(\beta)^{1-\lambda} A_n(2)^\lambda.$$

Thus using the fact that $A_n(2) \rightarrow 0$ as $n \rightarrow \infty$ (we proved this above), we conclude that

$$\liminf_{n \rightarrow \infty} A_n(\beta) \geq \frac{(\lim_{n \rightarrow \infty} A_n(1))^{\frac{1}{1-\lambda}}}{(\lim_{n \rightarrow \infty} A_n(2))^{\frac{\lambda}{1-\lambda}}} = \frac{1}{0} = \infty.$$

6. (Fall 2021) Let $f \in C^1[0, 1]$. Show that for every $\varepsilon > 0$, there exists a polynomial p such that

$$\|f - p\|_{L^\infty[0,1]} + \|f' - p'\|_{L^\infty[0,1]} \leq \varepsilon.$$

Solution: Since $f' \in C[0, 1]$, applying the *Stone-Weierstrass theorem*, for any $\varepsilon > 0$, we can find a polynomial q such that $\|f' - q\|_{L^\infty[0,1]} \leq \varepsilon/2$. But then if we consider the polynomial p such that $p(0) = f(0)$, and $p' = q$, then the fundamental theorem of calculus shows that for $x \in [0, 1]$,

$$|f(x) - p(x)| = \left| \int_0^x f'(x) - p'(x) \right| \leq \int_0^x |f'(x) - p'(x)| \leq (\varepsilon/2) \cdot x \leq (\varepsilon/2).$$

Thus we have proved that

$$\|f - p\|_{L^\infty[0,1]} + \|f' - p'\|_{L^\infty[0,1]} \leq \varepsilon.$$

Day 2: Warm Up Question

3 Day 2: Basic Analysis

7. (Spring 2017, Spring 2011, and Spring 2007) Show that the sequence of functions

$$S_n(x) = \sum_{k=1}^n \frac{\sin(kx)}{k}, \quad n = 1, 2, 3, \dots,$$

is uniformly bounded in \mathbb{R} .

Hint: Summation by parts. Break the sum into two parts for $kx \leq 1$ and $kx > 1$ respectively.

Solution: The idea is to use *Summation by Parts*, also known as *Abel's lemma*, which says the following: Let $\{a_k\}$ and $\{b_k\}$ be two sequences of real numbers, and let

$$T_n = \sum_{k=1}^n a_k$$

be the partial sum of the terms $\{a_k\}$. Then

$$\sum_{k=m}^n a_k b_k = T_n b_n - T_{m-1} b_m + \sum_{k=m}^{n-1} T_k (b_k - b_{k+1}).$$

Summation by parts can be useful in many different scenarios, but the most useful scenario occurs when the sequence $\{a_k\}$ is oscillating, and the sequence $\{b_k\}$ is 'smooth' (i.e. changing gradually). The oscillation of $\{a_k\}$ causes the sequence $\{T_k\}$ to be small overall because of cancellation, and the smoothness of $\{b_k\}$ causes the terms $b_k - b_{k+1}$ to be small.

Let us use summation by parts to prove this result. It suffices to prove a uniform bound for $-\pi/2 < x < \pi/2$, because the absolute value of each of the functions S_n is periodic with degree π . First, we break the sum into terms with little oscillation, but whose terms are overall relatively small, and terms with large oscillation to which we can apply summation by parts. Thus for each $x \in (-\pi/2, \pi/2)$, find a non-negative integer such that $N < 1/|x| < N+1$. Write

$$S_n(x) = L_n(x) + H_n(x),$$

where

$$L_n(x) = \sum_{k=1}^N \frac{\sin(kx)}{k} \quad \text{and} \quad H_n(x) = \sum_{k=N+1}^n \frac{\sin(kx)}{k}.$$

Utilizing the inequality $|\sin(a)| \leq |a|$ and the triangle inequality, we find that

$$|L_n(x)| = \left| \sum_{k=1}^N \frac{\sin(kx)}{k} \right| \leq \sum_{k=1}^N \frac{|kx|}{k} \leq N|x| < 1.$$

To estimate H_n , we apply summation by parts. Let

$$T_k(x) = \sum_{j=1}^k \sin(jx).$$

The triangle inequality gives the trivial estimate $|T_k(x)| \leq k$ for each $k > 0$ and all $x \in \mathbb{R}$. A more robust estimate follows via the calculation that for $x \in (-\pi/2, \pi/2)$,

$$|T_k(x)| = \left| \sum_{j=1}^k \sin(jx) \right| = \left| \operatorname{Im} \left(\sum_{j=1}^k e^{ijx} \right) \right| \leq \left| e^{ix} \frac{e^{kix} - 1}{e^{ix} - 1} \right| \leq \frac{2}{|\sin x|} \lesssim \frac{1}{|x|},$$

where in the second last inequality, we have used

$$\left| e^{ix} \frac{e^{kix} - 1}{e^{ix} - 1} \right| \leq \frac{2}{|e^{ix} - 1|} = \frac{2}{\sqrt{(\cos(x) - 1)^2 + \sin^2(x)}} = \sqrt{\frac{2}{1 - \cos x}} = \frac{1}{|\sin(x/2)|}.$$

Thus

$$\begin{aligned} |H_n(x)| &= \left| \sum_{k=N+1}^n \frac{\sin(kx)}{k} \right| \\ &= \left| \frac{T_n(x)}{n} - \frac{T_N(x)}{N+1} + \sum_{k=N+1}^{n-1} T_k(x) \left(\frac{1}{k} - \frac{1}{k+1} \right) \right| \\ &\leq 2 + \left| \sum_{k=N+1}^{n-1} \frac{T_k(x)}{k(k+1)} \right| \\ &\lesssim 1 + \frac{1}{|x|} \sum_{k=N+1}^{n-1} \frac{1}{k^2} \\ &\lesssim 1 + \frac{1}{N|x|} \lesssim 1. \end{aligned}$$

This shows that both L_n and H_n are uniformly bounded in n , and thus the same is true of S_n .

An alternate approach is to employ *Fourier analysis* together with a *Tauberian theorem*. Tauberian theorems use tools that are beyond the scope of the prerequisites of the qualifying exam, though the proof may be of interest for those who know some basic Fourier analysis. The sums here are the partial Fourier series corresponding to the Fourier coefficients of a bounded, 2π -periodic function $f \in L^\infty[-\pi, \pi]$, i.e. the function f such that

$$f \sim \sum_{k=1}^{\infty} \frac{\sin(kx)}{k}.$$

It is not necessary to know the function f , but it is a constant multiple of the function

$$x \mapsto \begin{cases} -\pi/2 - x/2 & : -\pi < x < 0 \\ +\pi/2 - x/2 & : 0 < x < \pi \end{cases}$$

It follows that $S_n = D_n * f$, where D_n is the *Dirichlet kernel*. If $\|D_n\|_{L^1([0, 2\pi])}$ was uniformly bounded in n , we could apply Young's inequality, implying

$$\|S_n\|_{L^\infty([0, 2\pi])} \leq \|D_n\|_{L^1([0, 2\pi])} \|f\|_{L^\infty([0, 2\pi])} \lesssim 1.$$

Unfortunately, we have $\|D_n\|_{L^1([0, 2\pi])} \sim \log n$, so this approach doesn't work completely. But there is a general result for functions $f \in L^1([0, 2\pi])$ such that

$$|\hat{f}(n)| \lesssim 1/n,$$

known as a *Tauberian theorem*, which shows that $D_n * f$ is uniformly bounded in n if and only if $P_r * f$ is bounded as $r \rightarrow 1$, where $\{P_r\}$ is the *Poisson kernel*. Since $\|P_r\|_{L^1([0, 2\pi])} \lesssim 1$ for $0 < r < 1$, we can use Young's inequality to conclude that $\|P_r * f\|_{L^\infty([0, 2\pi])} \lesssim 1$. Thus $P_r * f$ is uniformly bounded in r , and thus $D_n * f$ is uniformly bounded in n , completing the proof.

8. (Spring 2021)

$$\sum_{k=1}^{\infty} \frac{\cos(\sqrt{k})}{k}$$

converges.

Solution: Since the sum of $1/k$ diverges, this convergence could only possibly happen from the oscillation present in the $\cos(\sqrt{k})$ causes enough cancellation. Thus trying to control the oscillation is key to understanding this series. In particular, let us try and understand how often $\cos(\sqrt{k})$ changes sign. We can use conjugation, an often useful trick to understand square roots, to write

$$\begin{aligned} \sqrt{k+n} - \sqrt{k} &= (\sqrt{k+n} - \sqrt{k}) \left(\frac{\sqrt{k+n} + \sqrt{k}}{\sqrt{k+n} + \sqrt{k}} \right) \\ &= \frac{n}{\sqrt{k+n} + \sqrt{k}} \\ &= \frac{n}{2\sqrt{k} + O(n/\sqrt{k})} \\ &= \frac{n}{2\sqrt{k}} + O\left(\frac{n^2}{k^{3/2}}\right). \end{aligned}$$

In particular,

$$\sqrt{k + 2\pi\sqrt{k} + k} = \pi + O(1/\sqrt{k}).$$

Thus for each n , the sign of $\cos(\sqrt{k})$ stays roughly the same for $k = n \pm \sqrt{n}$. More precisely, let $\{m_1, m_2, \dots\}$ be an increasing family of integers with $m_1 = 1$, and such that for each i , and $k \in \{m_i, m_i + 1, \dots, m_{i+1} - 1\}$, $\cos(\sqrt{k})$ has the same sign. Then $m_{i+1} - m_i \sim \sqrt{m_i}$. For more than half of the values in $\{m_i, \dots, m_{i+1} - 1\}$, $|\cos(\sqrt{k})| \geq 1/2$, and it follows that

$$\left| \sum_{k=m_i}^{m_{i+1}-1} \frac{\cos(\sqrt{k})}{k} \right| \sim \frac{1}{\sqrt{m_i}}.$$

Roughly speaking, an application of Leibnitz's alternating series test then implies that this series converges.

9. (Fall 2015) Consider the series

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} \sin(x/n).$$

(a) Show that the series converges pointwise to some function f on \mathbb{R} .

Solution: The main idea to understanding the sum is to use the bound $|\sin(y)| \leq |y|$, which is tight when $|y| \leq 1$. Thus little is lost in our estimates if we apply this bound in the sum above for $|n| \geq |x|$, since then $|x/n| \leq 1$. We will actually find this sum converges absolutely, locally uniformly in x . Indeed, plugging in the bound $|\sin(x)| \leq |x|$, we find that

$$\sum_{n=1}^{\infty} \left| \frac{1}{\sqrt{n}} \sin(x/n) \right| \leq \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} \frac{|x|}{n} = |x| \sum_{n=1}^{\infty} \frac{1}{n^{3/2}} \lesssim |x|.$$

Since the sum converges pointwise absolutely, it must converge pointwise.

(b) Is f continuous on \mathbb{R} ? Does $f'(x)$ exist for all $x \in \mathbb{R}$?

Solution: To see that f is continuous, it suffices to show that the partial sums

$$f_N(x) = \sum_{n=1}^N \frac{1}{\sqrt{n}} \sin(x/n)$$

converge *locally uniformly* in x to f , since the limit of a sequence of continuous functions converging locally uniformly is continuous. But employing the bound $|\sin(x)| \leq |x|$ as above, we find that for $M > N$,

$$|f_M(x) - f_N(x)| \leq \sum_{n=N+1}^M \left| \frac{1}{\sqrt{n}} \sin(x/n) \right| \leq |x| \sum_{n=N+1}^M 1/n^{3/2} \lesssim |x|/N^{1/2}.$$

But this equation implies locally uniform convergence, i.e. for any interval $I = [-K, K]$, we have proved that

$$\|f_M - f_N\|_{L^\infty(I)} \lesssim K/N^{1/2}.$$

Thus f is a continuous function.

Now to understand the differentiability of f , it makes sense to look at the derivatives

$$f'_N(x) = - \sum_{n=1}^N (1/n^{3/2}) \cos(x/n).$$

Since the cosine term is uniformly bounded in x and n , it is fairly negligible to studying the limit of this quantity as $N \rightarrow \infty$. Indeed, we find that for $M > N$, we can plug in the bound $|\cos(y)| \leq 1$ to conclude that

$$|f'_M(x) - f'_N(x)| \leq \sum_{n=N+1}^M 1/n^{3/2} \lesssim N^{-1/2}.$$

Thus $\|f'_M - f'_N\|_{L^\infty(\mathbb{R})} \lesssim N^{-1/2}$, which implies the sequence is uniformly Cauchy, and thus converges to some function g . But it is a general rule that if $\{f_N\}$ converges locally uniformly to some function f , and $\{f'_N\}$ converges locally uniformly to some function g , then f is differentiable, and $f' = g$. Thus f is differentiable.

An alternative, but less clean way to show f is differentiable is using the definition of the derivative, i.e. by proving that for each $x \in \mathbb{R}$, the limit

$$\frac{f(x+h) - f(x)}{h}$$

exists. To see this, write

$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{1}{h} \sum_{n=1}^N \frac{1}{\sqrt{n}} \left[\sin\left(\frac{x+h}{n}\right) - \sin\left(\frac{x}{n}\right) \right] \\ &\quad \underbrace{\hspace{10em}}_{\text{Call it } A_{N,h}(x)} \\ &\quad + \frac{1}{h} \sum_{n=N+1}^{\infty} \frac{1}{\sqrt{n}} \left[\sin\left(\frac{x+h}{n}\right) - \sin\left(\frac{x}{n}\right) \right]. \end{aligned} \tag{1}$$

Call it $B_N(x)$

We consider the two terms on the right hand side individually. By the mean value theorem, $|\sin(\frac{x+h}{n}) - \sin(\frac{x}{n})| \leq \frac{|h|}{n}$, and hence

$$|B_N(x)| \leq \sum_{n=N+1}^{\infty} \frac{1}{n^{3/2}} \leq \frac{C}{\sqrt{N}} \quad (2)$$

for some constant $C > 0$ not depending on x .

Next we consider $A_{N,h}$. Let

$$g_N(x) = \sum_{n=1}^N \frac{1}{n^{3/2}} \cos(x/n)$$

and let

$$g_{\infty}(x) = \sum_{n=1}^{\infty} \frac{1}{n^{3/2}} \cos(x/n).$$

Then (it is easy to check that)

$$\lim_{h \rightarrow \infty} A_{N,h}(x) = g_N(x) \text{ for each } x \quad (3)$$

and

$$\lim_{N \rightarrow \infty} g_N(x) = g_{\infty}(x) \text{ uniformly in } x. \quad (4)$$

Therefore by Eq. (1),

$$\frac{f(x+h) - f(x)}{h} - g_{\infty}(x) = g_N(x) - g_{\infty}(x) + A_{N,h}(x) - g_N(x) + B_N$$

so that

$$\left| \frac{f(x+h) - f(x)}{h} - g_{\infty}(x) \right| \leq \sup_{x \in \mathbb{R}} |g_N(x) - g_{\infty}(x)| + |A_{N,h}(x) - g_N(x)| + \frac{C}{\sqrt{N}}$$

Let $\epsilon > 0$ be arbitrary. By (2) and (4), we may choose N sufficiently large that $C/\sqrt{N} < \epsilon/2$ and $\sup_{x \in \mathbb{R}} |g_N(x) - g_{\infty}(x)| < \epsilon/2$, which implies

$$\left| \frac{f(x+h) - f(x)}{h} - g_{\infty}(x) \right| \leq |A_{N,h}(x) - g_N(x)| + \epsilon.$$

Taking the limit supremum of both sides as $h \rightarrow 0$ and using (3), we obtain

$$\limsup_{h \rightarrow 0} \left| \frac{f(x+h) - f(x)}{h} - g_{\infty}(x) \right| \leq \epsilon$$

Since $\epsilon > 0$ was arbitrary, it holds that

$$\limsup_{h \rightarrow 0} \left| \frac{f(x+h) - f(x)}{h} - g_{\infty}(x) \right| = 0.$$

Therefore

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = g_{\infty}(x) \in \mathbb{R}.$$

(c) Does the series converge uniformly on \mathbb{R} ?

Solution: The bounds in part (a) of the problem, which depend on x , should hint to you that the series does *not* converge uniformly in x , in particular, as we take x to be very large. To show that the sum does not converge uniformly, we need lower bounds, and to do this we again employ the heuristic that $|\sin(y)| \leq |y|$ is a good estimate for $|y| \leq 1$, so we should get a similar estimate which lower bounds $\sin(y)$ in this range. In fact, for $0 \leq y \leq 1$ we actually have $\sin(y) \geq y/2$. Concentrating on the case where x is a positive integer for simplicity, we find that

$$\sum_{n \geq x} \frac{1}{\sqrt{n}} \sin(x/n) \gtrsim \sum_{n \geq x} \frac{x}{n^{3/2}} \gtrsim \sqrt{x}.$$

Thus if $N = x$, we can reword this inequality as saying that

$$\lim_{M \rightarrow \infty} f_M(x) - f_N(x) \gtrsim \sqrt{x}.$$

Thus

$$\limsup_{M \rightarrow \infty} \|f_M - f_N\|_{L^\infty(\mathbb{R})} \gtrsim \sqrt{N}.$$

In particular, this implies $\{f_N\}$ is not a Cauchy sequence in $L^\infty(\mathbb{R})$, and thus does not converge uniformly.

Alternatively, we can use the fact that if $\pi/3 \leq y \leq 2\pi/3$, then $\sin(y) \geq 1/2$. Thus, for any N , if we pick $x = (\pi/3)N$, then for $1 \leq n \leq N$, $\sin(x/n)$ is non-negative, and for $N/2 \leq n \leq N$, $\sin(x/n) \geq 1/2$, so

$$|f_N(x)| \gtrsim \sum_{N/2 \leq n \leq N} 1/\sqrt{n} \gtrsim \sqrt{N}.$$

Thus $\|f_N\|_{L^\infty(\mathbb{R})} \gtrsim \sqrt{N}$, which implies $\{f_N\}$ is not bounded in $L^\infty(\mathbb{R})$, and so $\{f_N\}$ is not a Cauchy sequence in $L^\infty(\mathbb{R})$.

10. (Fall 2019) Show that if $K \subset \mathbb{R}^n$, and every continuous function on K is bounded, then K is compact.

Solution: The *Heine-Borel theorem* implies that it suffices to show that K is bounded and closed.

If K was not closed, then we could find $y \in \overline{K} - K$. The function $f(x) = 1/|x - y|$ would then be continuous, but not bounded, on K . Thus $\overline{K} - K$ is empty, so $\overline{K} = K$.

If K was not bounded, then $f(x) = |x|$ would be a continuous, unbounded function. Thus K is bounded.

Thus K is compact.

11. (Spring 2015) Prove that the integral

$$f(a) = \int_0^\infty \frac{\sin(x^2 + ax)}{x} dx$$

converges for $a \geq 0$, and f is continuous on $[0, \infty)$.

Solution: Let

$$f_0(a) = \int_0^1 \frac{\sin(x^2 + ax)}{x} dx.$$

We perform a *dyadic decomposition*, writing, for $n \in \mathbb{Z}$,

$$f_n(a) = \int_{2^n}^{2^{n+1}} \frac{\sin(x^2 + ax)}{x} dx.$$

Then $f = \sum_{n=0}^{\infty} f_n$. Since the integrand is continuous, and each integral is on a finite interval, all of the functions $\{f_n\}$ are continuous.

We claim this sum converges locally uniformly. To do this we *must* exploit the oscillation in the integrand, for $1/x$ is not summable. To do this, we perform an *integration by parts*. Indeed, for $a \geq 0$,

$$\begin{aligned} \int_{2^n}^{2^{n+1}} \frac{\sin(x^2 + ax)}{x} &= \int_{2^n}^{2^{n+1}} \frac{d}{dx} (\cos(x^2 + ax)) \cdot \frac{-1}{x(2x + a)} \\ &= \left(\int_{2^n}^{2^{n+1}} \frac{\cos(x^2 + ax)}{x(2x + a)} \right) + O(1/2^n) \\ &\leq \int_{2^n}^{2^{n+1}} \frac{1}{x(2x + a)} + O(1/2^n) \\ &\lesssim 1/2^n. \end{aligned}$$

Thus $\|f_n\|_{L^\infty[0, \infty)} \lesssim 1/2^n$. Thus $\{f_n\}$ is uniformly summable, and so $f = \sum f_n$ exists, and is continuous for $a \geq 0$.

One can also work without a dyadic decomposition, i.e. using integration by parts to write

$$\begin{aligned} &= \int_A^B \frac{\sin(x^2 + ax)}{x} dx \\ &= \int_A^B \frac{\sin(x^2 + ax)(2x + a)}{x(2x + a)} dx \\ &= \int_A^B \frac{d}{dx} [-\cos(x^2 + ax)] \cdot \frac{1}{x(2x + a)} dx \\ &= \frac{-\cos(x^2 + ax)}{x(2x + a)} \Big|_{x=A}^{x=B} + \int_A^B \cos(x^2 + ax) \left(\frac{-1}{x^2(2x + a)} - \frac{2}{x(2x + a)^2} \right) dx \end{aligned}$$

Therefore if

$$f_C(a) = \int_0^C \frac{\sin(x^2 + ax)}{x} dx,$$

then we have shown that

$$|f_B(a) - f_A(a)| \leq \frac{1}{2A^2} + \frac{1}{2B^2} + \int_A^B \frac{1}{2x^3} + \frac{2}{x(2x)^2} dx \leq 10 \left(\frac{1}{A^2} + \frac{1}{B^2} \right).$$

Thus

$$\|f_B - f_A\|_{L^\infty[0, \infty)} \leq 10(1/A^2 + 1/B^2) \rightarrow 0,$$

which shows $\{f_A\}$ converges uniformly to a continuous function as $A \rightarrow \infty$.

12. (Fall 2017) Consider the sequence of functions $f_n : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f_n(x) = \int_0^n \frac{\sin(sx)}{\sqrt{s}} ds.$$

- (a) Show that $\{f_n\}$ converges locally uniformly on $(0, \infty)$.

Solution: This integral can only converge because of the oscillation of $\sin(sx)$ for large x . Thus we *integrate by parts*, writing

$$\begin{aligned} \int_{n_0}^{n_1} \frac{\sin(sx)}{\sqrt{s}} ds &= \int_{n_0}^{n_1} \frac{-d/ds(\cos(sx))}{x\sqrt{s}} ds \\ &= \frac{-\cos(sx)}{xs^{1/2}} \Big|_{n_0}^{n_1} + \int_{n_0}^{n_1} \frac{\cos(sx)}{xs^{3/2}} (-1/2) ds \\ &= O(1/xn_0^{1/2}) + O(1/x \int_{n_0}^{n_1} 1/s^{3/2} ds) \\ &= O(1/xn_0^{1/2}). \end{aligned}$$

Thus for any fixed $\alpha > 0$, we get uniform convergence for $\alpha < x < \infty$.

- (b) Show that $\{f_n\}$ does *not* converge uniformly on $(0, 1]$.

Solution: If the convergence was uniform, there would be n_0 such that for any $n_1 \geq n_0$, and $x \in [0, 1]$,

$$\left| \int_{n_0}^{n_1} \frac{\sin(sx)}{\sqrt{s}} ds \right| \leq 1.$$

Now pick $x \in [0, 1]$ with $x \sim 1/n_1$, such that for $s \in [n_0, n_1]$, $\sin(sx) \geq sx/2$. Then

$$1 \geq \int_{n_0}^{n_1} \frac{\sin(sx)}{\sqrt{s}} ds \geq \frac{x}{2} \int_{n_0}^{n_1} \sqrt{s} ds \gtrsim n_1^{3/2} x \gtrsim n_1^{1/2}.$$

This gives a contradiction for sufficiently large n_1 .

- (c) Does the sequence $\{f_n\}$ converge uniformly on $[1, \infty)$ as $n \rightarrow \infty$?

Solution: Yes, as implied by the calculations in part (a).

13. (Fall 2021) Does the improper integral

$$\int_2^\infty \frac{x \sin(e^x)}{x + \sin(e^x)} dx$$

converge?

Solution: Here's one solution. Our first intuition is to break the integrand into an oscillatory part, and a fairly well behaved part. A good option seems to be breaking the sum into the product

$$\frac{x}{x + \sin(e^x)} \cdot \sin(e^x) = A(x) \cdot B(x).$$

For large x , the well behaved part of the integrand is close to one. So let's do a Taylor expansion. A first order expansion yields $A(x) = 1 + O(1/x)$, but the error term here is not good enough (it's not integrable in x). Thus we perform a second order expansion, i.e. writing $A(x) = 1 - \sin(e^x)/x + O(1/x^2)$. Now the error term is integrable, so we conclude that the convergence of the improper integral above is equivalent to the convergence of the integral

$$\int_2^\infty \left(1 - \frac{\sin(e^x)}{x} \right) \sin(e^x) dx.$$

A change of variables might help us out here. We write $y = e^x$. Then $dy = ydx$, and so this integral is equal to

$$\int_c^\infty \frac{\sin(y)}{y} \left(1 - \frac{\sin(y)}{\ln y}\right).$$

A simple integration by parts (antidifferentiate the oscillating term $\sin(y)$, differentiate the term going to zero $1/y$) yields

$$\int_c^\infty \frac{\sin(y)}{y} = C + \int_{O(1)}^\infty \frac{\cos(y)}{y^2}.$$

The integral on the right hand side is summable, and thus the integral converges. Thus the convergence of the overall integral is equivalent to

$$\int_c^\infty \frac{\sin(y)^2}{y \ln(y)}.$$

This integral is positive, and *not integrable*, i.e. a dyadic decomposition (a tool for breaking a sum into parts on powers of two, which helps us out in an analogous to the Cauchy condensation theorem) shows that

$$\int_c^\infty \frac{\sin(y)^2}{y \ln(y)} \gtrsim \sum_{k=1}^\infty \int_{2^k}^{2^{k+1}} \frac{\sin(y)^2}{y \ln(y)} \gtrsim \sum_{k=1}^\infty 2^k \frac{1}{k 2^k} = \sum_{k=1}^\infty \frac{1}{k} = \infty.$$

Thus we conclude that the integral diverges, and thus the same must be true for the overall integral.

Here is an alternate solution. If $y = e^x$, then $dy = e^x dx = y dx$, and so

$$\int_2^\infty \frac{x \sin(e^x)}{x + \sin(e^x)} dx = \int_{\log 2}^\infty \frac{\sin y}{y} \frac{\log(y)}{\log(y) + \sin(y)}$$

The integrand oscillates between positive and negative values, so it suffices to decompose the integral onto the intervals where the integrand is positive or negative. For suitably large n , write

$$C_n = \left| \int_{\pi n}^{\pi(n+1)} \frac{\sin y}{y} \frac{\log y}{\log y + (-1)^n \sin y} \right|$$

The integral will then converge if and only if the alternating series

$$\sum_{n=1}^\infty (-1)^n C_n$$

converges. For $\pi n \leq y \leq \pi(n+1)$, a Taylor series expansion shows that

$$\begin{aligned} \frac{1}{y} \frac{\log y}{\log y + (-1)^n \sin y} &= \frac{1}{y} \frac{1}{1 + (-1)^n \sin y / \log y} \\ &= \left(\frac{1}{\pi n} + O(1/n^2) \right) \left(1 - (-1)^n \frac{\sin y}{\log y} + O\left(\frac{1}{(\log n)^2} \right) \right) \\ &= \left(\frac{1}{\pi n} + O(1/n^2) \right) \left(1 - (-1)^n \frac{\sin y}{\log(\pi n)} + O\left(\frac{1}{(\log n)^2} \right) \right) \\ &= \left(\frac{1}{\pi n} - (-1)^n \frac{\sin y}{\pi n \log \pi n} + O\left(\frac{1}{n(\log n)^2} \right) \right). \end{aligned}$$

Thus

$$\begin{aligned} C_n &= \left| \int_0^\pi \frac{\sin y}{\pi n} - (-1)^n \frac{(\sin y)^2}{\pi n \log \pi n} dy \right| + O\left(\frac{1}{n(\log n)^2}\right) \\ &= \left| \frac{2}{\pi n} - (-1)^n \frac{1}{2n \log(\pi n)} \right| + O\left(\frac{1}{n(\log n)^2}\right) \\ &= C'_n + O\left(\frac{1}{n(\log n)^2}\right). \end{aligned}$$

For large n , we have

$$C'_n = \frac{2}{\pi n} - (-1)^n \frac{1}{2n \log(\pi n)}.$$

Thus

$$\sum (-1)^n C'_n = \sum (-1)^n (2/\pi n) - \frac{1}{2n \log(\pi n)}.$$

The sum $\sum (-1)^n (2/\pi n)$ converges by the *Leibnitz test*. But the other part of the series diverges, i.e.

$$\sum_n \frac{1}{2n \log(\pi n)} = \infty.$$

One can prove this using the Cauchy condensation theorem. Thus the alternating series *diverges*, and so the entire integral diverges.

Day 3: Warm Up Question

14. (Fall 2018) Prove that for $1 \leq p \leq 2$ and $0 < b < a$,

$$(a+b)^p + (a-b)^p \geq 2a^p + p(p-1)a^{p-2}b^2.$$

Solution: [Hint: Apply Taylors Theorem.](#)

By subtracting the right-hand side, factoring out a^p , and noting that $0 < b/a < 1$, it suffices to show that the function

$$f(x) = (1+x)^p + (1-x)^p - 2 - p(p-1)x^2$$

is nonnegative for all $x \in (0, 1)$. Let $g(x) = (1+x)^p$, so that

$$f(x) = g(x) + g(-x) - 2 - p(p-1)x^2.$$

We next Taylor expand g about the point 0. For any $x > -1$, we have

$$g(x) = 1 + px + \frac{1}{2}p(p-1)x^2 + R_2(x)$$

where

$$R_2(x) = \frac{p(p-1)(p-2)(1+\xi)^{p-3}x^3}{6}$$

for some number ξ between 0 and x . Substituting the Taylor expansions for $g(x)$ and $g(-x)$ into the formula for f gives

$$f(x) = R_2(x) + R_2(-x)$$

or equivalently

$$f(x) = \frac{p(p-1)(p-2)}{6} [(1+\xi_+)^{p-3} - (1+\xi_-)^{p-3}] x^3$$

for some $\xi_+ \in (0, x)$ and $\xi_- \in (-x, 0)$. Since $\xi_- < \xi_+$ and $p < 3$, the part in brackets is negative, and since $0 \leq p \leq 2$, it follows that $f(x)$ is nonnegative, as required.

15. (Spring 2018 and Spring 2021) Determine if

$$\sum_{k=1}^{\infty} \frac{\cos(\sqrt{k})}{k}$$

converges.

Solution: [Hint: Integration by parts, Mean value theorem](#)

Yes, the series converges. We first claim that the integral

$$\int_1^{\infty} \frac{\cos(x^{1/2})}{x} dx$$

converges. To see this, we use integration by parts:

$$\begin{aligned} \int_1^{\infty} \frac{\cos(x^{1/2})}{x} dx &= \int_1^{\infty} 2x^{-1/2} \frac{d}{dx} [\sin(x^{1/2})] dx \\ &= -2\sin(1) + \int_1^{\infty} \frac{\sin(x^{1/2})}{x^{3/2}} dx < \infty. \end{aligned}$$

This proves the claim. Next we claim that

$$\sum_{k=1}^n \frac{\cos(\sqrt{k})}{k}$$

is a Cauchy sequence (and therefore converges as $n \rightarrow \infty$). To prove this, it suffices by the previous claim to show that for every $\epsilon > 0$ there exists N such that $m, n > N$ implies

$$\left| \sum_{k=m}^n \frac{\cos(\sqrt{k})}{k} - \int_m^{n+1} \frac{\cos(\sqrt{x})}{x} dx \right| < \epsilon. \quad (5)$$

Let $1 \leq n \leq m$. Then

$$\sum_{k=m}^n \frac{\cos(\sqrt{k})}{k} - \int_m^{n+1} \frac{\cos(\sqrt{x})}{x} dx = \sum_{k=m}^n \int_k^{k+1} \left(\frac{\cos(\sqrt{k})}{k} - \frac{\cos(\sqrt{x})}{x} \right) dx. \quad (6)$$

By the Mean Value Theorem, since

$$\frac{d}{dx} \left[\frac{\cos(x^{1/2})}{x} \right] = -\frac{\cos(x^{1/2})}{x^2} - \frac{\sin(x^{1/2})}{2x^{3/2}},$$

it follows that for every $x \in [k, k+1]$, there exists some $\xi \in (x, k+1)$ such that

$$\frac{\cos(\sqrt{k})}{k} - \frac{\cos(\sqrt{x})}{x} \leq \frac{1}{\xi^2} + \frac{1}{2\xi^{3/2}}$$

and since $\xi \geq k$,

$$\frac{\cos(\sqrt{k})}{k} - \frac{\cos(\sqrt{x})}{x} \leq \frac{1}{k^2} + \frac{1}{2k^{3/2}}.$$

Therefore

$$\left| \int_k^{k+1} \left(\frac{\cos(\sqrt{k})}{k} - \frac{\cos(\sqrt{x})}{x} \right) dx \right| \leq \frac{1}{k^2} + \frac{1}{2k^{3/2}}$$

Therefore using equation (6),

$$\left| \sum_{k=m}^n \frac{\cos(\sqrt{k})}{k} - \int_m^{n+1} \frac{\cos(\sqrt{x})}{x} dx \right| \leq \sum_{k=m}^n \frac{1}{k^2} + \frac{1}{2k^{3/2}}$$

Since the series on the right hand side is summable, it tends to zero as $m, n \rightarrow \infty$, which establishes (5).

4 Day 3: Basic Analysis

16. (Spring 2015) Let g be a non-constant differentiable real function on a finite interval $[a, b]$, with $g(a) = g(b) = 0$. Show that there exists $c \in (a, b)$ such that

$$|g'(c)| > \frac{4}{(b-a)^2} \int_a^b |g(t)| dt.$$

Solution: My main intuition is that the fundamental theorem of calculus enables us to relate g' and g . If we are clever about using the fundamental theorem, we obtain the bound, namely

$$\begin{aligned} \int_a^b |g(t)| dt &= \int_a^b \min \left(\left| \int_a^t g'(s) ds \right|, \left| \int_t^b g'(s) ds \right| \right) dt \\ &\leq \max_{a < c < b} |g'(c)| \cdot \int_a^b \min(|t-a|, |t-b|) dt \\ &= \max_{a < c < b} |g'(c)| \cdot \left(\int_a^{(a+b)/2} (t-a) dt + \int_{(a+b)/2}^b (b-t) dt \right) \\ &= \max_{a < c < b} |g'(c)| \cdot ((b-a)^2/8 + (b-a)^2/8) \\ &= \max_{a < c < b} |g'(c)| \cdot (b-a)^2/4. \end{aligned}$$

Rearranging this inequality completes the proof, except in the case that all the inequalities in this calculation are equalities (so we cannot get the strict inequality required). But this only holds if g' is a constant function, hence g is linear, which is impossible since g is non constant, and $g(0) = g(1) = 0$.

(Thanks to Karthik for this proof) There is a geometric argument for this result as well, though it is difficult to make it rigorous. If g is a function which maximizes

$$\int_a^b |g(t)| dt$$

subject to the constraint that $\|g'\|_{L^\infty[a,b]}$ and that $g(a) = g(b) = 0$, then we must have $|g'(x)| = 1$ for all $x \in (a, b)$ wherever g is differentiable, since otherwise we could modify g near this point so that $|g(t)|$ is larger at every point in this interval. In this case it is also true that g must be increasing on $[0, 1/2]$, and decreasing on $[1/2, 1]$, since if g' is negative at some point in $[0, 1/2]$, then we can flip the sign of the derivative near this point, and flip the sign later on, and this will increase the value of $\int_a^b |g(t)| dt$. But then $g(t)$ is a triangle function, i.e. obtained by linearly interpolating the points $g(a) = 0$, $g(b) = 0$, and $g((a+b)/2) = (b-a)/2$, and then

$$\int_a^b |g(t)| dt = \frac{(b-a)^2}{4}.$$

If g is differentiable at every point, it cannot be the function g above which maximizes the integral, so we must have

$$\int_a^b |g(t)| dt < \frac{(b-a)^2}{4}.$$

17. (Fall 2019) If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable on $\mathbb{R}^n - \{0\}$, continuous at 0, and

$$\lim_{x \rightarrow 0} \frac{\partial f}{\partial x^i}(x) = 0,$$

for $1 \leq i \leq n$, then f is differentiable at 0.

Solution: We claim that $\nabla f(0) = 0$, i.e. that $|f(x) - f(0)| = o(|x|)$ as $x \rightarrow 0$. To see this, we apply the mean value theorem. Assume without loss of generality that $f(0) = 0$. Fix $\varepsilon > 0$. For any $x \in \mathbb{R}^n - \{0\}$, since f is continuous at 0, there is $\lambda_0 \leq 1/2$ depending only on $|x|$ such that

$$|f(\lambda_0 x)| \leq |x|^2.$$

Now we apply the mean value theorem, writing $g(\lambda) = f(\lambda x)$. The mean value theorem implies that there is $\lambda \in (\lambda_0, 1)$ such that

$$g'(\lambda) = \frac{f(x) - f(\lambda_0 x)}{1 - \lambda_0}.$$

But

$$g'(\lambda) = \sum_{i=1}^n \frac{\partial f}{\partial x^i}(\lambda x) \cdot x_i.$$

If $|x|$ is suitably small, then $|\frac{\partial f}{\partial x^i}(\lambda x)| \leq \varepsilon$, which implies that

$$|f(x) - f(\lambda_0 x)| \leq 2 \left| \frac{f(x) - f(\lambda_0 x)}{1 - \lambda_0} \right| \lesssim \varepsilon |x|.$$

Thus

$$|f(x)| \leq |f(\lambda_0 x)| + 2\varepsilon |x| \leq |x|^2 + \varepsilon |x|.$$

Thus if $|x|$ is suitably small, we conclude that $|f(x)| \lesssim \varepsilon |x|$. Since ε was arbitrary, this shows that f is differentiable at zero, and $\nabla f(0) = 0$.

18. (Spring 2017) For any pair of sequences $\{a_k\}$ and $\{b_n\}$, show that

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{a_n b_m}{n+m} \lesssim \left(\sum_{n=1}^{\infty} a_n^2 \right)^{1/2} \left(\sum_{m=1}^{\infty} b_m^2 \right)^{1/2}.$$

Solution: This is a hard problem, with the simplest proof relying on some clever applications of Cauchy-Schwartz. Fix λ , and write

$$a_{nm} = \frac{a_n}{\sqrt{m+n}} (n/m)^\lambda \quad \text{and} \quad b_{nm} = \frac{b_m}{\sqrt{m+n}} (m/n)^\lambda.$$

Then Cauchy-Schwartz implies that

$$\sum_{n,m} \frac{a_n b_m}{n+m} = \sum_{n,m} a_{nm} b_{nm} \leq \left(\sum_{n,m} a_{nm}^2 \right)^{1/2} \left(\sum_{n,m} b_{nm}^2 \right)^{1/2}.$$

Now

$$\begin{aligned}\sum_{n,m} a_{nm}^2 &= \sum_{n=1}^{\infty} a_n^2 \sum_{m=1}^{\infty} \frac{(n/m)^{2\lambda}}{n+m} \\ &= \sum_{n=1}^{\infty} a_n^2 n^{2\lambda} \sum_{m=1}^{\infty} \frac{1}{m^{2\lambda}(n+m)}.\end{aligned}$$

Now

$$\begin{aligned}\sum_{m=1}^{\infty} \frac{1}{m^{2\lambda}(n+m)} &\lesssim \frac{1}{n} \sum_{m=1}^n \frac{1}{m^{2\lambda}} + \sum_{m=n+1}^{\infty} \frac{1}{m^{1+2\lambda}} \\ &\lesssim \frac{1}{n^{2\lambda}}\end{aligned}$$

Thus

$$\left(\sum_{n,m} a_{nm}^2 \right) \lesssim \sum_{n=1}^{\infty} a_n^2.$$

Similarly, one can show

$$\left(\sum_{n,m} b_{nm}^2 \right) \lesssim \sum_{n=1}^{\infty} b_n^2.$$

Combining these estimates completes the proof.

If one is less sneaky, but knows more analysis, there is an alternate, much more technical proof (using techniques that do not appear often in basic analysis problems). Consider the operator T , mapping a sequence $a(n)$ to a sequence $(Ta)(m)$ such that

$$(Ta)(m) = \sum_{n=1}^{\infty} \frac{a_n}{n+m},$$

The Marcinkiewicz interpolation theorem (i.e. see Section 8 of the first part of Terence Tao's notes for Math 247A) implies that to show T is bounded from $l^2(\mathbf{N})$ to $l^2(\mathbf{N})$, it suffices to show that the operator is of *restricted weak type* from $l^p(\mathbf{N})$ to $l^p(\mathbf{N})$ for *some value* $p < 2$, and *some value* $p > 2$. By restricted weak type, we mean that T satisfies a bound of the form

$$\|T\chi_E\|_{l^{p,\infty}(\mathbf{N})} \lesssim_p N^{1/p}$$

where χ_E is the indicator function of some finite set $E \subset \{1, \dots, \infty\}$ of cardinality N . And by this result, we mean that for any $\alpha > 0$, we have a bound of the form

$$\#\{m : T\chi_E(m) \geq \alpha\} \lesssim_p N/\alpha^p.$$

We will actually find that T is of restricted weak type for any $p \in [1, \infty)$. To obtain this result, it is simple to see that $T\chi_E \leq T\chi_N$, where χ_N is the indicator function of $\{1, \dots, N\}$, and so it suffices to prove that for any $\alpha > 0$, we have a bound

$$\#\{m : T\chi_N(m) \geq \alpha\} \lesssim_p N/\alpha^p.$$

We first claim that for all $\alpha > 0$,

$$\#\{m \geq N : T\chi_N(m) \geq \alpha\} \leq N/\alpha^p.$$

To get this result, we rely on the crude estimate

$$T_{\chi_N}(m) = \sum_{n=1}^N \frac{1}{n+m} \leq N/m.$$

If $\alpha > 1$, then if $m \geq N$, $T_{\chi_N}(m) \leq 1 \leq \alpha$, and thus

$$\#\{m \geq N : T_{\chi_N}(m) \geq \alpha\} = 0 \leq N/\alpha^p$$

For $\alpha \leq 1$,

$$\#\{m \geq N : T_{\chi_N}(m) \geq \alpha\} \leq \#\{m : N \leq m \leq N/\alpha\} \leq N/\alpha \leq N/\alpha^p.$$

Thus it suffices to show that for all $\alpha > 0$,

$$\#\{m < N : T_{\chi_N}(m) \geq \alpha\} \lesssim_p N/\alpha^p.$$

But now we use the more tight estimates, that for $m < N$,

$$T_{\chi_N}(m) \lesssim 1 + \ln(N/m).$$

Thus if $m < N$, and $T_{\chi_N}(m) \geq \alpha$, then $1 + \ln(N/m) \geq c\alpha$ for some small universal constant $c > 0$, which implies that $m \leq Ne^{1-c\alpha}$, which gives that for $\alpha \geq 2/c$,

$$\#\{m < N : T_{\chi_N}(m) \geq \alpha\} \leq Ne^{1-c\alpha} \leq Ne^{-\alpha/2} \lesssim N/\alpha^p.$$

For $\alpha \leq 2/c$,

$$\#\{m < N : T_{\chi_N}(m) \geq \alpha\} \leq N \lesssim N/\alpha^p.$$

This completes the proof that T has a restricted weak type bound for any $1 \leq p < \infty$.

5 Measure Theory Notes

It is useful to keep in mind *Littlewood's Three Principles*:

- Every finite measure set is *nearly* the union of a finite collection of disjoint sets. One instance of this principle is that if $E \subset \mathbb{R}^d$, then for any $\varepsilon > 0$, there is a disjoint family of cubes $\{Q_i\}$ such that $|E \Delta Q_i| \leq \varepsilon$.
- Every measurable function is nearly continuous. One instance of this principle is that $C_0(\mathbb{R}^d)$ is dense in $L^1(\mathbb{R}^d)$. A more technical instance is *Lusin's theorem*, that for every measurable function $f : E_0 \rightarrow \mathbb{C}$, where $|E_0| < \infty$, and for any $\varepsilon > 0$, there is $E \subset E_0$ with $|E \Delta E_0| \leq \varepsilon$, such that $f|_E$ is a continuous function.
- Every almost everywhere convergent sequence of measurable functions is nearly uniformly convergent. One instance of this principle is *Egorov's theorem*, that if $\{f_n\}$ is a sequence of measurable functions on a finite measure set $E_0 \subset \mathbb{R}^d$ converging pointwise to some function $f : E_0 \rightarrow \mathbb{C}$, then for any $\varepsilon > 0$, we can find $E \subset E_0$ with $|E \Delta E_0| < \varepsilon$ such that $\{f_n\}$ converges uniformly to f on E .

Here are some other results which are often useful:

- The *monotone convergence theorem*: If $\{f_n\}$ is a monotone sequence of non-negative measurable functions converging to some function f pointwise almost everywhere as $n \rightarrow \infty$, then

$$\lim_n \int f_n(x) dx = \int f(x) dx.$$

- The *dominated convergence theorem*: If $\{f_n\}$ is a sequence of measurable functions converging pointwise almost everywhere to some function f , and $|f_n| \leq g$ for some integrable function g , then

$$\lim_n \int f_n(x) dx = \int f(x) dx.$$

- *Fatou's lemma*, which says that if $\{f_n\}$ are a family of non-negative, measurable functions, then

$$\int \left(\liminf_{n \rightarrow \infty} f_n(x) \right) dx \leq \liminf_{n \rightarrow \infty} \left(\int f_n(x) dx \right).$$

- The *Borel-Cantelli Lemma*: If $\{E_n\}$ is a sequence of measurable sets such that

$$\sum_{n=1}^{\infty} |E_n| < \infty,$$

then $\limsup_n E_n = \bigcap_{n=1}^{\infty} \bigcup_{m \geq n} E_m$ is a set of measure zero.

One should also have an aptitude for manipulating sets to calculate their measure, but it is difficult to summarize these techniques here: they come from practice in various problems.

Day 4: Warm Up Problems

6 Day 4: Measure Theory

1. (Spring 2021 and Spring 2016) Let $E \subset \mathbb{R}$ be a Lebesgue measurable set with $|E| < \infty$. Prove that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(r) = |E \cap (E + r)|$ is continuous

Solution: It suffices to prove that for all $s \in \mathbb{R}$,

$$\lim_{r \rightarrow s} f(r) = f(s).$$

We write

$$f(r) = \int \mathbf{I}_E(x) \mathbf{I}_E(x + r) dx.$$

Now

$$f(r) - f(s) = \int \mathbf{I}_E(x) [\mathbf{I}_E(x + r) - \mathbf{I}_E(x + s)] dx.$$

The result then follows from the fact that

$$\lim_{r \rightarrow s} \int |\mathbf{I}_E(x + r) - \mathbf{I}_E(x + s)| dx = 0,$$

because for any $f \in L^1(\mathbb{R})$,

$$\lim_{r \rightarrow 0} \int |f(x + r) - f(x)| dx = 0.$$

To see that this equation is true, we note it is certainly true for any $f \in C_c^\infty(\mathbb{R})$, which is a dense subfamily of $L^1(\mathbb{R})$. But the operators $A_r f(x) = f(x + r) - f(x)$ are uniformly bounded in $L^1(\mathbb{R})$, i.e. $\|A_r f\|_{L^1(\mathbb{R})} \leq 2\|f\|_{L^1(\mathbb{R})}$. And thus the result follows by the uniform boundedness theorem.

Alternatively, write $g = \mathbf{I}_E$. Then g lies in L^1 and L^∞ , and $f(r) = (g * g)(-r)$. But Young's convolution inequality implies that $g * g$ is continuous, so that f is continuous.

2. (Spring 2015) Does there exist a Borel measurable function $f : \mathbb{R} \rightarrow [0, \infty)$ such that

$$\int_a^b f(x) dx = \infty$$

for all real numbers $a < b$. Find an example or show that no such function exists.

Solution: Here I construct such a function. There are many ways of doing this. We note that it suffices to construct such a function $f : [0, 1] \rightarrow [0, \infty)$, since one can piece together a Borel measurable function on \mathbb{R} by patching together translates of this function.

For each rational number $x \in [0, 1]$ expressed in the simplest form as p/q , let $E_x = \{y \in \mathbb{R} : |y - p/q| \leq 1/q^3\}$. Then

$$\sum_{x \in \mathbf{Q} \cap [0, 1]} |E_x| \leq \sum_{q=1}^{\infty} \sum_{p=1}^q 2/q^3 = \sum_{q=1}^{\infty} 2/q^2 < \infty.$$

The Borel-Cantelli lemma implies that $E = \limsup_x E_x$ is a set of measure zero. We set

$$f(x) = \mathbf{I}(x \notin E) \sum_{p/q \in \mathbf{Q} \cap [0, 1]} q^4 \cdot \mathbf{I}(x \in (p/q - 1/q^3, p/q + 1/q^3)).$$

If $x \notin E$, then x lies in at most finitely many of the intervals $(p/q - 1/q^3, p/q + 1/q^3)$, so $f(x)$ is finite. Moreover, f is a limit of simple functions, and is therefore measurable. And if $a < b$, there

exists arbitrarily large denominators q for which there is a simple fraction p/q with

$$a < p/q - 1/q^3 < p/q + 1/q^3 < b.$$

On $(p/q - 1/q^3, p/q + 1/q^3) \cap E^c$, we have $f(x) \geq q^4$, and thus

$$\int_a^b f(x) dx \geq \int_{p/q-1/q^3}^{p/q+1/q^3} q^4 = 2q.$$

Taking $q \rightarrow \infty$ shows that

$$\int_a^b f(x) dx = \infty.$$

3. (Fall 2018) Two parts:

(a) Give an example, with explanation, of each of the following:

- A sequence of functions on \mathbb{R} that converges to zero in $L^1(\mathbb{R})$, but it does not converge almost anywhere on \mathbb{R} to any function.
- A sequence of functions in $L^1(\mathbb{R})$ that converges almost everywhere to zero, but it does not converge in measure to any function.

Solution: For the first part of (a), It suffices to construct such a function on $L^1([0, 1])$, since one can patch together functions onto all of \mathbb{R} . Given $a \in \mathbb{R}$, let $[a] \in [0, 1)$ be the unique value such that $a - [a] \in \mathbb{Z}$. Define

$$H_n = \sum_{k=1}^n \frac{1}{k}.$$

Define $I_n = \{[x] \in [0, 1) : x \in [H_n, H_{n+1}]\}$ and then let $f_n = \chi_{I_n}$. These functions are known as the *typewriter sequence*, for some reason. Then

$$\|f_n\|_{L^1(\mathbb{R})} = H_{n+1} - H_n \lesssim 1/n.$$

Thus the sequence $\{f_n\}$ converges to zero in $L^1(\mathbb{R})$. However, since $H_n \rightarrow \infty$, $\limsup I_n = [0, 1)$, which means that for any $x \in [0, 1)$,

$$\liminf_{n \rightarrow \infty} f_n(x) = 1.$$

Thus for any $x \in [0, 1]$, $\{f_n(x)\}$ does not converge as $n \rightarrow \infty$. Thus the sequence of functions $\{f_n\}$ does not converge pointwise to zero at any point in $[0, 1)$.

For the second part, let $I_n = [n, n+1]$, and let $f_n = \chi_{I_n}$. The sequence $\{f_n\}$ is then the required example. This is the classic “moving bump”.

(b) Prove that a sequence of functions on \mathbb{R} that converges to zero in measure must have a subsequence that converges to zero almost everywhere. Do not quote any theorems that trivialize the problem.

Solution: By definition of convergence in measure, there exists a subsequence (n_k) such that

$$|\{x : f_{n_k}(x) > 2^{-k}\}| < 2^{-k}$$

for every $k \geq 1$. Therefore by the Borel-Cantelli lemma, it follows that

$$|\{x : f_{n_k}(x) > 2^{-k} \text{ i.o.}\}| = 0.$$

In other words, for a.e. $x \in \mathbb{R}$, there exists an integer k_0 (possibly depending on x) such that $f_{n_k}(x) < 2^{-k}$ whenever $k \geq k_0$. Therefore $\lim_{k \rightarrow \infty} f_{n_k}(x) = 0$ a.e.

4. (Spring 2017) Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a continuously differentiable function for which $\|f'\|_\infty < \infty$. Define, for $x > 0$,

$$F(x) = \int_0^\infty f(x + yx)\psi(y) dy,$$

where ψ satisfies

$$\int_0^\infty |\psi(y)| dy \quad \text{and} \quad \int_0^\infty y \cdot |\psi(y)| dy < \infty.$$

Show that $F(x)$ is well defined for all $x \geq 0$, and that F is continuously differentiable.

Solution: The fundamental theorem of calculus implies that

$$|f(x + yx)| = \left| f(x) + \int_x^{x+yx} f'(t) dt \right| = f(x) + O(yx).$$

Thus combined with the fact that $\int_0^\infty |\psi(y)| dy < \infty$ and $\int_0^\infty y \cdot |\psi(y)| dy < \infty$, this implies that

$$\int_0^\infty |f(x + yx)| |\psi(y)| dy \lesssim |f(x)| + |x| < \infty,$$

so F is well defined.

To show that F is continuously differentiable, we note that for a fixed $x \in [0, \infty)$, we calculate that, for h with $0 \leq x + h$,

$$\frac{F(x + h) - F(x)}{h} = \int_0^\infty \frac{|f(x + yx + h(1 + y)) - f(x + yx)|}{h} \psi(y) dy.$$

Now

$$\left| \frac{|f(x + yx + h(1 + y)) - f(x + yx)|}{h} \psi(y) \right| \lesssim (1 + y) |\psi(y)|.$$

Since $(1 + y)\psi(y)$ is integrable, the dominated convergence theorem implies that F is differentiable at x , and

$$F'(x) = \int_0^\infty f'(x + yx)(1 + y)\psi(y) dy.$$

Finally, we show F' is continuous. We note that for any $\varepsilon > 0$, there exists $R > 0$ such that

$$\int_R^\infty (1 + y) |\psi(y)| dy \leq \varepsilon.$$

Since f' is continuous, it is uniformly continuous on $[0, 2R(1 + x)]$. Thus there exists $\delta > 0$ such that for $|h| \leq \delta$, and $0 \leq y \leq R$, $|f'((x + yx) + h(1 + y)) - f'(x + yx)| \leq \varepsilon$, and so

$$\begin{aligned} F'(x + h) - F'(x) &\lesssim \varepsilon + \int_0^R [f'((x + yx) + h(1 + y)) - f'(x + yx)](1 + y)\psi(y) dy \\ &\lesssim \varepsilon + \int_0^R \varepsilon(1 + y)\psi(y) dy \lesssim \varepsilon. \end{aligned}$$

5. (Fall 2016) Let $f : [0, 1] \rightarrow \mathbb{R}$ be continuous with $\min_{0 \leq x \leq 1} f(x) = 0$. Assume that for any $0 \leq a \leq b \leq 1$ we have

$$\int_a^b [f(x) - \min_{a \leq y \leq b} f(y)] dx \leq \frac{|b-a|}{2}.$$

Prove that for any $\lambda \geq 0$, we have

$$|\{x : f(x) > \lambda + 1\}| \leq \frac{1}{2} |\{f(x) > \lambda\}|.$$

Solution: Since f is continuous, $\{x : f(x) > \lambda\}$ is open. Therefore we may write $\{x : f(x) > \lambda\} = \bigcup_{i=1}^{\infty} I_i$ where $(I_i)_{i=1}^{\infty}$ is a countable pairwise disjoint sequence of intervals $I_i = (a_i, b_i)$ with $a_i \leq b_i$.

We claim that for each i ,

$$\inf_{t \in \bar{I}_i} f(t) \leq \lambda.$$

If $I_i = \emptyset$, there is nothing to show. If $a_i = 0$ and $b_i = 1$, then by hypothesis, $\min_{[a_i, b_i]} f = \min_{[0, 1]} f = 0 \leq \lambda$. So assume that is not the case; i.e., that at least one of a_i, b_i is in $(0, 1)$. Without loss of generality, assume that $a_i \in (0, 1)$. Then $a_i \in \partial \{x : f(x) \leq \lambda\}$. Hence by continuity of f , we have $f(a_i) \leq \lambda$.

Now that this has been proved, we have:

$$\begin{aligned} \frac{1}{2} |I_i| &\geq \int_{I_i} \left\{ f(x) - \min_{t \in \bar{I}_i} f(t) \right\} dx && \text{by assumption} \\ &\geq \int_{I_i \cap \{f > \lambda + 1\}} \left\{ f(x) - \min_{t \in \bar{I}_i} f(t) \right\} dx && \text{by nonnegative of the integrand} \\ &\geq \int_{I_i \cap \{f > \lambda + 1\}} (f(x) - \lambda) dx && \text{by the claim} \\ &\geq \int_{I_i \cap \{f > \lambda + 1\}} ((\lambda + 1) - \lambda) dx && \text{by domain of integration} \\ &= |I_i \cap \{f > \lambda + 1\}|. \end{aligned}$$

By countable additivity for disjoint sets, summing over i and using $\{f > \lambda + 1\} \subseteq \{f > \lambda\}$ gives

$$\frac{1}{2} |\{f > \lambda\}| \geq |\{f > \lambda\} \cap \{f > \lambda + 1\}| = |\{f > \lambda + 1\}|.$$

7 Functional Analysis Notes

s

Day 5: Warm Up Problems

6. (Spring 2015) Let $f \in L^2[0, 1]$ satisfy $\int_0^1 t^n f(t) dt = (n+2)^{-1}$ for $n = 0, 1, \dots$. Must then $f(t) = t$ for almost every $t \in [0, 1]$?

Solution: The identity $\int_0^1 t^n f(t) dt = (n+2)^{-1}$ shows that for any polynomial $p(t)$,

$$\int_0^1 p(t)[f(t) - t] dt = 0.$$

Applying some type of density theorem (the Stone-Weierstrass theorem, and its variants), we find that the family of all polynomials is dense in $L^2[0, 1]$. Applying the continuity of the map $g \mapsto \int_0^1 g(t)[f(t) - t]$, we conclude that for all $g \in L^2[0, 1]$,

$$\int_0^1 g(t)[f(t) - t] dt = 0.$$

But setting $g(t) = \overline{f(t) - t}$, we conclude that

$$\int_0^1 |f(t) - t|^2 dt = 0.$$

This implies $f(t) = t$ for almost every t .

7. (Fall 2021) Let $\{f_n\}$ be a sequence of monotonic functions on $[0, 1]$ converging to a function f in measure. Show that f coincides almost everywhere with a monotonic function f_0 , and that $f_n(x) \rightarrow f_0(x)$ at every point of continuity of f_0 .

Solution: If the sequence $\{f_n\}$ converges uniformly, then for each $0 \leq x < y \leq 1$, we would have

$$f(y) - f(x) = \lim_n f_n(y) - f_n(x) \geq 0.$$

Thus f is monotonic. This sequence does *not* converge uniformly. But we utilize one of *Littlewood's three principles*, namely, that every convergent sequence of functions is nearly uniformly convergent.

We do this by first giving us a ε of room. It suffices to show that for each ε , there exists a set $E(\varepsilon)$ such that if $x, y \in E(\varepsilon)$ and $x < y$, then

$$f(y) - f(x) \geq -2\varepsilon.$$

If we set

$$E_n(\varepsilon) = \{x \in [0, 1] : |f_n(x) - f(x)| \geq \varepsilon\}$$

then $|E_n(\varepsilon)| \rightarrow 0$. If $x < y$, with $x, y \in E_n$ for some n , then

$$f(y) - f(x) \geq [f_n(y) - f_n(x)] - 2\varepsilon \geq -2\varepsilon.$$

Thus if $E(\varepsilon) = \bigcup_n E_n(\varepsilon)$, then $|E(\varepsilon)^c| = 0$, and for $x, y \in E(\varepsilon)$ with $x < y$,

$$f(y) - f(x) \geq -2\varepsilon.$$

The set $E = \bigcap_k E(1/2^k)$ is also a set with $|E^c| = 0$, and for any $x, y \in E$ with $x < y$,

$$f(y) - f(x) \geq -2/2^k$$

for all $k > 0$, so taking $k \rightarrow \infty$ gives

$$f(y) - f(x) \geq 0.$$

Thus f is monotonic when restricted to E , and thus agrees with a monotonic function $f_0 : [0, 1] \rightarrow \mathbb{R}$ on E .

It now suffices to show that $f_n(x_0) \rightarrow f_0(x_0)$, whenever x_0 is a continuity point of f_0 . Here we use the same Littlewood's principle more explicitly, i.e. using Egorov's theorem: for any $\delta > 0$, there is $E_\delta \subset [0, 1]$ with $|E_\delta|^c < \delta/2$, such that $f_n \rightarrow f_0$ uniformly on E_δ . If x_0 is a continuity point of f_0 , then for any $\varepsilon > 0$, there exists $\delta > 0$ such that if $|x - x_0| \leq \delta$, $|f(x) - f(x_0)| \leq \varepsilon$. Since $|E_\delta|^c < \delta/2$, there must exist

$$x_0 - \delta < x_1 < x_0 < x_2 < x_0 + \delta$$

with $x_1, x_2 \in E_\delta$. But this implies that

$$\limsup_n f_n(x_0) \leq \limsup_n f_n(x_2) = f_0(x_2) \leq f_0(x_0) + \varepsilon$$

and

$$\liminf_n f_n(x_0) \geq \liminf_n f_n(x_1) = f_0(x_1) \geq f_0(x_0) - \varepsilon.$$

Since ε was arbitrary, it is clear from these inequalities that $f_n(x_0)$ converges to $f_0(x_0)$.

8 Day 5: Functional Analysis

8. (Fall 2015) Find all $f \in L^2[0, \pi]$ such that

$$\int_0^\pi |f(x) - \sin x|^2 dx \leq \frac{4\pi}{9}$$

and

$$\int_0^\pi |f(x) - \cos x|^2 dx \leq \frac{\pi}{9}$$

Solution: We note that $\sin x$ and $\cos x$ are orthogonal in $L^2[0, 1]$. Thus we may consider a family of orthogonal vectors $\{e_n\}$ such that $\{\sin x, \cos x\} \cup \{e_n\}$ is an orthogonal basis for $L^2[0, 1]$. Thus there exists constants a , b , and $\{c_n\}$ such that

$$f(x) = a \sin x + b \cos x + \sum c_n e_n,$$

where the convergence of this sum is in the L^2 norm. Applying Parseval's theorem, we find that

$$\begin{aligned} \int_0^\pi |f(x) - \sin x|^2 dx &= |a - 1|^2 \int_0^\pi |\sin x|^2 dx + |b|^2 \int_0^\pi |\cos x|^2 dx + \sum |c_n|^2 \\ &= |a - 1|^2(\pi/2) + |b|^2(\pi/2) + \sum |c_n|^2. \end{aligned}$$

Similarly,

$$\int_0^\pi |f(x) - \cos x|^2 dx = |a|^2(\pi/2) + |b - 1|^2(\pi/2) + \sum_n |c_n|^2.$$

Next, we note that

$$\int_0^\pi |\sin x - \cos x|^2 dx = \pi.$$

The triangle inequality thus implies that for any f satisfying the requirements above, we actually have *equality*, i.e.

$$\int_0^\pi |f(x) - \sin x|^2 dx = \frac{4\pi}{9}$$

and

$$\int_0^\pi |f(x) - \cos x|^2 dx = \frac{\pi}{9}$$

Thus we must find all families of coefficients a , b , and $\{c_n\}$ such that

$$|a - 1|^2(\pi/2) + |b|^2(\pi/2) + \sum |c_n|^2 = 4\pi/9$$

and

$$|a|^2(\pi/2) + |b - 1|^2(\pi/2) + \sum |c_n|^2 = \pi/9.$$

Subtracting the second identity from the first, then multiplying by $2/\pi$, we conclude that

$$(|a - 1|^2 - |a|^2) + (|b|^2 - |b - 1|^2) = 2/3.$$

This equation can be simplified to

$$\operatorname{Re}(b) = \operatorname{Re}(a) + 1/3.$$

Substituting this into the first equation, we find that if $a = x + iy$, $b = (x + 1/3) + iz$, and $C = 2/\pi \sum c_n^2$, then

$$2x^2 - (4/3)x + (y^2 + z^2 + 2/9 + C) = 0.$$

The quadratic formula implies that there exists x satisfying this equation for a fixed y and z if

$$16/9 - 8(y^2 + z^2 + 2/9 + C) \geq 0,$$

which can be rearranged to read

$$8(y^2 + z^2 + C) \leq 0.$$

This can only occur if $y = z = C = 0$, and it then follows from the quadratic formula that $x = 1/6$. Thus the only function f satisfying this result is the function

$$f(x) = (1/6) \sin x + (1/2) \cos x.$$

This completes the argument.

Alternatively, let $u(x) = \sin(x) - f(x)$ and $v(x) = f(x) - \cos(x)$. Then

$$\|u + v\|_{L^2}^2 = \int_0^\pi (\sin(x) + \cos(x))^2 dx = \int_0^\pi \sin^2(x) + \cos^2(x) dx = \pi.$$

By the triangle inequality and the hypotheses of the problem,

$$\sqrt{\pi} = \|u + v\|_{L^2} \leq \|u\|_{L^2} + \|v\|_{L^2} \leq \frac{2}{3}\sqrt{\pi} + \frac{1}{3}\sqrt{\pi} = \sqrt{\pi}.$$

Therefore $\|u + v\|_{L^2} = \|u\|_{L^2} + \|v\|_{L^2}$. Therefore

$$(\|u\|_{L^2} + \|v\|_{L^2})^2 = \|u + v\|_{L^2}^2 = \langle u + v, u + v \rangle = \|u\|_{L^2}^2 + \|v\|_{L^2}^2 + 2\langle u, v \rangle.$$

Therefore since $(\|u\|_{L^2} + \|v\|_{L^2})^2 = \|u\|_{L^2}^2 + \|v\|_{L^2}^2 + 2\|u\|_{L^2}\|v\|_{L^2}$, it follows that $\|u\|_{L^2}\|v\|_{L^2} = \langle u, v \rangle$. Therefore by the Cauchy Schwarz inequality, u and v are linearly dependent (viz. the condition for equality in the statement of Cauchy Schwarz). That is, there exists $\alpha \in \mathbb{R}$ such that $\sin(x) - f(x) = \alpha(f(x) - \cos(x))$ almost everywhere. Solving for f gives

$$f(x) = \frac{1}{1 + \alpha} \sin(x) + \frac{\alpha}{1 + \alpha} \cos(x). \quad (7)$$

It remains to determine α . Since $\|u\|_{L^2} + \|v\|_{L^2} = \sqrt{\pi}$, $\|u\|_{L^2} \leq 2\sqrt{\pi}/3$, and $\|v\|_{L^2} \leq \sqrt{\pi}/3$, it follows that $\|u\|_{L^2}^2 = 4\pi/9$. Therefore

$$\begin{aligned} \frac{4}{9}\pi &= \int_0^\pi \left(\sin(x) - \frac{1}{1 + \alpha} \sin(x) - \frac{\alpha}{1 + \alpha} \cos(x) \right)^2 dx \\ &= \left(\frac{\alpha}{1 + \alpha} \right)^2 \int_0^\pi (\sin(x) - \cos(x))^2 dx \\ &= \left(\frac{\alpha}{1 + \alpha} \right)^2 \pi. \end{aligned}$$

This implies $\alpha = 2$ or $\alpha = -2/5$. A similar calculation using $\|v\|_{L^2}^2 = \pi/9$ implies that $\alpha = 2$ or $\alpha = -4$. The only value for α consistent with both of these conditions is $\alpha = 2$. Therefore by (7),

$$f(x) = \frac{1}{3} \sin(x) + \frac{2}{3} \cos(x).$$

9. (Spring 2017) Let $l^1(\mathbf{N})$ be the space of summable sequences, i.e.

$$l^1(\mathbf{N}) = \{x : \sum_{n=1}^{\infty} |x_n| < \infty\}.$$

Let $\{a_n\}$ be a sequence with $a_n \geq 0$ for all $n \in \mathbf{N}$ and consider the subset $K \subset l^1(\mathbf{N})$ defined by

$$K = \{x \in l^1(\mathbf{N}) : 0 \leq x_n \leq a_n \text{ for all } n\}.$$

Show that K is compact if and only if the sequence $\{a_n\}$ itself belongs to $l^1(\mathbf{N})$.

Solution: First suppose K is compact. Then consider the family of sequences

$$\{x(k) = (a_1, \dots, a_k, 0, \dots)\}.$$

Each sequence $x(k)$ belongs to K . Thus by compactness, this sequence has a convergent subsequence $x(k_i)$ converging to some sequence in $l^1(\mathbf{N})$. But $x(k_i)$ converges pointwise to a , and so $x(k_i)$ can only converge to a , which implies $a \in l^1(\mathbf{N})$.

On the other hand, suppose $a \in l^1(\mathbf{N})$. Consider an arbitrary sequence $\{x(k)\}$ in K . For any $N > 0$, the projection of K onto the first N elements of the sequence is compact, and thus we may inductively find subsequences $x(k_{N,i})$ for each N such that $\{k_{N,i}\}$ is a subsequence of $\{k_{M,i}\}$ for each $M \leq N$, and the first N elements of the sequences $x(k_{N,i})$ converge pointwise. Consider the subsequence of sequences $y(n) = x(k_{n,n})$. For each $N > 0$, this sequence is a subsequence of the sequence $\{x(k_{N,i})\}$ and so we can conclude from this that $\{y(n)\}$ converges pointwise to some sequence y . Since K is defined pointwise, it is simple to see that $y \in K$. For each $\varepsilon > 0$, we pick $N > 0$ such that

$$\sum_{k>N} a_k < \varepsilon.$$

Then exists M such that for $n > M$, and $1 \leq k \leq N$, $|y(n)_k - y_k| \leq \varepsilon/N$. And thus

$$\|y(n) - y\|_1 \leq \sum_{k=1}^N |y(n)_k - y_k| + \sum_{n>N} |y(n)_k| + |y_k| \leq \sum_{k=1}^N (\varepsilon/N) + \sum_{n>N} 2a_n \leq 3\varepsilon.$$

Taking $\varepsilon \rightarrow 0$ completes the proof.

10. (Spring 2018, Spring 2021) Let K be a continuous function on $[0, 1] \times [0, 1]$. Suppose that g is a continuous function on $[0, 1]$. Show that there exists a unique continuous function f on $[0, 1]$ such that

$$f(x) = g(x) + \int_0^x f(y)K(x, y)dy$$

Solution: Define an integral operator T by setting

$$Tf(x) = g(x) + \int_0^x f(y)K(x, y)dy.$$

Since g is continuous, $T : X \rightarrow X$ where $X = C[0, 1]$, a Banach space with respect to the norm $\|f\|_{L^\infty[0, 1]} = \max_{x \in [0, 1]} |f(x)|$. It suffices to show there is a unique f such that $Tf = f$. Since

K is continuous and $[0, 1] \times [0, 1]$ is compact, there exists M such that $|K(x, y)| \leq M$ for all $x, y \in [0, 1] \times [0, 1]$. By induction we can show that for any $f_1, f_2 \in C[0, 1]$ and any $x \in [0, 1]$,

$$\begin{aligned} |T^n f_1(x) - T^n f_2(x)| &\leq \|f_1 - f_2\|_{L^\infty[0,1]} M^n \int_0^x \int_0^{x_1} \cdots \int_0^{x_{n-1}} x_n dx_n \cdots dx_1 \\ &= \|f_1 - f_2\|_{L^\infty[0,1]} \frac{M^n x^n}{n!} \\ &\leq \|f_1 - f_2\|_{L^\infty[0,1]} \frac{M^n}{n!}. \end{aligned}$$

Thus for sufficiently large n , we conclude T^n is a contraction map. Thus by the Banach fixed point theorem, we conclude that there exists a unique $f \in C[0, 1]$ such that $T^n f = f$. Clearly, this shows that there exists at most one f such that $Tf = f$, since if $Tf = f$, then $T^n f = f$. TODO: Does it show existence?

In the previous solution, we wished to apply the contraction mapping theorem, but could not do so directly with T because T is not a contraction on $C[0, 1]$. We got around this difficulty by considering the operator T^n for some positive integer n , which fortuitously turned out to be a contraction when n is large enough. Another approach is to consider T as an operator on “smaller” Banach spaces, e.g. $C[0, x_1], C[x_1, x_2], \dots$ etc, where $x_i - x_{i-1}$ are small enough that T is a contraction on $C[x_{i-1}, x_i]$, and then patch the solutions together. We detail this second proof here. If $|K| \leq M$ for all x, y , define

$$T_{x_0 x_1}(f) = g(x) + \int_0^x f(y)K(x, y)dy$$

on $C([x_0, x_1])$. Then

$$\|T_{x_0 x_1} f_1 - T_{x_0 x_1} f_2\|_{L^\infty[x_0, x_1]} \leq \int_{x_0}^{x_1} |K(x, y)| dy \cdot \|f_1 - f_2\|_{L^\infty[x_0, x_1]} \leq M(x_1 - x_0) \|f_1 - f_2\|_{L^\infty[x_0, x_1]}.$$

Thus if $x_1 - x_0 < 1/2M$, then we conclude that $T_{x_0 x_1}$ is a contraction map on $C([x_0, x_1])$. This means that for any such interval, there exists a unique $f_{x_0 x_1} \in C[x_0, x_1]$ such that $T_{x_0 x_1} f = f$. But clearly if f satisfies this property, then its restriction to any subinterval also satisfies this property. This implies that any two solutions $f_{x_0 x_1}$ and $f_{x_2 x_3}$ agree on $[x_0, x_1] \cap [x_2, x_3]$. But the locality of continuity means we can patch these functions together to find $f \in C[0, 1]$ such that $T_{x_0 x_1} f = f$ for any x_0 and x_1 with $x_1 - x_0 < 1/2M$. But this implies that $Tf = f$.

11. Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function with compact support.

(a) Prove that there exists a constant A such that

$$\|f * \phi\|_q \leq A \|f\|_p$$

for $1 \leq p \leq q \leq \infty$.

Solution: Let us begin by showing that

$$\|f * \phi\|_\infty \leq A \|f\|_p.$$

Indeed, Hölder's inequality implies that if p^* is the conjugate to p , then

$$\begin{aligned}\|f * \phi\|_{L_x^\infty} &= \left\| \int f(x-y)\phi(y) dy \right\|_{L_x^\infty} \\ &\leq \left\| \left(\int |\phi(y)|^{p^*} dy \right)^{1/p^*} \cdot \left(\int |f(x-y)|^p dy \right)^{1/p} \right\|_{L_x^\infty} \\ &\leq \|\phi\|_{L^{p^*}} \|f\|_{L^p}.\end{aligned}$$

Next, we calculate using Minkowski's integral inequality that

$$\|f * \phi\|_{L_x^p} = \left\| \int f(x-y)\phi(y) dy \right\|_{L_x^p} \leq \int |\phi(y)| \|f\|_{L^p} dy = \|\phi\|_{L^1} \|f\|_{L^p}.$$

Thus we have found a constant $A = \max_{1 \leq r \leq \infty} (\|\phi\|_{L^r})$ such that the result holds for (p, q) of the form (p, ∞) , and of the form (p, p) . The remaining estimates lie 'in-between' this range. One approach here is to apply the Riesz-Thorin interpolation theorem, which gives all the intermediary bounds (p, q) for $p \leq q \leq \infty$. Alternatively, a sneaky application of Hölder's inequality shows that for $q \geq p$, if we fix $\alpha \in (0, 1)$ such that $\alpha q = p$, then

$$\begin{aligned}\|f * \phi\|_{L_x^q}^q &= \int |(f * \phi)(x)|^{\alpha q} |(f * \phi)(x)|^{(1-\alpha)q} dx \\ &\leq \left(\int |(f * \phi)(x)|^p |(f * \phi)(x)|^{(1-\alpha)q} dx \right) \\ &\leq \int |(f * \phi)(x)|^p dx \cdot \sup_x |(f * \phi)(x)|^{(1-\alpha)q} \\ &\leq \|f * \phi\|_{L_x^p}^p \|f * \phi\|_{L_x^\infty}^{(1-\alpha)q} \\ &\leq (A\|f\|_{L^p})^p (A\|f\|_{L^p})^{(1-\alpha)q} \\ &= A\|f\|_{L^p}^q.\end{aligned}$$

- (b) Show by example that such general inequality cannot hold for $p > q$.

Solution: It suffices, for any $R > 0$, to construct a function $f \in L^p$ such that

$$\|f * \phi\|_{L^q} \geq R\|f\|_{L^p}.$$

One should think of bounds on L^r for large r as controlling how 'peaked' a function can be, whereas bounds on L^r for small r control how 'long' a function can be. Thus, intuitively, we must construct a function f which is not peaked, but can be long (small in L^p norm), into a function $f * \phi$ which is long (has large L^q norm).

Here is one approach that 'almost' works for all ϕ . A good choice for a non-peaked, long function is just a constant function supported on a large set. For instance, if $f(x) = \mathbf{I}(|x| \leq R)$, then $\|f\|_p \sim R^{1/p}$. If we assume that ϕ is supported on $|x| \leq S$, then for $|x| \leq R - S$,

$$(\phi * f)(x) = \int \phi(x) dx.$$

Provided that $\int \phi(x) dx \neq 0$, this implies that $|(\phi * f)(x)| \gtrsim_\phi 1$ for $|x| \leq R - S$, which implies

that for $R \geq 2S$,

$$\|\phi * f\|_{L^q} \gtrsim_\phi \left(\int_{|x| \leq R-S} 1 \right)^{1/q} \gtrsim (R-S)^{1/q} \gtrsim R^{1/q}.$$

Thus

$$\|\phi * f\|_{L^q} \gtrsim_\phi R^{1/q-1/p} \|f\|_{L^p}.$$

If $q < p$, then $1/q - 1/p > 0$, and so we may take $R \rightarrow \infty$ to show that the operator $Tf = \phi * f$ cannot be bounded. The only problem is that this idea cannot approach functions such that

$$\int \phi(x) dx = 0.$$

If this is the case, in our examples above, $f * \phi = 0$ on $|x| \leq R - S$, which means $\|f * \phi\|_q \sim 1$, so this approach doesn't work. In order to approach this case, we must choose f more carefully so that $f * \phi$ cannot average out to zero on a large set.

Fix a large integer $N > 0$, and set

$$f(x) = \sum_{n=-N}^N \overline{\phi(2R \cdot n - x)},$$

where we assume $\phi(x) = 0$ for $|x| \geq R$. Then f is not 'peaked' (because ϕ is continuous, and not peaked), but supported on a very wide set. Indeed, we find that

$$\|f\|_p \sim_\phi N^{1/p}.$$

For $|n| \leq N$, we compute that

$$(\phi * f)(2Rn) = \int \phi(x) f(2Rn - x) dx = \int \phi(x) \overline{\phi(x)} dx = \|\phi\|_{L^1}.$$

By continuity of the quantities involved, there exists δ such that for $|x| \leq \delta$, $|(\phi * f)(x)| \geq \|\phi\|_{L^1}/2$. Thus $\phi * f$ is non-zero on a very wide set. More precisely, $|(\phi * f)| \gtrsim \|\phi\|_1$ on N intervals of radius δ , so

$$\|\phi * f\|_q \gtrsim (N\delta)^{1/q} \|\phi\|_{L^1} \gtrsim_\delta N^{1/q}.$$

Thus

$$\|\phi * f\|_q \gtrsim_{\phi, \delta} N^{1/q-1/p} \|f\|_p.$$

Taking $N \rightarrow \infty$ thus shows that the operator is unbounded.

12. (Fall 2016) Give an example of a non-empty closed subset of $L^2([0, 1])$ that does not contain a vector of smallest norm. Prove your assertion.

Solution: Let $\{e_n\}$ be an orthonormal basis for $L^2[0, 1]$, and define

$$x_n = (1 + 1/n)e_n.$$

Then $\{x_n\}$ is a closed subset of $L^2[0, 1]$ (it is actually *discrete*), and has no element of smallest norm.

Day 6: Warm Up Problems

9 Day 6: Functional Analysis

13. (Fall 2019) Show that there is no sequence $\{a_n\}$ of positive numbers such that $\sum a_n |c_n| < \infty$ if and only if $\{c_n\}$ is a bounded sequence. Hint: Suppose that there exists a sequence and consider the map $T : l^\infty \rightarrow l^1$ given by $Tf(n) = a_n f(n)$. The set of f such that $f(n) = 0$ for all but finitely many n is dense in l^1 but not in l^∞ .

Solution: Consider the operator T defined in the hint. Clearly T is linear. Also, T is injective, since all elements of the sequence $\{a_n\}$ are positive. We also claim that T is surjective, given the property assumed of the sequence. Indeed, if $\sum b_n < \infty$, and we define $c_n = b_n/a_n$, then $\sum a_n |c_n| < \infty$, and thus $\{c_n\}$ is a bounded sequence, and $Tc = b$. Finally, we claim that T is bounded. Indeed, we have

$$\|Tf\|_{l^1} = \sum |Tf(n)| = \sum |a_n| |f(n)| \leq \left(\sum |a_n| \right) \|f\|_{l^\infty}.$$

Thus we conclude that T is an isomorphism (a consequence of the open mapping theorem is that any bounded, bijective linear operator is an isomorphism). This means that there exists constants $C_1, C_2 > 0$ such that $C_1 \|c\|_{l^\infty} \leq \|Tc\|_{l^1} \leq C_2 \|c\|_{l^\infty}$.

This leads to a contradiction via one of several approaches. Plugging in the bounded sequence $(1, 1, 1, \dots)$ into the operator T gives the sequence $\{a_n\}$, and so this sequence lies in l^1 . Clearly any sequence $\{a_n\}$ in l^1 will give rise to a bounded operator from l^∞ to l^1 , so we wish to disprove that the lower bound exists, i.e. there does not exist a sequence $\{a_n\}$ in l^1 such that $\|Tc\|_{l^1} \gtrsim \|c\|_{l^\infty}$.

To do this, we must construct a non-negative sequence c where $\|c\|_{l^\infty}$ is large, but such that $\|Tc\|_{l^1}$ is small. The best way to do this is to concentrate the sequence c at a single point, i.e. we consider the sequence $c = (0, \dots, 0, 1, \dots)$, where the 1 is in the N th term in the sequence. Then $Tc = (0, \dots, 0, a_N, \dots)$, and so we have

$$\|c\|_{l^\infty} = 1 \quad \text{and} \quad \|Tc\|_{l^1} = |a_N|.$$

As $N \rightarrow \infty$, $|a_N| \rightarrow 0$ since the sequence is integrable. But this is impossible since the bound $\|Tc\|_{l^1} \gtrsim \|c\|_{l^\infty}$ implies that $|a_N| \gtrsim 1$ for all N .

Alternatively, it suffices to argue more abstractly that l^1 and l^∞ cannot be isomorphic to one another. For instance, one can show that l^1 is separable, but l^∞ is not separable, so the spaces cannot be isomorphic.

14. (Fall 2020)

Suppose that X, Y and Z are Banach spaces, and $T : X \times Y \rightarrow Z$ is a mapping such that:

- (a) For each $x \in X$, the map $y \mapsto T(x, y)$ is a bounded linear map $Y \rightarrow Z$.
- (b) For each $y \in Y$, the map $x \mapsto T(x, y)$ is a bounded linear map $X \rightarrow Z$.

Prove there exists a constant C such that

$$\|T(x, y)\|_Z \leq C \|x\|_X \|y\|_Y$$

Solution: Without loss of generality we may assume that x, y are nonzero since if either $x = 0$ or $y = 0$, the inequality holds for any C . Further, we claim that it suffices to show the inequality holds

for all x, y satisfying $\|x\|_X = 1$ and $\|y\|_Y = 1$. To see this, suppose $\|T(\tilde{x}, \tilde{y})\|_Z \leq C$ for all $\tilde{x} \in X$ and $\tilde{y} \in Y$ with $\|\tilde{x}\|_X = 1, \|\tilde{y}\|_Y = 1$. Then for any nonzero $x \in X, y \in Y$,

$$\|T(x, y)\|_Z = \|x\|_X \|y\|_Y \left\| T\left(\frac{x}{\|x\|_X}, \frac{y}{\|y\|_Y}\right) \right\|_Z \leq C \|x\|_X \|y\|_Y.$$

This proves the claim.

Next, let for each $x \in X$, let $\phi_x : Y \rightarrow Z$ denote the map $y \mapsto T(x, y)$, and for each $y \in Y$, let $\psi_y : X \rightarrow Z$ denote the map $x \mapsto T(x, y)$. Let $B = \{x \in X : \|x\|_X = 1\}$. Then since $\phi_x(y) = T(x, y) = \psi_y(x)$ and since ψ_y is assumed to be a bounded operator, it holds for all $y \in Y$ that

$$\sup_{x \in B} \|\phi_x(y)\|_Z = \sup_{x \in B} \|\psi_y(x)\|_Z \leq \sup_{x \in B} \|\psi_y\| \|x\|_X = \|\psi_y\| < \infty.$$

Therefore by the Uniform Boundedness Principle,

$$\sup_{x \in B} \|\phi_x\| = C < \infty.$$

Therefore for any x, y with $\|x\|_X = 1$ and $\|y\|_Y = 1$,

$$\|T(x, y)\|_Z = \|\phi_x(y)\|_Z \leq \|\phi_x\| \|y\|_Y = \|\phi_x\| \leq C.$$

15. (Fall 2015) For $p \in (1, \infty)$, and for $f \in L^p(\mathbb{R})$ define

$$Tf(x) = \int_0^1 f(x+y) dy.$$

(a) Show that $\|Tf\|_p \leq \|f\|_p$ and equality holds if and only if $f = 0$ almost everywhere.

Solution: We use Minkowski's inequality. For those unfamiliar, recall the triangle inequality, which for $1 \leq p \leq \infty$ says that

$$\left\| \sum_n f_n \right\|_p \leq \sum_n \|f_n\|_p.$$

Minkowski's inequality occurs when we replace the sum with an integral, i.e. for $1 \leq p \leq \infty$,

$$\left\| \int f_y dy \right\|_p \leq \int \|f_y\|_p dy.$$

In this case, we set $f_y(x) = f(x+y)$. Then

$$Tf(x) = \int_0^1 f_y(x) dy.$$

Thus, using the fact that $\|f_y\|_p = \|f\|_p$, we conclude

$$\|Tf\|_p = \left\| \int_0^1 f_y dy \right\|_p \leq \int_0^1 \|f_y\|_p dy = \int_0^1 \|f\|_p dy = \|f\|_p.$$

To address the case of equality, we recall the case of the triangle inequality, i.e. that

$$\left\| \sum_n f_n \right\|_p = \sum_n \|f_n\|_p$$

if and only if $\{f_n\}$ are all scalar multiples of one another. Similarly, *Minkowski's inequality* is an equality if and only if $\{f_y\}$ are all scalar multiples of one another *almost everywhere*. Thus if $\|Tf\|_p = \|f\|_p$, then the application of Minkowski's inequality is an equality, which implies that there exists $c(y)$ such that $f_y(x) = c(y)f(x)$, i.e. $f(x+y) = c(y)f(x)$.

Now suppose that

$$\int_{-\infty}^{\infty} |f(x)|^p dx \neq 0.$$

Then we can find some $x_0 \in \mathbb{R}$ with $f(x_0) \neq 0$. The dominated convergence theorem implies that for any $x \in \mathbb{R}$,

$$\begin{aligned} \|f\|_p^p &= \lim_{n \rightarrow \infty} \sum_{k=-n}^n \int_0^1 |f(x_0 + ky)|^p dy \\ &= \lim_{n \rightarrow \infty} |f(x_0)|^p \int_0^1 \sum_{k=-n}^n |c(y)|^k dy. \end{aligned}$$

The sum

$$\sum_{k=-n}^n |c(y)|^k$$

converges monotonically in n to ∞ everywhere except when $c(y) = 0$, where it is constantly zero in n . Thus by the monotone convergence theorem, if c does not equal zero almost everywhere on $[0, 1]$, then

$$\|f\|_p^p = \infty,$$

which gives a contradiction. Thus $c = 0$ almost everywhere on $[0, 1]$, and so as a result, $f = 0$ almost everywhere.

- (b) (Fall 2015) Prove that the map $f \mapsto Tf - f$ does not map $L^p(\mathbb{R})$ onto $L^p(\mathbb{R})$.

Solution: First, we claim that $Tf - f$ is injective. Indeed, if $Tf = f$, then $\|Tf\|_p = \|f\|_p$, and by the last part of the problem, this implies that $f = 0$. If $Tf - f$ was surjective, then the open mapping theorem would therefore imply that there would exist a constant $C > 0$ such that for all $f \in L^p(\mathbb{R})$, $\|Tf - f\|_p \geq C\|f\|_p$. Our goal is to show this is not true.

Intuitively, we wish to find a function f such that $Tf - f$ is small. This would be possible if $Tf \approx f$, which would hold if f was approximately constant. So we consider the ‘almost’ constant function $f(x) = \mathbf{I}(|x| \leq R)$. Then $Tf(x)$ is supported on $|x| \leq R + 1$, $Tf(x) = f(x)$ for $|x| \leq R - 1$, and $\|Tf\|_\infty \leq 1$. Thus $Tf - f$ is supported on a set of total mass $O(1)$, and $\|Tf - f\|_\infty = O(1)$, so

$$\|Tf - f\|_p \lesssim 1,$$

independently of R . On the other hand, $\|f\|_p \sim R^{1/p}$, so it is impossible for a bound of the form $\|Tf - f\|_p \geq C\|f\|_p$ to hold.

16. (Fall 2014)

- (a) For any $n \geq 1$ an integer, there exists two positive measures μ_1^n, μ_2^n supported on $[0, 1]$ such that for any polynomial $P(x)$ with $\deg P(x) \leq n$ it holds:

$$P'(0) = \int_0^1 P(x) d\mu_1^n(x) - \int_0^1 P(x) d\mu_2^n(x).$$

Solution: Let X_n be the space of all real-valued polynomials on $[0, 1]$ with degree at most n . Then X_n is a finite dimensional linear subspace of $C[0, 1]$. Let $T : X_n \rightarrow \mathbb{R}$ be the operator $TP = P'(0)$. Then T is bounded on X_n since X_n is finite dimensional. Thus by the Hahn-Banach theorem, we can find an extension of T to a bounded functional on $C[0, 1]$. The *Reisz Representation Theorem* says that $C[0, 1]^*$ is the space of all finite signed Borel measures on $[0, 1]$. Thus there exists a finite signed Borel measure μ on $[0, 1]$ such that for any $f \in C[0, 1]$,

$$Tf = \int f(x) d\mu(x).$$

In particular, if $P \in X_n$,

$$P'(0) = \int P(x) d\mu(x).$$

Then we perform a *Jordan decomposition* to write $\mu = \mu_1 - \mu_2$, where μ_1 and μ_2 are finite non-negative Borel measures, and this completes the construction.

One can also define the measures explicitly, though the argument becomes more technical. This is Chun's solution from the 2016 SEP, problem 39. We will show that there exists constants $c_j \in \mathbb{R}, x_j \in (0, 1)$ with $0 \leq j \leq n$ such that the measure μ^n defined by

$$\mu^n := \sum_{j=0}^n c_j \delta_{x_j} \quad (8)$$

satisfies

$$\int_0^1 P(x) d\mu^n(x) = P'(0)$$

for any polynomial $P \in X_n$. The result will then follow by taking

$$\mu_1^n = \sum_{j=0}^n c_j^+ \delta_{x_j} \quad \text{and} \quad \mu_2^n = \sum_{j=0}^n c_j^- \delta_{x_j}$$

where $c_j^+ = \max\{c_j, 0\}$ and $c_j^- = \max\{-c_j, 0\}$.

Let us determine what conditions the measure μ^n must satisfy. If $P(x) = \sum_{k=0}^n a_k x^k$, then $P'(0) = a_1$. By (8), $\int_0^1 x^k d\mu^n(x) = \sum_{j=0}^n c_j x_j^k$ whenever $0 \leq k \leq n$, and hence

$$\int_0^1 P(x) d\mu^n(x) = \int_0^1 \sum_{k=0}^n a_k x^k d\mu^n(x) = \sum_{k=0}^n a_k \int_0^1 x^k d\mu^n(x) = \sum_{k=0}^n \sum_{j=0}^n a_k c_j x_j^k.$$

Therefore $\int_0^1 f(x) d\mu^n(x) = P'(0)$ holds if and only if

$$\sum_{k=0}^n a_k \sum_{j=0}^n x_j^k = a_1$$

or equivalently,

$$\sum_{j=0}^n c_j x_j^k = \begin{cases} 1 & : k = 1 \\ 0 & : k \neq 1 \end{cases}$$

or equivalently

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_0 & x_1 & \cdots & x_n \\ x_0^2 & x_1^2 & \cdots & x_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_0^n & x_1^n & \cdots & x_n^n \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}. \quad (9)$$

Define

$$Q(x_0, \dots, x_n) := \det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_0 & x_1 & \cdots & x_n \\ x_0^2 & x_1^2 & \cdots & x_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_0^n & x_1^n & \cdots & x_n^n \end{pmatrix}.$$

The matrix equation necessarily has a solution $\vec{c} = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}$ if and only if we have $Q(x_0, \dots, x_n) \neq 0$. Therefore to prove the existence of the desired μ^n , it will suffice to show there exists $(x_0, \dots, x_n) \in (0, 1)^{n+1}$ such that $Q(x_0, \dots, x_n) \neq 0$. But this follows because Q is a polynomial, hence an entire function on \mathbb{C}^{n+1} , and because its derivatives do not vanish at the origin, i.e.

$$\partial_{x_1} \partial_{x_2}^2 \cdots \partial_{x_n}^n Q(0) \neq 0.$$

Thus $Q \neq 0$, so Q cannot vanish on any open set. In particular, it cannot vanish on $(0, 1)^{n+1}$.

- (b) Does there exist two finite positive measures μ_1, μ_2 supported on $[0, 1]$ such that for any polynomial $P(x)$, it holds

$$P'(0) = \int_0^1 P(x) d\mu_1(x) - \int_0^1 P(x) d\mu_2(x)?$$

Solution: There does not exist such a pair of measures. If the measures existed, and if $\mu = \mu_1 - \mu_2$, then μ corresponds to a continuous linear functional T on $C[0, 1]$, such that $Tf = f'(0)$ for any polynomial f . But then T cannot be continuous, for if $f_n = (1 - x)^n$, then $\|f_n\|_{L^\infty[0,1]} = 1$ for all n , whereas $Tf_n = n$.

10 Baire Category Notes

The *Baire Category Theorem* says that if X is a complete metric space, then for any sequence $\{U_n\}$ of open, dense subsets of X , $\bigcap_n U_n$ is dense in X . To make things less abstract, I like to think through this result in terms of logical properties:

- A logical property P of points in X is *stable* if, whenever $P(x_0)$ is true for some $x_0 \in X$, then $P(x)$ is true for x in a neighborhood of x_0 .
- A logical property P of points in X is *unstable* if, whenever $P(x_0)$ is true for some $x_0 \in X$, then for any $\varepsilon > 0$, we can find $x \in X$ with $d(x, x_0) < \varepsilon$, such that $P(x)$ is false. One advantage of being quantitative here is that it suffices to show that the property is unstable for x_0 in a *dense* subspace X_0 of X .

The Baire category theorem then says that, given a countable family of properties $\{P_n\}$, such that P_n is a stable property, and the *negation* of P_n is an unstable property, then the set of points $x \in X$ such that $P_n(x)$ is true for all n is a dense subset of X .

To see an example of this formulation of the theorem, let us go through the classic Baire category proof that there exists a continuous function on $[0, 1]$, which is differentiable nowhere. The first challenge is to find countably many properties $\{P_n\}$ of functions in $C[0, 1]$ such that a function f is differentiable nowhere if and only if $P_n(f)$ is true for all n , though we must also be careful to choose these properties to be stable. This leads to the family of properties $\{P_{N,M}\}$, where N and M are positive integers, and $P_{N,M}(f)$ is true if, for any $x_0 \in [0, 1]$, there exists $x_1 \in [0, 1]$ with $0 < |x_1 - x_0| < 1/M$ and with

$$|f(x_0) - f(x_1)| > N|x_0 - x_1|.$$

Let us now show that the properties $\{P_{N,M}\}$ are both stable, and their complement is unstable:

- Since $[0, 1]$ is compact, and for any fixed f , x_0 and x_1 such that the above properties hold, we can keep x_1 constant as we vary x_0 in a small neighborhood, we can find a finite collection of points $\{x_1(1), \dots, x_1(K)\}$ such that given any $x_0 \in [0, 1]$, there is k such that $0 < |x_0 - x_1(k)| < 1/M$ and $|f(x_0) - f(x_1)| > N|x_0 - x_1|$. Combined with the fact that we use a *strict inequality* above, this can be used to show $P_{N,M}$ is a stable property.
- Conversely, the negation of the property $P_{N,M}$ is unstable. To see this, we note that $C^\infty[0, 1]$ is dense in $C[0, 1]$. Thus to show the property is unstable, it suffices to show it is unstable at any such function $f \in C^\infty[0, 1]$. Fix $T > 0$ such that f is Lipschitz of order T for some large quantity $T \geq 1$. For any $R \geq 2M$, define a function $g_R \in C[0, 1]$ such that for $1 \leq k \leq R$, $g(k/R) = (-1)^k(10NT/R)$, and then linearly interpolating between these values. Then $\|g_R\|_{L^\infty} \leq 10NT/R$. Now consider the function $f_R = f + g_R$. Now if $1/R \leq x \leq 2/R$,

$$\begin{aligned} |f_R(k/R + x) - f_R(k/R)| &\geq |g_R(k/R + x) - g_R(k/R)| - |f(k/R + x) - f(k/R)| \\ &\geq 10NT/R - |f(k/R + x) - f(k/R)| \\ &\geq 10TN/R - 2T/R \\ &\geq 5TN/R > N|x|. \end{aligned}$$

Since $R \geq 2M$, this justifies that $P_{N,M}(f_R)$ is true. For any $\varepsilon > 0$, if $R \geq 10NT/\varepsilon$, then $\|f_R - f\|_{L^\infty} \leq \varepsilon$. Thus we see that the property $P_{N,M}$ is unstable.

The Baire category theorem then implies that the set of $f \in C[0, 1]$ such that $P_{N,M}(f)$ is true for all N and M is dense in $C[0, 1]$, which proves the existence of continuous functions differentiable at no point.

Day 7: Warm Up Problems

11 Day 7: Baire Category

17. (Spring 2020) A **Hamel basis** for a vector space X is a collection $\mathcal{H} \subset X$ of vectors such that $x \in X$ can be written uniquely as a finite linear combination of elements in \mathcal{H} . Prove that an infinite dimensional Banach space cannot have a countable Hamel basis. (Hint: otherwise the Banach space would be first category in itself.)

Solution: Denote the norm of X by $\|\cdot\|$. Suppose \mathcal{H} is countable. Then there exists an enumeration f_1, f_2, \dots of \mathcal{H} . Let $E_n = \text{span}\{f_1, \dots, f_n\}$. By definition of Hamel basis, $X = \cup_{n=1}^{\infty} E_n$. It will suffice to show that E_n is nowhere dense, as this will imply that X —which is a complete metric space with respect to the norm topology—is a countable union of nowhere dense sets, contradicting the *Baire Category theorem*. But this follows because *any* finite dimensional subspace of a Banach space is closed, and if E_n itself had nonempty interior, then it would contain a neighborhood of the origin, which is absorbing, and thus we would conclude that $E_n = X$.

Let us argue in slightly more detail some steps of this argument:

Lemma 1: Any finite dimensional subspace Y of a Banach space X is closed.

Proof. Since Y is finite dimensional, it has some basis $\{e_1, \dots, e_n\}$. If we define a norm

$$\|\sum a_i e_i\|_Y = \left(\sum |a_i|^2\right)^{1/2},$$

then Y is a Banach space with this norm, because it is easily seen to be isometric to \mathbb{R}^n or \mathbb{C}^n (depending on the underlying field). But all norms on a finite dimensional vector space are equivalent, so that $\|\cdot\|_Y$ is comparable to $\|\cdot\|_X$, and this shows Y is a Banach space under the norm induced by X . But this implies that Y is closed in X , since any sequence $\{y_n\}$ converging to an element $x \in X$ is Cauchy, and thus converges in Y . \square

Lemma 2: If Y is a proper subspace of a Banach space X , then Y has nonempty interior in X .

Proof. For each $x_0 \in X$ and $\varepsilon > 0$, let

$$B_\varepsilon(x_0) = \{x \in X : \|x - x_0\| < \varepsilon\}.$$

Suppose that there exists $y_0 \in Y$ and $\varepsilon > 0$ such that $B_\varepsilon(y_0) \subset Y$. Then

$$B_\varepsilon(0) = B_\varepsilon(y_0) - y_0 \subset Y - y_0 = Y.$$

If $r > 0$, then

$$B_r(0) = (r/\varepsilon) \cdot B_\varepsilon(0) \subset (r/\varepsilon) \cdot Y = Y.$$

Thus Y contains every ball centered at the origin. But this clearly means $X = Y$. \square

18. (Fall 2017) Let f_n be a sequence of real functions on \mathbb{R} such that each f'_n is continuous on \mathbb{R} . Suppose that as $n \rightarrow \infty$, f_n converges to a function $f : \mathbb{R} \rightarrow \mathbb{R}$ pointwise, and f'_n converges to a function g pointwise.

Prove that there exists a non-empty interval (a, b) and a constant $L < \infty$ such that

$$|f(x) - f(y)| \leq L|x - y|.$$

Hint: Consider the sets $K_c = \{x : \sup_n |f'_n(x)| \leq c\}$.

Solution: For each x , $f'_n(x) \rightarrow f(x)$, so that the sequence of quantities $\{f'_n(x)\}$ is bounded. It follows that if

$$K_c = \{x : \sup_n |f'_n(x)| \leq c\},$$

then $\bigcup_{c=1}^{\infty} K_c = \mathbb{R}$. Since the supremum of continuous functions is lower semi-continuous, each K_c is a closed set. And thus the *Baire category theorem* implies that there exists L such that K_L contains an interval. Thus there is an interval (a, b) such that for all $x \in (a, b)$, $\sup_n |f'_n(x)| \leq L$. Thus by the mean value theorem

$$|f(x) - f(y)| = \limsup_{n \rightarrow \infty} |f_n(x) - f_n(y)| \leq \limsup_{n \rightarrow \infty} |x - y| \|f_n\|_{L^\infty(a,b)} \leq L|x - y|.$$

19. (Fall 2016) Show that there is a continuous real valued function on $[0, 1]$ that is not monotone on any open interval $(a, b) \subset [0, 1]$.

Solution: Suppose not. Then we can write $C([0, 1])$ as the following countable union:

$$C([0, 1]) = \bigcup_{a,b \in \mathbb{Q}, a < b} E_{a,b}. \quad (10)$$

where $E_{a,b} = \{f \in C([0, 1]) : f \text{ monotone on } (a, b)\}$.

We claim that each $E_{a,b}$ is closed. Indeed suppose (f_k) is a sequence of monotone function on (a, b) such that $f_k \rightarrow f$ uniformly. Then there exists a subsequence (k_j) with $k_j \rightarrow \infty$ and such that f_{k_j} is an increasing function for all j or f_{k_j} is a decreasing function for all j . Without loss suppose that they are increasing functions. If $x \leq y$ then $f_{k_j}(x) - f_{k_j}(y) \leq 0$ for all j , and hence sending $j \rightarrow \infty$, we have $f(x) \leq f(y)$. Thus f is increasing. Thus $E_{a,b}$ is closed. This proves the claim.

The $C[0, 1]$ is a complete metric space which is the countable union of closed sets. It follows by Baire's theorem that there exists some $a, b \in \mathbb{Q}$ with $a < b$ such that $E_{a,b}$ has nonempty interior. That is, there exists $f^* \in E_{a,b}$ and $\epsilon > 0$ such that $\{f \in C([0, 1]) : \|f^* - f\|_\infty < \epsilon\} \subset E_{a,b}$. Therefore every f with $\|f^* - f\|_\infty < \epsilon$ is monotone on the interval (a, b) . But this is absurd, since any function can be approximated uniformly by “zig-zag functions”.

20. (Spring 2014) Does there exist a sequence of continuous functions $f_n : [0, 1] \rightarrow \mathbb{R}$ such that $f_n \rightarrow \chi_Q$ pointwise?

Solution: No. Suppose such a sequence (f_n) exists. Then we can write

$$[0, 1] = \bigcup_{k \geq 1} \left\{ x \in [0, 1] : |f_n(x) - f_m(x)| \leq \frac{1}{4} \text{ for all } m, n \geq k \right\}.$$

Since the right hand side is a countable union of closed sets, it follows by the Baire Category theorem that there exists some k_0 and some (a, b) such that for all $x \in (a, b)$,

$$|f_n(x) - f_m(x)| \leq \frac{1}{4} \quad \text{for all } n, m \geq k_0.$$

Sending $m \rightarrow \infty$,

$$|f_n(x) - f(x)| \leq \frac{1}{4} \quad \text{for all } n \geq k_0.$$

Let $n \geq k_0$. If $x \in (a, b) \setminus \mathbb{Q}$ then $|f_n(x)| \leq \frac{1}{4}$. But if $x \in (a, b) \cap \mathbb{Q}$ then $|f_n(x) - 1| \geq \frac{1}{4}$, so that $|f_n(x)| \geq \frac{3}{4}$. But this contradicts the continuity of f_n .

21. (Fall 2014) Let X, Y be Banach spaces and $\{T_{j,k} : j, k \in \mathbb{N}\}$ be a set of bounded linear transformations $X \rightarrow Y$. Suppose for each k , there exists $x \in X$ such that $\sup \{\|T_{j,k}x\| : j \in \mathbb{N}\} = \infty$. Then there is an $x \in X$ such that $\sup \{\|T_{j,k}x\| : j \in \mathbb{N}\} = \infty$ for all k .

Solution: Proof by contrapositive. Suppose that the conclusion does not hold: that is, suppose that for all $x \in X$ there exists some k such that $\sup \{\|T_{j,k}x\| : j \in \mathbb{N}\} < \infty$. Then we can write

$$X = \bigcup_{k=1}^{\infty} \bigcup_{N=1}^{\infty} \bigcap_{j=1}^{\infty} \{x \in X : \|T_{j,k}x\| \leq N\}.$$

By the Baire Category theorem, there exists $k_0, N_0 \geq 1$ and some ball $\bar{B}_r(x_0) = \{x \in X : \|x - x_0\| \leq r\}$ with $r > 1$ such that

$$\sup_{j \geq 1} \|T_{j,k_0}x\| \leq N_0$$

for all $x \in \bar{B}_r(x_0)$.

It will suffice to show that for this choice of k_0 , we have $\sup \{\|T_{j,k_0}x\| : j \in \mathbb{N}\} < \infty$ for all $x \in X$.

Let $x \in X$. There are two cases:

Case 1. Suppose $\|x\| \leq 1$. Then $rx + x_0 \in \bar{B}_r(x_0)$, and hence

$$\begin{aligned} \|T_{j,k_0}x\| &= \frac{1}{r} \|T_{j,k_0}(rx)\| \\ &= \frac{1}{r} \|T_{j,k_0}(rx + x_0) - T_{j,k_0}x_0\| \\ &\leq \frac{1}{r} \|T_{j,k_0}(rx + x_0)\| + \frac{1}{r} \|T_{j,k_0}x_0\| \\ &\leq 2N_0/r \end{aligned}$$

for all j .

Case 2. Suppose $\|x\| > 1$. Let $x_1 = x/\|x\|$. By Case 1,

$$\|T_{j,k_0}x\| = \|x\| \|T_{j,k_0}x_1\| \leq 2\|x\| N_0/r$$

for all j .

Combining the two cases, the result follows.

12 Distribution Theory Notes

Here we detail the very basics of distribution theory. The hope is that provided one knows these basics, then without having to study much distribution theory, one can turn many problems on the exam involving distributions into more basic analysis problems, to which one can apply the tools of basic analysis, measure theory, or functional analysis.

If $U \subset \mathbb{R}^d$ is an open set, a distribution on U is a *continuous linear functional* u on the vector space $\mathcal{D}(U) := C_c^\infty(U)$ of all smooth, compactly supported functions on U . Thus with each $\phi \in C_c^\infty(U)$, the functional u associates a quantity $\langle u, \phi \rangle$, which we might also denote as

$$\int u(x)\phi(x) dx.$$

The set of all distributions on U is denoted $\mathcal{D}(U)^*$. There is an abstract theory of topological vector spaces that allows us to define what it means for u to be continuous, but it is not necessary to learn this theory if one remembers a more practical definition. In order for u to be continuous, one needs to show that for any compact set $K \subset U$, there exists an integer $N > 0$, possibly depending on K , such that for all $\phi \in C_c^\infty(U)$ with $\text{supp}(\phi) \subset K$,

$$|\langle u, \phi \rangle| \lesssim_K \sup_{x \in K} \sup_{|\alpha| \leq N} |D^\alpha \phi(x)|.$$

To check you understand this definition, prove that the linear functional u defined by setting

$$\langle u, \phi \rangle = \sum_{n=1}^{\infty} e^{e^n} D^{n!} \phi(n)$$

is a distribution. As other examples, for any locally integrable function $f : U \rightarrow \mathbb{C}$, or any locally finite measure μ , one can view f and μ as distributions by defining

$$\langle f, \phi \rangle = \int f(x)\phi(x) dx \quad \text{and} \quad \langle \mu, \phi \rangle = \int \phi(x) d\mu(x).$$

Thus many mathematical objects in analysis are special cases of distributions. An important example is the Dirac delta distribution δ_x , for any $x \in U$, such that $\langle \delta_x, \phi \rangle = \phi(x)$.

Surprisingly, one can take the formal derivative of *any distribution*, by applying integration by parts. If u is a distribution on U , we define its partial derivatives $D^i u$, which are also distributions, by the formal definition

$$\langle D^i u, \phi \rangle = -\langle u, D^i \phi \rangle.$$

As an example, we calculate that

$$\langle D^i \delta_x, \phi \rangle = -\langle \delta_x, D^i \phi \rangle = -D^i \phi(x).$$

Thus the derivative of the Dirac delta at a point x is the distribution that measures the derivatives of a given function at the point x . For more well behaved functions, the distributional derivative of a function will be equal to the derivative of the function.

A sequence of distributions $\{u_n\}$ converges *distributionally* to another distribution u if, for any $\phi \in \mathcal{D}(U)$, the quantities $\langle u_n, \phi \rangle$ converge to $\langle u, \phi \rangle$. If this is the case, it also follows that the derivatives $\{D^i u_n\}$ converge distributionally to $D^i u$.

Roughly speaking, if you start to get intuition about how to work with distributions, these facts will allow you to convert most problems about distributions to more basic problems about the convergence of numbers, or integrals, and so on, and so you can focus on trying to apply the more basic analytical tools.

Day 8: Warm Up Problems

13 Day 8: Intro to Distribution

22. (Fall 2020) Consider the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = |x|$. Find $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)f$, where the derivative is taken in the sense of distributions.

Solution: Let $\phi \in C_c^\infty(\mathbb{R}^2)$. By definition of distributional derivative,

$$\begin{aligned} \left\langle \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)f, \phi \right\rangle &= \left\langle f, \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)\phi \right\rangle \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |x| \phi_{xx}(x, y) + |x| \phi_{yy}(x, y) dx dy \\ &= \underbrace{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |x| \phi_{xx}(x, y) dx dy}_A + \underbrace{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |x| \phi_{yy}(x, y) dx dy}_B. \end{aligned}$$

Integrating by parts,

$$\begin{aligned} A &= \int_{-\infty}^{\infty} \left[\int_0^{\infty} x \phi_{xx}(x, y) dx - \int_{-\infty}^0 x \phi_{xx}(x, y) dx \right] dy \\ &= \int_{-\infty}^{\infty} \left[- \int_0^{\infty} \phi_x(x, y) dx + \int_{-\infty}^0 \phi_x(x, y) dx \right] dy \\ &= \int_{-\infty}^{\infty} -[\phi(x, y)]_{x=0}^{x=\infty} + [\phi(x, y)]_{x=-\infty}^{x=0} dy \\ &= 2 \int_{-\infty}^{\infty} \phi(0, y) dy \end{aligned}$$

Next, by Fubini's theorem,

$$B = \int_{-\infty}^{\infty} |x| \int_{-\infty}^{\infty} \phi_{yy}(x, y) dy dx = 0$$

because $\int_{-\infty}^{\infty} \phi_{yy}(x, y) dy = [\phi_y(x, y)]_{y=-\infty}^{y=\infty} = 0$ for every x . It follows from our calculations for A and B that

$$\left\langle \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)f, \phi \right\rangle = 2 \int_{-\infty}^{\infty} \phi(0, y) dy$$

for every ϕ . Therefore $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)f = 2\delta(x)$ in the sense of distributions, where δ is the point mass at $x = 0$.

23. (Spring 2017)

- (a) Let $f \in L^1(\mathbb{R})$ and consider the sequence of distributions $T_n(x) = \sin(nx^2)f(x)$. Show that $\lim_{n \rightarrow \infty} T_n = 0$ in the sense of distributions.

Solution: We perform an integration by parts, which is always handy when dealing with

oscillatory things. If $\Lambda_n(x) = \sin(nx^2)$, then

$$\begin{aligned} \int \Lambda_n(x) \phi(x) dx &= \int \Lambda'_n(x) \frac{\phi(x)}{2nx} \\ &= - \int \Lambda_n(x) \frac{d}{dx} \left(\frac{\phi(x)}{2nx} \right) \\ &= - \int \Lambda_n(x) \left(\frac{\phi'(x)}{2nx} - \frac{\phi(x)}{2nx^2} \right). \end{aligned}$$

Now performing a Taylor expansion, we can write

$$\frac{\phi'(x)}{2nx} - \frac{\phi(x)}{2nx^2} = \frac{\phi'(0)}{2nx} - \frac{\phi(0)}{2nx^2} - \frac{\phi'(0)}{2nx} + O(1/n) = \frac{\phi'(0)}{2nx} + O(1/n).$$

But since $1/x$ is odd, and $\sin(nx^2)$ is even, we have

$$\int \Lambda_n(x) \frac{\phi'(0)}{2nx} dx = 0.$$

Thus

$$\int \Lambda_n(x) \phi(x) \lesssim_\phi O(1/n),$$

Taking $n \rightarrow \infty$ shows that Λ_n converges to 0 distributionally.

- (b) Find a distribution $T \in \mathcal{D}'(\mathbb{R})$ such that $T_n = \sin(nx^2)T$ does not converge to 0 in the sense of distributions as $n \rightarrow \infty$.

Solution: Let T be the distribution defined by

$$T(\phi) = \phi\left(\sqrt{\pi/2}\right)$$

for every test function ϕ . Then

$$T_n(\phi) = T(\sin(nx^2)\phi(x)) = \sin\left(\frac{n\pi}{2}\right) \phi\left(\sqrt{\frac{\pi}{2}}\right) = (-1)^n \phi\left(\sqrt{\frac{\pi}{2}}\right),$$

which does not converge as $n \rightarrow \infty$ if $\phi(\sqrt{\pi/2}) \neq 0$.

24. (Spring 2015)

- (a) If $f \in C[0, 1]$, and the distributional derivative f' of f on $(0, 1)$ is in $L^1((0, 1))$, prove that

$$f(1) - f(0) = \int_0^1 f'(x) dx.$$

Solution: In order to relate f' and f , since f' is a distributional derivative, we need to integrate it against a test function, which reduces the question to a basic analysis problem. Fix $\delta > 0$ and consider $\phi_\delta \in C_c^\infty[0, 1]$ with $\|\phi_\delta\|_\infty \leq 1$, $\phi_\delta(x) = 1$ for $\delta \leq x \leq 1 - \delta$, and supported on $\delta/2 \leq x \leq 1 - \delta/2$, and with $\|\phi'_\delta\|_\infty \leq 10/\delta$. Then ϕ'_δ is supported on $\delta/2 \leq x \leq \delta$ and

$1 - \delta/2 \leq x \leq 1 - \delta$. Since f' is integrable, we have

$$\int_0^1 f'(x) dx = \lim_{\delta \rightarrow 0} \int_0^1 f'(x) \phi_\delta(x) dx.$$

Since f' is a distributional derivative, we can apply an integration by parts, from which we conclude that

$$\int_0^1 f'(x) \phi_\delta(x) dx = - \int_0^1 f(x) \phi'_\delta(x) dx.$$

We can write

$$\int_0^1 f(x) \phi'_\delta(x) dx = \int_0^\delta f(x) \phi'_\delta(x) dx + \int_{1-\delta}^1 f(x) \phi'_\delta(x) dx.$$

Since $f \in C[0, 1]$, for each $\varepsilon > 0$, there exists $\delta_0 > 0$ such that for $x, y \in [0, 1]$ with $|x - y| \leq \delta_0$, $|f(x) - f(y)| \leq \varepsilon$. If $\delta \leq \delta_0$, this implies that

$$\left| \int_0^\delta [f(x) - f(0)] \phi'_\delta(x) dx \right| \leq \int_0^\delta \varepsilon |\phi'_\delta(x)| dx \leq \varepsilon.$$

Similarly,

$$\left| \int_{1-\delta}^1 [f(x) - f(1)] \phi'_\delta(x) dx \right| \leq \int_{1-\delta}^1 \varepsilon |\phi'_\delta(x)| dx \leq \varepsilon.$$

The fundamental theorem of calculus implies that

$$\int_0^\delta f(0) \phi'_\delta(x) dx = f(0) [\phi_\delta(\delta) - \phi_\delta(0)] = f(0),$$

and

$$\int_{1-\delta}^1 f(1) \phi'_\delta(x) dx = f(1) [\phi_\delta(1) - \phi_\delta(1 - \delta)] = -f(1).$$

Putting these calculations together shows that

$$\lim_{\delta \rightarrow 0} \int_0^1 f(x) \phi'_\delta(x) dx = f(0) - f(1),$$

and thus

$$\int_0^1 f'(x) dx = - \lim_{\delta \rightarrow 0} \int_0^1 f(x) \phi'_\delta(x) dx = f(1) - f(0).$$

- (b) Let $p \in [1, \infty)$ and let $F \subset C[0, 1]$ be such that for each $f \in F$ we have $\|f\|_{L^1[0,1]} \leq 1$ and $\|f'\|_{L^p[0,1]} \leq 1$, where f' is the distributional derivative of f . Prove that F is precompact in $C[0, 1]$, or find a counter-example.

Solution: Problems about pre-compactness in $C[0, 1]$ normally want us to apply the Arzela-Ascoli theorem, so let's try that here. We wish to show there is a constant $C > 0$ such that for each $f \in C[0, 1]$,

$$\|f\|_\infty \leq C,$$

(uniform boundedness), and that for each $\varepsilon > 0$, there is $\delta > 0$ such that for $|x - y| \leq \delta$ and $f \in F$, $|f(x) - f(y)| \leq \varepsilon$ (equicontinuity).

Using the last part of the problem, together with Hölder's inequality, if p and p^* are conjugates,

then for $x < y$,

$$|f(x) - f(y)| = \left| \int_x^y f'(x) dx \right| \leq \|f'\|_p \left(\int_x^y 1 \right)^{1/p^*} = |y - x|^{1/p^*} \|f'\|_p \leq |y - x|^{1/p^*}.$$

Thus f satisfies a Hölder condition, uniformly over elements of F . This gives equicontinuity.

To show F is uniformly bounded, we must exploit the fact that $\|f\|_1 \leq 1$. Suppose that $\|f\|_\infty \geq M$. Then there exists $x_0 \in [0, 1]$ such that $|f(x_0)| \geq M$. Using the Hölder condition, we find that $|f(x_0)| \geq M/2$ for $|y - x_0| \leq (M/2)^{p^*}$. And so we find that $\|f\|_1 \geq (M/2) \cdot 2(M/2)^{p^*} \gtrsim_p M^{1+p^*}$. Since $\|f\|_1 \leq 1$, this means that $M^{1+p^*} \lesssim_p 1$, and so $M \lesssim_p 1$. This means we have shown that there exists a constant C_p such that for all $f \in F$, $\|f\|_\infty \leq C_p$.

Thus we have shown that the hypothesis of the Arzela-Ascoli theorem applies, and so F is precompact.

25. (Fall 2021, Spring 2016) Prove or disprove:

(a) There exists a distribution $u \in \mathcal{D}'(\mathbb{R})$ so that the restriction to $(0, \infty)$ is given by

$$\langle u, f \rangle = \int_0^\infty e^{1/x^2} f(x) dx$$

for all $f \in C^\infty(\mathbb{R})$ which are compactly supported in $(0, \infty)$.

Solution: Part (a): Does not exist. Idea: if such a distribution exists, then there exists N, C such that

$$|\langle u, \phi \rangle| \leq C \|\phi\|_N.$$

But we will construct a sequence ϕ_n so that the left hand side blows up faster than the right hand side. We will want the functions ϕ_n to get more and more concentrated near 0.

Let $\phi \in C_c^\infty(0, 2)$ such that $\phi(x) = 1$ for all $x \in [1/2, 1]$. Define $\phi_n(x) = \phi(2^n x)$. Then $\phi_n(x) = 1$ for all $x \in [2^{-(n+1)}, 2^{-n}] =: I_n$. We have

$$\langle u, \phi_n \rangle \geq \int_{I_n} e^{x^{-2}} dx \geq e^{4^n} 2^{-(n+1)}$$

On the other hand,

$$\|\phi_n\|_N = \sum_{k \leq N} \left\| \phi_n^{(k)} \right\|_\infty = \sum_{k \leq N} \left\| 2^{kn} \phi^{(k)}(2^n x) \right\|_\infty \leq 2^{Nn} \|\phi\|_N$$

Therefore,

$$\frac{\langle u, \phi_n \rangle}{\|\phi_n\|_N} \geq \frac{e^{4^n} 2^{-(n+1)}}{2^{Nn}} \rightarrow \infty \text{ as } n \rightarrow \infty.$$

(b) There exists a distribution $u \in \mathcal{D}'(\mathbb{R})$ so that its restriction to $(0, \infty)$ is given by

$$\langle u, f \rangle = \int_0^\infty x^{-2} e^{i/x^2} f(x) dx$$

for all $f \in C^\infty$ which are compactly supported in $(0, \infty)$.

Solution: Part (b): Yes, such a distribution exists. Let $f \in \mathcal{D}(\mathbb{R})$ with $\text{supp}(f) \subset (0, \infty)$. Write $x^{-2}e^{ix^{-2}} = \frac{ix}{2} \frac{d}{dx} \left[e^{ix^{-2}} \right]$, so that

$$\begin{aligned} \langle u, f \rangle &= \int_0^\infty \frac{d}{dx} \left[e^{ix^{-2}} \right] \frac{ix}{2} f(x) dx \\ &= \int_0^\infty e^{ix^{-2}} \left(\frac{i}{2} f(x) + \frac{ix}{2} f'(x) \right) dx \\ &= \frac{i}{2} \int_0^\infty e^{ix^{-2}} (f(x) + x f'(x)) dx. \end{aligned}$$

and it is easy to verify that this formula defines a distribution for all $f \in \mathcal{D}(\mathbb{R})$.

26. (Fall 2017) A distribution $T \in \mathcal{S}'(\mathbb{R}^n)$ is said to be *nonnegative* if $\langle T, \phi \rangle \geq 0$ for every test function $\phi \in \mathcal{S}(\mathbb{R}^n)$ with $\phi(x) \geq 0$ for all $x \in \mathbb{R}^n$.

(a) Suppose $f \in L^1_{\text{loc}}(\mathbb{R}^n)$, and let T_f be the distribution defined by f . Show that $T_f \geq 0$ if and only if $f \geq 0$ for almost all $x \in \mathbb{R}^n$.

Solution: Suppose $f < 0$ on a set of positive measure. Then we may find $R > 0$ and $c > 0$ such that the set $E = \{|x| \leq R : f(x) < -c\}$ has positive measure. Fix $\delta > 0$ and find an open set U contained E with $|U - E| \leq \delta$. Since U is pre-compact, we can find a smooth function ϕ compactly supported on a set V such that $f(x) = 1$ for $x \in U$, and $|V - U| \leq \delta$. Then

$$\left| \int f(x)\phi(x) dx - \int_E f(x) dx \right| \leq \int_{V-U} |f(x)| dx.$$

Since f is integrable, and $|V - E| \leq 2\delta$, for any fixed $\varepsilon > 0$, if δ is suitably small, then

$$\int_{V-U} |f(x)| dx \leq \varepsilon.$$

On the other hand,

$$\int_E f(x) dx \leq -c|E|,$$

and so

$$0 \leq \int f(x)\phi(x) \leq \varepsilon - c|E|.$$

Taking $\varepsilon \rightarrow 0$ gives a contradiction.

(b) Show that if $T_n \rightarrow T$ in the sense of distributions, and if $T_n \geq 0$ for all n , then $T \geq 0$.

Solution: If $\phi \geq 0$, then $\langle T_n, \phi \rangle \geq 0$ for all n since $T_n \geq 0$. But $\langle T, \phi \rangle = \lim_n \langle T_n, \phi \rangle$, which implies that $\langle T, \phi \rangle \geq 0$.

(c) Suppose $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a C^2 function with $\Phi'' \geq 0$ in \mathbb{R} , and let $f \in C^2(\mathbb{R}^n)$ have compact support. Show that $\Delta(\Phi(f(x))) \geq \Phi'(f(x))\Delta f(x)$.

Solution: We calculate directly that

$$\Delta(\Phi(f(x))) = \Phi'(f(x))\Delta f + \sum_{i=1}^n \Phi''(f(x)) \left(\frac{\partial f}{\partial x^i} \right)^2.$$

Since $\Phi''(f(x))(\partial f/\partial x^i)^2 \geq 0$, this easily implies the inequality.

- (d) Suppose $f \in C^2(\mathbb{R}^n)$ has compact support. Show that $\Delta|f| \geq \text{sign}(f(x))\Delta f(x)$ holds in the sense of distributions. (Hint use (c) with $\Phi(t) = \sqrt{\varepsilon + t^2}$).

Solution: For each $\varepsilon > 0$, if $\Phi_\varepsilon(t) = \sqrt{\varepsilon + t^2}$, then $\Phi_\varepsilon \in C^\infty(\mathbb{R})$, and the part (c) implies that

$$\Delta(\sqrt{\varepsilon + f^2}) \geq \frac{f \cdot \Delta f}{\sqrt{\varepsilon + f^2}}$$

Our goal is to take $\varepsilon \rightarrow 0$ to show that $\Delta|f| \geq \text{sign}(f)\Delta f$.

As $\varepsilon \rightarrow 0$, $\sqrt{\varepsilon + f^2}$ converges distributionally to $|f|$ by the dominated convergence theorem. Given $\phi \in C_c^\infty(\mathbb{R}^d)$, $\phi(x)\sqrt{\varepsilon + f(x)^2} \rightarrow \phi(x)|f(x)|$ pointwise for every x , and is upper bounded by $2|\phi(x)|(1 + |f(x)|)$, which is integrable since ϕ is compactly supported. Thus

$$\int \sqrt{\varepsilon + f(x)^2} \phi(x) dx \rightarrow \int |f(x)| \phi(x) dx.$$

Similarly, one can argue using the dominated convergence theorem that $f(x)\Delta f(x)/\sqrt{\varepsilon + f(x)^2}$ converges distributionally to $\text{sign}(f(x))\Delta f(x)$, since it converges pointwise to this function, and is uniformly upper bounded in absolute value by $\Delta f(x)$. Applying (b) shows that $\Delta|f(x)| \geq \text{sign}(f(x)) \cdot \Delta f(x)$.

27. (Spring 2020)

- (a) Suppose Λ is a distribution on \mathbb{R}^n such that $\text{supp}(\Lambda) = \{0\}$. If $f \in C^\infty(\mathbb{R}^n)$ satisfies $f(0) = 0$, does it follow that $f\Lambda = 0$ as a distribution?

Solution: No. Let $n = 1$, $f(x) = x$, and $\Lambda = D\delta$. It is clear that Λ has support $\{0\}$. But $\langle f\Lambda, \phi \rangle = \langle \Lambda, x\phi \rangle = \langle \delta, -(x\phi)' \rangle = \langle \delta, -\phi' - x\phi \rangle = -\phi'(0)$. Therefore $f\Lambda = -\delta$ in the sense of distributions, and hence is not the zero distribution.

- (b) Suppose Λ is a distribution on \mathbb{R}^n such that $\text{supp}(\Lambda) \subseteq K$, where $K = \{x \in \mathbb{R}^n : |x| \leq 1\}$. If $f \in C^\infty(\mathbb{R}^n)$ vanishes on K , does it follow that $f\Lambda = 0$ as a distribution?

Solution: This part is true. Since $\text{supp}(\Lambda)$ is compact, it follows that Λ is of finite order. Then $\Lambda = \sum_{\alpha \in \mathcal{A}} D^\alpha g_\alpha$ where g_α is a continuous function for each $\alpha \in \mathcal{A}$ and \mathcal{A} is a finite set of multi-indices. Let $N = \max\{|\alpha| : \alpha \in \mathcal{A}\}$.

Let $\phi \in C_c^\infty(\mathbb{R}^n)$ and let $M > 0$ such that $\text{supp}(\phi) \subseteq \{x : |x| \leq M\}$. Then

$$\begin{aligned} \langle f\Lambda, \phi \rangle &= \langle \Lambda, f\phi \rangle \\ &= \int_{\mathbb{R}^n} \sum_{\alpha \in \mathcal{A}} (-1)^{|\alpha|} g_\alpha(x) D^\alpha(f\phi)(x) dx \\ &= \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{\mathbb{R}^n} g_\alpha(x) (D^{\alpha-\beta} f)(D^\beta \phi) dx && \text{by Leibniz Formula} \\ &= \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{|x| > 1} g_\alpha(x) (D^{\alpha-\beta} f)(D^\beta \phi) dx && \text{since } D^{\alpha-\beta} f(x) = 0 \text{ for all } x \in K \\ &= \lim_{\epsilon \rightarrow 0^+} \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{|x| > 1+\epsilon} g_\alpha(x) (D^{\alpha-\beta} f)(D^\beta \phi) dx \end{aligned}$$

We claim this limit is zero. To see this, we first define a cutoff function $\eta_1(x) \in C_c^\infty(\mathbb{R}^n)$ with $0 \leq \eta_1 \leq 1$ such that $\eta_1(x) = 0$ if $|x| < 1$, and $\eta_1(x) = 1$ if $2 \leq |x| \leq M$. Next, for each $\epsilon \in (0, 1)$, define $\eta_\epsilon \in C_c^\infty$ satisfying

$$\eta_\epsilon(x) = \begin{cases} 0 & : |x| \leq 1 + \epsilon/2 \\ \eta_1\left(\frac{2x}{\epsilon} - \frac{2}{\epsilon}\right) & : 1 + \epsilon/2 < |x| \leq 1 + \epsilon \\ 1 & : 1 + \epsilon < |x| \leq M \\ \text{smooth} & : |x| > M \end{cases}$$

(Draw a picture for the $n = 1$ case. This isn't as complicated as it looks.)

By the chain rule, $|D^\alpha \eta_\epsilon| \leq C \cdot 2^{|\alpha|} \epsilon^{-|\alpha|}$ where $C < \infty$ is an upper bound for the first order derivatives of η_1 . Then taking $\phi_\epsilon = \eta_\epsilon \phi$,

$$\begin{aligned} \langle f\Lambda, \phi_\epsilon \rangle &= \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} g_\alpha(x) (D^{\alpha-\beta} f) (D^\beta \phi_\epsilon) dx \\ &\quad + \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{|x| > 1+\epsilon} g_\alpha(x) (D^{\alpha-\beta} f) (D^\beta \phi) dx \end{aligned}$$

On the other hand, $\text{supp}(f\phi_\epsilon) \subset \mathbb{R}^n \setminus B(0, 1 + \frac{\epsilon}{2})$. Therefore $f\phi_\epsilon \in \mathcal{D}(K^c)$ and hence by definition of distribution support, $\langle f\Lambda, \phi_\epsilon \rangle = \langle \Lambda, f\phi_\epsilon \rangle = 0$. Therefore

$$\langle f\Lambda, \phi \rangle = \lim_{\epsilon \rightarrow 0^+} - \sum_{\alpha \geq \beta} (-1)^{|\alpha|} c_{\alpha\beta} \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} g_\alpha(x) (D^{\alpha-\beta} f) (D^\beta \phi_\epsilon) dx$$

Therefore

$$|\langle f\Lambda, \phi \rangle| \leq \limsup_{\epsilon \rightarrow 0^+} \sum_{\alpha \geq \beta} |c_{\alpha\beta}| \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} |g_\alpha(x) (D^{\alpha-\beta} f) (D^\beta \phi_\epsilon)| dx. \quad (11)$$

In particular, for each $\epsilon > 0$, using the Leibniz formula again

$$\begin{aligned} &\sum_{\alpha \geq \beta} |c_{\alpha\beta}| \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} |g_\alpha(D^{\alpha-\beta} f) (D^\beta \phi_\epsilon)| dx \\ &\leq \sum_{\alpha \geq \beta} |c_{\alpha\beta}| \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} |g_\alpha(D^{\alpha-\beta} f)| \left| \sum_{\beta \geq \gamma} c_\gamma D^\gamma \phi D^{\gamma-\beta} \eta_\epsilon \right| dx \\ &\leq \sum_{\alpha \geq \beta} \sum_{\beta \geq \gamma} |c_{\alpha\beta}| |c_\gamma| \frac{C}{(\epsilon/2)^N} \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} |g_\alpha(D^{\alpha-\beta} f)| |D^\gamma \phi| dx. \end{aligned}$$

Therefore as $\epsilon \rightarrow 0^+$, we have $\frac{1}{(\epsilon/2)^N} \int_{1+\epsilon \geq |x| > 1+\frac{\epsilon}{2}} |g_\alpha(D^{\alpha-\beta} f)| |D^\gamma \phi| dx \rightarrow 0$ since $(D^{\alpha-\beta} f)(x) = 0$ for all x with $|x| = 1$. Then it follows by (11) that $\langle f\Lambda, \phi \rangle = 0$. Since ϕ was arbitrary, $f\Lambda = 0$ in the sense of distributions.

Day 9: Warm Up Problems

28. (Fall 2019) Let $s \in \mathbb{R}$, and let $H^s(\mathbb{R})$ be the Sobolev space on \mathbb{R} with the norm

$$\|u\|_{(s)} = \left(\int_{\mathbb{R}} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi \right)^{1/2}$$

where \hat{u} is the Fourier transform of u . Let $r < s < t$ be real numbers. Prove that for every $\varepsilon > 0$ there is $C > 0$ such that

$$\|u\|_{(s)} \leq \varepsilon \|u\|_{(t)} + C \|u\|_{(r)}$$

for every $u \in H^t(\mathbb{R})$.

Solution: Intuitively, we only need the H^t norm to control H^s for large frequencies $|\xi| \gtrsim 1$, since the H^r norm gives enough control over low frequencies since $(1 + |\xi|^2)^s \approx 1 \approx (1 + |\xi|^2)^r$ when $|\xi| \lesssim 1$. Thus we fix $R > 0$, and write

$$\int (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi = \int_{|\xi| \leq R} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi + \int_{|\xi| > R} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi.$$

Now

$$\int_{|\xi| \leq R} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi \leq R^{s-r} \int_{|\xi| \leq R} (1 + |\xi|^2)^r |\hat{u}(\xi)|^2 d\xi \leq R^{s-r} \|u\|_{(r)}^2,$$

and

$$\int_{|\xi| \geq R} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi \leq R^{s-t} \int_{|\xi| \geq R} (1 + |\xi|^2)^t |\hat{u}(\xi)|^2 d\xi \leq R^{s-t} \|u\|_{(t)}^2.$$

Thus we find that

$$\|u\|_{(s)} \leq \sqrt{R^{s-r} \|u\|_{(r)}^2 + R^{s-t} \|u\|_{(t)}^2} \leq (2R^{s-r})^{1/2} \|u\|_{(r)} + (2R^{s-t})^{1/2} \|u\|_{(t)}.$$

Given $\varepsilon > 0$, the proof is completed by picking $R \geq (2/\varepsilon^2)^{1/(t-s)}$.

There is an alternate solution using Young's inequality. Let $A = 1 + |\xi|^2$.

Claim 1: Suppose $r < s < t$. Then for each $\epsilon > 0$, there exists $C > 0$ such that $A^s \leq \epsilon A^t + CA^r$.

To prove this claim, we will use Young's inequality for products with ϵ . Recall Young's inequality states that if $a, b \geq 0$ and $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ then

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

For any $\epsilon > 0$ we may write $ab = ((p\epsilon)^{1/p} a) \cdot \frac{b}{(p\epsilon)^{1/p}}$, which gives

$$ab \leq \epsilon a^p + (p\epsilon)^{-q/p} \frac{b^q}{q}. \quad (12)$$

Let $\beta = \frac{t(s-r)}{t-r}$, $p = t/\beta$ and $q = \frac{p}{p-1} = \frac{t}{t-\beta} = \frac{t-r}{t-s}$. Then $p, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$. Applying inequality (12) with $a = A^\beta$ and $b = A^{s-\beta}$, we have:

$$\begin{aligned} A^s &= A^\beta A^{s-\beta} \\ &\leq \epsilon A^{\beta t/\beta} + \frac{1}{q(p\epsilon)^{q/p}} A^{(s-\beta)q} \\ &= \epsilon A^t + \frac{1}{q(p\epsilon)^{q/p}} A^{(s-\beta)q} \\ &= \epsilon A^t + \frac{1}{q(p\epsilon)^{q/p}} A^r. \end{aligned}$$

This proves Claim 1.

Next we apply Claim 1 with $r/2 < s/2 < t/2$, which gives

$$A^{s/2} \leq \epsilon A^{t/2} + C A^{r/2}.$$

Multiplying both sides by $|\hat{u}|$ and making the substitution $A = 1 + |\xi|^2$, we have

$$(1 + |\xi|^2)^{s/2} |\hat{u}(\xi)| \leq \epsilon (1 + |\xi|^2)^{t/2} |\hat{u}(\xi)| + C (1 + |\xi|^2)^{r/2} |\hat{u}(\xi)|.$$

Therefore using Minkowski's inequality,

$$\begin{aligned} \left\| (1 + |\xi|^2)^{s/2} \hat{u}(\xi) \right\|_{L^2} &\leq \left\| \epsilon (1 + |\xi|^2)^{t/2} \hat{u}(\xi) + C (1 + |\xi|^2)^{r/2} \hat{u}(\xi) \right\|_{L^2} \\ &\leq \epsilon \left\| (1 + |\xi|^2)^{t/2} \hat{u}(\xi) \right\|_{L^2} + C \left\| (1 + |\xi|^2)^{r/2} \hat{u}(\xi) \right\|_{L^2}. \end{aligned}$$

This is exactly the inequality we needed to show, so we are done.

29. (Fall 2015) Let f be a tempered distribution on \mathbb{R} with Fourier transform

$$\hat{f}(\xi) = 1 + \xi^{12} + \sin \xi + \text{sign}(\xi).$$

Find f and f' (specify the definition of the Fourier transform you are using).

Solution: I will be using the Fourier transform defined by

$$\hat{f}(\xi) = \int f(x) e^{-2\pi i \xi \cdot x} dx.$$

The problem is really asking us to compute the inverse Fourier transform of 1 , ξ^{12} , $\sin \xi$, and $\text{sign}(\xi)$ individually.

If δ is the Dirac delta function at the origin, then

$$\hat{\delta}(\xi) = \int e^{-2\pi i \xi \cdot x} \delta(x) dx = 1.$$

Thus $\hat{\delta} = 1$.

To find a function f whose Fourier transform is ξ^{12} , we recall the applying linear differential operators with constant coefficients on one side of the Fourier transform is the same as multiplying by a polynomial on the other side. Since ξ^{12} is equal to $\xi^{12} \cdot \hat{\delta}$, which implies that f is a constant multiple of the distributional derivative $D^{12}\delta$. If we compute constants carefully, we find that

$$f = \frac{1}{(2\pi i)^{12}} D^{12}\delta = \frac{D^{12}\delta}{(2\pi)^{12}}.$$

Thus we have found the inverse Fourier transform of ξ^{12} .

Next, we find a function whose Fourier transform is $\sin \xi$. A simple way to compute this is to write

$$\sin \xi = \frac{e^{i\xi} - e^{-i\xi}}{2i}.$$

Since $e^{i\xi}$ is the Fourier transform of the Dirac delta $\delta_{-1/2\pi}$ at $-1/2\pi$, and $e^{-i\xi}$ is the Fourier transform of the Dirac delta $\delta_{1/2\pi}$ at $1/2\pi$, if we write $\Gamma(x) = \delta_{-1/2\pi}/2i - \delta_{1/2\pi}/2i$, then $\hat{\Gamma}(\xi) = \sin \xi$.

Finally, we find a function f such that $\hat{f}(\xi) = \text{sign}(\xi)$. Taking distributional derivatives, we find that

$$D\hat{f}(\xi) = 2\delta.$$

Taking inverse Fourier transforms, we find that

$$-2\pi i x f(x) = 2.$$

In particular, we might guess that

$$f(x) = \frac{-1}{i\pi x}.$$

We must be slightly more careful, however, since integration against the singular function $1/\xi$ can be interpreted in several different ways. One choice is the *principal value*

$$\int \frac{\phi(\xi)}{\xi} d\xi = \lim_{\varepsilon \rightarrow 0} \int_{|\xi| \geq \varepsilon} \frac{\phi(\xi)}{\xi},$$

which converges because ϕ is smooth. Any other choice differs from this principal value by a value supported at the origin, and thus differs by a constant multiple of the Dirac delta function (the only distribution supported at the origin which is homogeneous of degree -1), so we might as well guess that f is given by the principal value formula above, and fix the function by adding or subtracting Dirac deltas if necessary. Thus we assume that

$$\int f(x)\phi(x) = \lim_{\varepsilon \rightarrow 0} \frac{-1}{i\pi x}.$$

But we now calculate that

$$\begin{aligned} \int \hat{f}(\xi)\phi(\xi) d\xi &= \frac{-1}{i\pi} \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \frac{\hat{\phi}(x)}{x} dx \\ &= \frac{1}{\pi} \int \phi(\xi) \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^{\infty} \frac{\sin(2\pi\xi x)}{x} dx d\xi. \end{aligned}$$

Note that we may assume without loss of generality that ϕ is smooth and compactly supported, which justifies that we can apply Fubini's theorem, and to bring the limit into the integral, since then the inner integrals converge uniformly on the support of ϕ . It is easy to see via a change of variables this limiting quantity is radial and odd in ξ , and thus this quantity is equal to

$$\left(\frac{1}{\pi} \int \phi(\xi) \text{sign}(\xi) d\xi \right) \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^{\infty} \frac{\sin(2\pi x)}{x} dx d\xi.$$

One can explicitly calculate this integral, e.g. using contour integration techniques. But we can also now rely on a little bit of a cheat: we *already know* from prior calculations that our choice of f differs from the right answer by a distribution supported at the origin, which implies the limiting integral must be equal to π , for otherwise our answer will differ by a quantity that is supported on the entire real line. Thus we have chosen the right answer.

Alternately, to verify that our choice of f above is correct, it suffices only to plug in a Gaussian $\phi(x) = e^{-\pi|x|^2}$, which has the property that $\hat{\phi}(\xi) = \phi(\xi)$, and then because the Gaussian is even, we find that

$$\begin{aligned} \int \hat{f}(\xi) e^{-\pi|\xi|^2} &= \int f(x) e^{-\pi|x|^2} \\ &= \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \frac{-1}{i\pi x} e^{-\pi|x|^2} \\ &= \lim_{\varepsilon \rightarrow 0} 0 = 0. \end{aligned}$$

Since we know $\widehat{f}(\xi) = \text{sgn}(\xi) + C\delta$ for some constant C , we find that

$$0 = \int \widehat{f}(\xi) e^{-\pi|\xi|^2} = \int \text{sgn}(\xi) e^{-\pi|\xi|^2} + C = C.$$

Thus f is actually the right answer.

We finish by giving the rough idea of how to calculate the distributional derivative of f . One approach is to multiply by a power of ξ on the Fourier transform side, and then take a Fourier transform, but we choose to work directly from the definition of the distributional derivative.

First, we calculate directly using the definition of the distributional derivative that for any test function ϕ ,

$$\int \delta'_0(x) \phi(x) dx = - \int \delta_0(x) \phi'(x) dx = -\phi'(0).$$

Thus δ'_0 maps a test function ϕ to $-\phi'(0)$. Similarly, the derivative of $\delta_{-1/2\pi}$ maps ϕ to $-\phi'(-1/2\pi)$, and the derivative of $\delta_{1/2\pi}$ maps ϕ to $-\phi'(1/2\pi)$. This allows us to calculate the distributional derivative of Γ simply. For the distribution $u : \phi \mapsto D^{12}\phi(0)$ we calculate

$$\int u'(x) \phi(x) dx = - \int u(x) \phi'(x) dx = -D^{12}\phi'(0) = -D^{13}(0).$$

This enables us to calculate the distributional derivative of Λ . Finally,

$$\int \text{p.v}(1/x)' \cdot \phi(x) dx = - \int \text{p.v}(1/x) \phi'(x) dx = - \lim_{\delta \rightarrow 0} \int_{|x| \geq \delta} (1/x) \phi'(x) dx.$$

Integration by parts shows that

$$\int_{\delta}^{\infty} (1/x) \phi'(x) dx = -\phi(\delta)/\delta + \int_{\delta}^{\infty} (1/x^2) \phi(x) dx$$

and

$$\int_{-\infty}^{-\delta} (1/x) \phi'(x) dx = -\phi(-\delta)/\delta + \int_{-\infty}^{-\delta} (1/x^2) \phi(x) dx.$$

Thus

$$\begin{aligned} \lim_{\delta \rightarrow 0} - \int_{|x| \geq \delta} (1/x) \phi'(x) dx &= \lim_{\delta \rightarrow 0} \frac{\phi(\delta) + \phi(-\delta)}{\delta} - \int_{|x| \geq \delta} (1/x^2) \phi(x) dx \\ &= \lim_{\delta \rightarrow 0} \frac{2\phi(0)}{\delta} - \int_{|x| \geq \delta} (1/x^2) \phi(x) dx. \end{aligned}$$

Thus if we define the distribution $\text{f.p}(1/x^2)$ (the finite part distribution) as mapping a test function ϕ to

$$\lim_{\delta \rightarrow 0} \int_{|x| \geq \delta} (1/x^2) \phi(x) - 2\phi(0)/\delta,$$

then the derivative of $\text{p.v}(1/x)$ is $-\text{f.p}(1/x^2)$. This allows us to calculate the distributional derivative of h .

14 Fourier Analysis / Tempered Distribution Notes

With distribution theory, there are a couple simple facts that allow one to convert technical sounding problems into more basic problems. Fourier analysis consists of some more deep tools than this, so I'd recommend making a study of these tools separately. But the theory of Fourier analysis applied to tempered distributions is very similar to the theory of distributions, in the sense that if you know Fourier analysis, you can often convert problems involving tempered distributions into more basic problems about Fourier analysis / other analysis techniques.

To begin with, let's review the Fourier transform. If $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is integrable, we define the Fourier transform $\hat{f} : \mathbb{R}^d \rightarrow \mathbb{C}$ by the integral formula

$$\hat{f}(\xi) = \int f(x) e^{-2\pi i \xi \cdot x} dx.$$

To avoid technical assumptions, from now on we assume we are working with functions in the *Schwartz class* $\mathcal{S}(\mathbb{R}^d)$, which consists of smooth functions $f \in C^\infty(\mathbb{R}^d)$ which are rapidly decaying, i.e. such that for any $N > 0$,

$$|f(x)| \lesssim_N \frac{1}{(1 + |x|)^N}.$$

The Fourier transform is then a bijection from $\mathcal{S}(\mathbb{R}^d)$ to itself, with inverse given by

$$\check{g}(x) = \int g(\xi) e^{2\pi i \xi \cdot x} d\xi.$$

One can then verify that the *multiplication formula*

$$\int \hat{f}(\xi) g(\xi) d\xi = \int f(x) \hat{g}(x) dx$$

holds. This will enable us to use duality to extend the definition of the Fourier transform to a family of distributions.

Distributions on \mathbb{R}^d were defined as continuous linear functionals on $\mathcal{D}(\mathbb{R}^d) = C_c^\infty(\mathbb{R}^d)$. Some of these distributions extend to continuous linear functionals on $\mathcal{S}(\mathbb{R}^d)$, which contains $\mathcal{D}(\mathbb{R}^d)$ as a subclass. These distributions are called *tempered*, and the class of all such tempered distributions is denoted $\mathcal{S}'(\mathbb{R}^d)$. Again, there is an abstract theory which determines when a linear functional on $\mathcal{S}(\mathbb{R}^d)$ is continuous. But in practice, it suffices to verify the following: a tempered distribution u is a functional that associates with each $\phi \in \mathcal{S}(\mathbb{R}^d)$ a quantity $\langle u, \phi \rangle$, such that there exists $N > 0$ and $M > 0$ such that

$$|\langle u, \phi \rangle| \leq \sup_{|\alpha| \leq N} \sup_{x \in \mathbb{R}^d} \frac{|D^\alpha \phi(x)|}{(1 + |x|)^M}.$$

If u is a tempered distribution, we formally define the Fourier transform of u , a tempered distribution \hat{u} , by the formula

$$\langle \hat{u}, \phi \rangle = \langle u, \hat{\phi} \rangle,$$

which behaves like the multiplication formula above. The Fourier transform is then a bijection of $\mathcal{S}'(\mathbb{R}^d)$, where the inverse Fourier transform is given by

$$\langle \check{u}, \phi \rangle = \langle u, \check{\phi} \rangle,$$

where the inverse Fourier transform $\check{\phi}$ is defined above.

We now summarize several useful properties of the Fourier transform:

- Because the Fourier transform is a bijection, if $\hat{u} = 0$, then $u = 0$.

- If f is an integrable function, then \widehat{f} is a continuous function with $\lim_{|x| \rightarrow \infty} |f(x)| = 0$ (the *Riemann-Lebesgue Lemma*), and $\|\widehat{f}\|_{L^\infty} \leq \|f\|_{L^1}$.
- The Fourier transform is a bijection of $L^2(\mathbb{R}^d)$, and for any $f \in L^2(\mathbb{R}^d)$, $\|\widehat{f}\|_{L^2} = \|f\|_{L^2}$ (*Parseval's Formula*). In particular, if u is a tempered distribution, and $\widehat{u} \in L^2(\mathbb{R}^d)$, then $u \in L^2(\mathbb{R}^d)$.
- The Fourier transform of $D^\alpha u$ is equal to $(2\pi i \xi)^\alpha \widehat{u}$.
- The Fourier transform of $(-2\pi i x)^\alpha u$ is equal to $D^\alpha \widehat{u}$.
- If we define the convolution of $\phi_1, \phi_2 \in \mathcal{S}(\mathbb{R}^d)$ by setting

$$(\phi_1 * \phi_2)(x) = \int \phi_1(y) \phi_2(x - y) dy,$$

then $\phi_1 * \phi_2$ is Schwartz, and its Fourier transform is $\widehat{\phi_1 * \phi_2}$. There is a way to define the convolution of a distribution u with a Schwartz function ϕ , denoted $u * \phi$, which will be a tempered distribution. The important thing to remember is that the Fourier transform of this convolution is also equal to $\widehat{u} \widehat{\phi}$.

15 Day 9: Fourier Analysis + Distribution Theory

30. (Spring 2017) Let $f \in L^1(\mathbb{R}^n)$ be a function all of whose distributional derivatives $D^\alpha f$ of order $|\alpha| = m$ also belong to $L^1(\mathbb{R}^n)$. Show that if $m > n$, then $f \in C(\mathbb{R}^n)$.

Solution: The duality between differentiation and multiplication by polynomials in the Fourier transform, and the fact that the Fourier transform of an L^1 function is in L^∞ , implies that for each α with $|\alpha| = m$,

$$|\xi^\alpha \hat{f}| \in L^\infty(\mathbb{R}^n).$$

We have $\sum_{|\alpha|=m} |\xi^\alpha| \sim |\xi|^m$, so we conclude that $|\xi^m \hat{f}| \in L^\infty$. Since $f \in L^1$, $\hat{f} \in L^\infty$, so that we have a bound

$$|\hat{f}(\xi)| \lesssim \frac{1}{1 + |\xi|^m}.$$

Since $m > n$, this means \hat{f} is also integrable. The Fourier inversion formula thus implies that f agrees almost everywhere with a bounded, continuous function.

31. (Fall 2019) Let $f \in L^2(\mathbb{R})$. Define

$$g(x) = \int_{-\infty}^{\infty} f(x-y)f(y) dy$$

Show that there exists a function $h \in L^1(\mathbb{R})$ such that

$$g(\xi) = \int_{-\infty}^{\infty} e^{-i\xi x} h(x) dx,$$

i.e. g is a Fourier transform of a function in $L^1(\mathbb{R})$. Hint: The following formal argument may be helpful:

$$\hat{g}(x) = \widehat{f * f}(x) = \hat{f}(x)^2,$$

where $*$ denotes convolution, and $\hat{\cdot}$ denotes the Fourier transform.

Solution: If one only knows Fourier analysis in $L^1(\mathbb{R})$, the argument in the hint may be justified using the fact that $L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ is dense in $L^2(\mathbb{R})$, together with some continuity arguments.

Young's convolution inequality implies that g is a continuous function in $L^\infty(\mathbb{R})$. One cannot take the Lebesgue integral defining the Fourier transform for g , but one may take the distributional Fourier transform. If we consider a family of functions $\{f_n\}$ lying in $L^2(\mathbb{R}) \cap L^1(\mathbb{R})$, which converge to f in $L^2(\mathbb{R})$ as $n \rightarrow \infty$, then Young's convolution inequality implies that $f_n * f_n$ converges to $f * f$ in $L^\infty(\mathbb{R})$. In particular, $f_n * f_n$ converges to $f * f$ distributionally. By continuity of the Fourier transform, $\widehat{f_n * f_n}$ converges to $\widehat{f * f}$, and $\widehat{f_n * f_n} = \widehat{f_n}^2$, which converges distributionally to \hat{f}^2 (by Cauchy Schwartz it actually converges in $L^1(\mathbb{R})$, which is a stronger statement). Thus we conclude that $\hat{g} = \widehat{f * f} = \hat{f}^2$. But since the Fourier transform of g lies in L^1 , we know that g acts as a distribution in exactly the same way as the distribution

$$\xi \mapsto \int \hat{f}(x)^2 e^{2\pi i \xi \cdot x}.$$

Since g is a continuous function, and the integral above defines a continuous function, it follows that for all ξ ,

$$g(\xi) = \int \hat{f}(x)^2 e^{2\pi i \xi \cdot x}.$$

32. (Fall 2015) Recall that $H^s(\mathbb{R}^n)$ is the Sobolev space consisting of all tempered distributions g on \mathbb{R}^n for which the Fourier transform \widehat{g} of g is locally integrable and satisfies

$$\int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\widehat{g}(\xi)|^2 d\xi < \infty.$$

Let u be a Schwartz function on \mathbb{R}^n and for $a \in \mathbb{C}$, let

$$f_a(x) = |x|^a u(x).$$

Show that if $\operatorname{Re}(a) > -n/2$ and $s \in [0, \operatorname{Re}(a) + n/2)$, then $f_a \in H^s(\mathbb{R}^n)$.

Solution: We wish to compute the Fourier transform of f_a , in order to apply the H^s norm. f_a is the product of a Schwartz function with $|x|^a$, so a useful avenue is to study the distributional derivative of $\Lambda(x) = |x|^a$, since $\widehat{f_a} = \widehat{\Lambda} * \widehat{u}$. The important properties of Λ we will use is that Λ acts as a C^∞ functions on $\mathbb{R}^n - \{0\}$, is a radial function, and is *homogeneous* of degree a . It follows from general Fourier analysis that $\widehat{\Lambda}$ acts as a C^∞ function on $\mathbb{R}^n - \{0\}$, is a radial function, and is homogeneous of degree $-a - n$. Thus we can write $\widehat{\Lambda}(\lambda\theta) = \lambda^{-a-n} g(\theta)$ where g is a C^∞ function defined for $|\theta| = 1$, and $\lambda > 0$. Since $\operatorname{Re}(a) > -n/2$, $\operatorname{Re}(-a-n) < -n/2$. This implies that $\widehat{\Lambda}$ decays away from the origin. In particular, $\widehat{\Lambda}$ is L^2 away from the origin.

Now we write

$$|x|^a u(x) = T_1 u(x) + T_2 u(x),$$

where

$$\widehat{T_1 u}(\xi) = \int_{|\eta| \leq 1} |\eta|^{-a-n} g(\eta/|\eta|) \widehat{u}(\xi - \eta) d\eta,$$

and

$$\widehat{T_2 u}(\xi) = \int_{|\eta| \geq 1} |\eta|^{-a-n} g(\eta/|\eta|) \widehat{u}(\xi - \eta) d\eta.$$

Since u is Schwartz, \widehat{u} is also Schwartz. In particular, for $|\eta| \geq 2|\xi|$, $|\widehat{u}(\xi - \eta)| \lesssim_N |\eta|^{-N}$ for all $N > 0$, which implies that

$$\left| \int_{|\eta| \geq 2|\xi|} |\eta|^{-a-n} g(\eta/|\eta|) \widehat{u}(\xi - \eta) d\eta \right| \lesssim_M |\xi|^{-M}.$$

For $1 \leq |\eta| \leq |\xi|/2$, $|\widehat{u}(\xi - \eta)| \lesssim_N |\xi|^{-N}$, which implies that

$$\left| \int_{1 \leq |\eta| \leq |\xi|/2} |\eta|^{-a-n} g(\eta/|\eta|) \widehat{u}(\xi - \eta) d\eta \right| \lesssim_N |\xi|^{-N}.$$

For $|\xi|/2 \leq |\eta| \leq 2|\xi|$, we can really only use the bound $|\widehat{u}(\xi - \eta)| \lesssim 1$, which implies that

$$\left| \int_{|\xi|/2 \leq |\eta| \leq 2|\xi|} |\eta|^{-a-n} g(\eta/|\eta|) \widehat{u}(\xi - \eta) d\eta \right| \lesssim |\xi|^{-\operatorname{Re}(a)}.$$

Putting these bounds together shows that

$$|\widehat{T_2 u}(\xi)| \lesssim |\xi|^{-\operatorname{Re}(a)},$$

which is sufficient to show that $T_2 u \in H^s$ for $0 \leq s \leq \operatorname{Re}(a) + n/2$.

To show $T_1 u \in H^s$, we ‘cheat’ a little, since this is the average of u over ‘low frequencies’ (and a good intuition to have is that low frequencies are very smooth). More precisely, $\widehat{T_1 u} = \widehat{\Psi} * \widehat{u}$, where $\widehat{\Psi}$ is a compactly supported distribution. The Paley-Wiener theorem implies that Ψ is a C^∞ function, and there exists some $N > 0$ such that $|\Psi(x)| \lesssim_N 1 + |x|^N$. And $T_1 u = \Psi \cdot u$, so $T_1 u$ is therefore actually a *Schwartz function*, and thus certainly in $H^s(\mathbb{R}^n)$.

Day 10: Warm Up Problems

33. (Fall 2021) Let σ be a Borel probability measure on $[0, 1]$ satisfying

1. $\sigma([1/3, 2/3]) = 0$.
2. $\sigma([a, b]) = \sigma([1 - b, 1 - a])$ for any $0 \leq a < b \leq 1$.
3. $\sigma([3a, 3b]) = 2\sigma([a, b])$ for any a, b such that $0 \leq 3a < 3b \leq 1$.

σ is called the $1/3$ Cantor measure on $[0, 1]$.

(a) Find $\sigma([0, 1/8])$.

Solution: Property 2 implies that

$$\sigma([0, 1/3]) = \sigma([2/3, 1]).$$

Since σ is a probability measure, and $\sigma([1/3, 2/3]) = 0$, this implies that σ gives mass $1/2$ to each of these sets. Property 1 and 3 implies that

$$\sigma([1/9, 2/9]) = (1/2)\sigma([1/3, 2/3]) = 0.$$

Since $1/9 < 1/8 < 2/9$, we find that

$$\sigma([0, 1/8]) = \sigma([0, 1/9]).$$

But now Property 1 implies that

$$\sigma([0, 1/9]) = (1/2)\sigma([0, 1/3]) = 1/4.$$

Thus $\sigma([0, 1/8]) = 1/4$.

(b) Calculate the second moment of σ , i.e. the integral

$$M = \int_0^1 x^2 d\sigma(x).$$

Solution: The trick to calculating M is to exploit the recursive nature of the Cantor measure to write M in terms of itself. First, we note that

$$\int_0^1 x d\sigma(x) = 1/2.$$

To see why this is true, we note that Property 2. implies that $d\sigma(1 - x) = d\sigma(x)$, and so a change of variables gives that

$$\int_0^1 x d\sigma(x) = \int_0^1 (1 - x) d\sigma(1 - x) = \int_0^1 (1 - x) d\sigma(x) = 1 - \int_0^1 x d\sigma(x).$$

Rearranging this equation gives the result.

Now we have done this, let's address the second moment. Property 1. implies that we have $d\sigma(x/3) = (1/2)d\sigma(x)$, and so a change of variables gives that

$$\int_0^{1/3} x^2 d\sigma(x) = \int_0^1 (x/3)^2 d\sigma(x/3) = (1/18) \int_0^1 x^2 d\sigma(x) = M/18.$$

Since $\sigma([1/3, 2/3]) = 0$,

$$\int_{1/3}^{2/3} x^2 d\sigma(x) = 0.$$

Finally, using Property 2. of the integral, we have $d\sigma(1-x) = d\sigma(x)$, which gives that

$$\int_{2/3}^1 x^2 d\sigma(x) = \int_0^{1/3} (1-x)^2 d\sigma(1-x) = \int_0^{1/3} (1-2x+x^2) d\sigma(x).$$

Now

$$\int_0^{1/3} d\sigma(x) = \sigma([0, 1/3]) = 1/2.$$

Another change of variables gives that

$$\int_0^{1/3} x d\sigma(x) = \int_0^1 (x/3) d\sigma(x/3) = (1/6) \int_0^1 x d\sigma(x) = (1/12).$$

Finally, we have already calculated that

$$\int_0^{1/3} x^2 d\sigma(x) = M/18.$$

Putting everything together, we conclude that

$$\int_{2/3}^1 x^2 d\sigma(x) = 1 - 2/12 + M/18 = 1 - 1/6 + M/18.$$

Thus

$$M = \int_0^1 x^2 d\sigma(x) = (M/18) + (0) + (1 - 1/6 + M/18) = M/9 + (5/6).$$

Rearranging gives

$$M = 15/16.$$

34. (Fall 2021) Find the spectrum of the linear operator A on $L^2(\mathbb{R})$ defined as

$$Af(x) = \int_{-\infty}^{\infty} \frac{f(y)}{1 + (x-y)^2} dy$$

(The spectrum of a linear operator T is the closure of the set of all complex numbers λ such that the operator $T - \lambda$ does not have a bounded inverse). Hint: It may be useful to find the Fourier transform of $1/(1+x^2)$.

Solution: That Fourier analysis might help us in this problem is suggest by the fact that $Af = K*f$, where

$$K(x) = \frac{1}{1+x^2}.$$

Thus the operator is a convolution operator. If $m = \widehat{K}$, then we can view A as a *Fourier multiplier operator*, i.e. such that

$$\widehat{Af}(\xi) = m(\xi)\widehat{f}(\xi).$$

There are several ways to compute K . The first is to calculate it explicitly, writing

$$m(\xi) = \int_{-\infty}^{\infty} \frac{e^{-2\pi i \xi \cdot x}}{1+x^2} dx$$

and apply contour integration. The second is to note that since

$$\left(1 - \frac{1}{4\pi^2}(-2\pi i x)^2\right) K = (1+x^2)K = 1,$$

taking Fourier transforms on both sides gives that m satisfies the differential equation

$$D^2 m = (4\pi^2)(m - \delta).$$

Solving this equation gives

$$m(\xi) = Ae^{2\pi\xi} + Be^{-2\pi\xi} + \pi[e^{2\pi x}H(x) - e^{-2\pi x}H(x)],$$

Since m is tempered, the constants must be chosen to avoid exponential increase near infinity. Thus we must set $A = \pi$, and $B = 0$, which gives

$$m(\xi) = \pi e^{-2\pi|\xi|}.$$

This viewpoint will help us find *approximate eigenvalues* for A , which, by virtue of the fact that A is self-adjoint, are the only barriers to $A - \lambda$ being invertible. Morally speaking, if $e_{\xi_0}(x) = e^{2\pi i \xi_0 \cdot x}$, then $\widehat{e_{\xi_0}} = \delta_{\xi_0}$, and so the Fourier transform of Ae_{ξ_0} *should* be equal to $m(\xi_0)\delta_{\xi_0}$, and thus we should have

$$Ae_{\xi_0} = m(\xi_0)e_{\xi_0}.$$

Thus e_{ξ_0} is an eigenfunction of A with eigenvalue $m(\xi_0)$. *However*, e_{ξ_0} does not lie in $L^2(\mathbb{R})$, and in fact, A has no eigenfunctions. To fix this, we simply approximate e_{ξ_0} by elements of $L^2(\mathbb{R})$, i.e. considering a Schwartz function ϕ with $\|\phi\|_{L^2(\mathbb{R})} = 1$, with $\widehat{\phi}$ supported on $[-1, 1]$, and with $\widehat{\phi} \geq 0$, and consider the functions

$$e_{\xi_0, R}(x) = R^{-1/2} e^{2\pi i \xi_0 \cdot x} \phi(x/R)$$

for $R > 0$. Then $\|e_{\xi_0, R}\|_{L^2(\mathbb{R})} = 1$ for all $R > 0$, and by applying Plancherel's formula,

$$\begin{aligned} \|Ae_{\xi_0, R} - m(\xi_0)e_{\xi_0, R}\|_{L^2(\mathbb{R})} &= \|[m(\xi) - m(\xi_0)\widehat{e_{\xi_0, R}}](\xi)\|_{L^2_{\xi}(\mathbb{R})} \\ &= R^{1/2} \left(\int |m(\xi) - m(\xi_0)|^2 |\widehat{\phi}(R(\xi - \xi_0))|^2 d\xi \right)^{1/2}. \end{aligned}$$

This integral is supported on $|\xi - \xi_0| \leq 1/R$, and on this region, by continuity of m , we have $|m(\xi) - m(\xi_0)| \lesssim_{\xi_0} 1/R$. Putting this into the integral above gives that

$$\|Ae_{\xi_0, R} - m(\xi_0)e_{\xi_0, R}\|_{L^2(\mathbb{R})} \leq R^{-1/2} = R^{-1/2} \|e_{\xi_0, R}\|_{L^2(\mathbb{R})}.$$

Taking $R \rightarrow \infty$ shows that $A - m(\xi_0)$ cannot be an isomorphism, since if T is an isomorphism, we have $\|Tf\|_{L^2(\mathbb{R})} \gtrsim \|f\|_{L^2(\mathbb{R})}$. Since the range of m includes all positive numbers less than or equal to π , we have shown that the spectrum contains $[0, \pi]$.

We claim this is *all* of the spectrum of A . Indeed, if $\alpha \notin [0, \pi]$, then $|m(\xi) - \alpha| \gtrsim 1$ for all $\xi \in \mathbb{R}$, and so

$$\|(A - \alpha)f\|_{L^2(\mathbb{R})} = \|[m(\xi) - \alpha]\widehat{f}(\xi)\|_{L^2(\mathbb{R})} \gtrsim \|\widehat{f}\|_{L^2(\mathbb{R})} = \|f\|_{L^2(\mathbb{R})}.$$

Thus $A - \alpha$ is an isomorphism onto its image, which must contain all of $L^2(\mathbb{R})$ because it has a continuous inverse, namely the operator $B_{\alpha} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ defined such that for $f \in L^2(\mathbb{R})$,

$$\widehat{B_{\alpha}f}(\xi) = \frac{\widehat{f}(\xi)}{m(\xi) - \alpha}.$$

16 Day 10: Bonus Questions

35. (Fall 2017) Let $a_1, \dots, a_n > 0$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be defined by

$$f(x) = \frac{1}{1 + \sum |x_i|^{\alpha_i}}.$$

Determine for each $p > 0$ whether

$$\int |f(x)|^p dx < \infty.$$

Solution: Let $E_i = \{x : x_i^{\alpha_i} \geq x_1^{\alpha_1}, \dots, x_n^{\alpha_n}\}$. On E_i , we have

$$|f(x)|^p \sim \frac{1}{1 + |x_i|^{p\alpha_i}}.$$

Let us determine what values of $p > 0$ we have

$$\int_{E_i} \frac{1}{1 + |x_i|^{p\alpha_i}} < \infty.$$

Since $\mathbb{R}^n = \bigcup E_i$, this will suffice to determine the range of p required in the problem. Set $E_i(n) = E_i \cap \{x : 2^{n-1} \leq x_1 \leq 2^n\}$. For $n > 0$, on $E_i(n)$ we have $|f(x)|^p \sim_p 1/2^{p\alpha_i n}$, and for $n < 0$ we have $|f(x)|^p \sim 1$. It is simple to calculate that $|E_i(n)| \sim 2^n 2^{n(\alpha_i/\alpha_2 + \dots + \alpha_i/\alpha_n)} = 2^{n\alpha_i(1/\alpha_1 + \dots + 1/\alpha_n)}$. For any value of p , we thus have

$$\sum_{n \leq 0} \int_{E_i(n)} |f(x)|^p \sim \sum_{n < 0} 2^{n\alpha_i(1/\alpha_1 + \dots + 1/\alpha_n)} < \infty.$$

On the other hand, we have

$$\sum_{n > 0} \int_{E_i(n)} |f(x)|^p \sim \sum_{n=1}^{\infty} 2^{-p\alpha_i n} 2^{n\alpha_i(1/\alpha_1 + \dots + 1/\alpha_n)} = \sum_{n=1}^{\infty} 2^{n\alpha_i(1/\alpha_1 + \dots + 1/\alpha_n - p)}.$$

This sum is finite if and only if $1/\alpha_1 + \dots + 1/\alpha_n < p$. Thus this is the range for which

$$\int_{E_i} |f(x)|^p < \infty.$$

But this condition is invariant of i , so this is precisely the range for which

$$\int |f(x)|^p < \infty.$$

Solution:

Let $\gamma = \frac{1}{\alpha_1} + \dots + \frac{1}{\alpha_n}$. We claim the integral converges if and only if $p > \gamma$.

Case 1: Suppose $p > \gamma$. Then

$$\int_{\mathbb{R}^n} |f(x)|^p dx = \int_0^\infty p t^{p-1} |E_t| dt$$

where $E_t = \{x : f(x) > 1\}$. Since $E_t = \emptyset$ if $t \geq 1$, it follows that

$$\int_{\mathbb{R}^n} |f(x)|^p dx = \int_0^1 p t^{p-1} |E_t| dt. \quad (13)$$

Observe that

$$\begin{aligned} E_t &= \left\{ x \in \mathbb{R}^n : \sum_{i=1}^n |x_i|^{\alpha_i} < \frac{1-t}{t} \right\} \\ &\subseteq \bigcap_{i=1}^n \left\{ x \in \mathbb{R}^n : |x_i| < \left(\frac{1-t}{t} \right)^{\frac{1}{\alpha_i}} \right\} \\ &= \left\{ x \in \mathbb{R}^n : - \left(\frac{1-t}{t} \right)^{\frac{1}{\alpha_i}} < x_i < \left(\frac{1-t}{t} \right)^{\frac{1}{\alpha_i}} \text{ for all } i \right\} \end{aligned}$$

Since the set on the right hand side is a rectangle with Lebesgue measure $2^n \left(\frac{1-t}{t} \right)^\gamma$, it follows that

$$|E_t| \leq 2^n \left(\frac{1-t}{t} \right)^\gamma.$$

Combining this with equation (13) gives

$$\int_{\mathbb{R}^n} |f(x)|^p dx \leq 2^n p \int_0^1 t^{p-1} (1-t)^\gamma t^{-\gamma} dt \leq 2^n p \int_0^1 t^{p-1-\gamma} dt$$

which is integrable since $p > \gamma$.

Case 2: Suppose $p \leq \gamma$. We will prove that $\int_{\mathbb{R}^n} |f|^p dx = +\infty$ by induction on n . If $n = 1$,

$$\int_{\mathbb{R}} \left(\frac{1}{1+|x|^{\alpha_1}} \right)^p dx \geq \frac{1}{2} \int_{|x|>1} |x|^{-\alpha_1 p} dx \geq \frac{1}{2} \int_{|x|>1} \frac{1}{|x|} dx = +\infty.$$

Next assume $n > 1$. For convenience of notation we will write $R = 1 + \sum_{i=2}^n |x_i|^{\alpha_i}$. We have

$$\begin{aligned} \int_{\mathbb{R}^n} |f(x)|^p dx &= \int_{\mathbb{R}^{n-1}} \left[\int_{\mathbb{R}} \left(\frac{1}{|x_1|^{\alpha_1} + 1 + \sum_{i=2}^n |x_i|^{\alpha_i}} \right)^p dx_1 \right] dx_2 \cdots dx_n \\ &= \int_{\mathbb{R}^{n-1}} \left[\int_{\mathbb{R}} \left(\frac{1}{|x_1|^{\alpha_1} + R} \right)^p dx_1 \right] dx_2 \cdots dx_n \\ &\geq \int_{\mathbb{R}^{n-1}} \left[\int_{|x_1| > R^{1/\alpha_1}} \left(\frac{1}{|x_1|^{\alpha_1} + R} \right)^p dx_1 \right] dx_2 \cdots dx_n \\ &\geq \frac{1}{2^p} \int_{\mathbb{R}^{n-1}} \left[\int_{|x_1| > R^{1/\alpha_1}} \left(\frac{1}{|x_1|^{\alpha_1}} \right)^p dx_1 \right] dx_2 \cdots dx_n \\ &= \frac{1}{2^{p-1}} \int_{\mathbb{R}^{n-1}} \int_{R^{1/\alpha_1}}^{\infty} x_1^{-p\alpha_1} dx_1 \cdots dx_n. \end{aligned}$$

If $p\alpha_1 \leq 1$ then $\int_{\mathbb{R}^{n-1}} \int_R^{\infty} x_1^{-p\alpha_1} dx_1 = +\infty$ and we are done. On the other hand, if $p\alpha_1 < 1$, then by

the above, we have

$$\begin{aligned} \int_{\mathbb{R}^n} |f(x)|^p dx &= \frac{1}{2^{p-1}(1-p\alpha_1)} \int_{\mathbb{R}^{n-1}} R^{\frac{1}{\alpha_1}-p} dx_2 \cdots dx_n \\ &= \frac{1}{2^{p-1}(1-p\alpha_1)} \underbrace{\int_{\mathbb{R}^{n-1}} \left(\frac{1}{1 + \sum_{i=2}^n |x_i|^{\alpha_i}} \right)^{p-\frac{1}{\alpha_1}} dx_2 \cdots dx_n}_{\text{Call this } A} \end{aligned}$$

However, since $p \geq \gamma$, it follows that $p - \frac{1}{\alpha_1} \geq \frac{1}{\alpha_2} + \dots + \frac{1}{\alpha_n}$. Therefore by the induction hypothesis, the integral $A = +\infty$. Therefore $\int_{\mathbb{R}^n} |f(x)|^p dx = +\infty$.

36. Let $f \in L^1(\mathbb{R})$. Let

$$G(\lambda) = \int_{-\infty}^{\infty} e^{i\lambda t^2} f(t) dt.$$

Prove that G is a continuous function and that $\lim_{\lambda \rightarrow \infty} G(\lambda) = 0$.

Solution: It is simple to see that the operator $Tf = G$ is continuous from $L^1(\mathbb{R})$ to $L^\infty(\mathbb{R})$, with $\|Tf\|_\infty \leq \|f\|_1$. We claim that it thus suffices to show the result for $\phi \in C_c^\infty(\mathbb{R})$ supported away from the origin. Indeed, if $f \in L^1(\mathbb{R})$, we can find $\phi_n \in C_c^\infty(\mathbb{R})$ supported away from the origin, converging in $L^1(\mathbb{R})$ to f . Then $T\phi_n$ converges to Tf uniformly, and $T\phi_n$ is continuous for each n , so Tf is continuous. Moreover, if $\varepsilon > 0$, there is n large enough that $\|f - \phi_n\|_1 \leq \varepsilon$. This means that $\|Tf - T\phi_n\|_\infty \leq \varepsilon$, and so

$$\limsup_{\lambda \rightarrow \infty} |Tf(\lambda)| \leq \limsup_{\lambda \rightarrow \infty} |Tf(\lambda) - T\phi_n(\lambda)| + \limsup_{\lambda \rightarrow \infty} |T\phi_n(\lambda)| \leq \varepsilon + 0 \leq \varepsilon.$$

Taking $\varepsilon \rightarrow 0$ completes the argument. So now we show the result for $\phi \in C_c^\infty(\mathbb{R})$. Continuity here is easy because ϕ is compactly supported (in this case, G is actually a smooth function). And to show that $T\phi(\lambda)$ converges to zero as $\lambda \rightarrow \infty$, we integrate by parts, writing

$$T\phi(\lambda) = \int_{-\infty}^{\infty} e^{i\lambda t^2} \phi(t) dt = \int_{-\infty}^{\infty} \frac{\phi(t)}{2i\lambda t} \frac{d}{dt} (e^{i\lambda t^2}) = -\frac{1}{2i\lambda} \int_{-\infty}^{\infty} \frac{d}{dt} \left(\frac{\phi(t)}{t} \right) e^{i\lambda t^2} dt.$$

Since ϕ lies in C_c^∞ and is supported away from the origin, so too is $\phi(t)/t$, and thus we find

$$|T\phi(\lambda)| \leq \frac{1}{2\lambda} \int_{-\infty}^{\infty} \left| \frac{d}{dt} \left(\frac{\phi(t)}{t} \right) \right| dt \lesssim 1/\lambda.$$

Thus $T\phi(\lambda) \rightarrow 0$ as $\lambda \rightarrow \infty$.

37. Let (X, μ) be a σ -finite measure space. Let $\{f_n\}$ be a sequence of measurable functions and assume that $f_n \rightarrow f$ almost everywhere. Prove that there exists measurable $A_1, A_2, \dots \subset X$ such that $\mu(X - \bigcup_i A_i) = 0$, and such that $f_n|_{A_i} \rightarrow f|_{A_i}$ uniformly for each i .

Solution: We may assume without loss of generality that X is actually a *finite* measure space, since if $\bigcup X_n = X$, with X_n a finite measure space, then we can find A_{n1}, A_{n2}, \dots such that $\mu(X_n - \bigcup_i A_{ni}) = 0$, such that f_k converges uniformly on each A_{ni} , and then $\mu(X - \bigcup_{n,i} A_{ni}) = 0$. Normalizing, we assume $|X| = 1$.

Now let

$$E_n(k) = \{x \in X : \sup_{m \geq n} |f_m(x) - f(x)| \leq 1/2^k\}.$$

Since $f_n \rightarrow f$ almost everywhere, as $n \rightarrow \infty$ and for a fixed $k > 0$, $\{E_n(k)\}$ is an increasing sequence, and $X - \bigcup_n E_n(k)$ is a set of measure zero. Thus for each $r > 0$, we can pick n_1 such that $|X - E_{n_1}(1)| \leq 1/2^r$. Similarly, we can for each k pick n_k such that $|E_{n_{k-1}}(k-1) - E_{n_k}(k)| \leq 1/2^{k+r}$. If we define $A_r = \bigcap_k E_{n_k}(k)$, then $|X - A_r| \leq 1/2^{r-1}$, and by definition, f_k converges uniformly on A_r for each r since for $n \geq n_k$, $|f_n(x) - f(x)| \leq 1/2^k$. Thus we have found the required sequence $\{A_r\}$.

38. (Spring 2017) Given for each function $f \in C^0(\mathbb{R}^2)$ we define for each $y \in \mathbb{R}$ a function $f_y \in C[0, 1]$ by $f_y(x) = f(x, y)$. Assume that for each fixed y , the distributional derivative of $f_y \in \mathcal{D}'(\mathbb{R})$ defines a function $a_y \in L^p(\mathbb{R})$. Assume further that

$$\|a_y\|_p \leq C < \infty$$

for some constant C independent of y . Show that the distributional derivative $\partial_x f \in \mathcal{D}'(\mathbb{R}^2)$ is in $L^p_{\text{loc}}(\mathbb{R}^2)$, provided $1 < p \leq \infty$.

Solution: We apply duality, which says that $\partial_x f$ is in L^p_{loc} if and only if for each $R > 0$, and $\phi \in C_c^\infty(\mathbb{R}^d)$ supported on $|z| \leq R$,

$$\left| \int_{\mathbb{R}^2} (\partial_x f)(z) \phi(z) dz \right| \lesssim_R \|\phi\|_{p^*},$$

where p^* is the conjugate to p . Now

$$\int_{\mathbb{R}^2} (\partial_x f)(z) \phi(z) dz = - \int f(z) (\partial_x \phi)(z) dz.$$

Since f is continuous, and $\partial_x \phi$ is compactly supported on $|z| \leq R$, we can apply Fubini's theorem, writing

$$\int f(z) (\partial_x \phi)(z) dz = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) (\partial_x \phi)(x, y) dx dy.$$

But applying integration by parts in just the x -variable, we have

$$\int_{-\infty}^{\infty} f(x, y) (\partial_x \phi)(x, y) dx = - \int a_y(x) \phi(x, y) dx.$$

Thus we conclude that

$$\int (\partial_x f)(z) \phi(z) dz = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} a_y(x) \phi(x, y) dx dy.$$

But by Hölder's inequality, if p^* is the dual to p ,

$$\begin{aligned}
 \left| \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} a_y(x) \phi(x, y) \, dx \, dy \right| &= \left| \int_{-R}^R \int_{-\infty}^{\infty} a_y(x) \phi(x, y) \, dx \, dy \right| \\
 &\leq \left(\int_{-R}^R \int_{-\infty}^{\infty} |a_y(x)|^p \, dx \, dy \right)^{1/p} \|\phi\|_{p^*} \\
 &= \left(\int_{-R}^R \|a_y\|_p^p \, dy \right)^{1/p} \|\phi\|_{p^*} \\
 &\leq C \cdot (2R)^{1/p} \|\phi\|_{p^*} \\
 &\lesssim_R \|\phi\|_{p^*}.
 \end{aligned}$$

Thus we conclude that $\partial_x f \in L_{\text{loc}}^p$.

39. (Spring 2020) Let $E \subset [0, 1]$ be a measurable set with positive Lebesgue measure. Moreover, it satisfies the following property: As long as x and y belong to E , we know $\frac{x+y}{2}$ belongs to E . Prove that E is an interval.

Solution: Let $m = \inf E$ and $M = \sup E$. Since $|E| > 0$, E contains more than one element and hence $m < M$. We first prove two claims:

Claim 1: E is dense in (m, M) .

Proof of Claim 1: Let $z \in (m, M)$. By definition of infimum and supremum, there exists $u, v \in E$ such that $m \leq u < z < v \leq M$. Let $\delta > 0$. It will suffice to show there exists an element $w \in E$ such that $|w - z| < \delta$. Define a sequence of intervals inductively by $I_0 = [u, v]$ and if $I_{n-1} = [x, y]$, then

$$I_n = \begin{cases} \left[\frac{x+y}{2}, y \right] & \text{if } w \geq \frac{x+y}{2} \\ \left[x, \frac{x+y}{2} \right] & \text{if } w < \frac{x+y}{2} \end{cases}$$

By construction, $w \in I_n$, and using the midpoint convexity of E , the endpoints of I_n are elements of E . Choosing N sufficiently large that $|I_N| = 2^{-N}(v - u) < \delta$, it follows that $|w - \max I_N| < |\max I_N - \min I_N| = 2^{-N}(v - u) < \delta$. This proves the claim. \square

Claim 2: E contains a nonempty interval.

Proof of Claim 2: Denote by $E/2$ the set $\{x/2 : x \in E\}$. First observe that $|E/2| > 0$. (Indeed, if $|E/2| = 0$, then by absolute continuity of the function $x \mapsto 2x$, $|E| = 0$, a contradiction.) We will show that $\frac{E}{2} + \frac{E}{2}$ contains a nonempty interval. The proof of the claim will then follow since $E \supset \frac{E+E}{2} = \frac{E}{2} + \frac{E}{2}$.

Let $h(x) = \int_{\mathbb{R}} \chi_A(x) \chi_A(x - y) dy$ where $A = E/2$. If $x \notin A + A$, then $h(x) = 0$. Therefore by contrapositive, if $h(x) > 0$ then $x \in A + A$. Moreover, h is continuous, so the set $\{x : h(x) > 0\}$ is open and therefore contains an interval provided it is nonempty. Therefore it remains to show that $\{x : h(x) > 0\}$ is nonempty. Indeed, by the Fubini-Tonelli theorem, $\int_{\mathbb{R}} h(x) dx = \int_A \int_{\mathbb{R}} \chi_A(x - y) dx dy = \int_A \int_{\mathbb{R}} \chi_A(x) dx dy = |A|^2 > 0$. Since h is nonnegative function whose integral is positive, it follows that $h(x) > 0$ for some x , as desired. \square

We now proceed with the proof. By Claim 2, there exists a nonempty interval $(a, b) \subset E$. Let $b_0 = \sup \{r > 0 : (a, r) \subset E\}$ and $a_0 = \inf \{s > 0 : (s, b) \subset E\}$. It will suffice to show that $a_0 = m$ and $b_0 = M$, since this will imply that $(m, M) \subset E$, which is enough to conclude that E is an interval. We shall show that $b_0 = M$ as the proof that $a_0 = m$ is similar.

Suppose $b_0 < M$. By Claim 1, there exists $b_1 \in E$ such that $b_0 < b_1 < 2b_0 - a$. Since $(a, b_0) \subset E$ and $b_1 \in E$, it follows that

$$E \supset \left\{ \frac{b_1 + s}{2} : a < s < b_0 \right\} = \left\{ t : \frac{a + b_1}{2} < t < \frac{b_0 + b_1}{2} \right\} = \left(\frac{a + b_1}{2}, \frac{b_0 + b_1}{2} \right).$$

Since $b_1 < 2b_0 - a$, we have $\frac{a+b_1}{2} < b_0$. Therefore $(a, \frac{b_0+b_1}{2}) = (a, b_0) \cup (\frac{a+b_1}{2}, \frac{b_0+b_1}{2}) \subset E$. Moreover, since $b_0 < b_1$, we have $\frac{b_0+b_1}{2} > b_0$. Therefore b_0 is not an upper bound for the set $\{r > 0 : (a, r) \subset E\}$. But this is a contradiction since we defined b_0 to be the supremum of that set. Therefore $b_0 = M$.

40. (a) Does $p_N = \prod_{n=2}^N (1 + (-1)^n/n)$ tend to a nonzero limit as $N \rightarrow \infty$.

Solution: We have

$$\sum_{n=2}^N \log(1 + (-1)^n/n) = \sum_{n=2}^N (-1)^n/n + O(1/n^2).$$

The error terms converge absolutely, since $1/n^2$ is summable, and Leibniz's test implies since $(-1)^n/n$ is an alternating sequence decreasing in absolute value, that

$$\sum_{n=2}^N (-1)^n/n$$

converges as $N \rightarrow \infty$, and that the limiting value exceeds

$$1/2 - 1/3 > 0.$$

Thus taking exponentials shows that p_N converges as $N \rightarrow \infty$.

- (b) Does $q_N = \prod_{n=2}^N (1 + (-1)^n/\sqrt{n})$ tend to a nonzero limit as $N \rightarrow \infty$.

Solution: We apply Taylor series, writing

$$\log(1 + (-1)^n/\sqrt{n}) = (-1)^n/\sqrt{n} - 1/2n + O(1/n^{3/2}).$$

The error term is absolutely summable, and the sum of $(-1)^n/\sqrt{n}$ converges by Leibniz test as above. Thus the convergence of the sum

$$\sum_{n=2}^N \log(1 + (-1)^n/\sqrt{n})$$

is equivalent to the convergence of the sum

$$\sum_{n=2}^N -1/2n.$$

But this sum converges to $-\infty$, so we find that, taking exponentials, $q_N \rightarrow 0$ as $N \rightarrow \infty$.

Day 11: Warm Up Question

41. (Fall 2015) Let $\chi \in C^\infty(\mathbb{R})$ have a compact support and define

$$f_n(x) = n^2 \chi'(nx).$$

- (a) Does f_n converge in the sense of distributions as $n \rightarrow \infty$? If so, what is the limit?

Solution: Applying an integration by parts, we calculate that

$$\int f_n(x) \phi(x) dx = \int n^2 \chi'(nx) \phi(x) dx = - \int n \chi(nx) \phi'(x) dx.$$

As $n \rightarrow \infty$, each function $x \mapsto n \chi(nx)$ has total mass one, but is concentrated in small and smaller neighborhoods of the origin. In particular, these functions operate as an approximation to the identity, so that

$$- \int n \chi(nx) \phi'(x) dx = -\phi'(0).$$

Thus as $n \rightarrow \infty$, f_n converges distributionally to the distribution $\phi \mapsto -\phi'(0)$.

- (b) Let $p \in [1, \infty)$ and $g \in L^p(\mathbb{R})$ be such that the distributional derivative of g also lies in $L^p(\mathbb{R})$. Does $f_n * g$ converge in $L^p(\mathbb{R})$ as $n \rightarrow \infty$? If so, what is the limit?

Solution: As mentioned before, integration by parts shows that

$$(f_n * g)(x) = \int_{-\infty}^{\infty} f_n(y) g(x-y) dy = \int n \chi(ny) g'(x-y) dy = (n \chi(ny) * g')(x).$$

The function $n \chi(ny)$ is an approximation to the identity, and so as $n \rightarrow \infty$, $(f_n * g)$ converges in L^p to g' .

42. (Fall 2018) Prove or Disprove that in an infinite dimensional Banach space,

- (a) every norm bounded set is weakly bounded,
- (b) every norm closed set is weakly closed
- (c) a norm bounded set has empty interior in the weak topology

Solution: We first prove part (a). Let X be a Banach space and suppose $E \subset X$ is norm bounded; i.e., that $\sup_{x \in E} \|x\| = C < \infty$. Recall that to show E is weakly bounded, we need to show that $\sup_{x \in E} |\phi(x)| < \infty$ for all $\phi \in X^*$. This follows immediately by definition of norm:

$$\sup_{x \in E} |\phi(x)| \leq \sup_{x \in E} \|\phi\| \|x\| \leq C \|\phi\| < \infty.$$

Next we prove (c). Let X be an infinite-dimensional Banach space, and let $A \subset X$ be a norm-bounded set. Then there exists $\rho > 0$ such that $A \subset B := \{x \in X : \|x\| < \rho\}$. It suffices to show that B has empty interior in the weak topology.

By definition, the weak topology has a base consisting of sets of the form

$$N(x_0, F, r) = \{x \in X : |\phi(x - x_0)| < r \text{ for all } \phi \in F\} \quad (14)$$

where $x_0 \in X$, $r > 0$, and F is a finite subset of X^* . Letting $K := \bigcap_{\phi \in F} \ker(\phi)$, observe that

$$x_0 + K \subset N(x_0, F, r).$$

Since $\ker(\phi)$ is of co-dimension 1 in X for every $\phi \in X^*$ and since F is a finite set, it follows that K is of finite codimension in X . Therefore since X is infinite-dimensional, it follows that $\dim K > 0$, and therefore K contains a nonzero element y . Since K is a subspace, it follows that $\alpha y \in K$ for any $\alpha \in \mathbb{R}$. This implies that K —and therefore $N(x_0, F, r)$ —contains elements of arbitrarily large norm. Therefore $N(x_0, F, r) \not\subset B$. Therefore B has empty interior in the weak topology. This proves statement (c).

Incidentally, we have also shown that statement (b) is false: while B^c is clearly norm closed, it is not weakly closed because B is not weakly open.

Finally we claim that the converse of statement (b)—i.e., that every weakly closed set is norm closed—is true. To see this, let F be weakly closed. Then F^c is weakly open, and hence can be written as a union of sets of the form given in equation (18). Moreover, since $x \mapsto |\phi(x - x_0)|$ is continuous with respect to the norm topology, it follows that $N(x_0, F, r)$ is open in the norm topology. Therefore F^c is a union of norm open sets. Therefore F^c is norm open, and hence F is norm closed.

Another way to prove the converse of statement (b) is the following. Suppose $F \subset X$ is weakly closed, and suppose that $x_n \in F$ converges in norm to $x \in X$. To show that F is strongly closed, we need to show that $x \in F$. Using the linearity of ϕ and the definition of operator norm,

$$|\phi(x_n) - \phi(x)| = |\phi(x_n - x)| \leq \|\phi\| \|x_n - x\|$$

for every $\phi \in X^*$. Since the right-hand side converges to zero as $n \rightarrow \infty$, it follows that $\phi(x_n) \rightarrow \phi(x)$ as well, i.e. that x_n converges weakly to x . Therefore since F is weakly closed, $x \in F$, as required.

17 Day 11: Bonus Questions

43. Suppose $f : [0, 1]^2 \rightarrow \mathbb{R}$ is continuous, and $\partial^2 f / \partial x^2$ exists pointwise on $[0, 1]$, is continuous in the x variable, and is bounded. Then $\partial f / \partial x$ is continuous.

Solution: If $\partial_x f$ was not continuous, we could find $z \in [0, 1]^2$ and $R > 0$ such that for any $\delta > 0$, we can find $w \in [0, 1]^2$ with $|z - w| \leq \delta$ and with $|\partial_x f(w) - \partial_x f(z)| \geq R$. Applying Taylor's formula and the boundedness and continuity of $\partial_x^2 f$, there exists $C > 0$, such that if $z + te_1 \in [0, 1]^2$ and $w + te_1 \in [0, 1]^2$

$$|f(z + te_1) - f(z) - t\partial_x f(z)| \leq Ct^2.$$

and

$$|f(w + te_1) - f(w) - t\partial_x f(w)| \leq Ct^2.$$

Thus

$$|f(z + te_1) - f(w + te_1)| \geq |t||\partial_x f(z) - \partial_x f(w)| - |f(z) - f(w)| - 2Ct^2 \geq |t|R - |f(z) - f(w)| - 2Ct^2$$

Since f is continuous, it is uniformly continuous. Thus for any $\varepsilon > 0$, there is $\delta > 0$ such that if $|v - w| \leq \delta$, then $|f(v) - f(w)| \leq \varepsilon$. But

$$|(z + te_1) - (w + te_1)| = |z - w| \leq \delta,$$

so we conclude that $|f(z + te_1) - f(w + te_1)| \leq \varepsilon$ and $|f(z) - f(w)| \leq \varepsilon$, and thus that

$$|t|R - 2Ct^2 \leq 2\varepsilon$$

It is possible to pick $|t| = \min(1/10, R/4C)$. If $R/4C \geq 1/10$, then $R \geq 2C/5$, and so setting $t = 1/10$ gives

$$2\varepsilon \geq R/10 - C/50 \geq C/50.$$

Thus ε cannot be arbitrarily small, which gives a contradiction. If $R \leq 2C/5$, we can pick $|t| = R/4C$, which gives

$$2\varepsilon \geq R^2/8C.$$

This again prevents ε from being arbitrarily small.

44. Let

$$s_N(x) = \sum_{n=1}^N (-1)^n \frac{x^{3n}}{n^{2/3}}.$$

Prove that $s_N(x)$ converges to a limit $s(x)$ on $[0, 1]$, and that there is a constant $C > 0$ so that for all $N \geq 1$ the inequality

$$\sup_{x \in [0, 1]} |s_N(x) - s(x)| \leq CN^{-2/3}$$

holds.

Solution: The convergence here becomes highly singular near $x = 1$, because the power series diverges for $|x| > 1$. Thus we cannot use normal power series techniques here. It seems we must exploit the oscillation of the series here to get explicit bounds since removing the $(-1)^n$ in the definition of the series causes the sum to diverge at $x = 1$. If we define

$$S_N(x) = \sum_{n=1}^N (-x^3)^n$$

for $N > 0$, and $S_0 = 0$, then the geometric series formula shows that there is $C > 0$ such that $|S_N(x)| \leq C$ for all N and $x \in [0, 1]$, and a summation by parts shows that

$$\sum_{n=1}^N (-1)^n \frac{x^{3n}}{n^{2/3}} = \sum_{n=1}^N (S_n - S_{n-1}) \frac{1}{n^{2/3}} = \sum_{n=1}^{N-1} S_n \left(\frac{1}{n^{2/3}} - \frac{1}{(n+3)^{2/3}} \right) + S_N/n^{2/3}.$$

Now a mean value theorem / Taylor series expansion implies that

$$\left| \frac{1}{n^{2/3}} - \frac{1}{(n+3)^{2/3}} \right| \lesssim \frac{1}{n^{5/3}},$$

which is summable, and so for $M \geq N$,

$$|s_M(x) - s_N(x)| \lesssim \sum_{n=N}^{M-1} S_n/n^{5/3} + S_M/M^{2/3} \lesssim C/N^{2/3}.$$

This is sufficient to show the existence of the point wise limit (since it shows s_N is a Cauchy sequence), and also shows the required inequality.

45. (Spring 2016) Let $1 < p < \infty$, and let $\chi_{[1-\frac{1}{n}, 1]}$ denote the characteristic function of $[1-\frac{1}{n}, 1]$. For which $\alpha \in \mathbb{R}$ does the sequences $n^\alpha \chi_{[1-\frac{1}{n}, 1]}$ converge weakly to 0 in $L^p(\mathbb{R})$?

Solution: Case 1: Suppose $\alpha > 1/p$. Then $\|f_n\|_p^p = \int_{1-\frac{1}{n}}^1 n^{\alpha p} dx = n^{\alpha p-1} \rightarrow \infty$ as $n \rightarrow \infty$. Since weakly convergent sequences are bounded in norm, it follows that f_n does not weakly converge if $\alpha > 1/p$.

Case 2: Suppose $\alpha < 1/p$. Let $g \in L^q(\mathbb{R})$ where $\frac{1}{p} + \frac{1}{q} = 1$. Using Holder's inequality,

$$\int f_n g = \int_{1-\frac{1}{n}}^1 n^\alpha g \leq \left(\int_{\mathbb{R}} n^{\alpha p} \chi_{[1-\frac{1}{n}, 1]} \right)^{\frac{1}{p}} \|g\|_q = n^{\alpha-\frac{1}{p}} \|g\|_q \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since $g \in L^q$ was arbitrary, it follows that f_n converges weakly to 0 in L^p .

Case 3: Suppose $\alpha = 1/p$. Let $g \in L^q(\mathbb{R})$ where $\frac{1}{p} + \frac{1}{q} = 1$. By density of $C_c^\infty(\mathbb{R})$ in $L^q(\mathbb{R})$, there exists $\phi \in C_c^\infty(\mathbb{R})$ with $\|\phi - g\|_q < \epsilon$. Then

$$\begin{aligned} \left| \int_{\mathbb{R}} f_n g \right| &= \int_{\mathbb{R}} |f_n(g - \phi)| + \int_{\mathbb{R}} |f_n \phi| \\ &\leq \|f_n\|_p \underbrace{\|g - \phi\|_q}_{< \epsilon} + \int_{1-\frac{1}{n}}^1 n^{\frac{1}{p}} |\phi(x)| dx && \text{by Holder's inequality} \\ &\leq \epsilon + \int_{1-\frac{1}{n}}^1 n^{\frac{1}{p}} |\phi(x)| dx && \text{since } \|f_n\|_p = 1 \text{ for all } n \\ &\leq \epsilon + \|\phi\|_\infty \int_{1-\frac{1}{n}}^1 n^{\frac{1}{p}} dx && \text{since } \phi \in C_c^\infty(\mathbb{R}) \\ &= \epsilon + \|\phi\|_\infty n^{\frac{1}{p}-1}. \end{aligned}$$

Since $\frac{1}{p} < 1$, it follows that $\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}} f_n g \right| \leq \epsilon$. Since $\epsilon > 0$ and $g \in L^q(\mathbb{R})$ were arbitrary, it follows that $\lim_{n \rightarrow \infty} \int f_n g = 0$ for all $g \in L^q$. So f_n converges weakly to 0 in L^p .

46. (Spring 2018) Let x_n be a sequence in a Hilbert space H . Suppose that x_n converges to x weakly. Prove that there is a subsequence x_{n_k} such that

$$\frac{1}{N} \sum_{k=1}^N x_{n_k}$$

converges to x (in norm) as $N \rightarrow \infty$.

Solution: (Note: this result is called the Banach-Saks theorem). Without loss of generality, assume that $x = 0$. Since x_n converges weakly, it follows that x_n is bounded in norm (this is a consequence of the uniform boundedness principle). Let $C = \max_{n \geq 1} \|x_n\|$. We construct a subsequence in the following manner. Let $x_{n_1} = x_1$. Having chosen x_{n_N} , choose $x_{n_{N+1}}$ so that

$$\left\langle \sum_{i=1}^N x_{n_i}, x_{n_{N+1}} \right\rangle \leq 2^{-(N+1)}.$$

Such an $x_{n_{N+1}}$ exists since $\langle y, x_n \rangle \rightarrow 0$ as $n \rightarrow \infty$ for all $y \in H$. We need to show that

$$\left\| N^{-1} \sum_{k=1}^N x_{n_k} \right\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

To see this, we have:

$$\begin{aligned} N^{-2} \left\langle \sum_{k=1}^N x_{n_k}, \sum_{j=1}^N x_{n_j} \right\rangle &= N^{-2} \sum_{j,k=1}^N \langle x_{n_k}, x_{n_j} \rangle \\ &= N^{-2} \left(\sum_{j=1}^N \|x_{n_j}\|^2 + \sum_{k=2}^N \sum_{j=1}^{k-1} \langle x_{n_j}, x_{n_k} \rangle \right) \\ &= N^{-2} \left(\sum_{j=1}^N \|x_{n_j}\|^2 + \sum_{k=2}^N \underbrace{\left\langle \sum_{j=1}^{k-1} x_{n_j}, x_{n_k} \right\rangle}_{\leq 2^{-k}} \right) \\ &\leq N^{-2} (CN + 1) \rightarrow 0 \text{ as } N \rightarrow \infty. \end{aligned}$$

47. (Fall 2015) Let $E \subset \mathbb{R}$ be a measurable set, such that $E + r = E$ for all $r \in \mathbf{Q}$. Show that $|E| = 0$ or $|E^c| = 0$.

Solution: We apply the Lebesgue density theorem. Suppose $|E| \neq 0$. Fix $\delta > 0$. Then there exists $\varepsilon_0 > 0$ such that for $\varepsilon < \varepsilon_0$, we can find $a, b \in \mathbf{Q}$ with $b - a < \varepsilon$ with

$$|E \cap (a, b)| \geq (1 - \delta)(b - a)$$

Set $\varepsilon_1 = b - a$. Because E is invariant under translations in \mathbf{Q} ,

$$|E \cap (0, \varepsilon_1)| \geq (1 - \delta)\varepsilon_1.$$

Then $|E \cap (a, a + \varepsilon_1)| \geq (1 - \delta)\varepsilon_1$ for all $a \in \mathbf{Q}$. This implies that for any $x \in \mathbb{R}$,

$$\limsup_{\substack{x \in I \\ |I| \rightarrow 0}} \frac{|E \cap I|}{|I|} \geq 1.$$

The Lebesgue density theorem thus implies that $|E^c| = 0$.

48. (Spring 2015) Let $\{r_n\} \in [0, 1]$ be an arbitrary sequence, and define the function

$$f(x) = \sum_{r_n < x} \frac{1}{2^n}$$

Show that f is Borel measurable, find all its points of discontinuity, and find $\int_0^1 f(x) dx$.

Solution: Write

$$f_n(x) = \sum_{k=1}^n \mathbf{I}_{(r_n, \infty)} \cdot 2^{-n}.$$

Then for each n , f_n is a simple function, and is thus measurable. Moreover, $f = \lim_{n \rightarrow \infty} f_n$, and this limit is monotone. The pointwise limit of measurable functions is measurable, so f is measurable.

Next, we look at the discontinuity points. We claim that the set of discontinuity points is *precisely* the set of values $\{r_n\}$. To see this, fix $\varepsilon > 0$, and note that

$$f(x + \varepsilon) - f(x) = \sum_{x \leq r_n < x + \varepsilon} 2^{-n}$$

Suppose there exists some n_0 such that $x = r_{n_0}$. Then

$$f(x + \varepsilon) - f(x) \geq 1/2^{n_0},$$

and thus

$$\limsup_{\varepsilon \rightarrow 0^+} f(x + \varepsilon) - f(x) \geq 1/2^{n_0},$$

implying f is not continuous at x . On the other hand, if $x \neq r_n$ for any n , then for any $N > 0$, there exists ε_N such that for $\varepsilon < \varepsilon_N$, $\{r_1, \dots, r_N\}$ is disjoint from $[x, x + \varepsilon)$. Therefore, for $\varepsilon < \varepsilon_0$,

$$f(x + \varepsilon) - f(x) \leq \sum_{k=N+1}^{\infty} 2^{-k} = 2^{-N}.$$

Thus we conclude that

$$\lim_{\varepsilon \rightarrow 0^+} f(x + \varepsilon) - f(x) = 0.$$

On the other hand, I claim that the left hand limit does not impact continuity, since for all $x \in [0, 1]$,

$$\lim_{\varepsilon \rightarrow 0^-} f(x) - f(x - \varepsilon) = 0.$$

Indeed, we have

$$f(x) - f(x - \varepsilon) = \sum_{x - \varepsilon \leq r_n < x} 2^{-n}.$$

If $N > 0$, there is ε_0 such that for $\varepsilon < \varepsilon_0$, $[x - \varepsilon, x)$, $\{r_1, \dots, r_N\}$ is disjoint from $[x - \varepsilon, x)$, and thus for such ε ,

$$f(x) - f(x - \varepsilon) \leq 2^{-N}.$$

Taking $N \rightarrow \infty$ gives the limit, and completes the classification of discontinuity points.

Finally, we note the monotone convergence theorem implies that

$$\int_0^1 f(x) dx = \lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx.$$

Since f_n is a simple function, we easily evaluate

$$\int_0^1 f_n(x) \, dx = \sum_{k=1}^n (1 - r_k) \cdot 2^{-k}.$$

Thus

$$\int_0^1 f(x) \, dx = \sum_{k=1}^{\infty} (1 - r_k) \cdot 2^{-k}.$$

Day 12: Warm Up Question

49. (Spring 2021) Let f be a C^1 function on $[0, \infty)$. Suppose that

$$\int_0^\infty t|f'(t)|^2 dt < \infty$$

and

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(t) dt = L.$$

Show that $f(t) \rightarrow L$ as $t \rightarrow \infty$.

Solution: To relate f and f' , we apply the fundamental theorem of calculus. Suppose that

$$\int_R^\infty t|f'(t)|^2 dt \leq \varepsilon.$$

Then if $S = Re^{1/\varepsilon^{1/2}}$, then for $s \in [R, S]$,

$$|f(s) - f(R)| = \left| \int_R^s f'(t) dt \right| \leq \left(\int_R^s t|f'(t)|^2 dt \right)^{1/2} \left(\int_R^s 1/t dt \right)^{1/2} \leq \varepsilon^{1/2} \log(s/R)^{1/2} \leq \varepsilon^{1/4}.$$

If R is suitably large, then

$$\left| \frac{1}{R} \int_0^R f(t) dt \right| \leq 2L.$$

Thus

$$\left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_0^R f(t) dt \right| \leq 2Le^{-1/\varepsilon^{1/2}}.$$

Next,

$$\begin{aligned} \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_R^{Re^{1/\varepsilon^{1/2}}} f(t) dt &= \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_R^{Re^{1/\varepsilon^{1/2}}} f(R) + [f(t) - f(R)] dt \\ &= \left(1 - e^{-1/\varepsilon^{1/2}}\right) f(R) + \int_R^{Re^{1/\varepsilon^{1/2}}} [f(t) - f(R)] dt. \end{aligned}$$

Again, for any $\delta > 0$, if R is suitably large, then

$$\left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_0^{Re^{1/\varepsilon^{1/2}}} f(t) dt - L \right| \leq \delta$$

Combining all these inequalities yields that

$$\begin{aligned} |L - f(R)| &= \left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_0^{Re^{1/\varepsilon^{1/2}}} f(t) dt - f(R) \right| + \delta \\ &= \left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_R^{Re^{1/\varepsilon^{1/2}}} f(t) dt - f(R) \right| + \delta + 2Le^{-1/\varepsilon^{1/2}} \\ &= \left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_R^{Re^{1/\varepsilon^{1/2}}} [f(t) - f(R)] dt - e^{-1/\varepsilon^{1/2}} f(R) \right| + \delta + 2Le^{-1/\varepsilon^{1/2}} \\ &\leq e^{-1/\varepsilon^{1/2}} |f(R)| + \delta + 2Le^{-1/\varepsilon^{1/2}} + \varepsilon^{1/4}. \end{aligned}$$

The fact that

$$|f(R)| \leq 2 \left| \frac{1}{Re^{1/\varepsilon^{1/2}}} \int_R^{Re^{1/\varepsilon^{1/2}}} f(t) dt \right| + 2 \left| \int_R^{Re^{1/\varepsilon^{1/2}}} [f(t) - f(R)] dt \right| \leq 2L + 2\varepsilon^{1/4}$$

shows f is bounded, which is sufficient to complete the proof.

Solution: (Due to Anuk D.) Integration by parts with $u = f(t)$, $dv = 1$ gives

$$\frac{1}{T} \int_0^T f(t) dt = f(T) - \frac{1}{T} \int_0^T t f'(t) dt.$$

Therefore it suffices to show that $\frac{1}{T} \int_0^T t f'(t) dt$ converges as $T \rightarrow \infty$. Indeed, by the Cauchy Schwarz inequality,

$$\begin{aligned} \left| \frac{1}{T} \int_0^T t f'(t) dt \right| &= \left| \frac{1}{T} \int_0^{\sqrt{T}} \sqrt{t} \cdot \sqrt{t} f'(t) dt + \frac{1}{T} \int_{\sqrt{T}}^T \sqrt{t} \cdot \sqrt{t} f'(t) dt \right| \\ &\leq \frac{1}{T} \left(\int_0^{\sqrt{T}} t dt \right)^{\frac{1}{2}} \left(\int_0^{\sqrt{T}} t |f'(t)|^2 dt \right)^{\frac{1}{2}} + \frac{1}{T} \left(\int_{\sqrt{T}}^T t dt \right)^{\frac{1}{2}} \left(\int_{\sqrt{T}}^T t |f'(t)|^2 dt \right)^{\frac{1}{2}} \\ &\leq \frac{1}{2\sqrt{T}} \left(\int_0^{\infty} t |f'(t)|^2 dt \right)^{\frac{1}{2}} + \frac{1}{T} \left(\frac{T^2 - T}{2} \right)^{1/2} \left(\int_{\sqrt{T}}^{\infty} t |f'(t)|^2 dt \right)^{\frac{1}{2}} \\ &\leq \frac{C}{2\sqrt{T}} + \left(\int_{\sqrt{T}}^{\infty} t |f'(t)|^2 dt \right)^{\frac{1}{2}} \rightarrow 0 \text{ as } T \rightarrow \infty \end{aligned}$$

50. (Fall 2013) Let $E = \{(x_1, x_2) : x_1, x_2 \in \mathbb{R}, x_1 - x_2 \in \mathbb{Q}\}$. Is it possible to find Lebesgue measurable sets $A_1, A_2 \subset \mathbb{R}$ such that $|A_1|, |A_2| > 0$, and $A_1 \times A_2 \subset E^c$?

Solution: No, it is not possible. We will show that $|A_1| \cdot |A_2| = 0$ whenever $A_1 \times A_2 \subset E^c$.

Case 1: Suppose $A_1, A_2 \subset [-n, n]$ for some positive integer n . Let $h(y) = \int_{\mathbb{R}} \chi_{A_1}(x) \chi_{A_2}(x+y) dx$. By Tonelli's Theorem,

$$\begin{aligned} \int_{\mathbb{R}} h(y) dy &= \int_{\mathbb{R}} \chi_{A_1}(x) \int_{\mathbb{R}} \chi_{A_2}(y+x) dy dx \\ &= \int_{\mathbb{R}} \chi_{A_1}(x) \int_{\mathbb{R}} \chi_{A_2}(y) dy dx \\ &= \left(\int_{\mathbb{R}} \chi_{A_1}(x) dx \right) \left(\int_{\mathbb{R}} \chi_{A_2}(y) dy \right) \\ &= |A_1| \cdot |A_2|. \end{aligned}$$

Next we claim that h is also continuous. This is by average continuity of the L^1 norm. For a detailed proof, let $\epsilon > 0$. Since $\chi_{A_2} \in L^1(\mathbb{R})$, there exists $\phi \in C_c(\mathbb{R})$ with compact support $K \subset \mathbb{R}$ such that

$\|\chi_{A_2} - \phi\|_{L^1} < \epsilon/3$. Since ϕ is continuous on a compact set, it is uniformly continuous, so we may choose $\delta > 0$ such that $|\phi(s) - \phi(t)| \leq \epsilon/3|K|$ whenever $|s - t| \leq \delta$. Then

$$\begin{aligned}
 |h(y + \delta) - h(y)| &= \left| \int_{A_1} \chi_{A_2}(x + y + \delta) dx - \int_{A_1} \chi_{A_2}(x + y) dx \right| \\
 &\leq \int_{A_1} |\chi_{A_2}(x + y + \delta) - \phi(x + y + \delta)| dx + \int_{A_1} |\phi(x + y + \delta) - \phi(x + y)| dx \\
 &\quad + \int_{A_1} |\phi(x + y) - \chi_{A_2}(x + y)| dx \\
 &\leq 2 \int_{\mathbb{R}} |\chi_{A_2} - \phi| dx + \int_K |\phi(x + \delta) - \phi(x)| dx \\
 &\leq 2\epsilon/3 + \epsilon/3 \\
 &= \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ was arbitrary, h is continuous.

Finally we claim that $h(y) = 0$ for all $y \in \mathbb{Q}$. Indeed, if $y \in \mathbb{Q}$ then for any x , we have $x - (y + x) \in \mathbb{Q}$ and hence $(x, y + x) \in E$. Therefore since $A_1 \times A_2 \subset E^c$, $(x, y + x) \notin A_1 \times A_2$. Therefore

$$\chi_{A_1}(x)\chi_{A_2}(y + x) = 0.$$

Since this holds for all $x \in \mathbb{R}$, it follows that $h(y) = 0$. We have shown that

1. $\int_{\mathbb{R}} h = |A_1| \cdot |A_2|$
2. h is continuous
3. $h(y) = 0$ for all $y \in \mathbb{Q}$.

By density of \mathbb{Q} in \mathbb{R} , (2) and (3) imply that $h \equiv 0$, so that by (1), $|A_1| \cdot |A_2| = 0$.

Case 2: Let $A_1, A_2 \subset \mathbb{R}$ be arbitrary measurable sets. By case 1, $|A_1 \cap [-n, n]| \cdot |A_2 \cap [-n, n]| = 0$ for all n . Therefore by continuity of Lebesgue measure,

$$|A_1| \cdot |A_2| = \lim_{n \rightarrow \infty} |A_1 \cap [-n, n]| \cdot |A_2 \cap [-n, n]| = \lim_{n \rightarrow \infty} 0 = 0.$$

51. (Spring 2021)

- (a) Let H_1 and H_2 be Hilbert spaces, and let $T : H_1 \rightarrow H_2$ be a continuous linear operator. Give a precise definition of the adjoint operator T^* .

Solution: The adjoint operator T^* is a bounded operator from H_2 to H_1 . For each $x \in H_2$, T^*x is the unique element of H_1 such that for any $y \in H_1$, $\langle T^*x, y \rangle = \langle x, Ty \rangle$.

- (b) Let $(a, b) \subset \mathbb{R}$ be a (possibly infinite) open interval. If $f \in L^2(a, b)$, explain what it means that the distributional derivative f' is also in $L^2(a, b)$.

Solution: This means that there exists $g \in L^2(a, b)$, such that for any $\phi \in C_c^\infty(a, b)$,

$$\int f(X)\phi'(x) dx = - \int g(x)\phi(x) dx.$$

- (c) Let \mathbb{R}_+ denote the positive real axis $[0, \infty)$. Let $H^1(\mathbb{R})$ (respectively $H^1(\mathbb{R}_+)$) be the space of real-valued functions $f \in L^2(\mathbb{R})$ (respectively $f \in L^2(\mathbb{R}_+)$) such that the distributional derivative f' is also in $L^2(\mathbb{R})$ (respectively $L^2(\mathbb{R}_+)$). Then $H^1(\mathbb{R})$ and $H^1(\mathbb{R}_+)$ are Hilbert spaces with inner product given by

$$\begin{aligned}\langle f, g \rangle_{H^1(\mathbb{R})} &= \int_{\mathbb{R}} f(x)g(x)dx + \int_{\mathbb{R}} f'(x)g'(x)dx, \\ \langle f, g \rangle_{H^1(\mathbb{R}_+)} &= \int_{\mathbb{R}_+} f(x)g(x)dx + \int_{\mathbb{R}_+} f'(x)g'(x)dx\end{aligned}$$

Let $T : H^1(\mathbb{R}) \rightarrow H^1(\mathbb{R}_+)$ be the mapping given by the restriction. Compute exactly the adjoint operator T^* .

Solution:

18 Day 12: Bonus Questions

52. (Fall 2015) Let (X, μ) be a measure space, and let $f : X \rightarrow \mathbb{R}$ be measurable. Then if $1 \leq p < r < q < \infty$ and there is $C < \infty$ such that

$$\mu(\{x : |f(x)| > \lambda\}) \leq \frac{C}{\lambda^p + \lambda^q}$$

for every $\lambda > 0$. Then $f \in L^r(\mu)$.

Solution: We perform a dyadic decomposition, writing

$$\begin{aligned} \int |f(x)|^r dx &= \sum_{k=-\infty}^{\infty} \int_{2^k < |f(x)| \leq 2^{k+1}} |f(x)|^r dx \\ &\leq \sum_{k=-\infty}^{\infty} \mu(\{2^k < |f(x)|\}) \cdot 2^{kr}. \end{aligned}$$

For $\lambda < 1$, $\lambda^p \leq \lambda^q$, so

$$\mu(\{2^k < |f(x)|\}) \leq C/2^{kp}.$$

For $\lambda \geq 1$, $\lambda^q \leq \lambda^p$, so

$$\mu(\{2^k < |f(x)|\}) \leq C/2^{kq}.$$

Thus we conclude

$$\int |f(x)|^r dx \leq C \left(\sum_{k=-\infty}^0 2^{k(r-p)} + \sum_{k=1}^{\infty} 2^{k(r-q)} \right).$$

Both of these sums converge, so

$$\int |f(x)|^r dx \lesssim_{p,q} C < \infty.$$

Solution: *Case 1:* Suppose that (X, \mathcal{M}, μ) is σ -finite. This assumption of σ -finiteness allows us to apply the Fubini-Tonelli Theorem in the following calculation:

$$\begin{aligned} \int_X |f(x)|^r dx &= \int_X \int_0^{|f(x)|} r t^{r-1} dt dx \\ &= \int_X \int_0^{\infty} r t^{r-1} \chi_{[|f(x)| > t]} dt dx \\ &= \int_0^{\infty} r t^{r-1} \int_X \chi_{[|f(x)| > t]} dx dt && \text{by Tonelli's Theorem} \\ &= \int_0^{\infty} r t^{r-1} \mu\{x : |f(x)| > t\} dt \\ &\leq Cr \int_0^1 \frac{t^{r-1}}{t^p + t^q} dt + Cr \int_1^{\infty} \frac{t^{r-1}}{t^p + t^q} dt \\ &\leq Cr \int_0^1 t^{r-1-p} dt + Cr \int_1^{\infty} t^{r-1-q} dt \end{aligned}$$

and the right-hand side is finite since $1 < p < r < q < \infty$.

Case 2: Suppose (X, \mathcal{M}, μ) is not σ -finite. Let $X' = \{x \in X : |f(x)| > 0\}$. Also define $\mathcal{M}' = \{M \cap X' : M \in \mathcal{M}\}$ (it is easy to see this is a σ -algebra) and define μ' as the restriction of μ to \mathcal{M}' . Then (X', \mathcal{M}', μ') is σ -finite because

$$X' = \bigcup_{n=1}^{\infty} \left\{ x \in X' : |f(x)| > \frac{1}{n} \right\}$$

and

$$\mu \left\{ x \in X' : |f(x)| > \frac{1}{n} \right\} \leq \frac{C}{n^{-p} + n^{-q}} < \infty$$

for all n .

Therefore by Case 1, $\int_{X'} |f(x)|^r dx < \infty$. And since $\int_X |f(x)|^r dx = \int_{X'} |f(x)|^r dx$, we are done.

53. (Spring 2020) Let $\sum_{n=1}^{\infty} a_n$ be a convergent series. Let $b_n \in \mathbb{R}$ be an increasing sequence with $\lim_{n \rightarrow \infty} b_n = \infty$. Show that

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{k=1}^n b_k a_k = 0.$$

Solution: Let $k_n = \max \left\{ k \geq 1 : \left| \sum_{i=1}^k a_i b_i \right| < \sqrt{b_n} \text{ and } k \leq n \right\}$. Since $\sqrt{b_n} \rightarrow \infty$ monotonically as $n \rightarrow \infty$, k_n is defined for sufficiently large n , is increasing, and $k_n \rightarrow \infty$. For n sufficiently large, we may write

$$\frac{1}{b_n} \sum_{k=1}^n b_k a_k = \frac{1}{b_n} \sum_{k=1}^{k_n-1} b_k a_k + \frac{1}{b_n} \sum_{k=k_n}^n b_k a_k$$

By choice of k_n , $\left| \frac{1}{b_n} \sum_{k=1}^{k_n-1} b_k a_k \right| \leq \frac{1}{\sqrt{b_n}} \rightarrow 0$ as $n \rightarrow \infty$, so it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{k=k_n}^n b_k a_k = 0.$$

Let $T_k = \sum_{i=1}^k a_i$ and let $a = \sum_{n=1}^{\infty} a_n$. Using the summation by parts formula,

$$\begin{aligned} \frac{1}{b_n} \sum_{k=k_n}^n b_k a_k &= \frac{1}{b_n} \left[T_n b_n - T_{k_n-1} b_{k_n} + \sum_{k=k_n}^{n-1} T_k (b_k - b_{k+1}) \right] \\ &= \frac{1}{b_n} \left[(T_n - a) b_n + a b_n - (T_{k_n-1} - a) b_{k_n} - a b_{k_n} \right. \\ &\quad \left. + \sum_{k=k_n}^{n-1} (T_k - a) (b_k - b_{k+1}) + a \sum_{k=k_n}^{n-1} (b_k - b_{k+1}) \right] \\ &= \frac{1}{b_n} \left[(T_n - a) b_n - (T_{k_n-1} - a) b_{k_n} + \sum_{k=k_n}^{n-1} (T_k - a) (b_k - b_{k+1}) \right] \end{aligned}$$

where for the last equality we used the fact that $\sum_{k=k_n}^{n-1} (b_k - b_{k+1}) = b_{k_n} - b_n$. Let $M_n =$

$\max_{k_n \leq k < n} |T_k - a|$. By the triangle inequality, and using $b_{k_n} \leq b_n$,

$$\begin{aligned} \left| \frac{1}{b_n} \sum_{k=k_n}^n b_k a_k \right| &\leq 2 |T_n - a| + \frac{1}{b_n} \left| \sum_{k=k_n}^{n-1} (T_k - a)(b_k - b_{k+1}) \right| \\ &\leq 2 |T_n - a| + \frac{M_n}{b_n} \sum_{k=k_n}^{n-1} |b_k - b_{k+1}| \\ &= 2 |T_n - a| + \frac{M_n}{b_n} \sum_{k=k_n}^{n-1} b_{k+1} - b_k \\ &= 2 |T_n - a| + \frac{M_n}{b_n} (b_n - b_{k_n}) \\ &\leq 2 |T_n - a| + 2M_n \end{aligned}$$

The result then follows since $T_n \rightarrow a$ and $M_n \rightarrow 0$ as $n \rightarrow \infty$.

54. (Spring 2021) Let $f_n \rightarrow f$ weakly in $L^2(\mathbb{R})$ and $\|f_n\|_2 \rightarrow \|f\|_2$ as $n \rightarrow \infty$. Show that $f_n \rightarrow f$ strongly in $L^2(\mathbb{R})$.

Solution: Since $f_n \rightarrow f$ weakly, therefore $\langle f_n, g \rangle \rightarrow \langle f, g \rangle$ for any $g \in L^2$. Since $\|f\|_2 \leq \sup_n \|f_n\|_2 < \infty$, we have $f \in L^2$. Therefore taking $g = f$, we have $\langle f_n, f \rangle \rightarrow \|f\|_2^2$, so that

$$\|f_n - f\|_2^2 = \langle f_n - f, f_n - f \rangle = \|f_n\|_2^2 - 2\langle f_n, f \rangle + \|f\|_2^2 \rightarrow 0$$

as $n \rightarrow \infty$.

55. (Spring 2017) Let $E \subset \mathbb{R}^n$ be a set of finite, positive measure, and let $\{t_k\}$ be a sequence with $\{t_k\} > 0$ and $\lim_k t_k = 0$. Define, for $f \in L^p(\mathbb{R}^n)$,

$$Mf(x) = \sup_k \int_{t_k E} |f(x - y)| dy.$$

Suppose furthermore that there is $C > 0$ such that

$$|\{x : Mf(x) > \lambda\}| \leq C\lambda^{-p} \|f\|_p^p.$$

Show that for every $f \in L^p(\mathbb{R}^n)$,

$$\lim_k \int_{t_k E} f(x - y) dy = f(x).$$

for almost every $x \in \mathbb{R}^d$.

Solution: We note that the result is true for any $f \in C(\mathbb{R}^n)$, i.e. for such functions, and any $x \in \mathbb{R}$,

$$\limsup_k \left| \int_{t_k E} f(x) - f(x - y) \right| = 0.$$

Now given any $f \in L^p(\mathbb{R}^n)$, for $1 \leq p < \infty$, fix $\varepsilon > 0$, and find $g \in C(\mathbb{R}^n)$ with $\|f - g\|_p \leq \varepsilon$. Then

$$|\{M(f - g) > \lambda\}| \leq C\varepsilon^p \lambda^{-p}.$$

Thus

$$\begin{aligned} & \left\{ \limsup_k \left| \int_{t_k E} f(x) - f(x-y) dy \right| \geq \delta \right\} \\ & \subset \left\{ \limsup_k \left| \int_{t_k E} f(x) - g(x) + g(x-y) - f(x-y) dy \right| \geq \delta \right\} \\ & \subset \{x : |f(x) - g(x)| \geq \delta/2\} \cup \left\{ \limsup_k \left| \int_{t_k E} g(x-y) - f(x-y) dy \right| \geq \delta/2 \right\}. \end{aligned}$$

Now Markov's inequality implies that

$$|\{x : |f(x) - g(x)| \geq \delta/2\}| \leq \|f - g\|_p (\delta/2)^{-p} \lesssim_p \varepsilon \delta^{-p}.$$

and

$$\left| \left\{ x : \limsup_k \left| \int_{t_k E} g(x-y) - f(x-y) dy \right| \geq \delta/2 \right\} \right| \leq |\{x : |M(g-f)(x)| \geq \delta/2\}| \lesssim \varepsilon^p \delta^{-p}.$$

Thus we conclude that

$$\left| \left\{ \limsup_k \left| \int_{t_k E} f(x) - f(x-y) dy \right| \geq \delta \right\} \right| \lesssim (\varepsilon + \varepsilon^p) \delta^{-p}.$$

Taking $\varepsilon \rightarrow 0$ shows that for each $\delta > 0$,

$$\left| \left\{ \limsup_k \left| \int_{t_k E} f(x) - f(x-y) dy \right| \geq \delta \right\} \right| = 0.$$

Taking a union bound over $\delta = 1, 1/2, 1/4, \dots$ completes the proof.

56. (Spring 2020) Let $f_n : [0, 1] \rightarrow \mathbb{R}$ be a sequence of Lebesgue measurable functions such that f_n converges to f almost everywhere on $[0, 1]$ and such that $\|f_n\|_{L^2([0,1])} \leq 1$ for all n . Show that

$$\lim_{n \rightarrow \infty} \|f_n - f\|_{L^1([0,1])} = 0.$$

Solution:

[This proof may border on trivializing the problem by invoking the Vitali convergence theorem, but I seem to have gotten away with a proof like this on my qual, so... -m]. Since $|E| < \infty$, it follows trivially that $\{f_n\}$ is tight. Since $f_n \rightarrow f$ a.e., the result will then follow by the Vitali Convergence Theorem provided that the sequence $\{f_n\}$ is uniformly integrable. We wish to show that $\{f_n\}$ is uniformly integrable. Let $\epsilon > 0$ and let $A \subset [0, 1]$ be measurable. We need to show there exists a $\delta > 0$ such that $\int_A |f_n| < \epsilon$ for all n whenever $|A| < \delta$. By Cauchy-Schwarz inequality and the assumption that $\|f_n\|_{L^2} \leq 1$ for all n ,

$$\int_A |f_n| = \int_{[0,1]} \chi_A |f_n| \leq |A|^{1/2} \|f_n\|_{L^2} \leq |A|^{1/2}.$$

It follows that $\int_A |f_n| < \epsilon$ for all n whenever $|A| < (\epsilon/2)^2$. Therefore $\{f_n\}$ is uniformly integrable. This completes the proof.

Another (similar) proof is given:

Proof. We first prove the following claim:

Claim 1: $f_n, f \in L^1([0, 1])$ and $\|f_n\|_{L^1}, \|f\|_{L^1} \leq 1$.

Proof of Claim 1: Since $x \mapsto x^2$ is convex and $[0, 1]$ has measure 1, it follows by Jensen's inequality that

$$\left(\int_0^1 |f_n(x)| dx \right)^2 \leq \int_0^1 |f_n(x)|^2 dx \leq 1.$$

Therefore $\|f_n\|_{L^1} \leq 1$ and hence $f_n \in L^1([0, 1])$ for all n . Since $f_n \rightarrow f$ a.e., it follows by Fatou's lemma that

$$\int_0^1 |f(x)| dx \leq \liminf_{n \rightarrow \infty} \int_0^1 |f_n(x)| dx \leq 1.$$

Therefore $f \in L^1([0, 1])$ with $\|f\|_{L^1} \leq 1$ as well. \square

By Egorov's Theorem, since $[0, 1]$ has finite measure, there exists $E \subset [0, 1]$ with $|E| < (\epsilon/4)^2$ such that $f_n \rightarrow f$ uniformly on $[0, 1] \setminus E$. By uniform convergence, we may choose a positive integer n_0 such that $|f_n(x) - f(x)| < \epsilon/2$ for all $x \in [0, 1] \setminus E$ and all $n \geq n_0$. Then for $n \geq n_0$,

$$\begin{aligned} \int_0^1 |f_n - f| &= \int_E |f_n - f| + \int_{[0,1] \setminus E} |f_n - f| \\ &< \int_E |f_n(x) - f(x)| dx + \epsilon/2 \\ &= \int_{[0,1]} \chi_E |f_n(x)| dx + \int_{[0,1]} \chi_E |f(x)| dx + \epsilon/2 \\ &\leq |E|^{1/2} \|f_n\|_{L^2} + |E|^{1/2} \|f\|_{L^2} + \epsilon/2 && \text{by Cauchy Schwarz} \\ &\leq 2|E|^{1/2} + \epsilon/2 && \text{by Claim 1} \\ &< \epsilon/2 + \epsilon/2 \\ &= \epsilon. \end{aligned}$$

It follows that $\lim_{n \rightarrow \infty} \|f_n - f\|_{L^1([0,1])} = 0$. \square

57. (Fall 2017, Spring 2021) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a compactly supported function that satisfies the Hölder condition with exponent $\beta \in (0, 1)$, i.e. that there exists a constant $A < \infty$ such that for all $x, y \in \mathbb{R}$, $|f(x) - f(y)| \leq A|x - y|^\beta$. Consider the function g defined by

$$g(x) = \int_{-\infty}^{\infty} \frac{f(y)}{|x - y|^\alpha} dy,$$

where $\alpha \in (0, \beta)$.

- (a) Prove that g is a continuous function at zero.

Solution: One way to see this immediately is to apply Young's inequality for convolution. More directly, we can write

$$g(x) - g(0) = \int_{-\infty}^{\infty} f(y) \left(\frac{1}{|x - y|^\alpha} - \frac{1}{|y|^\alpha} \right) dy$$

For $|y| \leq |x|/2$, we have

$$\left(\frac{1}{|x - y|^\alpha} - \frac{1}{|y|^\alpha} \right) \lesssim 1/|x|^\alpha.$$

Thus

$$\left| \int_{|y| \leq |x|/2} f(y) \left(\frac{1}{|x-y|^\alpha} - \frac{1}{|y|^\alpha} \right) dy \right| \lesssim \|f\|_\infty |x|^{1-\alpha}$$

For $|y| \geq 2|x|$, we can apply Taylor's theorem / the mean value theorem to conclude that

$$\left| \frac{1}{|x-y|^\alpha} - \frac{1}{|y|^\alpha} \right| \lesssim \frac{|x|}{|y|^{\alpha+1}}.$$

This bound implies that

$$\left| \int_{|y| \geq 2|x|} f(y) \left(\frac{1}{|x-y|^\alpha} - \frac{1}{|y|^\alpha} \right) dy \right| \lesssim \|f\|_\infty |x|^{1-\alpha}.$$

For $|x|/2 \leq |y| \leq 2|x|$, we have

$$\left| \int_{|x|/2 \leq |y| \leq 2|x|} f(y) \left(\frac{1}{|x-y|^\alpha} - \frac{1}{|y|^\alpha} \right) dy \right| \leq 2\|f\|_\infty \int_{|y| \leq 3|x|} \frac{1}{|y|^\alpha} \lesssim \|f\|_\infty |x|^{1-\alpha}.$$

These bounds together imply that

$$|g(x) - g(0)| \lesssim \|f\|_\infty |x|^{1-\alpha}.$$

Thus $g(x) \rightarrow g(0)$ as $x \rightarrow 0$.

(b) Prove that g is differentiable at zero. (Hint: Try the dominated convergence theorem).

Solution: Assume first that $f(0) = 0$. We have

$$\frac{g(x) - g(0)}{x} = \int_{-\infty}^{\infty} f(y) \frac{1}{x} \left(\frac{1}{|y-x|^\alpha} - \frac{1}{|y|^\alpha} \right) dy.$$

The integrand here converges pointwise as $x \rightarrow 0$ to

$$f(y)\alpha/|y|^{\alpha+1},$$

which is integrable since the function is compactly supported and continuous away from the origin, and $|f(y)| \lesssim |y|^\beta$ near the origin. The mean value theorem implies that if $x \leq 1$,

$$\left| \frac{1}{x} \left(\frac{1}{|y-x|^\alpha} - \frac{1}{|y|^\alpha} \right) \right| \lesssim 1/|y|^{\alpha+1}$$

so we might expect that

$$g'(0) = \int_{-\infty}^{\infty} f(y)\alpha/|y|^{\alpha+1},$$

at least if the dominated convergence theorem

58. (Fall 2018) Let $1 < p \leq \infty$. Let (X, \mathcal{M}, μ) be a finite measure space. Let $\{f_n\}$ be a sequence of measurable functions converging μ -a.e. to the function f . Assume further that $\|f_n\|_p \leq 1$ for all n . Prove that $f_n \rightarrow f$ as $n \rightarrow \infty$ in L^r for all $1 \leq r < p$.

Solution: *Case 1.* If $p = \infty$ then since $\|f\|_\infty \leq 1$ and $f_n \rightarrow f$ a.e., it follows that $\|f\|_\infty \leq 1$. Therefore by the triangle inequality and using the fact that $r \geq 1$, we have $|f_n - f|^r \leq (|f_n| + |f|)^r \leq 2^r$. Since the constant function 2^r is integrable on X with respect to μ , it follows by the dominated convergence theorem that $\lim_{n \rightarrow \infty} \int_X |f_n - f|^r d\mu = 0$. Therefore $f_n \rightarrow f$ in L^r .

Case 2. Suppose $1 < p < \infty$. Let $\epsilon > 0$, and let $\epsilon_0 = 2^{-r} \epsilon^{\frac{p}{p-r}}$ (here we use the assumption that $p > r$ so that $\epsilon_0 > 0$ is defined). By Egorov's theorem there exists $A \subset X$ such that $\mu(A^c) < \epsilon_0$ and $f_n \rightarrow f$ uniformly on A . Then

$$\int_X |f_n - f|^r = \int_{A^c} |f_n - f|^r d\mu + \int_A |f_n - f|^r d\mu$$

Since $r \geq 1$, it follows that $|f_n - f|^r \rightarrow 0$ uniformly. Therefore $\lim_{n \rightarrow \infty} \int_A |f_n - f|^r d\mu = 0$. Therefore

$$\limsup_{n \rightarrow \infty} \int_X |f_n - f|^r \leq \limsup_{n \rightarrow \infty} \int_{A^c} |f_n - f|^r d\mu \quad (15)$$

By Holder's inequality,

$$\begin{aligned} \int_{A^c} |f_n - f|^r d\mu &= \int_X |f_n - f|^r \chi_{A^c} d\mu \\ &\leq \left(\int_X |f_n - f|^p d\mu \right)^{r/p} \left(\int_X \chi_{A^c} d\mu \right)^{\frac{p-r}{p}} \\ &= \|f_n - f\|_p^r \mu(A^c)^{\frac{p-r}{r}} \\ &= \epsilon \|f_n - f\|_p^r \\ &\leq \epsilon (\|f_n\|_p + \|f\|_p)^r \\ &\leq \epsilon \left(1 + \|f\|_p \right)^r \end{aligned}$$

Also, by Fatou's lemma, $\|f\|_p \leq \liminf_{n \rightarrow \infty} \|f_n\|_p \leq 1$. Therefore

$$\int_{A^c} |f_n - f|^r d\mu \leq \epsilon 2^r$$

Therefore by (15),

$$\limsup_{n \rightarrow \infty} \int_X |f_n - f|^r d\mu \leq \epsilon 2^r$$

Since $\epsilon > 0$ was arbitrary, it follows that $\limsup_{n \rightarrow \infty} \int_X |f_n - f|^r d\mu = 0$. Therefore $f_n \rightarrow f$ in L^r .

19 Day 14: Analysis Qualifying Exam 2022

59. Let x_1, \dots, x_{n+1} be pairwise distinct real numbers. Prove that there exists $C > 0$ such that if $P : \mathbb{R} \rightarrow \mathbb{R}$ is a polynomial with degree at most n , then

$$\|P\|_{L^\infty[0,1]} \leq C \max(|P(x_1)|, \dots, |P(x_{n+1})|).$$

Solution: Let V_n be the vector space of degree $\leq n$ polynomials. The theory of polynomial interpolation shows that the operator $T : V_n \rightarrow \mathbb{R}^n$ given by

$$TP = (p(x_1), \dots, p(x_{n+1}))$$

is a bijection. Thus $T^{-1} : \mathbb{R}^n \rightarrow V_n$ is well defined. As a linear map between two finite dimensional vector spaces, T^{-1} is continuous from any norm on \mathbb{R}^n , to any norm on V_n . If we equip \mathbb{R}^n with the l^∞ norm, and V_n with the $L^\infty[0,1]$ norm, then the result immediately follows.

60. Given a real number x , let $\{x\}$ denote the fractional part of x . Suppose α is an irrational number and define $T : [0, 1] \rightarrow [0, 1]$ by

$$T(x) = \{x + \alpha\}.$$

Prove: If $A \subset [0, 1]$ is measurable and $T(A) = A$ then $|A| \in \{0, 1\}$.

Solution: Define $f : \mathbb{R} \rightarrow \{0, 1\}$ by setting $f(x) = \mathbf{I}(x \in A)$. Since $T(A) = A$, it follows that $f(x + \alpha) = f(x)$, i.e. f is both $[0, 1]$ periodic and α periodic. But applying the Fourier transform to both sides of this formula (viewing f as a tempered distribution) yields that

$$\widehat{f}(\xi)e^{2\pi i\xi\alpha} = \widehat{f}(\xi),$$

i.e. that \widehat{f} is supported on integer multiples of $2\pi\alpha$. But the Poisson summation formula says precisely that \widehat{f} is supported on the integer multiples of 2π , and the intersection of these two supports lies only at the origin. Thus \widehat{f} is supported at the origin. But the only locally integrable tempered function with this property is a constant function, so either $f(x) = 1$ almost everywhere, or $f(x) = 0$ almost everywhere, which yields the result.

61. Let $\{f_n\}$ be a sequence of measurable, real-valued functions on a measure space X such that $f_n \rightarrow f$ pointwise as $n \rightarrow \infty$, where $f : X \rightarrow \mathbb{R}$, and suppose that for some constant $M > 0$,

$$\int |f_n| d\mu \leq M \text{ for all } n \in \mathbb{N}.$$

(a) Prove that

$$\int |f| d\mu \leq M.$$

Solution: The result follows immediately from Fatou's lemma, i.e. that

$$\int |f| d\mu = \int \liminf_n |f_n| d\mu \leq \liminf_n \int |f_n| d\mu \leq M.$$

- (b) Give an example to show that we may have $\int |f_n| d\mu = M$ for every n , but $\int |f| d\mu < M$.

Solution: Let $X = \mathbb{R}$, and let $f_n(x) = M\mathbf{I}(x \in [n, n+1])$. Then $f = 0$.

- (c) Prove that

$$\lim_{n \rightarrow \infty} \int ||f_n| - |f| - |f_n - f|| = 0.$$

Solution: The integrand converges pointwise to zero. Moreover, we have

$$||f_n| - |f| - |f_n - f|| = |f_n - f| + |f| - |f_n|$$

by virtue of the fact that

$$|f_n - f| \geq |f_n| - |f|.$$

Now the triangle inequality gives that

$$|f_n - f| + |f| - |f_n| \leq |f_n| + |f| + |f| - |f_n| \leq 2|f|.$$

Thus the integrands are uniformly dominated by an *integrable function*. The dominated convergence thus immediately gives the result.

62. Consider the following equation for an unknown function $f : [0, 1] \rightarrow \mathbb{R}$:

$$f(x) = g(x) + \lambda \int_0^1 (x-y)^2 f(y) dy + \frac{1}{2} \sin(f(x)).$$

Prove that there exists a number $\lambda_0 > 0$ such that for all $\lambda \in [0, \lambda_0)$ and all continuous functions g on $[0, 1]$, the equation has a unique continuous solution.

Solution: Define an integral operator T_g by setting

$$T_g f(x) = g(x) + \frac{\sin(f(x))}{2} + \lambda \int_0^1 (x-y)^2 f(y) dy.$$

The problem is to find, and prove the uniqueness, of a fixed point for T_g . We calculate that

$$\begin{aligned} \|T_g f_1 - T_g f_2\|_{L^\infty[0,1]} &\leq \left\| \frac{\sin \circ f_1 - \sin \circ f_2}{2} \right\|_{L^\infty[0,1]} + \lambda \|f_1 - f_2\|_{L^\infty[0,1]} \sup_{x \in [0,1]} \int_0^1 (x-y)^2 dy \\ &\leq (1/2) \|f_1 - f_2\|_{L^\infty[0,1]} + \lambda \|f_1 - f_2\|_{L^\infty[0,1]} \left(\sup_{x \in [0,1]} x^2 - x + 1/3 \right) \\ &\leq (1/2 + \lambda/3) \|f_1 - f_2\|_{L^\infty[0,1]}. \end{aligned}$$

If $\lambda < 1/6$, we conclude that T_g is a contraction map, and thus has a unique fixed point.

63. Given $\alpha \geq 0$, the α -dimensional Hausdorff measure of a set $X \subset \mathbb{R}^n$ is

$$H^\alpha(X) = \liminf_{r \rightarrow 0} \left\{ \sum_{i=1}^{\infty} r_i^\alpha : X \subset \bigcup_{i=1}^{\infty} B(x_i, r_i), r_i < r \text{ for all } i \right\}$$

(where $B(x, r)$ is the Euclidean ball with center x and radius r) and the Hausdorff dimension is

$$\dim_{\mathbf{H}}(X) = \inf\{\alpha \geq 0 : H^\alpha(X) = 0\}$$

Prove:

- (a) If $X \subset \mathbb{R}^n$ and μ is a finite Borel measure on X such that $\mu(X) > 0$ and $\mu(B(x, r)) \leq r^\alpha$ for all open balls $B(x, r)$, then $\dim_{\mathbf{H}}(X) \geq \alpha$.

Solution: Suppose X is covered by a family of balls $\{B(x_i, r_i)\}$. Then

$$1 \lesssim \mu(X) = \mu\left(\bigcup B(x_i, r_i)\right) \leq \sum_i \mu(B(x_i, r_i)) \leq \sum_i r_i^\alpha.$$

Thus

$$H^\alpha(X) = \liminf \sum_i r_i^\alpha \gtrsim 1,$$

which implies that $\dim_{\mathbf{H}}(X) \geq \alpha$.

- (b) If $\mathbf{S}^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$, then $\dim_{\mathbf{H}}(\mathbf{S}^1) = 1$.

Solution: For any r , we can cover \mathbf{S}^1 by $O(1/r)$ balls of radius r . Thus for $\alpha > 1$,

$$H^\alpha(X) = \liminf_{r \rightarrow 0} O(1/r)r^\alpha = \liminf_{r \rightarrow 0} r^{\alpha-1} = 0.$$

Thus $\dim_{\mathbf{H}}(\mathbf{S}^1) \leq 1$. We use the first part to prove that $\dim_{\mathbf{H}}(\mathbf{S}^1) \geq 1$, which will complete the proof. Namely, we let μ denote the angle measure on \mathbf{S}^1 divided by 1000. Then it is easy to prove that $\mu(B(x, r)) \lesssim r$ for all balls B - since the angle measure has mass $2\pi/1000$ we may assume $r \leq 1/10$, and then the intersection of $B(x, r)$ with \mathbf{S}^1 is an arc with angular length $O(r)$. Thus the above part of the problem gives the lower bound.

64. Let $X = [0, 1]$ with Lebesgue measure and $Y = [0, 1]$ with counting measure. Give an example of an integrable function $f : X \times Y \rightarrow [0, \infty)$ for which Fubini's theorem does not apply.

Solution: Let $f(x, y) = 1$ if $x = y$, and $f(x, y) = 0$ otherwise. Then

$$\int_Y \int_X f(x, y) dx dy = \int_Y 0 dy = 0,$$

whereas

$$\int_X \int_Y f(x, y) dy dx = \int_X 1 dx = 1.$$

65. For $s > 1/2$ let $H^s(\mathbb{R}^n)$ denote the Sobolev space

$$H^s(\mathbb{R}^n) = \{f \in L^2(\mathbb{R}^n) : \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{f}(\xi)|^2 d\xi < \infty\}.$$

Use the Fourier transform to prove that if $u \in H^s(\mathbb{R}^n)$ for $s > n/2$ then $u \in L^\infty(\mathbb{R}^n)$ with the boundary

$$\|u\|_{L^\infty(\mathbb{R}^n)} \leq C \|u\|_{H^s(\mathbb{R}^n)}.$$

for a constant C depending only on s and n .

Solution: The Fourier inversion formula tells us that

$$\|u\|_{L^\infty(\mathbb{R}^n)} \leq \|\hat{u}\|_{L^1(\mathbb{R}^n)}.$$

But for $s > n/2$, we have

$$\|\hat{u}\|_{L^1(\mathbb{R}^n)} \lesssim_s \|u\|_{H^s(\mathbb{R}^n)},$$

by virtue of Cauchy Schwartz, i.e. that

$$\int |\hat{u}(\xi)| d\xi = \int \left[|\hat{u}(\xi)|(1 + |\xi|^2)^{s/2} \right] \left[(1 + |\xi|^2)^{-s/2} \right] \leq \left(\int |\hat{u}(\xi)|^2 (1 + |\xi|^2)^s \right)^{1/2} \left(\int (1 + |\xi|^2)^{-s} \right)^{1/2}.$$

The first integral is precisely the H^s norm, and for large ξ , the right integrand is bounded, and essentially equal to $|\xi|^{-2s}$ for $|\xi| \geq 1$, which proves the second integral is finite, completing the proof.

66. Assume that X is a compact metric space and $T : X \rightarrow X$ is a continuous map. Let $M_1(T)$ denote the set of Borel probability measures on X such that $T_*\mu = \mu$. Prove

(a) $M_1(T) \neq \emptyset$.

Solution: Fix any probability measure ν . Then $T_*^k \nu$ is a probability measure for all $k \geq 0$. Define a probability measure μ_n such that

$$\mu_n = \frac{1}{n} \sum_{k=0}^{n-1} T_*^k \nu$$

Applying weak * compactness, there exists a subsequence of these measures that converges in the weak * topology to some probability measure μ , i.e. $\mu = \lim_m \mu_{n_m}$. Then by continuity of the pushforward,

$$T_*\mu = \lim_{m \rightarrow \infty} \frac{1}{n_m} \sum_{k=0}^{n_m-1} T_* T_*^k \nu = \lim_{m \rightarrow \infty} \frac{1}{n_m} \sum_{k=0}^{n_m-1} T_*^{k+1} \nu = \lim_{m \rightarrow \infty} [(n_m)/(n_m + 1)](\mu_{n_m+1} - \nu).$$

The limit point of this

Let $\{B_i\}$ be a countable basis of balls for X . It suffices to show that for any finite, disjoint collection of balls $\{B_1, \dots, B_n\}$, we can find a Borel probability measure μ_n such that $\mu(T^{-1}(B_i)) = \mu(B_i)$, for then if we take a limit of maximal collections of balls whose radius tends to zero, and consider a subsequence of probability measures which converge weakly to some probability measure μ , then it follows that $T_*\mu = \mu$. It does not take too much work to find a μ_n which works.

We can break down the sets $\{B_1, \dots, B_n, T^{-1}B_1, \dots, T^{-1}B_n\}$ into finitely many atomic parts $\{A_1, \dots, A_m\}$, i.e. such that each set is a union of these parts, and such that the intersection of any subcollection of the sets is a finite union of these sets. The problem then just becomes a linear programming problem, i.e. each of the equations $\mu(T^{-1}(B_i)) = \mu(B_i)$ breaks down into an equation $\sum_{j \in S'_i} c_j = \sum_{j \in S_i} c_j$ where $c_i = 0$, subject to the constraint that $c_i \geq 0$.

- (b) If $M_1(T) = \{\mu\}$ consists of a single measure μ , then

$$\int_X f d\mu = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f \circ T^n(x)$$

for every continuous function $f : X \rightarrow \mathbb{R}$ and $x \in X$.

Solution: The result above shows that every subsequence of the measures

$$\mu_n = \frac{1}{N} \sum_{n=0}^{N-1} f \circ T^n$$

has a further subsequence converging in the weak $*$ topology to the invariant measure μ . But this is only possible if the sequence itself converges to μ .

67. Find the Fourier transform of the following function f in \mathbb{R}^2 :

$$f(x) = e^{ix\xi_0} |x - x_0|^{-1}.$$

Solution: First, we note that if $g(x) = |x - x_0|^{-1}$, then

$$\hat{f}(\xi) = \hat{g}(\xi - \xi_0/2\pi).$$

But if $h(x) = |x|^{-1}$, then

$$\hat{g}(\xi) = \hat{h}(\xi) e^{-2\pi i \xi \cdot x_0}.$$

Thus it suffices to find the Fourier transform of h . But h is radial, C^∞ away from the origin, and homogeneous of degree -1 . It follows that the Fourier transform of h is radial, C^∞ away from the origin, and homogeneous of degree -1 . Since there are no such homogeneous functions supported at the origin, we conclude that there must exist a quantity C such that

$$\hat{h}(\xi) = C|\xi|^{-1}.$$

To calculate C , it suffices to test h against a Gaussian. Namely, if $\Phi(x) = e^{-\pi|x|^2}$, then Φ is its own Fourier transform, and so the multiplication formula shows that

$$C \int |\xi|^{-1} \Phi(\xi) = \int \hat{h}(\xi) \Phi(\xi) = \int h(x) \Phi(x) = \int |x|^{-1} \Phi(x).$$

Thus we conclude that $C = 1$, i.e. that h is its own Fourier transform. Thus

$$\begin{aligned} \hat{f}(\xi) &= \hat{g}(\xi - \xi_0/2\pi) \\ &= \hat{h}(\xi - \xi_0/2\pi) e^{-2\pi i \xi \cdot x_0} e^{i \xi_0 \cdot x} \\ &= \frac{e^{-2\pi i \xi \cdot x_0} e^{i \xi_0 \cdot x}}{|\xi - \xi_0/2\pi|}. \end{aligned}$$

20 Day 13: Interchanging limits and integrals + Arzela-Ascoli Theorem

20.1 Limits and Integrals (DCT and friends)

68. (Rice, Winter 2011) Let $\{f_n\}$ be a sequence of Lebesgue measurable functions defined on $[0, 1]$ such that $\|f(x)\| \leq 1$ for all $n \geq 1$ and all $0 < x \leq 1$, and

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

exists for each $0 \leq x \leq 1$. Prove that

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{f_n(x)}{\sqrt{|x - 1/n|}} = \int_0^1 \frac{f(x)}{\sqrt{x}} dx$$

Solution: Let $g_n(x) = f_n(x)/\sqrt{|x - 1/n|}$. We wish to apply the Vitali convergence theorem. To do this, let us first show that the sequence $\{g_n\}$ is uniformly integrable. Now for a fixed $M > 0$ and $n > 0$, since $|f_n(x)| \leq 1$, if $|g_n(x)| \geq M$, then $|x - 1/n| \leq 1/M^2$, and so

$$\int_{|g_n(x)| \geq M} |g_n(x)| \leq \int_{|x - 1/n| \leq 1/M^2} \frac{1}{|x - 1/n|^{1/2}} \leq \int_{|x| \leq 1/M^2} \frac{1}{|x|^{1/2}} \lesssim M^{-1}.$$

Thus

$$\lim_{M \rightarrow \infty} \sup_{|g_n(x)| \geq M} |g_n(x)| = 0.$$

To show g_n converges to $g(x) = f(x)/\sqrt{x}$, it thus suffices to show that g_n converges to g in measure. Now g_n clearly converges to g except when $x = 0$, and thus g_n converges to g almost everywhere. But this implies g_n converges to g in measure (since we are working over a finite measure space), and thus g_n converges to g in $L^1[0, 1]$.

69. (Rice, Spring 2005) Compute

(a) $\lim_{n \rightarrow \infty} \int_0^\infty \frac{x^{n-2}}{1+x^n}.$

Solution: We apply the dominated convergence theorem. For $0 \leq x \leq 1$, $x^{n-2}/(1+x^n) \leq 1$, and for $1 \leq x \leq \infty$, $x^{n-2}/(1+x^n) \leq x^{n-2}/x^n \leq 1/x^2$. Thus the family of integrands $x^{n-2}/(1+x^n)$ is dominated by an integrable function, namely the function

$$\mathbf{I}(0 \leq x \leq 1) + \mathbf{I}(1 \leq x) \cdot 1/x^2.$$

The sequence converges pointwise to zero for $0 \leq x < 1$, and for $x > 1$,

$$\lim_{n \rightarrow \infty} x^{n-2}/(1+x^n) = \lim_{n \rightarrow \infty} 1/(x^2 + 1/x^{n-2}) = 1/x^2.$$

Thus the dominated convergence theorem implies that

$$\lim_{n \rightarrow \infty} \int_0^\infty \frac{x^{n-2}}{1+x^n} = \int_1^\infty \frac{1}{x^2} = 1.$$

(b) $\lim_{n \rightarrow \infty} n \int_0^\infty \frac{\sin y}{y(1+n^2 y)} dy$ (Hint: Substitute $x = ny$).

Solution: We have

$$n \int_0^\infty \frac{\sin y}{y(1+n^2 y)} dy = n \int_0^\infty \frac{\sin(y/n)}{y(1+ny)} dy.$$

For $y \geq 1$,

$$n \frac{\sin(y/n)}{y(1+ny)} \leq n \frac{1}{y(ny)} \leq 1/y^2.$$

For $y \leq 1$, we employ the bound $\sin(y/n) \leq y/n$ to conclude that

$$n \frac{\sin(y/n)}{y(1+ny)} \leq \frac{1}{1+ny} \leq 1.$$

These bounds imply that the family of integrands is uniformly integrable. Now for all $0 \leq x \leq 1$,

$$\lim_{n \rightarrow \infty} n \frac{\sin(y/n)}{y(1+ny)} = \lim_{n \rightarrow \infty} \frac{1}{1+ny} + O\left(n \frac{(y/n)^2}{y(1+ny)}\right) = \lim_{n \rightarrow \infty} \frac{1}{1+ny} + O\left(\frac{y}{n(1+ny)}\right) = 0.$$

Thus the dominated convergence theorem implies that

$$\lim_{n \rightarrow \infty} n \int_0^\infty \frac{\sin y}{y(1+n^2 y)} dy = 0.$$

70. (Spring 2017) Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a continuously differentiable function for which $\|f'\|_\infty < \infty$. Define, for $x > 0$,

$$F(x) = \int_0^\infty f(x+yx)\psi(y) dy,$$

where ψ satisfies

$$\int_0^\infty |\psi(y)| dy \quad \text{and} \quad \int_0^\infty y \cdot |\psi(y)| dy < \infty.$$

Show that $F(x)$ is well defined for all $x \geq 0$, and that F is continuously differentiable.

Solution: The fundamental theorem of calculus implies that

$$|f(x+yx)| = \left| f(x) + \int_x^{x+yx} f'(t) dt \right| = f(x) + O(yx).$$

Thus combined with the fact that $\int_0^\infty |\psi(y)| dy < \infty$ and $\int_0^\infty y \cdot |\psi(y)| dy < \infty$, this implies that

$$\int_0^\infty |f(x+yx)| |\psi(y)| dy \lesssim |f(x)| + |x| < \infty,$$

so F is well defined.

To show that F is continuously differentiable, we note that for a fixed $x \in [0, \infty)$, we calculate that, for h with $0 \leq x+h$,

$$\frac{F(x+h) - F(x)}{h} = \int_0^\infty \frac{|f(x+yx+h(1+y)) - f(x+yx)|}{h} \psi(y) dy.$$

Now

$$\left| \frac{|f(x+yx+h(1+y)) - f(x+yx)|}{h} \psi(y) \right| \lesssim (1+y) |\psi(y)|.$$

Since $(1+y)\psi(y)$ is integrable, the dominated convergence theorem implies that F is differentiable at x , and

$$F'(x) = \int_0^\infty f'(x+yx)(1+y)\psi(y) dy.$$

Finally, we show F' is continuous. We note that for any $\varepsilon > 0$, there exists $R > 0$ such that

$$\int_R^\infty (1+y)|\psi(y)| dy \leq \varepsilon.$$

Since f' is continuous, it is uniformly continuous on $[0, 2R(1+x)]$. Thus there exists $\delta > 0$ such that for $|h| \leq \delta$, and $0 \leq y \leq R$, $|f'((x+yx) + h(1+y)) - f'(x+yx)| \leq \varepsilon$, and so

$$\begin{aligned} F'(x+h) - F'(x) &\leq \varepsilon + \int_0^R [f'((x+yx) + h(1+y)) - f'(x+yx)](1+y)\psi(y) dy \\ &\leq \varepsilon + \int_0^R \varepsilon(1+y)\psi(y) dy \leq \varepsilon. \end{aligned}$$

20.2 Arzela-Ascoli

The key part of the Arzela-Ascoli theorem to know for the qual is the following: If $\{f_n\} \subset C[0,1]$ is a sequence which is uniformly bounded and equicontinuous, then $\{f_n\}$ has a uniformly convergent subsequence.

(Note that we can replace $[0,1]$ by any compact subset of \mathbb{R}^d . Also, there is a converse to the theorem, but I haven't seen it used in any qual problems.)

By *uniformly bounded*, we mean that $|f_n(x)| \leq C$ for all $x \in [0,1]$, $n \in \mathbb{N}$.

By *equicontinuous*, we mean that for all $\epsilon > 0$, there exists δ such that $|f_n(x) - f_n(y)| < \epsilon$ whenever $|x - y| < \delta$ for all $n \in \mathbb{N}$.

71. (From a UBC Math 321 Midterm) Let $\{f_n\}$ be a sequence of functions in $C[a,b]$ with no uniformly convergent subsequence. Define

$$F_n(x) = \int_a^x \sin(f_n(t)) dt.$$

Does $\{F_n\}$ has a uniformly convergent subsequence.

Solution: The sequence $\{F_n\}$ does have a uniformly convergent subsequence, by Arzela-Ascoli. Indeed, it is simple to see that $\|F_n\|_\infty \leq b - a$ for all $n > 0$, so the sequence $\{F_n\}$ is uniformly bounded. The sequence is also equicontinuous, because

$$|F_n(x) - F_n(y)| \leq |x - y|$$

uniformly in n . Thus Arzela-Ascoli implies the existence of a uniformly convergent subsequence.

72. (Fall 2004) Let $f_n : [0,1] \rightarrow \mathbb{R}$ be a sequence of continuous functions whose derivatives f'_n in the sense of distributions belong to $L^2(0,1)$. The functions also satisfy $f_n(0) = 0$.

(a) Assume that

$$\lim_{n \rightarrow \infty} \int_0^1 f'_n(x)g(x)dx$$

exists for all $g \in L^2(0, 1)$. Show that the f_n converge uniformly as $n \rightarrow \infty$.

Solution: Define $T_n(g) = \int_0^1 f'_n g dx$ for $g \in L^2(0, 1)$. Then T_n is clearly linear and by Cauchy-Schwarz inequality, T_n is bounded. Then by the part (a) assumption, we have

$$\sup\{T_n(g) : n \geq 1\} < \infty.$$

Therefore by the Uniform Boundedness Principle,

$$C := \sup_n \|f'_n\|_{L^2} < \infty$$

Uniform boundedness principle to obtain a uniform bound on f'_n . Then apply Arzela ascoli to a subsequence to obtain convergence of whole sequence. [Note the uniform boundedness of the derivatives suggests that we have uniform equicontinuity – and hence maybe we can use the Arzela-Ascoli Theorem.] In particular, since $f' \in L^2[0, 1]$, it follows that $f' \in L^1[0, 1]$. Therefore

$$f_n(x) - f_n(y) = \int_y^x f'_n(t) dt \quad (16)$$

Taking $y = 0$ and using $f_n(0) = 0$, we have

$$|f_n(x)| \leq \int_0^x |f'_n(t)| dt \leq \int_0^1 |f'_n(t)| dt \leq \left(\int_0^1 |f'_n(t)|^2 dt \right)^{\frac{1}{2}} \leq C.$$

Therefore $\{f_n\}_n$ is uniformly bounded. In addition, by (16) and Cauchy-Schwarz inequality, we have

$$|f_n(x) - f_n(y)| \leq \left| \int_y^x f'_n(t) dt \right| \leq \sqrt{x-y} \left(\int_y^x |f'_n|^2 dt \right)^{\frac{1}{2}} \leq C \sqrt{x-y}.$$

Therefore $\{f_n\}$ is uniformly equi-continuous. Therefore by the Arzela-Ascoli theorem, $\{f_n\}_{n \in \mathbb{N}}$ is a relatively compact subset of $C([0, 1])$.

Recall that by assumption $\lim_{n \rightarrow \infty} \int_0^1 f'_n(x) g(x) dx$ exists for any $g \in L^2[0, 1]$. Taking $g(x) = \chi_{[0, a]}$, we have $\lim_{n \rightarrow \infty} \int_0^a f'_n(x) dx = \lim_{n \rightarrow \infty} f'_n(a)$ exists. That is, there exists some real-valued function f such that $f_n \rightarrow f$ pointwise.

Suppose that f_n does not converge uniformly to f . Then there exists a subsequence n_k and $\epsilon_0 > 0$ such that

$$\sup_{x \in [0, 1]} |f(x) - f_{n_k}(x)| \geq \epsilon_0$$

for all k . However, since $\{f_n\}_n$ is relatively compact, there exists a subsequence $f_{n_{k_j}}$ which is uniformly Cauchy—and therefore must converge uniformly to f . But this implies that

$$\sup_{x \in [0, 1]} |f(x) - f_{n_{k_j}}(x)| \rightarrow 0 \quad \text{as } j \rightarrow \infty$$

This is a contradiction.

(b) Assume that

$$\lim_{n \rightarrow \infty} \int_0^1 f'_n(x) g(x) dx$$

exists for all $g \in C([0, 1])$. Do we still have the f_n converge uniformly?

Solution: Maybe yes by density?

73. (Spring 2014)

Consider the following operator

$$Af = \frac{1}{x\sqrt{1+|\log x|}} \int_0^x f(t)dt$$

Is A bounded as an operator from $L^2[0, 1]$ to $L^2[0, 1]$. Is it compact?

Solution: We first prove the following lemma:

Lemma 1 (Hardy's Inequality): If $F(x) = \int_0^x f(t)dt$ then

$$\int_0^1 \left(\frac{F(x)}{x} \right)^2 dx \leq 4 \int_0^1 f^2(x)dx$$

Proof of Lemma:

$$\begin{aligned} \int_0^1 \frac{F(x)^2}{x^2} dx &= - \left. \frac{F(x)^2}{x} \right|_0^1 - \int_0^1 \left(\frac{-1}{x} \right) 2F(x)f(x)dx && \text{Integration by parts} \\ &\leq 2 \int_0^1 \frac{F(x)}{x} f(x)dx \\ &\leq 2 \left(\int_0^1 \frac{F(x)^2}{x^2} dx \right)^{\frac{1}{2}} \left(\int_0^1 f(x)^2 dx \right)^{\frac{1}{2}} && \text{Cauchy-Schwarz} \end{aligned}$$

Dividing by $\left(\int_0^1 \frac{F(x)^2}{x^2} dx \right)^{\frac{1}{2}}$ and squaring both sides gives the result. \square

We will use Hardy's inequality to prove that A is bounded. Letting $F(x) = \int_0^x f(t)dt$, we have

$$\|Af\|_2^2 = \int_0^1 \frac{|F(x)|^2}{|x|^2} \frac{1}{1+|\log x|} dx \leq \int_0^1 \frac{|F(x)|^2}{|x|^2} dx \leq 4 \|f\|_2^2$$

where the last inequality is due to Hardy's inequality. Since A is clearly linear, it follows that A is a bounded linear operator.

Next we show that A is a compact operator. Let $\{f_n\}_{n \geq 1}$ be a bounded sequence in $L^2[0, 1]$. To show that A is compact, it suffices to show that there exists a subsequence $\{f_{n_k}\}_{k \geq 1}$ such that $\{Af_{n_k}\}_{k \geq 1}$ converges in $L^2[0, 1]$.

Claim 2. Let $F_n(x) := \int_0^x f_n(t)dt$. Then $\{F_n\}_{n=1}^\infty$ is a relatively compact subset of $C[0, 1]$ (with respect to the topology induced by the uniform norm).

To prove claim 2, we apply the Arzela-Ascoli Theorem. First, since $\{f_n\}_n$ is bounded in $L^2[0, 1]$, there exists some $C > 0$ such that $\|f_n\|_{L^2} \leq C$ for all n . Then

$$|F_n(x)| \leq \int_0^x |f_n(t)| dt \leq \left(\int_0^1 |f_n(t)|^2 dt \right)^{\frac{1}{2}} \leq C < \infty.$$

Therefore $\{F_n\}$ is uniformly bounded. It remains to show that $\{F_n\}$ is equicontinuous. By the Cauchy-Schwarz inequality,

$$|F_n(x) - F_n(y)| = \left| \int_y^x f_n(t) dt \right| \leq \|f_n\|_{L^2} \sqrt{|x - y|} \leq C \sqrt{|x - y|}.$$

Therefore $\{F_n\}$ is equicontinuous. Therefore by the Arzela-Ascoli theorem, $\{F_n\}_{n=1}^\infty$ is a relatively compact. This proves Claim 2.

By Claim 2, there exists a subsequence F_{n_k} such that $\{F_{n_k}\}_k$ is uniformly Cauchy. We claim that $\{Af_{n_k}\}_{k \geq 1}$ converges in L^2 -norm, which will complete the proof. Indeed,

$$\begin{aligned} \|Af_{n_j} - Af_{n_k}\|_{L^2}^2 &\leq \int_0^1 \frac{|F_{n_j}(x) - F_{n_k}(x)|^2}{|x|^2 (1 + |\log x|)} dx \\ &= \int_0^\delta \frac{|F_{n_j}(x) - F_{n_k}(x)|^2}{|x|^2 (1 + |\log x|)} dx + \int_\delta^1 \frac{|F_{n_j}(x) - F_{n_k}(x)|^2}{|x|^2 (1 + |\log x|)} dx \\ &\leq \frac{1}{1 + |\log \delta|} \underbrace{\int_0^\delta \frac{|F_{n_j}(x) - F_{n_k}(x)|^2}{|x|^2} dx}_{\text{apply Hardy's inequality again}} + \|F_{n_j} - F_{n_k}\|_{L^\infty} \cdot \frac{1}{\delta^2} \\ &\leq \frac{1}{1 + |\log \delta|} \|f_{n_j} - f_{n_k}\|_{L^2} + \frac{1}{\delta^2} \|F_{n_j} - F_{n_k}\|_{L^\infty} \\ &\leq \frac{2C}{1 + |\log \delta|} + \frac{1}{\delta^2} \|F_{n_j} - F_{n_k}\|_{L^\infty}. \end{aligned}$$

Let $\epsilon > 0$. Choose δ sufficiently small that $\frac{2C}{1 + |\log \delta|} < \epsilon/2$, and choose $N \geq 0$ sufficiently large that $\|F_{n_j} - F_{n_k}\|_{L^\infty} \leq \delta^2 \epsilon/2$ whenever $j, k \geq N$. So $j, k \geq N$ implies $\|Af_{n_j} - Af_{n_k}\|_{L^2} \leq \epsilon/2 + \epsilon/2 = \epsilon$. We have shown that $\{Af_{n_k}\}_k$ is Cauchy in $L^2[0, 1]$, as required.

74. (Problem 36 from the 2017 SEP) Consider the Hilbert space $L^2([0, 1])$ with inner product $(f, g) := \int_0^1 f(t)\bar{g}(t)dt$. Let $\{e_n\}_{n=1}^\infty$ be an orthonormal system of functions in $L^2([0, 1])$.

(a) Suppose that $e_n \in L^2([0, 1])$ for all $n \in \mathbb{N}$. Show that

$$\sup_n \max_{x \in [0, 1]} |e'_n(x)| = \infty.$$

Solution: Suppose not. Then there exists some $C < \infty$ such that

$$\sup_n \max_{x \in [0, 1]} |e'_n(x)| \leq C.$$

Therefore $|e_n(x) - e_n(y)| \leq C|x - y|$. Hence $\{e_n\}$ is equicontinuous.

By Chebychev's inequality,

$$|\{x \in [0, 1] : |e_n(x)| > \alpha\}| \leq \frac{\int_0^1 e_n(x)^2 dx}{\alpha^2} = \frac{1}{\alpha^2}.$$

Taking $\alpha = 2$, we find that $|\{x \in [0, 1] : |e_n(x)| \leq 2\}| > \frac{1}{4}$. Since every set with positive measure has at least one element, there exists some $x_n \in [0, 1]$ such that $|e_n(x)| \leq 2$. Therefore for any $x \in [0, 1]$ and any $n \in \mathbb{N}$, we have

$$|e_n(x)| = \left| e_n(x_n) + \int_{x_n}^x e'_n(t) dt \right| \leq 2 + |x_n - x| \leq 3.$$

Therefore $\{e_n\}$ is uniformly bounded. Therefore by the Arzela-Ascoli theorem, there exists some subsequence (e_{n_k}) such that $e_{n_k} \rightarrow f$ uniformly in $C[0, 1]$ as $k \rightarrow \infty$. Therefore $e_{n_k} \rightarrow f$ in $L^2[0, 1]$. But this contradicts the orthonormality of the sequence $\{e_n\}_{n=1}^\infty$. (In particular, by orthonormality we have for all $m \neq n$:

$$\|e_n - e_m\|_{L^2}^2 = \langle e_n - e_m, e_n - e_m \rangle = \langle e_n, e_n \rangle - \langle e_m, e_n \rangle - \langle e_n, e_m \rangle + \langle e_m, e_m \rangle = 1 - 0 - 0 + 1 = 2$$

and this implies that $\{e_n\}$ can have no subsequence which is Cauchy in L^2 .)

- (b) Suppose that e_n is complete, which means $(g, e_n) = 0$ for all n implies $g = 0$ almost everywhere. Prove

$$\sum_{n=1}^{\infty} |e_n(x)|^2 = \infty, \quad \text{almost everywhere.}$$

Solution: We first prove the following claim:

Claim 1. If $E \subset [0, 1]$ is measurable and $|E| > 0$, then

$$\int_E \sum_{n=1}^{\infty} |e_n(x)|^2 dx \geq 1.$$

To prove the claim, observe that

$$|E| = \|\chi_E\|_{L^2}^2 = \sum_{n=1}^{\infty} (\chi_E, e_n)^2 = \sum_{n=1}^{\infty} \left(\int_0^1 \chi_E(x) e_n(x) dx \right)^2 \leq \sum_{n=1}^{\infty} \left(\int_E e_n^2 dx \right) |E|$$

where the second equality is by Completeness of (e_n) and the inequality is due to Cauchy-Schwarz. Dividing both sides by $|E|$ and applying the Fubini-Tonelli theorem proves the claim. Define

$$A_M = \left\{ x : \sum_{n=1}^{\infty} |e_n(x)|^2 \leq M \right\}.$$

Suppose there exists some $M > 0$ such that $|A_M| > 0$. Let $E_n \subset A_M$ with $0 < |E_n| < 1/n$. Then by Claim 1,

$$1 \leq \int_{E_n} \sum_{n=1}^{\infty} |e_n(x)|^2 dx \leq \frac{M}{n} \rightarrow 0$$

a contradiction. Therefore $|A_M| = 0$ for all M . Since $\{A_M\}_M$ is an ascending sequence of sets, it follows by continuity of measure that

$$\left| \left\{ x : \sum_{n=1}^{\infty} |e_n(x)|^2 dx < \infty \right\} \right| = \left| \bigcup_{M=1}^{\infty} A_M \right| = \lim_{M \rightarrow \infty} |A_M| = 0.$$

21 Day 14: Misc. Topics

75. (Rice, Winter 2008) Is it possible to construct a measurable set $E \subset \mathbb{R}$ of positive measure such that for any pair $a < b$, $|E \cap [a, b]| \leq 0.5(b - a)$?

Solution: No, since the Lebesgue differentiation theorem implies that if E has positive measure, then there exists x_0 such that

$$\limsup_{x_0 \in I} |E \cap I|/|I| = 1.$$

However, if E is as above, then $|E \cap I|/|I| \leq 0.5$ for any interval I , so x_0 cannot possibly exist. Thus any E satisfying the condition above must have measure zero.

76. (Spring 2010) For $\lambda > 0$, set

$$F(\lambda) = \int_0^1 e^{-10\lambda x^4 + \lambda x^6} dx$$

Prove there exists constants A and $C > 0$, such that $F(\lambda) = \frac{A}{\lambda^{\frac{1}{4}}} + E(\lambda)$ where $|E(\lambda)| \leq \frac{C}{\lambda^{\frac{1}{2}}}$.

Solution: Supposing A and C exist, then $A = \lim_{\lambda \rightarrow \infty} \lambda^{\frac{1}{4}} F(\lambda)$. To determine the value of A , we make the u -substitution $u = \lambda^{\frac{1}{4}} x$, which gives

$$\lambda^{\frac{1}{4}} F(\lambda) = \int_0^{\lambda^{\frac{1}{4}}} e^{-10u^4 + \lambda^{-\frac{1}{2}} u^6} du = \int_0^{\infty} \chi_{(0, \lambda^{\frac{1}{4}})} e^{-10u^4 + \lambda^{-\frac{1}{2}} u^6} du. \quad (17)$$

Note that for $0 < u < \lambda^{\frac{1}{4}}$, it holds that $\lambda^{-\frac{1}{2}} u^6 \leq u^4$. It follows that

$$\chi_{(0, \lambda^{\frac{1}{4}})} e^{-10u^4 + \lambda^{-\frac{1}{2}} u^6} \leq e^{-9u^4}.$$

Since the right-hand side is integrable and $\chi_{(0, \lambda^{\frac{1}{4}})} e^{-10u^4 + \lambda^{-\frac{1}{2}} u^6} \rightarrow \chi_{(0, \infty)} e^{-10u^4}$ pointwise as $\lambda \rightarrow \infty$, it follows by the Dominated Convergence Theorem that

$$\lim_{\lambda \rightarrow \infty} \lambda^{\frac{1}{4}} F(\lambda) = \int_0^{\infty} e^{-10u^4} du.$$

So let $A := \int_0^{\infty} e^{-10u^4} du$. We need to show that there exists some constant C such that

$$\left| \lambda^{\frac{1}{4}} F(\lambda) - A \right| \leq \frac{C}{\lambda^{\frac{1}{4}}}.$$

In particular, by equation (17) and the triangle inequality,

$$\left| \lambda^{\frac{1}{4}} F(\lambda) - A \right| \leq \underbrace{\left| \int_0^{\lambda^{\frac{1}{4}}} e^{-10u^4} \left(e^{\lambda^{-\frac{1}{2}} u^6} - 1 \right) du \right|}_{\text{Call this } g(\lambda)} + \underbrace{\left| \int_{\lambda^{\frac{1}{4}}}^{\infty} e^{-10u^4} du \right|}_{\text{Call this } h(\lambda)},$$

so it will suffice to show that $\lambda^{\frac{1}{4}} g(\lambda) \leq C_1$ and $\lambda^{\frac{1}{4}} h(\lambda) \leq C_2$ for all λ .

To estimate $\lambda^{\frac{1}{4}}g(\lambda)$, we use the inequality $|e^x - 1| \leq |x|e^x$ (which holds for all $x > 0$ since $|e^x - 1| = |\int_0^x e^t dt| \leq xe^x$). Using this inequality, we have:

$$\begin{aligned}\lambda^{\frac{1}{4}}g(\lambda) &\leq \lambda^{\frac{1}{4}} \int_0^{\lambda^{\frac{1}{4}}} e^{-10u^4} \lambda^{-\frac{1}{2}} u^6 e^{\lambda^{-\frac{1}{2}} u^6} du \\ &= \lambda^{-\frac{1}{4}} \int_0^{\lambda^{\frac{1}{4}}} u^6 e^{-10u^4 + \lambda^{-\frac{1}{2}} u^6} du \\ &\leq \int_0^{\lambda^{\frac{1}{4}}} u^5 e^{-9u^4} du && \text{using } u \leq \lambda^{\frac{1}{4}} \\ &\leq C_1\end{aligned}$$

where $C_1 := \int_0^\infty u^5 e^{-9u^4} du < \infty$.

It remains to estimate $\lambda^{\frac{1}{4}}h(\lambda)$. Using $\lambda^{\frac{1}{4}} \leq u$, we have

$$\lambda^{\frac{1}{4}}h(\lambda) = \lambda^{\frac{1}{4}} \int_{\lambda^{\frac{1}{4}}}^\infty e^{-10u^4} du < \int_{\lambda^{\frac{1}{4}}}^\infty u e^{-10u^4} du < C_2$$

where $C_2 := \int_0^\infty u e^{-10u^4} du < \infty$. This completes the proof.

77. (Fall 2010) Let $I = [0, 1]$ and define for $f \in L^2(I)$ the Fourier coefficients as $\hat{f}(k) = \int_0^1 f(t) e^{-2\pi i k t} dt$ for any $k \in \mathbb{Z}$.

(a) Let \mathcal{G} be the set of all $L^2(I)$ functions with the property that $|\hat{f}(0)| \leq 1$ and $|\hat{f}(k)| \leq |k|^{-3/5}$ for any $k \in \mathbb{Z}$, $k \neq 0$. Prove that \mathcal{G} is a compact subset of $L^2(I)$.

Solution:

Let $f_n \in \mathcal{G}$. It will suffice to show that there exists a subsequence f_{n_j} and an $f \in \mathcal{G}$ such that $f_{n_j} \rightarrow f$ in $L^2(I)$.

We will apply Cantor's diagonal sequence argument to obtain a subsequence n_j such that $\lim_{j \rightarrow \infty} \hat{f}_{n_j}(k)$ exists for all $k \in \mathbb{Z}$. The details of this argument are as follows:

First, since $\{\hat{f}_n(0)\}_{n \in \mathbb{N}}$ is a bounded sequence of real numbers, there exists a subsequence $\{s(n, 0)\}_{n \in \mathbb{N}}$ of \mathbb{N} such that $\lim_{n \rightarrow \infty} \hat{f}_{s(n, 0)}(0) = c_0$ for some $c_0 \in \mathbb{R}$.

Next, since $|\hat{f}_{s(n, 0)}(k)| \leq |k|^{-3/5}$ for any $k \in \mathbb{Z} \setminus \{0\}$, it follows that $\{\hat{f}_{s(n, 0)}(1)\}_{n \in \mathbb{N}}$ and $\{\hat{f}_{s(n, 0)}(-1)\}_{n \in \mathbb{N}}$ are both bounded sequences of real numbers. Therefore there exists a subsequence $\{s(n, 1)\}_{n \in \mathbb{N}}$ of $\{s(n, 0)\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} \hat{f}_{s(n, 1)}(1) = c_1$ and $\lim_{n \rightarrow \infty} \hat{f}_{s(n, 1)}(-1) = c_{-1}$ for some $c_1, c_{-1} \in \mathbb{R}$.

In general, suppose we have chosen the subsequence $s(n, m-1)$. Then by the inequality $|\hat{f}(k)| \leq |k|^{-3/5}$, it follows that both $\{\hat{f}_{s(n, m-1)}(m)\}_{n \in \mathbb{N}}$ and $\{\hat{f}_{s(n, m-1)}(-m)\}_{n \in \mathbb{N}}$ are bounded sequences of real numbers, and therefore we may choose a further subsequence $\{s(n, m)\}_{n \in \mathbb{N}}$ of $\{s(n, m-1)\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} \hat{f}_{s(n, m)}(m) = c_m$ and $\lim_{n \rightarrow \infty} \hat{f}_{s(n, m)}(-m) = c_{-m}$ for some $c_m, c_{-m} \in \mathbb{R}$.

We now have in our hands a nested sequence of subsequences

$$\{s(n, 0)\}_{n \in \mathbb{N}} \supset \{s(n, 1)\}_{n \in \mathbb{N}} \supset \{s(n, 2)\}_{n \in \mathbb{N}} \supset \cdots$$

Now consider the diagonal subsequence $\{s(n, n)\}_{n \in \mathbb{N}}$. By construction,

$$\lim_{n \rightarrow \infty} \hat{f}_{s(n, n)}(k) = c_k \in \mathbb{R} \tag{18}$$

for all $k \in \mathbb{Z}$.

Let $f(t) = \sum_{k \in \mathbb{Z}} c_k e^{2\pi i k t}$. By Parseval's identity and the inequality $|\hat{f}(k)| \leq |k|^{-3/5}$, we have

$$\|f\|_{L^2(I)}^2 = \sum_{k \in \mathbb{Z}} |\hat{f}(k)|^2 \leq \left(1 + 2 \sum_{k \geq 1} |k|^{-6/5}\right) < \infty$$

Therefore f is well defined and in $L^2(I)$.

It remains to show that $f_{s(n,n)} \rightarrow f$ in $L^2(I)$. To see this, observe that by Parseval's identity,

$$\begin{aligned} \|f_{s(n,n)} - f\|_{L^2}^2 &= \sum_{k \in \mathbb{Z}} |\hat{f}_{s(n,n)}(k) - c_k|^2 \\ &= \sum_{|k| \leq N} |\hat{f}_{s(n,n)}(k) - c_k|^2 + \sum_{|k| > N} |\hat{f}_{s(n,n)}(k) - c_k|^2 \\ &\leq \sum_{|k| \leq N} |\hat{f}_{s(n,n)}(k) - c_k|^2 + \sum_{|k| > N} |K|^{-6/5} \end{aligned}$$

Therefore

$$\limsup_{n \rightarrow \infty} \|f_{s(n,n)} - f\|_{L^2}^2 \leq \sum_{|k| > N} |K|^{-6/5} \rightarrow 0 \text{ as } N \rightarrow \infty$$

Therefore $f_{s(n,n)} \rightarrow f$ in $L^2(I)$.

- (b) Let \mathcal{E} be the set of all $L^2(I)$ functions with the property that $\sum_k |\hat{f}(k)|^{5/3} \leq 2016^{-2016}$. Is \mathcal{E} a compact subset of $L^2(I)$?

Solution: \mathcal{E} is not compact. To see this, let $f_n(t) = e^{2\pi i n t}$. Then $\hat{f}_n(k) = \delta(k - n)$ where δ is the Kronecker-delta function. If $\epsilon_0 \in (0, 2016^{-2016})$, then $\{\epsilon_0 f_n\}_{n=1}^\infty \subset \mathcal{E}$ but it is not compact because it contains no convergent subsequence.

78. (Fall 2011) Let $\ell^2(\mathbb{N})$ denote the Hilbert space of square summable sequences with inner product $(x, y) = \sum_{n=1}^\infty x_n y_n$, where $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$.

- (a) What are the necessary and sufficient conditions on $\lambda_n > 0$ for the set

$$S = \{(x_1, x_2, \dots) \in \ell^2(\mathbb{N}) : |x_n| \leq \lambda_n, \forall n\}$$

to be compact in $\ell^2(\mathbb{N})$?

Solution: For any sequence $\{x(1), x(2), \dots\}$ in S , and any fixed i , the sequence $\{x(1)_i, x(2)_i, \dots\}$ is bounded, and thus has a convergent sub-sequence. Assuming that

$$\lim_{N \rightarrow \infty} \sum_{i=N}^\infty \lambda_i^2 = 0,$$

we will use diagonalization to construct a convergent subsequence. This is necessary, because the sequence $\{(\lambda_1, 0, \dots), (\lambda_1, \lambda_2, 0, \dots), (\lambda_1, \lambda_2, \lambda_3, 0, \dots), \dots\}$ only has a convergent subsequence if this property was true.

We will define a family of elements $x(i, j) \in \ell^2(\mathbb{N})$ for $0 \leq i < \infty$ and $1 \leq j < \infty$, such that for $0 < k \leq i$, $x(i, j)_k$ converges as $j \rightarrow \infty$. Begin by setting $x(0, i) = x(i)$. Given $x(i_0, j)$,

the sequence $x(i_0, j)_{i_0+1}$ is bounded in j , so we can find a sequence $y(j)$ such that $y(j)_{i_0+1}$ converges. Let $x_{i_0+1, j}$ be this convergent sub-sequence. Now define $y(k) = x(k, k)$. For each i , the sequence $\{y(k) : k > i\}$ is a subsequence of $\{x(i, j)\}$, and so $y(k)_i$ converges as $i \rightarrow \infty$ for each i . Thus for any fixed K ,

$$\limsup_{N \rightarrow \infty} \sup_{M > N} \sum_{j \leq K} |y(N)_j - y(M)_j|^2 = 0.$$

But

$$\limsup_{N \rightarrow \infty} \sup_{M > N} \sum_{j > K} |y(N)_j - y(M)_j|^2 \leq 4 \sum_{j > K} \lambda_j^2.$$

Thus for any K ,

$$\limsup_{N \rightarrow \infty} \sum_{M > N} \|y(N) - y(M)\|_2 \leq 2 \left(\sum_{j > K} \lambda_j^2 \right)^{1/2}$$

Taking $K \rightarrow \infty$ gives

$$\limsup_{N \rightarrow \infty} \sum_{M > N} \|y(N) - y(M)\|_2 = 0,$$

and so $\{y(k)\}$ is a Cauchy sequence in $\ell^2(\mathbb{N})$, and is thus convergent.

- (b) What are the necessary and sufficient conditions on $\mu_n > 0$ for the set

$$\left\{ (x_1, x_2, \dots) \in \ell^2(\mathbb{N}) : \sum_n \frac{|x_n|^2}{\mu_n^2} \leq 1 \right\}$$

to be compact in $\ell^2(\mathbb{N})$?

Solution: The necessary and sufficient condition is that $\mu_n \rightarrow 0$ as $n \rightarrow \infty$.

Proof of Necessity: Suppose that $\mu_n \not\rightarrow 0$. Then there exists an $\epsilon > 0$ and a subsequence $\{\mu_{n_j}\}$ with $\mu_{n_j} \geq \epsilon$. Let $x^{(j)} = (0, \dots, \mu_{n_j}, 0, \dots)$. It is easy to check that this sequence has no convergent subsequence.

Proof of Sufficiency: Suppose $\mu_n \rightarrow 0$.

Let $\{x^{(n)}\}$ be a sequence with $x^{(n)} \in \left\{ (x_1, x_2, \dots) \in \ell^2(\mathbb{N}) : \sum_n \frac{|x_n|^2}{\mu_n^2} \leq 1 \right\}$.

By a Cantor diagonalization argument, there exists a subsequence $\{x^{(n_j)}\}_j$ such that $x_k^{n_j} \rightarrow y_k \in \mathbb{R}$ for each k as $j \rightarrow \infty$.

We then check that that $y = (y_1, y_2, \dots) \in \ell^2$ and that $x^{n_j} \rightarrow y$ in ℓ^2 .

79. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a convex function, let $E = \{x \in \mathbb{R} : f \text{ is not differentiable at } x\}$. Show that E is at most countable.

Solution: For each $x_0 \in \mathbb{R}$, define $f'_-(x_0) := \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}$ and $f'_+(x_0) := \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0}$.

Claim 1: Both $f'_-(x_0)$ and $f'_+(x_0)$ exist and are finite for all $x_0 \in \mathbb{R}$.

To prove this, we will demonstrate that $f'_-(x_0)$ exists and is finite, as the case for $f'_+(x_0)$ is similar. Recall the chordal slope property of convex functions: if f is convex on \mathbb{R} , then for any $x_1 < x_2 < x_3$, the following inequality holds (draw a picture of this)

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} \leq \frac{f(x_3) - f(x_1)}{x_3 - x_1} \leq \frac{f(x_3) - f(x_2)}{x_3 - x_2}. \quad (19)$$

The proof of (19) requires only the definition of convexity and a little bit of algebra. Suppose $x_1 < x_2 < x_3$. Recall f convex means that for all $a, b \in \mathbb{R}$ and all $0 \leq \lambda \leq 1$, we have

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y).$$

Taking $x = x_1, y = x_3$ and $\lambda = \frac{x_3 - x_2}{x_3 - x_1}$ (so that $x_2 = \lambda x_1 + (1 - \lambda)x_3$) in the above equation, we obtain

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} \leq \frac{f(x_3) - f(x_2)}{x_3 - x_2}.$$

Then, with a little more algebra, we can rewrite this inequality as the equivalent forms

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} \leq \frac{f(x_3) - f(x_1)}{x_3 - x_1} \quad \text{and} \quad \frac{f(x_1) - f(x_3)}{x_1 - x_3} \leq \frac{f(x_2) - f(x_3)}{x_2 - x_3}.$$

This proves (19)

From (19), we can deduce the following:

1. For each fixed $x_0 \in \mathbb{R}$, the function $x \mapsto \frac{f(x) - f(x_0)}{x - x_0}$ is increasing on $(-\infty, x_0)$.
2. The family $\left\{ \frac{f(x) - f(x_0)}{x - x_0} : x \in (-\infty, x_0) \right\}$ is bounded above by $\frac{f(x_0 + 1) - f(x_0)}{(x_0 + 1) - x_0} = f(x_0 + 1) - f(x_0) < \infty$.

These two facts together imply that the limit

$$f'_-(x_0) := \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}$$

is both increasing and bounded above, and therefore exists as a real number. The case for $f'_+(x_0)$ is similar. This proves Claim 1.

By Claim 1, it follows that f is not differentiable at a point $x \in \mathbb{R}$ if and only if $f'_+(x) \neq f'_-(x)$. Let $E = \{x \in \mathbb{R} : f'_-(x) \neq f'_+(x)\}$. We need to show that E is at most countable.

To see this, we first make the following observations, all of which follow from (19):

- (a) f'_+ and f'_- are both increasing functions
- (b) $f'_-(x) \leq f'_+(x)$ for all $x \in \mathbb{R}$
- (c) $f'_+(a) \leq \frac{f(b) - f(a)}{b - a} \leq f'_-(b)$ if $a < b$

By (a), f'_+ and f'_- are continuous at all but countably many points. That is, the set

$$D := \{x \in \mathbb{R} : f'_+ \text{ is not continuous at } x\} \cup \{x \in \mathbb{R} : f'_- \text{ is not continuous at } x\}$$

is countable. We will show that $E \subset D$, which will complete the proof.

Suppose $x \notin D$. We need to show that $x \notin E$. By (b), $E = \{x \in \mathbb{R} : f'_-(x) < f'_+(x)\}$, so it will suffice to show that $f'_-(x) \geq f'_+(x)$.

Let x_n be a sequence such that $x_n > x$ and $x_n \rightarrow x$. Since $x \notin D$,

$$f'_-(x) = \lim_{n \rightarrow \infty} f'_-(x_n) \geq f'_+(x)$$

where the final inequality is justified by (c) since $x_n > x$.

(This proof was adapted from Royden and Fitzpatrick's *Real Analysis* 4th ed.)

80. (Fall 2015) Identify all $\alpha \in \mathbb{R}$ such that $\lim_{n \rightarrow \infty} \sin(2\pi n\alpha)$ exists.

Solution: Define $f(\alpha) = \sin(2\pi\alpha)$. It will suffice to characterize the α for which $\lim_{n \rightarrow \infty} f(n\alpha)$ converges, since $\lim_{n \rightarrow \infty} f(n\alpha)$ exists if and only if $\lim_{n \rightarrow \infty} \sin(2\pi n\alpha)$ exists. Observe that f is periodic: if $k \in \mathbb{Z}$ then $f(x) = \sin(2\pi x) = \sin(2\pi x + 2\pi k) = f(x + k)$. By periodicity it suffices to consider $\alpha \in [0, 1)$, since $\lim_{n \rightarrow \infty} f(n\alpha)$ converges if and only if $\lim_{n \rightarrow \infty} f(n(\alpha + k))$ converges for all $k \in \mathbb{Z}$.

Case 1. Suppose $\alpha \notin \mathbb{Q}$. Let $\xi_n := n\alpha - [n\alpha]$. Then by *Weyl's equidistribution theorem*, the sequence $\{\xi_n\}$ is equidistributed on $[0, 1)$ and hence is dense. If we choose a sequence n_k such that $\xi_{n_k} \rightarrow 3/4$ as $k \rightarrow \infty$, and another sequence m_k such that $\xi_{m_k} \rightarrow 1/4$, then we conclude that by continuity,

$$\liminf_{n \rightarrow \infty} f(n\alpha) \leq \liminf_{k \rightarrow \infty} f(n_k\alpha) = -1$$

and

$$\limsup_{n \rightarrow \infty} \sin(2\pi n\alpha) \geq \limsup_{k \rightarrow \infty} \sin(2\pi m_k) = 1.$$

This shows the \liminf differs from the \limsup , so $\lim_{n \rightarrow \infty} \sin(2\pi n\alpha)$ cannot exist. Therefore $\sin(2\pi n\alpha)$ does not converge for any $\alpha \notin \mathbb{Q}$.

Case 2. Suppose $\alpha \in \mathbb{Q}$. Write $\alpha = p/q$ where p, q are integers with $\gcd(p, q) = 1$. Let $x = 2\pi p/q$. Then the sequence $\xi_n := n\alpha - [n\alpha]$ is periodic with q distinct values, one of which is zero. Indeed,

$$\xi_q = q\frac{p}{q} - \left\lfloor q\frac{p}{q} \right\rfloor = p - p = 0$$

and

$$\xi_{q+1} = (q+1)\frac{p}{q} - \left\lfloor \frac{(q+1)p}{q} \right\rfloor = \frac{p}{q} - \left\lfloor \frac{p}{q} \right\rfloor = \xi_1.$$

Since $f(n\alpha) = f(\xi_n)$, it follows that the sequence $\{f(n\alpha)\}_{n=1}^{\infty}$ is periodic and takes up to q distinct values. Therefore, in order for $\lim_{n \rightarrow \infty} f(n\alpha)$ to converge, we must have

$$f(\xi_1) = f(\xi_2) = \dots = f(\xi_q) = f(0) = 0.$$

By periodicity of f , this is equivalent to

$$f\left(\frac{p}{q}\right) = f\left(\frac{2p}{q}\right) = \dots = f\left(\frac{(q-1)p}{q}\right) = f(p)$$

and

$$f(p) = \sin(2\pi p) = 0.$$

Therefore $\sin(2\pi p/q) = 0$. This implies that $q = 1$ or $q = 2$. So either $\alpha = 0$ or $1/2 \pmod{1}$. That is, $x = k\pi$ for $k \in \mathbb{Z}$.

22 Day 15: Spring 2021 Final Qualifying Exam

81. (3) For a Lebesgue measurable subset E of \mathbb{R} , denote $\mathbf{1}_E$ the indicator function of E (i.e. $\mathbf{1}_E(x) = 1$ for $x \in E$ and $\mathbf{1}_E(x) = 0$ for $x \in E^c$).

Let $\{E_n : n \in \mathbb{N}\}$ be a family of Lebesgue measurable subsets of \mathbb{R} with finite measure and let f be a measurable function such that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} |f(x) - \mathbf{1}_{E_n}(x)| dx = 0.$$

Prove that f is the indicator function of a measurable set.

Solution: For each $\epsilon > 0$,

$$\epsilon \mu \{x : |f(x) - \mathbf{1}_{E_n}(x)| > \epsilon\} < \int_{[|f - \mathbf{1}_{E_n}| > \epsilon]} |f(x) - \mathbf{1}_{E_n}(x)| dx \leq \int_{\mathbb{R}} |f(x) - \mathbf{1}_{E_n}(x)| dx \rightarrow 0$$

as $n \rightarrow \infty$. Therefore $\mathbf{1}_{E_n} \rightarrow f$ in measure. By definition of convergence in measure, we may choose a subsequence n_k such that

$$\mu \{x : |f(x) - \mathbf{1}_{E_{n_k}}(x)| > \epsilon\} \leq 2^{-k}.$$

Therefore

$$\sum_{k \geq 1} \mu \{x : |f(x) - \mathbf{1}_{E_{n_k}}(x)| > \epsilon\} < \infty.$$

Therefore by the Borel-Cantelli lemma,

$$\mu \{x : |f(x) - \mathbf{1}_{E_{n_k}}(x)| > \epsilon \text{ for infinitely many } k\} = 0.$$

In other words, $\lim_{k \rightarrow \infty} \mathbf{1}_{E_{n_k}}(x) = f(x)$ for a.e. $x \in \mathbb{R}$. Therefore f is equal to zero or one almost everywhere, and as the a.e. limit of measurable functions, f is also measurable. Therefore f is the indicator function of a measurable set.

82. (6) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a compactly supported function that satisfies the Hölder condition with exponent $\beta \in (0, 1)$, i.e., there exists a constant $A < \infty$ such that $\forall x, y \in \mathbb{R} : |f(x) - f(y)| \leq A|x - y|^\beta$. Consider the function g defined by

$$g(x) = \int_{-\infty}^{\infty} \frac{f(y)}{|x - y|^\alpha} dy$$

where $\alpha \in (0, \beta)$.

- (a) Prove that g is a continuous function at 0.

Solution: We will show that $|g(x) - g(0)| \rightarrow 0$ as $|x| \rightarrow 0$. Without loss of generality, we may assume that $|x| \leq 1$. Performing the u -substitution $u = y - x$, we may write

$$g(x) = \int_{-\infty}^{\infty} \frac{f(x + y)}{|y|^\alpha} dy.$$

Therefore, taking $k > 0$ sufficiently large that $\text{supp}(f) \subseteq [-k, k]$, we have

$$\begin{aligned}
 |g(x) - g(0)| &= \left| \int_{-\infty}^{\infty} \frac{f(x+y) - f(y)}{|y|^\alpha} dy \right| \\
 &= \left| \int_{-(k+1)}^{k+1} \frac{f(x+y) - f(y)}{|y|^\alpha} dy \right| && \text{since } |x| \leq 1 \\
 &\leq \int_{-(k+1)}^{k+1} \frac{|f(x+y) - f(y)|}{|y|^\alpha} dy \\
 &\leq 2A|x|^\beta \int_0^{k+1} y^{-\alpha} dy \\
 &= 2A|x|^\beta \frac{(k+1)^{1-\alpha}}{1-\alpha} \rightarrow 0 \text{ as } |x| \rightarrow 0
 \end{aligned}$$

where we have used the fact that $\alpha < 1$ so that the integral is finite.

- (b) Prove that g is differentiable at 0. (Hint: Try the dominated convergence theorem).

Solution:

So the idea is to use the so called generalized dominated convergence theorem, whose proof is exactly the same as the usual dominant convergence theorem. And the statement is the following:

Let $\{h_n\}$ be a family of L^1 function which converge almost everywhere to a L^1 function h , let g_n be another family of L^1 functions such that $g_n \rightarrow g$ a.e., $|h_n| \leq g_n$ for each n and $|h| \leq g$, Suppose now we have $\lim \int g_n = \int g$, then $\lim \int h_n = \int h$.

Now come back to our problem. First attempt would be to use the standard test, which means we just naively take the derivative of the inside function and then check integrability. We immediately encounter a problem because we don't have the desired integrability condition. In fact, we have $f(y) \cdot \text{sgn}(x-y)|x-y|^{-\alpha-1}$, which is not integrable.

So our second attempt would be to use the Holder condition to balance this singularity. Assume that $\text{supp}(f) \subset [-K, K]$ for some $K > 1$. Let (x_n) be a sequence of real numbers converging to zero. Without loss of generality, we may assume $|x_n| \leq 1$ for all n . By the Mean Value Theorem, for each n there exists some ξ_n with $0 \leq |\xi_n| \leq |x_n|$ such that

$$\frac{1}{|y-x_n|^\alpha} - \frac{1}{|y|^\alpha} = \alpha \cdot \text{sgn}(y-\xi_n) \cdot |y-\xi_n|^{-\alpha-1} \cdot |x_n|.$$

Therefore

$$\begin{aligned}
 \frac{g(x_n) - g(0)}{x_n} &= \int_{-K}^K \frac{f(y) - f(\xi_n)}{|y-x_n|^\alpha \cdot x_n} - \frac{f(y) - f(\xi_n)}{|y|^\alpha \cdot x_n} dy + E_n \\
 &= \int_{-K}^K \frac{f(y) - f(\xi_n)}{x_n} \cdot \left(\frac{1}{|y-x_n|^\alpha} - \frac{1}{|y|^\alpha} \right) dy + E_n \\
 &= \int_{-K}^K (f(y) - f(\xi_n)) \cdot \alpha \cdot \text{sgn}(y-\xi_n) \cdot |y-\xi_n|^{-\alpha-1} dy + E_n \quad (20)
 \end{aligned}$$

where

$$E_n = \int_{-K}^K \frac{f(\xi_n)}{x_n} \left(\frac{1}{|y-x_n|^\alpha} - \frac{1}{|y|^\alpha} \right) dy.$$

In particular, we have

$$|E_n| = \left| \int_{-K-x_n}^{-K} \frac{f(\xi_n)}{x_n} \frac{1}{|y|^\alpha} dt - \int_{K-x_n}^K \frac{f(\xi_n)}{x_n} \frac{1}{|y|^\alpha} dy \right| = \left| \frac{f(\xi_n)}{x_n} \int_{K-x_n}^{K+x_n} \frac{1}{|y|^\alpha} dy \right| \leq \frac{C}{K^\alpha}.$$

for $C > 0$ sufficiently large. Therefore by the above estimate for $|E_n|$ and equation (20) we have:

$$\left| \frac{g(x_n) - g(0)}{x_n} - \int_{\mathbb{R}} h_n(y) dy \right| \leq \frac{C}{K^\alpha} \quad (21)$$

where

$$h_n(y) = \chi_{[-K, K]} \cdot (f(y) - f(\xi_n)) \cdot \alpha \cdot \operatorname{sgn}(y - \xi_n) \cdot |y - \xi_n|^{-\alpha-1}.$$

Let

$$h(y) := \lim_{n \rightarrow \infty} h_n(y) = \chi_{[-K, K]} \cdot (f(y) - f(0)) \cdot \alpha \cdot \operatorname{sgn}(y) \cdot |y|^{-\alpha-1}.$$

We wish to show that $\int_{\mathbb{R}} h_n(y) dy \rightarrow \int_{\mathbb{R}} h(y) dy$ as $n \rightarrow \infty$. We now justify this by applying the generalized dominant convergence theorem. Define

$$g_n = C \cdot |y - \xi_n|^{\beta-\alpha-1} \cdot \chi_{[-K, K]}$$

where C is some large constant. Similarly, define

$$g = C \cdot |y|^{\beta-\alpha-1} \cdot \chi_{[-K, K]}.$$

One easily sees that $|h_n| \leq g_n$ and $|h| \leq g$ by applying the Holder condition. Moreover, each g_n is in L^1 . Since $\xi_n \rightarrow 0$, we have $\lim \int g_n \rightarrow \int g$ and $g_n \rightarrow g$ a.e. Finally, invoke the generalized dominated convergence theorem, and it follows that $\lim_{n \rightarrow \infty} \int h_n = \int h$, so that by equation (21),

$$\limsup_{n \rightarrow \infty} \left| \frac{g(x_n) - g(0)}{x_n} - \int_{\mathbb{R}} h(y) dy \right| \leq \frac{C}{K^\alpha}$$

Taking $K \rightarrow \infty$, it follows that

$$\lim_{n \rightarrow \infty} \frac{g(x_n) - g(0)}{x_n} = \int_{\mathbb{R}} (f(y) - f(0)) \cdot \alpha \cdot \operatorname{sgn}(y) \cdot |y|^{-\alpha-1} dy.$$

Since (x_n) was an arbitrary sequence converging to zero, it follows that

$$g'(0) = \int_{\mathbb{R}} (f(y) - f(0)) \cdot \alpha \cdot \operatorname{sgn}(y) \cdot |y|^{-\alpha-1} dy$$

which is finite.

83. ((Spring 2021) points) A real valued function f defined on \mathbb{R} belongs to the space $C^{1/2}(\mathbb{R})$ if and only if

$$\sup_{x \in \mathbb{R}} |f(x)| + \sup_{x \neq y} \frac{|f(x) - f(y)|}{\sqrt{|x - y|}} < \infty.$$

Prove that a function $f \in C^{1/2}(\mathbb{R})$ if and only if there exists a constant C so that for every $\varepsilon > 0$, there is a bounded function $\varphi \in C^\infty(\mathbb{R})$ such that

$$\sup_{x \in \mathbb{R}} |f(x) - \varphi(x)| \leq C\varepsilon^{1/2} \quad \text{and} \quad \sup_{x \in \mathbb{R}} \varepsilon^{1/2} |\varphi'(x)| \leq C.$$

Solution: Suppose $f \in C^{1/2}(\mathbb{R})$. Let ϕ be a bump function symmetrical about the origin supported on $|x| \leq 1$, set $\phi_\varepsilon(y) = (1/\varepsilon)\phi(y/\varepsilon)$ and define $\varphi = f * \phi_\varepsilon$. Then φ is bounded, and lies in $C^\infty(\mathbb{R})$. Since $\phi_\varepsilon(y)$ is supported on $|y| \leq \varepsilon$, We have

$$f(x) - \varphi(x) = \int [f(x) - f(x-y)]\phi_\varepsilon(y) \lesssim \varepsilon^{1/2}.$$

We write (using the symmetry of ϕ)

$$\varphi'(x) = (1/\varepsilon)^2 \int_0^\infty \phi'(y/\varepsilon)[f(x-y) - f(x+y)] dy.$$

Now $|f(x-y) - f(x+y)| \lesssim \varepsilon^{1/2}$ for $|y| \leq \varepsilon$, the support of our integrand lies on $0 \leq y \leq \varepsilon$, and $|\phi'(y/\varepsilon)| \lesssim 1$. Thus it follows that $|\varphi'(x)| \lesssim \varepsilon^{1/2}$, which completes the proof.

Now suppose the second condition. Given φ satisfying the condition above, the derivative condition together with the fundamental theorem of calculus implies that

$$|f(x) - f(y)| \leq |f(x) - \varphi(x)| + |\varphi(x) - \varphi(y)| + |\varphi(y) - f(y)| \leq 2C\varepsilon^{1/2} + C|x-y|/\varepsilon^{1/2}.$$

If we choose $\varepsilon = |x-y|$, then we conclude that

$$|f(x) - f(y)| \leq 3C|x-y|^{1/2},$$

so that

$$\sup_{x \neq y} \frac{|f(x) - f(y)|}{|x-y|^{1/2}} \leq 3C.$$

The fact that φ is bounded shows that if the condition holds for any $\varepsilon > 0$, then f is bounded. Thus $f \in C^{1/2}(\mathbb{R})$.

84. (Fall 2021) Let $f \in L^1(\mathbb{R})$ satisfy $\int_a^b f(x)dx = 0$ for any two rational numbers a and b , $a < b$. Does it follow that $f(x) = 0$ for almost every x ?

Solution: Yes. We observe that this property follows immediately from the Lebesgue differentiation theorem, however this would trivialize the problem, so we present another solution here, using the $\pi - \lambda$ theorem. The $\pi - \lambda$ theorem states that if Π is a π -system and \mathcal{L} is a λ -system, then

$$\Pi \subseteq \mathcal{L} \quad \text{implies} \quad \sigma(\Pi) \subseteq \mathcal{L}.$$

Let $\mathcal{B}_{\mathbb{R}}$ denote the Borel σ -algebra on \mathbb{R} . Let

$$\mathcal{L} := \left\{ E \in \mathcal{B}_{\mathbb{R}} : \int_E f(x)dx = 0 \right\}$$

and let

$$\Pi := \{(a, b) : a \leq b, \text{ and } a, b \in \mathbb{Q}\}.$$

Next, we make the following observations

1. Π is a π -system. (This is obvious).
2. $\Pi \subseteq \mathcal{L}$. (This is given to us in the problem statement).

3. \mathcal{L} is a λ -system. (Just need to verify that \mathcal{L} satisfies the definition of λ -system. Establishing that \mathcal{L} is closed under countable unions requires use of the fact that f is integrable in order to justify use of either continuity of integration or the dominated convergence theorem).
4. $\mathcal{B}_{\mathbb{R}} \subseteq \sigma(\Pi)$. (Since $\mathcal{B}_{\mathbb{R}}$ is by definition the minimal σ -algebra containing the open sets, it suffices to show that $\sigma(\Pi)$ contains the open sets. Therefore, since each open set in \mathbb{R} can be written as a disjoint union of open intervals, it suffices to show that $\sigma(\Pi)$ contains the open intervals. Indeed, if $(x, y) \subseteq \mathbb{R}$ is an open interval, then there exists sequences of rational numbers (a_n) and (b_n) such that $x \leq a_n \leq b_n \leq y$ for all n and $a_n \rightarrow x$ and $b_n \rightarrow y$. Then $(x, y) = \cup_{n=1}^{\infty} (a_n, b_n)$. This implies $(x, y) \in \sigma(\Pi)$, as required.)

Observations (1)-(3) justify the use of the $\pi - \lambda$ theorem, which implies

$$\sigma(\Pi) \subseteq \mathcal{L}$$

and hence by observation (4), it follows that

$$\mathcal{B}_{\mathbb{R}} \subseteq \mathcal{L}.$$

But this implies that

$$\int_E f(x) dx = 0 \quad \text{for all } E \in \mathcal{B}_{\mathbb{R}}. \quad (22)$$

for all $E \in \mathcal{B}_{\mathbb{R}}$. In particular, let $E_+ := \{x : f(x) > 0\}$. Since f is measurable, it follows that $E_+ \in \mathcal{B}_{\mathbb{R}}$ and hence by (22) that $f(x) \leq 0$ a.e. A similar argument using the set $E_- := \{x : f(x) < 0\}$ implies that $f(x) \geq 0$ a.e. Therefore $f(x) = 0$ a.e.

23 Questions that need solutions

85. (Fall 2017) Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function that has continuous partial derivatives in \mathbb{R}^2 . Define $\chi : \mathbb{R}^3 \rightarrow \{0, 1\}$ by $\chi(x) = 1$ if $x_3 > g(x_1, x_2)$, and $\chi(x) = 0$ otherwise. Compute the derivatives $\partial\chi/\partial x^i$ for $i = 1, 2, 3$.

Solution: The function χ is constant away from the surface $x_3 = g(x_1, x_2)$, so we should expect the distributional derivatives of χ to be supported on this surface. Now for a test function ϕ , using the fundamental theorem of calculus, we have

$$\begin{aligned} \int \partial_3 \chi(x) \phi(x) &= - \int \int_{g(x_1, x_2)}^{\infty} \chi(x) \partial_3 \phi(x) \, dx_3 \, dx_1 \, dx_2 \\ &= \int \chi(x_1, x_2, g(x_1, x_2)) \phi(x_1, x_2, g(x_1, x_2)) \, dx_1 \, dx_2. \end{aligned}$$

Thus $\partial_3 \chi$ is equal to integration against the measure obtained by pulling back the Lebesgue measure on \mathbb{R}^2 along the projection π projecting the graph $x_3 = g(x_1, x_2)$ onto the x_3 axis. Next, if $S_{x_1} = \{(x_2, x_3) : g(x_1, x_2) < x_3\}$, we have

$$\begin{aligned} \int \partial_1 \chi(x) \phi(x) \, dx &= - \int \chi(x) \partial_1 \phi(x) \, dx \\ &= - \int \end{aligned}$$

86. Let $\alpha \in (0, 1)$, and for $f \in C[0, 1]$, and $x \in [0, 1]$, define

$$(T_\alpha f)(x) = \int_0^1 \sin(x-y) |x-y|^{-\alpha} f(y) \, dy.$$

- (a) Prove that T_α extends to a bounded operator on $L^2[0, 1]$.

Solution: We calculate using Minkowski's inequality that

$$\begin{aligned} \|T_\alpha f\|_{L^1[0,1]} &= \left\| \int_0^1 \sin(x-y) |x-y|^{-\alpha} f(y) \, dy \right\|_{L_x^1[0,1]} \\ &\leq \int_0^1 \| |\sin(x-y)| |x-y|^{-\alpha} f(y) \|_{L_x^1[0,1]} \, dy \\ &= \int_0^1 |f(y)| \left(\int_0^1 |\sin(x-y)| |x-y|^{-\alpha} \, dx \right) \, dy \\ &\leq \int_0^1 |f(y)| \int_0^1 |x-y|^{-\alpha} \, dx \, dy \\ &\leq \left(2 \int_0^1 t^{-\alpha} \, dt \right) \left(\int_0^1 |f(y)| \, dy \right) \\ &\leq \frac{2 \cdot 5^{1-\alpha}}{1-\alpha} \|f\|_{L^1[0,1]}. \end{aligned}$$

Thus T_α is bounded from $L^1[0, 1]$. Similarly, we verify that

$$\begin{aligned}\|T_\alpha f\|_{L^\infty[0,1]} &= \sup_x \left| \int_0^1 \sin(x-y)|x-y|^{-\alpha} f(y) dy \right| \\ &\leq \|f\|_{L^\infty[0,1]} \sup_x \int_0^1 |\sin(x-y)||x-y|^\alpha dy \\ &\leq \|f\|_{L^\infty[0,1]} \left(2 \int_0^1 t^{-\alpha} dt \right) \\ &\lesssim \|f\|_{L^\infty[0,1]}.\end{aligned}$$

Thus T_α is bounded from $L^\infty[0, 1]$ to itself. Interpolation theory then implies that T_α is bounded on $L^p[0, 1]$ for all $1 \leq p \leq \infty$. In particular, T_α is bounded on $L^2[0, 1]$.

- (b) For which $\alpha \in (0, 1)$ is $T_\alpha : L^2[0, 1] \rightarrow L^2[0, 1]$ a compact operator?

Solution: The operator T_α is an integral operator with kernel

$$K_\alpha(x, y) = \sin(x-y)|x-y|^{-\alpha}.$$

We calculate that

$$\begin{aligned}\|K_\alpha\|_{L^2_x[0,1]L^2_y[0,1]} &\leq \left(\int_0^1 \int_0^1 |x-y|^{-2\alpha} dx dy \right)^{1/2} \\ &\lesssim \left(\int_{|z| \leq 10} |z|^{-2\alpha} dz \right)^{1/2} \\ &\lesssim \left(\int_0^{10} t^{-2\alpha} dt \right)^{1/2}.\end{aligned}$$

For $\alpha < 1/2$, this quantity is finite, and thus T_α is a Hilbert-Schmidt operator, and thus compact. It is natural to guess this is also the range under which operator is compact, i.e. the operator is not compact for $\alpha \geq 1/2$.

The remaining part of the problem is quite hard. Here are some ideas on a proof, but the ideas are certainly not rigorous. Let me know if you find a solution, and I can add it to this document, crediting you. We let

$$K_\alpha(x) = \sin(x)|x|^{-\alpha}$$

Then $T_\alpha f = K_\alpha * f$, restricted to $[0, 1]$. The Fourier transform of $|x|^{-\alpha}$ is $|\xi|^{-(1-\alpha)}$ multiplied by a constant, and so, morally speaking, ignoring constants, we have

$$\widehat{K_\alpha} = (\delta_1 - \delta_{-1})|\xi|^{-(1-\alpha)} = |\xi - 1|^{-(1-\alpha)} - |\xi + 1|^{-(1-\alpha)}.$$

Since T_α acts by multiplying f on the frequency side of things by $\widehat{K_\alpha}$, we see that the function might behave the worst when f is oscillating at a frequency -1 or 1 . For the remainder of the proof, we will pretend like we are actually working on \mathbb{R} instead of $[0, 1]$, i.e. we are working with the operator $T_\alpha f = K_\alpha * f$ on $L^2(\mathbb{R})$. If ϕ is a Schwartz function whose Fourier transform is supported on $1/10 \leq |\xi| \leq 2/10$, and we let

$$f_N(x) = N^{-1/2} \phi(x/N) e^{ix},$$

then the Fourier transform of $T_\alpha f_N$ should be approximately equal to

$$\begin{aligned} & |\xi - 1|^{1-\alpha} N^{1/2} \widehat{\phi}(N(\xi - 1)) - |\xi + 1|^{-(1-\alpha)} N^{1/2} \widehat{\phi}(N(\xi - 1)) \\ & \approx |(1 + 1/10N) - 1|^{-(1-\alpha)} N^{1/2} \widehat{\phi}(N(x - 1)) - |1 + 1|^{-(1-\alpha)} N^{1/2} \widehat{\phi}(N(\xi - 1)) \\ & \approx (1/10N)^{-(1-\alpha)} N^{1/2} \widehat{\phi}(N(x - 1)) - 2^{-(1-\alpha)} N^{1/2} \widehat{\phi}(N(\xi - 1)) \\ & \approx N^{1/2-\alpha} \widehat{\phi}(N(x - 1)) - N^{1/2} \widehat{\phi}(N(\xi - 1)). \end{aligned}$$

Thus we should have

$$T_\alpha f_N \approx (N^{-\alpha} - 1) f_N$$

and so

$$\|T_\alpha f_N + f_N\|_{L^2[0,1]} \lesssim N^{-\alpha}.$$

This means $\{T_\alpha f_N\}$ has a convergent subsequence if and only if $\{f_N\}$ has a convergent subsequence. But this is clearly impossible since $\limsup |f_N(x)| = 0$ and so any subsequence must converge pointwise almost everywhere to zero.