

Fractals avoiding Fractal Sets

Jacob Denson

University of British Columbia

November 10, 2018

Examples of Configuration Problems

- ▶ How large can a subset of the \mathbf{R} be such that it contains no three term arithmetic progressions?

Examples of Configuration Problems

- ▶ How large can a subset of the \mathbf{R} be such that it contains no three term arithmetic progressions?
- ▶ How large can a subset of \mathbf{R}^d be such that the distances between any two points are distinct?

Examples of Configuration Problems

- ▶ How large can a subset of the \mathbf{R} be such that it contains no three term arithmetic progressions?
- ▶ How large can a subset of \mathbf{R}^d be such that the distances between any two points are distinct?
- ▶ How large can a subset of \mathbf{R}^d be such that the angles formed from any three points in X are irrational?

Two Observations

- ▶ These are ‘multi-scale’ configurations.

Two Observations

- ▶ These are ‘multi-scale’ configurations.
 - ▶ Every open set contains these configurations.

Two Observations

- ▶ These are ‘multi-scale’ configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.

Two Observations

- ▶ These are ‘multi-scale’ configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a ‘second order’ notion of size.

Two Observations

- ▶ These are ‘multi-scale’ configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a ‘second order’ notion of size.
 - ▶ We use Hausdorff dimension (Not too technically!).

Two Observations

- ▶ These are 'multi-scale' configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a 'second order' notion of size.
 - ▶ We use Hausdorff dimension (Not too technically!).
- ▶ The problems can be stated by avoiding values of a function.

Two Observations

- ▶ These are 'multi-scale' configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a 'second order' notion of size.
 - ▶ We use Hausdorff dimension (Not too technically!).
- ▶ The problems can be stated by avoiding values of a function.
 - ▶ x, y, z is in a three term arithmetic progression if and only if $f(x, y, z) = (x - y) - (y - z) = 0$.

Two Observations

- ▶ These are 'multi-scale' configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a 'second order' notion of size.
 - ▶ We use Hausdorff dimension (Not too technically!).
- ▶ The problems can be stated by avoiding values of a function.
 - ▶ x, y, z is in a three term arithmetic progression if and only if $f(x, y, z) = (x - y) - (y - z) = 0$.
 - ▶ The distances between two pairs x, y and z, w are non distinct when $f(x, y; z, w) = d(x, y) - d(z, w) = 0$.

Two Observations

- ▶ These are ‘multi-scale’ configurations.
 - ▶ Every open set contains these configurations.
 - ▶ Even every set of positive measure has these configurations.
 - ▶ We need a ‘second order’ notion of size.
 - ▶ We use Hausdorff dimension (Not too technically!).
- ▶ The problems can be stated by avoiding values of a function.
 - ▶ x, y, z is in a three term arithmetic progression if and only if $f(x, y, z) = (x - y) - (y - z) = 0$.
 - ▶ The distances between two pairs x, y and z, w are non distinct when $f(x, y; z, w) = d(x, y) - d(z, w) = 0$.
 - ▶ The angles formed by three points are irrational when $f(x, y, z) = (x - z)(y - z)/|x - z||y - z| \notin \cos(\mathbf{Q})$.

General Results

- ▶ **Configuration Avoidance Problem:** Given $f : [0, 1]^{nd} \rightarrow \mathbf{R}$, find $X \subset \mathbf{R}^d$ with high Hausdorff dimension such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.

General Results

- ▶ **Configuration Avoidance Problem:** Given $f : [0, 1]^{nd} \rightarrow \mathbf{R}$, find $X \subset \mathbf{R}^d$ with high Hausdorff dimension such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.
- ▶ Mathé (2017) proved that if f is a polynomial of degree r , we can find X with dimension d/r .

General Results

- ▶ **Configuration Avoidance Problem:** Given $f : [0, 1]^{nd} \rightarrow \mathbf{R}$, find $X \subset \mathbf{R}^d$ with high Hausdorff dimension such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.
- ▶ Mathé (2017) proved that if f is a polynomial of degree r , we can find X with dimension d/r .
- ▶ Pramanik and Fraser (2018) proved that if f is smooth and nonsingular, we can find X with dimension $d/(n-1)$.

What if the zero sets of the function are also fractals...

Main Result

Theorem

If the zero set of f is α dimensional, we can find X with

$$\dim_{\mathbf{H}}(X) = \frac{nd - \alpha}{n - 1} = \frac{\text{codim}_{\mathbf{H}}(f^{-1}(0))}{n - 1}$$

such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.

Main Result

Theorem

If the zero set of f is α dimensional, we can find X with

$$\dim_{\mathbf{H}}(X) = \frac{nd - \alpha}{n - 1} = \frac{\operatorname{codim}_{\mathbf{H}}(f^{-1}(0))}{n - 1}$$

such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.

- ▶ Extension of Pramanik and Fraser's result (2018), who proved the case where f is smooth and nonsingular, so $f^{-1}(0)$ is a smooth surface of dimension $nd - d$.

Main Result

Theorem


If the zero set of f is α dimensional, we can find X with

$$\dim_{\mathbf{H}}(X) = \frac{nd - \alpha}{n - 1} = \frac{\operatorname{codim}_{\mathbf{H}}(f^{-1}(0))}{n - 1}$$

such that for any distinct $x_1, \dots, x_n \in X$, $f(x_1, \dots, x_n) \neq 0$.


- ▶ Extension of Pramanik and Fraser's result (2018), who proved the case where f is smooth and nonsingular, so $f^{-1}(0)$ is a smooth surface of dimension $nd - d$.
- ▶ Shows smoothness is only needed to get a bound on the Hausdorff dimension. Not needed anywhere else.

Applications




- More robust result of Pramanik and Fraser, showing that we can 'thicken' or 'thin' the zero set of our function with stable effects on the Hausdorff dimension of X .

Applications



- ▶ More robust result of Pramanik and Fraser, showing that we can 'thicken' or 'thin' the zero set of our function with stable effects on the Hausdorff dimension of X .
- ▶ Uncountable unions of regular sets are allowed!

Applications



- ▶ More robust result of Pramanik and Fraser, showing that we can 'thicken' or 'thin' the zero set of our function with stable effects on the Hausdorff dimension of X .
- ▶ Uncountable unions of regular sets are allowed!
- ▶ General results in this direction assume f is regular. We don't assume anything about f .

Applications

- ▶ Given a continuous $\gamma : [0, 1] \rightarrow \mathbf{R}^d$, finding $X \subset [0, 1]$ such that $\gamma(x), \gamma(y), \gamma(z)$ do not form an ‘arithmetic progression’ on the curve, in the sense that

$$(\gamma(x) - \gamma(y)) - (\gamma(y) - \gamma(z)) = 0$$

Pramanik and Fraser can avoid smooth curves. We can now avoid any curve given a Hausdorff dimension bound.
Potentially even a Hilbert curve?

Applications

- ▶ Given a continuous $\gamma : [0, 1] \rightarrow \mathbf{R}^d$, finding $X \subset [0, 1]$ such that $\gamma(x), \gamma(y), \gamma(z)$ do not form an ‘arithmetic progression’ on the curve, in the sense that

$$(\gamma(x) - \gamma(y)) - (\gamma(y) - \gamma(z)) = 0$$

Pramanik and Fraser can avoid smooth curves. We can now avoid any curve given a Hausdorff dimension bound. Potentially even a Hilbert curve?

- ▶ Given a subset Y of \mathbf{R}^d with dimension α , we can find a subset X of dimension $d - \alpha$ such that $X + X$ and $X - X$ are disjoint from Y . I hope to extend this shortly with the additional criterion that the set X we construct is a rational vector space, so for instance $X + X + X$ is disjoint from Y . Then $X \oplus Y$ is a full dimensional subspace of \mathbf{R} .

Two key ideas:

Two key ideas:

- ▶ Discretization of Scales.

Two key ideas:

- ▶ Discretization of Scales.
- ▶ Random Dissection.

Discretization of Scales: Intuition

- ▶ Every number in the Cantor set has a trinary expansion which does not contain the digit 1.

Discretization of Scales: Intuition

- ▶ Every number in the Cantor set has a trinary expansion which does not contain the digit 1.
- ▶ We can view this as a 'fine scale property', since it involves the infinite digit expansion of a number.


Discretization of Scales: Intuition

- ▶ Every number in the Cantor set has a trinary expansion which does not contain the digit 1.
- ▶ We can view this as a 'fine scale property', since it involves the infinite digit expansion of a number.
- ▶ Scale discretization changes fine-scale configurations into an infinite sequence of discrete configurations which are easier to achieve.

Discretization of Scales: Intuition


- ▶ Every number in the Cantor set has a trinary expansion which does not contain the digit 1.
- ▶ We can view this as a 'fine scale property', since it involves the infinite digit expansion of a number.
- ▶ Scale discretization changes fine-scale configurations into an infinite sequence of discrete configurations which are easier to achieve.
- ▶ **Discrete Problem:** Given X_n , find X_{n+1} such that no element of X_{n+1} has 1 as the n 'th digit in its expansion.

Discretization of Scales




- We take $X = \lim X_n$, where X_{n+1} is obtained from X_n by interval dissection avoiding a discretized problem.

Discretization of Scales




- ▶ We take $X = \lim X_n$, where X_{n+1} is obtained from X_n by interval dissection avoiding a discretized problem.
- ▶ **Discrete Configuration Problem:** If $f(x_1, \dots, x_n) = 0$ for $x_1, \dots, x_n \in X_n$, then some x_i and x_j occur in a common interval.

Discretization of Scales




- ▶ We take $X = \lim X_n$, where X_{n+1} is obtained from X_n by interval dissection avoiding a discretized problem.
- ▶ **Discrete Configuration Problem:** If $f(x_1, \dots, x_n) = 0$ for $x_1, \dots, x_n \in X_n$, then some x_i and x_j occur in a common interval.
- ▶ If the interval lengths in X_n tend to zero as $n \rightarrow \infty$, and $x_1, \dots, x_n \in X$ have $f(x_1, \dots, x_n) = 0$, then there is i, j such that $|x_i - x_j| = o(1)$, so $x_i = x_j$.

Discrete Scale Problem




- For technical reasons, we only dissect subsets of X_n at a time.

Discrete Scale Problem




- ▶ For technical reasons, we only dissect subsets of X_n at a time.
- ▶ **Discrete Configuration Problem:** Given disjoint unions of intervals sets l_1, \dots, l_n , find $J_1 \subset l_1, \dots, J_n \subset l_n$ such that the cartesian product of the J_k avoids $f^{-1}(0)$.

Discrete Scale Problem




- ▶ For technical reasons, we only dissect subsets of X_n at a time.
- ▶ **Discrete Configuration Problem:** Given disjoint unions of intervals sets l_1, \dots, l_n , find $J_1 \subset l_1, \dots, J_n \subset l_n$ such that the cartesian product of the J_k avoids $f^{-1}(0)$.
- ▶ Using a queueing procedure, if the subsets we choose cover our entire set infinitely often, we still avoid the finite scale configuration. See Pramanik and Fraser (2018) for the same technique applied to their problem.

Exploiting Randomness




- How do we prove the discrete scale argument?

Exploiting Randomness




- ▶ How do we prove the discrete scale argument?
- ▶ Aside from the Hausdorff dimension of $f^{-1}(0)$, we have little knowledge of its structure.

Exploiting Randomness



- ▶ How do we prove the discrete scale argument?
- ▶ Aside from the Hausdorff dimension of $f^{-1}(0)$, we have little knowledge of its structure.
- ▶ Random choices of the J_k avoid $f^{-1}(0)$ effectively.

Exploiting Randomness



- ▶ How do we prove the discrete scale argument?
- ▶ Aside from the Hausdorff dimension of $f^{-1}(0)$, we have little knowledge of its structure.
- ▶ Random choices of the J_k avoid $f^{-1}(0)$ effectively.
- ▶ We obtain that for all but $o(1)$ of these sections, J_k contains a length $1/N^\beta$ section, where $\beta = d(nd - \alpha)/(n - 1)$. This ratio gives the Hausdorff dimension bound $(nd - \alpha)/(n - 1)$ for X .

So What's Next?

Extension to Hausdorff Dimension

- ▶ I left out something in the statement of my theorem.

Extension to Hausdorff Dimension

- ▶ I left out something in the statement of my theorem.
- ▶ Our theorem needs the assumption that $f^{-1}(0)$ is the countable union of sets with *lower* Minkowski dimension α .

Extension to Hausdorff Dimension

- ▶ I left out something in the statement of my theorem.
- ▶ Our theorem needs the assumption that $f^{-1}(0)$ is the countable union of sets with *lower* Minkowski dimension α .
- ▶ Our techniques ‘should’ work when $f^{-1}(0)$ has *Hausdorff dimension* α .

Extension to Hausdorff Dimension

- ▶ I left out something in the statement of my theorem.
- ▶ Our theorem needs the assumption that $f^{-1}(0)$ is the countable union of sets with *lower* Minkowski dimension α .
- ▶ Our techniques ‘should’ work when $f^{-1}(0)$ has *Hausdorff dimension* α .
- ▶ We are trying to use hyperdyadic coverings to obtain this.

Analogies with Hypergraphs

- ▶ Partition I_1, \dots, I_n into length $1/N^\beta$ sections. Form a hypergraph whose vertices are the sections, and add a hyperedge between $K_1 \subset I_1, \dots, K_n \subset I_n$ if $K_1 \times \dots \times K_n$ intersects $f^{-1}(0)$.

Analogies with Hypergraphs

- ▶ Partition I_1, \dots, I_n into length $1/N^\beta$ sections. Form a hypergraph whose vertices are the sections, and add a hyperedge between $K_1 \subset I_1, \dots, K_n \subset I_n$ if $K_1 \times \dots \times K_n$ intersects $f^{-1}(0)$.
- ▶ Our discrete configuration problem reduces to finding independent sets in hypergraphs.

Analogies with Hypergraphs

- ▶ Partition I_1, \dots, I_n into length $1/N^\beta$ sections. Form a hypergraph whose vertices are the sections, and add a hyperedge between $K_1 \subset I_1, \dots, K_n \subset I_n$ if $K_1 \times \dots \times K_n$ intersects $f^{-1}(0)$.
- ▶ Our discrete configuration problem reduces to finding independent sets in hypergraphs.
- ▶ The method we use essentially generalizes Turan's theorem to hypergraphs, and is tight?

Analogies with Hypergraphs

- ▶ Partition I_1, \dots, I_n into length $1/N^\beta$ sections. Form a hypergraph whose vertices are the sections, and add a hyperedge between $K_1 \subset I_1, \dots, K_n \subset I_n$ if $K_1 \times \dots \times K_n$ intersects $f^{-1}(0)$.
- ▶ Our discrete configuration problem reduces to finding independent sets in hypergraphs.
- ▶ The method we use essentially generalizes Turan's theorem to hypergraphs, and is tight?
- ▶ We are looking to using other methods on hypergraphs to improve the bound when $f^{-1}(0)$ has certain structural properties.

You're Interested in Learning More?

Read these to find out more about configuration problems:

- ▶ Read Keleti (1999) for a simple use of scale discretization on a particular configuration avoidance problem.

You're Interested in Learning More?

Read these to find out more about configuration problems:

- ▶ Read Keleti (1999) for a simple use of scale discretization on a particular configuration avoidance problem.
- ▶ Read Máthé (2017) for a construction where f is assumed to be a polynomial of bounded degree.

You're Interested in Learning More?

Read these to find out more about configuration problems:

- ▶ Read Keleti (1999) for a simple use of scale discretization on a particular configuration avoidance problem.
- ▶ Read Máthé (2017) for a construction where f is assumed to be a polynomial of bounded degree.
- ▶ Read Pramanik and Fraser (2018) for a general construction where f is smooth and nonsingular. We generalize their work.

You're Interested in Learning More?

Read these to find out more about configuration problems:

- ▶ Read Keleti (1999) for a simple use of scale discretization on a particular configuration avoidance problem.
- ▶ Read Máthé (2017) for a construction where f is assumed to be a polynomial of bounded degree.
- ▶ Read Pramanik and Fraser (2018) for a general construction where f is smooth and nonsingular. We generalize their work.

Thanks for listening!