

Complex Analysis

Jacob Denson

September 13, 2018

Table Of Contents

1	Introduction	2
1.1	The Complex Numbers	2
1.2	Geometry of the Complex Plane	4
1.3	Complex Functions and Holomorphicity	5
1.4	Power Series	7
1.5	Integration Along Curves	11
1.6	Harmonic Functions	15
2	Cauchy's Theorem	17
2.1	Consequences of Cauchy's Formula	20
2.2	Uniqueness of Holomorphic Functions	22
2.3	Meromorphic Functions and Residues	23
2.4	Sequences of Holomorphic Functions	23
2.5	The Reflection Principle	25
3	Geometric Function Theory	26
4	Riemann Surfaces	29
4.1	Derivations	32

Chapter 1

Introduction

The study of complex numbers is the analysis of a strange mathematical creation. Originally, these numbers were used as an ‘imaginary’ mathematical creation used to find solutions to polynomial equations. However, in the early 19th century these numbers began being viewed as a geometric construction, another way to describe the two dimensional plane \mathbf{R}^2 . The introduction of this number system on top of the plane has a much higher structure. And function differentiable with respect to this coordinate system has drastic differences to the study of complex numbers. It offers results many would described as miraculous and magical. What’s more, the theory of complex analysis introduces many concepts which have formed the building blocks of many 20th century mathematical fields in topology, geometry, algebra, and analysis. In this first section, we introduce the complex numbers formally, as well as discussing the fundamental tools we will use in the study of complex analysis.

1.1 The Complex Numbers

As a numerical calculation system, the real numbers have many advantages: they are complete, enabling us to use Newton’s calculus with precision, as well as forming an ordered system of numbers, giving us a standard tool for comparing magnitudes which occur throughout the sciences. Nevertheless, the number system has one deficiency: there are some mathematical equations we are unable to solve. Since the square of every real number is positive, we cannot find a number x which satisfies the equa-

tion $x^2 = -1$. The complex number system \mathbf{C} is obtained by ‘imagining’ a new number i , which does satisfy the equation $i^2 = -1$, and building a new number system containing i , and all real numbers, in such a way that most of the useful mathematical laws which hold for the real numbers continue to hold for all numbers in \mathbf{C} .

To be precise, we introduce the language of modern algebra. The algebraic rules in \mathbf{R} we tend to use the most often are those which make \mathbf{R} into a *field*. We define \mathbf{C} to be the smallest set of numbers containing \mathbf{R} , as well as containing a new i satisfying the equation $i^2 = -1$, while also forming a field. Since we must be able to add and multiply arbitrary numbers in \mathbf{C} , we must have numbers of the form $a + ib$, for each $a, b \in \mathbf{R}$, because we must be able to multiply each real number by i , and to add the result to a real number. Using the commutative and distributive law, we find that

$$(a + ib) + (x + iy) = (a + x) + i(b + y)$$

$$(a + ib)(x + iy) = (ax - by) + i(bx + ay)$$

so the operations of addition and multiplication are closed under these operations. What’s more, every nonzero number of the form $z = x + iy$ also has an inverse of this form, because if we define the **complex conjugate** $\bar{z} = x - iy$, we find that $z\bar{z} = x^2 + y^2$, which is a nonzero, positive real number if either x or y is nonzero, and then

$$z \frac{\bar{z}}{x^2 + y^2} = z \left[\frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2} \right] = 1$$

so the inverse of a complex number z is $\bar{z}(x^2 + y^2)^{-1}$. This means the set of numbers of the form $a + ib$ is precisely the *smallest* field containing \mathbf{R} and i . Thus every complex number is formally expandable in this form. This expansion must be unique, because if $x + iy = a + ib$, then $(x - a) + i(y - b) = 0$, and if $y \neq b$, we conclude that $i = (a - x)/(y - b)$ is a real number, which is impossible, so we must have $y = b$, and then $x = a$ follows automatically. Thus a complex number $z = a + ib$ is uniquely defined by its real and complex parts $\Re(z) = a$ and $\Im(z) = b$. Summing up these results using the language of field theory, we say that \mathbf{C} is the field extension of degree two obtained from \mathbf{R} by solving the polynomial equation $x^2 = -1$. We also note that since the standard laws of arithmetic apply to the complex numbers, we find that the equations

$$(a + ib) + (x + iy) = (a + x) + i(b + y)$$

$$(a + ib)(x + iy) = ax + (ay + bx)i + (by)i^2 = (ax - by) + (ay + bx)i$$

can be used to formally define the complex numbers; We don't need to 'assume' the existence of \mathbf{C} as an axiom. We can instead construct it by placing certain arithmetical operations on \mathbf{R}^2 , using the definitions above.

Of course, certain properties of the real numbers are lost when we extend to the complex plane. Most importantly, we lose the property of being an ordered field, which was used to define the topological structure of the real line. We cannot possibly give \mathbf{C} an ordered field structure, because in any ordered field we have $x^2 \geq 0$ for all numbers x . Nonetheless, in the next section we will show that we can use another algebraic construct to give \mathbf{C} a metric structure by viewing the complex numbers as a 'two dimensional' field, whereas the real line is one dimensional.

1.2 Geometry of the Complex Plane

Geometrically, the complex plane can be identified with \mathbf{R}^2 . This leads to useful insight into the theorems of complex analysis. First off, it gives a topological structure to the class of complex numbers, which when viewed geometrically is called the *complex plane*. The x and y axis of the plane are called the **real axis** and **imaginary axis**. A novel result of the introductions of arithmetical operations to \mathbf{R}^2 is that the norm has a pleasant form. If we denote the norm of a complex number z by $|z|$, then we have seen that $|z|^2 = x^2 + y^2 = z\bar{z}$, which has a pleasant complex algebraic form. We call $|z|$ the **modulus** of a complex number z . Since

$$|zw|^2 = zw\overline{zw} = z\bar{z}w\bar{w} = |z|^2|w|^2$$

we find that $|zw| = |z||w|$ and $|\bar{z}| = |z|$. The modulus gives a complete metric space structure to the complex plane. Just as for the plane, we have the useful inequalities

$$|z + w| \leq |z| + |w| \quad |\operatorname{Re}(z)|, |\operatorname{Im}(z)| \leq |z| \quad ||z| - |w|| \leq |z - w|$$

which follow from their real analytical theorems about \mathbf{R}^2 , once converted back to the appropriate language.

Geometrically, the arithmetical operations of the complex numbers become incredibly beautiful when we look at the numbers in their **polar form**. Recall that any point on the plane \mathbf{R}^2 can be expressed in terms

of an angle θ and a radius r . Looking ahead, we note that the complex exponential e^{it} satisfies Euler's formula $e^{it} = \cos t + i \sin t$ (we will prove this rigorously later), so that in polar coordinates, the number with coordinates r and θ is exactly $re^{i\theta}$. Because we have the addition formula $e^{i(x+y)} = e^{ix}e^{iy}$, we find that $re^{it}r'e^{it'} = (rr')e^{i(t+t')}$, so that multiplying two complex numbers dilates their radius multiplicatively, and rotates their angular coordinate additively. This means that for any $w \neq 0$, the function $z \mapsto wz$ is a rotation and dilation in two dimensions. In particular, this justifies the utility of the abstractly defined complex numbers; it has a purely geometric definition, which explains why complex analysis leads to such majestic geometrical results, and explains why complex numbers occur in the studies of rotation in physics.

1.3 Complex Functions and Holomorphicity

The really interesting part of complex analysis begins when we start looking at complex-valued functions defined on subsets of the complex plane. Classically, we write such a function in the form $w = f(z)$, referring to the domain as the 'z-plane', and the codomain as the 'w-plane'. We will mostly be analyzing maps defined on **domains**, that is, open, connected subsets of the plane. Unless otherwise specified, we will assume that all the domains of the functions we consider are open and connected.

The standard definition of differentiability of real-valued functions generalizes easily to complex-valued functions, but the definition hides the fact that complex differentiability is a much stronger property than standard differentiability. A complex-valued function f defined in a neighbourhood of a point z is **complex differentiable**, **holomorphic**, or **regular** at z if the limit

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$$

exists. Unlike differentiability on the real plane, we note that complex differentiability is a much stronger condition on the local behaviour of f around z . Whereas the real-valued limit can only be approached by two directions, complex differentiability is a limit approached from any approach trajectory. The set of all holomorphic functions defined on a domain D is denoted $C^\omega(D)$. If f is complex differentiable at z , then the limit

implies exactly that we can write $f(z+w) = f(z) + wf'(z) + o(w)$. In terms of complex multiplication, that means that f is locally well approximated by a translation, combined with a rotation and a dilation. We leave it to verify that the addition, multiplication, quotient, and chain rule continue to hold from normal calculus, and the proofs are algebraically the same.

Example. Every polynomial $f(z) = a_0 + a_1z + \cdots + a_nz^n$ is differentiable at any point in the complex plane, with derivative $f'(z) = a_1 + 2a_2z + \cdots + na_nz^{n-1}$. It is easy to see that the derivative of $w = z$ is $w' = 1$, and then the addition and product rule verifies the rule for all polynomials.

Example. The function $w = 1/z$ is differentiable on $\mathbf{C} - \{0\}$, because

$$\lim_{h \rightarrow 0} \frac{(z+h)^{-1} - z^{-1}}{h} = \lim_{h \rightarrow 0} -\frac{1}{z(z+h)} = -\frac{1}{z^2}$$

Using the product rule, this implies that for $w = z^{-n}$, $w' = -nz^{-(n+1)}$.

Example. The function $f(z) = \bar{z}$, though just a simple reflection in the complex plane, is not complex differentiable anywhere. This is because

$$\lim_{h \rightarrow 0} \frac{\overline{z+h} - \bar{z}}{h} = \lim_{h \rightarrow 0} \frac{\bar{h}}{h}$$

And this limit does not exist, because if we approach the limit from the imaginary axis, with $h = ti$, then $\bar{h}/h = -1$, and if we approach the limit from the real axis, with $h = t$, then $\bar{h}/h = 1$. A reflection is not locally a rotation and a dilation.

We previously noted that complex multiplication corresponds to a rotation and a dilation. In particular, for each w , the map $T_w(z) = wz$ is a linear map from \mathbf{C} to itself. Viewing \mathbf{C} as \mathbf{R}^2 , we see it has a matrix representation

$$T_w = \begin{pmatrix} w_1 & w_2 \\ -w_2 & w_1 \end{pmatrix}$$

The function $f(z) = \bar{z}$, viewed as a function from \mathbf{R}^2 to \mathbf{R}^2 , is certainly differentiable, and in fact, C^∞ . It is linear, with matrix representation

$$\begin{pmatrix} +1 & 0 \\ 0 & -1 \end{pmatrix}$$

This cannot be put in the form T_w for any w , which reflects the nonexistence of a complex derivative. Not all linear endomorphisms of the plane are homotheties. However, there are relations which guarantee a linear map to be a rotation and a dilation. In fact, the matrix representation above shows that a function $f = u + iv$ is complex differentiable at a point if and only if it is real differentiable, and

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

These are known as the **Cauchy-Riemann equations**. As an alternate viewpoint, we can also approach this purely from the complex perspective, something we will find often gives the most elegant arguments as we go deeper into complex analysis. If f is complex differentiable at z , then

$$f(z + w) = f(z) + f'(z)w + o(w)$$

so

$$\begin{aligned} \frac{\partial f}{\partial x} &= \lim_{t \rightarrow 0} \frac{f(z + t) - f(z)}{t} = \lim_{t \rightarrow 0} \frac{f'(z)t + o(t)}{t} = f'(z) \\ \frac{\partial f}{\partial y} &= \lim_{t \rightarrow 0} \frac{f(z + it) - f(z)}{t} = \lim_{t \rightarrow 0} \frac{itf'(z) + o(it)}{t} = if'(z) \end{aligned}$$

If f is complex differentiable, then we find

$$\frac{\partial f}{\partial y} = i \frac{\partial f}{\partial x}$$

This is precisely the Cauchy Riemann equations in complex form.

1.4 Power Series

Polynomials are the simplest examples of holomorphic functions. Our next family of holomorphic functions will be obtained by taking limits of polynomials. In fact, we will soon show that this family essentially contains all holomorphic functions. A **formal power series** about a point z_0 is an ‘infinite dimensional polynomial’

$$\sum_{n=0}^{\infty} c_n (z - z_0)^n$$

We may evaluate a power series to obtain a function f , defined as

$$f(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k$$

whose domain is the set where the series converges. A function f can be **expanded in a power series** at a point z_0 if we may write f as a power series locally around z_0 . An **analytic** function is a function which can be locally expanded as a power series at every point in its domain.

Standard tricks from real power series come into play here, and the proofs go unedited. The most useful is the root test, or Cauchy-Hadamard theorem, which states that, if we define

$$1/R = \limsup_{n \rightarrow \infty} |c_n|^{1/n}$$

Then the power series f converges locally uniformly for $|z - z_0| < R$, and diverges for $|z - z_0| > R$. It may sometimes be easier to apply the ratio test. If the limit

$$1/L = \lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right|$$

exists, then f converges for $|z - z_0| < L$, and diverges for $|z - z_0| > L$. These statements do not say anything about what happens on the boundary, when $|z - z_0|$ is equal to R or L .

Example. *The most important power series, and perhaps the most important function in mathematics, is the exponential function, defined at all points in the complex plane by the formula*

$$e^z = \exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

Which converges by the ratio test, since $(n+1)!/n! = n \rightarrow \infty$. The product formula shows

$$\begin{aligned} \exp(z+w) &= \sum_{k=0}^{\infty} \frac{(z+w)^k}{k!} = \sum_{k=0}^{\infty} \sum_{j=0}^k \binom{k}{j} \frac{z^j w^{k-j}}{k!} \\ &= \sum_{k=0}^{\infty} \sum_{j=0}^k \frac{z^j}{j!} \frac{w^{k-j}}{(k-j)!} = \exp(z) \exp(w) \end{aligned}$$

which gives us the multiplicative property of the function.

Example. The hypergeometric series is defined for $\alpha, \beta \in \mathbf{C}$, and γ not a negative integer by

$$\begin{aligned} F(\alpha, \beta, \gamma; z) &= \sum_{n=1}^{\infty} \frac{\alpha(\alpha+1)\dots(\alpha+n-1)\beta(\beta+1)\dots(\beta+n-1)}{\gamma(\gamma+1)\dots(\gamma+n-1)} \frac{z^n}{n!} \\ &= \sum_{n=1}^{\infty} \frac{(\alpha)_n(\beta)_n}{(\gamma)_n} \frac{z^n}{n!} \end{aligned}$$

The notation $(\cdot)_n$ is known as a Pockhammer symbol. We calculate

$$\begin{aligned} \log \left(\prod_{m=0}^{N-1} (\lambda + m) \right) &= \sum_{m=0}^{N-1} \log(\lambda + m) = \int_{\lambda}^{\lambda+N} \log x \, dx + O(\log(\lambda + N)) \\ &= N \log(\lambda + N) + O(\log(\lambda + N)) \\ &= N \log(N) + O(\log(\lambda + N)) \end{aligned}$$

In particular, $\log(N!) = N \log N + O(\log N)$. This means

$$\log \left(\left(\frac{(\alpha)_n(\beta)_n}{(\lambda)_n} \frac{1}{n!} \right)^{1/n} \right) = \frac{O(\log(\alpha + \beta + \lambda + n))}{n} = o(1)$$

so

$$\left(\frac{(\alpha)_n(\beta)_n}{(\lambda)_n} \frac{1}{n!} \right)^{1/n} \rightarrow 1$$

so the series has a radius of convergence of one.

Example. If we define the Bessel function

$$J_m(z) = (z/2)^m \sum_{n=0}^{\infty} \frac{(-1)^n}{n!(n+m)!}$$

Then we calculate

$$\begin{aligned} \log \left(\left(\frac{1}{2^{2n} n! (n+m)!} \right)^{1/2n} \right) &= \frac{-2n \log 2 - 2n \log n + O(\log(n+m))}{2n} \\ &= -\log 2 - \log n + o(1) \rightarrow -\infty \end{aligned}$$

so J_m is defined by a power series everywhere.

Calculus tells us that the derivative of the exponential function is the exponential function itself. If there is any justice in the world, this idea should extend to the complex case. We can formally calculate

$$\exp'(z) = \sum_{n=0}^{\infty} \left(\frac{z^n}{n!} \right)' = \sum_{n=1}^{\infty} \frac{z^{n-1}}{(n-1)!} = \sum_{n=0}^{\infty} \frac{z^n}{n!} = \exp(z)$$

But how do we know that the derivative of the power series can be obtained by differentiating each term in the series?

Theorem 1.1. *Let*

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

be a power series, convergent inside a disk D . Then f is holomorphic in D , and

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

Proof. The fact that the power series defining f' converges in D follows from the Hadamard formula. Now let us verify that this series, which defines a function g , approaches the derivative of f . Define S_n to be the finite polynomial of degree n defined by f , $S_n(z) = \sum_{k=0}^n a_k z^k$. Let E_n denote the error term, $E_n(z) = \sum_{k=n+1}^{\infty} a_k z^k$. Then $S'_n(z) = \sum_{k=1}^n k a_k z^{k-1}$, and

$$\frac{f(z+w) - f(z)}{w} = \left[\frac{S_n(z+w) - S_n(z)}{w} - S'_n(z) \right] + \left[\frac{E_n(z+w) - E_n(z)}{w} \right] + S'_n(z)$$

The first term converges to zero for small enough w . The third term converges to $g(z)$ as $n \rightarrow \infty$. The only tricky component is the second term. Since $a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + \dots + ab^{n-2} + b^{n-1})$. If we choose $|z|, |z+w| < r - \varepsilon$, then

$$\left| \frac{E_n(z+w) - E_n(z)}{w} \right| = \left| \sum_{k=n+1}^{\infty} a_k \frac{(z+w)^k - z^k}{w} \right| \leq \sum_{k=n+1}^{\infty} k |a_k| (r - \varepsilon)^{n-1}$$

The right side is a convergent power series, and hence as $n \rightarrow \infty$, the term converges to zero. By picking n large enough, and $|z - w|$ small enough, the value can be made as close to g as possible. \square

Corollary 1.2. *The power series f is infinitely differentiable in D , with*

$$f^{(m)}(z) = \sum_{n=m}^{\infty} \frac{n!}{(n-m)!} a_n z^{n-m}$$

Proof. All that remains to be proven is that f' has the same radius of convergence as f , so that we may successively apply the theorem. If f has radius of converge R , then $R^{-1} = \limsup |a_n|^{1/n}$, and so

$$\limsup |a_n|^{1/n} n^{1/n} = \lim n^{1/n} \limsup |a_n|^{1/n} = \limsup |a_n|^{1/n}$$

and so f' has the same radius of convergence as f . \square

Corollary 1.3. *Every analytic function is holomorphic throughout it's domain.*

1.5 Integration Along Curves

One distinguishing feature separating the study of complex analysis from Newton's one dimensional calculus is the reliance on line integration, which you may have encountered in multivariate calculus. Firstly, we will find ourselves not integrating over the whole complex plane, but instead integrating over curves lying in the complex plane. A **smooth curve** is an immersion of a closed interval into \mathbf{C} (viewing the interval as an oriented one dimensional manifold with boundary, and \mathbf{C} as a two dimensional manifold). More simply, we define a **smooth parameterization** $z : [a, b] \rightarrow \mathbf{C}$ to be a differentiable map whose derivative is continuous, and $z'(t) \neq 0$ for all $t \in [a, b]$, where we interpret $z'(a)$ and $z'(b)$ as the left and right hand limits

$$\lim_{h \rightarrow 0^+} \frac{z(a+h) - z(a)}{h} \quad \lim_{h \rightarrow 0^-} \frac{z(b+h) - z(b)}{h}$$

Two smooth curves $z_0 : [a_0, b_0] \rightarrow \mathbf{C}$ and $z_1 : [a_1, b_1] \rightarrow \mathbf{C}$ are **equivalent** if there is a continuously differentiable function $s : [a_0, b_0] \rightarrow [a_1, b_1]$ with $s'(t) > 0$ for all t , such that $z_0(t) = z_1(s(t))$ for all $t \in [a_0, b_0]$. We then define a smooth curve to be an equivalence class of smooth parameterizations. More generally, a **piecewise smooth curve** is an equivalence class of continuous parameterizations $z : [a, b] \rightarrow \mathbf{C}$, with values $a = a_0 < a_1 < \dots < a_n = b$ such that z is a smooth curve on $[a_i, a_{i+1}]$, where $z_0 : [a_0, b_0] \rightarrow \mathbf{C}$ and $z_1 : [c_0, d_0] \rightarrow \mathbf{C}$ are equivalent if there are $a = a_0 < \dots < a_n = b$ and

$c = c_0 < \cdots < b_n = d$ such that z_0 and z_1 are equivalent on each of the intervals $[a_i, a_{i+1}]$ and $[c_i, c_{i+1}]$. We rarely need to discuss classes of curves more general than this in complex analysis, and certainly not in elementary complex analysis.

Example. The standard example of a curve is the circle $C_r(z_0)$, of radius r centered at z_0 . The positive orientation of this curve is given by the parameterization

$$z(t) = z_0 + re^{it}$$

On $[0, 2\pi]$, and the negative orientation

$$z(t) = z_0 + re^{-it}$$

also defined on $[0, 2\pi]$. In general, we will interpret $C_r(z_0)$ as having the positive orientation. In particular, we denote $C_1(0)$ by S^1 , the unit circle.

We will also need to introduce slightly more terminology to manipulate curves. The **trace** of a curve C is the point set corresponding to the curve. It is the image of all the parameterizations of the curve. The **inverse orientation** of a curve with parameterization $z : [a, b] \rightarrow \mathbf{C}$, which is the curve with parameterization $z^- : [a, b] \rightarrow \mathbf{C}$ with $z^-(t) = z(a + b - t)$. A curve $z : [a, b] \rightarrow \mathbf{C}$ is **closed** if $z(a) = z(b)$, which means its image forms a ‘closed loop’ in the plane. Now given a smooth curve C with parameterization $z : [a, b] \rightarrow \mathbf{C}$, and a continuous function defined on the trace of C , we define the **line integral** of f on C as

$$\int_C f(z)dz = \int_a^b f(z(t))z'(t)dt$$

which we can remember by the mnemonic

$$f(z)dz = f(z) \left(\frac{dz}{dt} \right) dt$$

so we can obtain the formula by ‘multiplying by a dt/dt factor’. If z' is only piecewise smooth, with partition $a_0 < \cdots < a_n$, then we define

$$\int_C f(z)dz = \sum \int_{a_i}^{a_{i+1}} f(z(t))z'(t)dt$$

The change of variables formula for one dimensional integrals verifies that these definitions are independent of parameterization.

Theorem 1.4. *The line integral satisfies the following properties*

- For a given curve C on some domain D , for any two $f, g \in C(D)$, $\alpha, \beta \in \mathbf{C}$,

$$\int_C (\alpha f + \beta g)(z) dz = \alpha \int_C f(z) dz + \beta \int_C g(z) dz$$

- If C^- is the reverse orientation of C , then

$$\int_{C^-} f(z) dz = - \int_C f(z) dz$$

- One has

$$\left| \int_C f(z) dz \right| \leq l(C) \|f\|_\infty$$

where $\|f\|_\infty$ is the supremum of f over image of the curve C , and $l(C)$ is the **length** of the curve C , defined to be the integral $\int_a^b |z'(t)| dt$.

Proof. The first theorem follows by the linearity of one dimensional integration by expanding out formulas, and the second theorem follows because $(z^-)'(t) = -z'(a + b - t)$. Finally, the third theorem follows because if $z : [a, b] \rightarrow \mathbf{C}$ is smooth, then

$$\left| \int_C f(z) dz \right| = \left| \int_a^b f(z(t)) z'(t) dt \right| \leq \int_a^b |f(z(t))| |z'(t)| dt \leq \|f\|_\infty \int_a^b |z'(t)| dt$$

and the result for piecewise smooth curves follows by linearity. \square

Finally in our review of line integration, we recall a very useful result. We say a function $f : D \rightarrow \mathbf{C}$ on some domain D has a **primitive** if there is a function $F : D \rightarrow \mathbf{C}$ holomorphic on D such that $F'(z) = f(z)$.

Theorem 1.5. *If f is continuous, and has a primitive F , then a curve C beginning at z_0 and ending at z_1 has*

$$\int_C f(z) dz = F(z_1) - F(z_0)$$

In particular, if C is a closed curve, then $\int_C f(z) dz = 0$.

Proof. If C is smooth, then

$$\int_C f(z) dz = \int_a^b f(z(t)) z'(t) dt$$

Now if $F'(z) = f(z)$ for each $z \in D$, then the chain rule (which is easily proved for compositions of functions from intervals to \mathbf{C} and from holomorphic functions on \mathbf{C}) implies that $(F \circ z)'(t) = F'(z(t)) z'(t)$, and therefore the fundamental theorem of calculus (applied componentwise on the integral of the complex valued function $f(z(t)) z'(t)$), we conclude that

$$\int_a^b f(z(t)) z'(t) dt = F(z(b)) - F(z(a)) = F(w) - F(z)$$

The theorem is then proved for piecewise smooth curves by applying this theorem for the individual smooth curves and then applying a telescoping summation. \square

Corollary 1.6. *If f is holomorphic, with $f' = 0$, then f is constant on D .*

Proof. If z_0 and z_1 are two points, let C be a smooth curve beginning at z_0 and ending at z_1 . Then since $f' = 0$,

$$\int_C f'(z) dz = 0$$

But also since f' has a primitive f , we find that

$$\int_C f'(z) dz = f(z_1) - f(z_0)$$

and hence $f(z_0) = f(z_1)$. \square

Example. *The function $f(z) = 1/z$ does not have a primitive in $\mathbf{C} - \{0\}$, where it is defined, because*

$$\int_{S^1} \frac{dz}{z} = \int_0^{2\pi} \frac{ie^{it}}{e^{it}} = \int_0^{2\pi} i = 2\pi i$$

If $1/z$ did have a primitive, we would conclude this integral is zero. We shall find that this is the essential holomorphic function without a primitive, and we will use this theorem to calculate a great many contour integrals.

We know that if f has a primitive F , then $\int_{\gamma} f(z) dz = 0$ for all closed curves γ . But conversely, if $\int_{\gamma} f(z) dz = 0$ for all closed curves γ , then such a primitive exists. The idea is to fix a point z_0 in the domain, and to define $F(z_1) = \int_{\gamma} f(z) dz$, where γ is a curve from z_0 to z_1 . We find that

$$\frac{\partial F}{\partial x} = f(z) \quad \frac{\partial F}{\partial y} = if(z)$$

follows from the fundamental theorem of calculus. But this implies that F is holomorphic by the Cauchy Riemann equations, and the continuity of f , and $F' = f$.

1.6 Harmonic Functions

A function $f : \mathbf{R}^2 \rightarrow \mathbf{C}$ is known as harmonic if

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$$

We now discuss a connection which allows us to reduce their study to the study of holomorphic functions. We calculate

$$\partial_z \partial_{\bar{z}} = \frac{(\partial_x - i\partial_y)(\partial_x + i\partial_y)}{4} = \frac{\partial_x^2 + i\partial_x \partial_y - i\partial_y \partial_x + \partial_y^2}{4} = \frac{\Delta}{4}$$

So any holomorphic or antiholomorphic function is harmonic, as is its real and imaginary part. Conversely, suppose that g is a real-valued harmonic function. Then $\partial_{\bar{z}} \partial_z g = 0$, so $\partial_z g$ is holomorphic. We will later find that any holomorphic locally has an antiderivative, so in particular, we can find a holomorphic function f with $f' = 2\partial_z g$. But this means that

$$\partial_x f_0 - i\partial_y f_0 = f' = 2\partial_z g = \partial_x g - i\partial_y g$$

so $\nabla f_0 = \nabla g$, implying f_0 and g differ by a constant. Patching the f_0 together shows that g is the real part of a holomorphic function.

In particular, this implies that for any circle $C_r(z_0)$,

$$2\pi i f(z_0) = \int_{C_r(z_0)} \frac{f(z)}{z - z_0} dz = i \int_0^{2\pi} f(z_0 + re^{it}) dt$$

Taking the imaginary parts on both sides gives

$$2\pi g(z_0) = \int_0^{2\pi} f(z_0 + e^{it}) dt$$

So any harmonic function satisfies the mean value formula; the average value of the function around a circle is the same as the value of that function at the centre of the circle. Note that this is not necessarily true for the average of the function over every curve, since in the calculations above for curves other than circles the imaginary parts of f might contribute to the imaginary part of the integral.

Chapter 2

Cauchy's Theorem

Cauchy's theorem says that the integral of a holomorphic function about a closed curve is always zero. This is simple, but forms the backbone for the theory of holomorphic functions. Originally, certain regularity were assumed on holomorphic functions to obtain Cauchy's theorem. But it was later discovered that one can obtain Cauchy's theorem assuming only that the function is holomorphic at each point in the region, and it is this procedure we will use to show the regularity of holomorphic functions completely.

Theorem 2.1 (Goursat's lemma). *If T is a triangle, and f is holomorphic in an open set containing the triangle and its interior, then*

$$\int_T f(z) dz = 0$$

Proof. We can split the triangle T into four smaller triangles A , B , C , and D , each with half the diameter and length. Since

$$\int_T f(z) dz = \left(\int_A f(z) dz + \int_B f(z) dz + \int_C f(z) dz + \int_D f(z) dz \right)$$

We may choose one such triangle T_1 with

$$\left| \int_{T_1} f(z) dz \right| \geq \left| \int_T f(z) dz \right|$$

Continuing inductively, we can find a nested family of rectangles $T_1 \supset T_2 \supset \dots T_n$ with $\text{diam}(T_n) = \text{diam}(T)/2^n$, $l(T_n) = l(T)/2^n$, and

$$\left| \int_{T_n} f(z) dz \right| \geq 4^{-n} \left| \int_T f(z) dz \right|$$

As $n \rightarrow \infty$, $\text{diam}(T_n) \rightarrow 0$, so $\bigcap T_n$ contains a unique point z_0 . We can write $f(z) = f(z_0) + f'(z_0)(z - z_0) + \psi(z - z_0)(z - z_0)$, with $\psi(w) \rightarrow 0$ as $w \rightarrow 0$. Now it is easy to calculate that

$$\int_{T_n} f(z_0) dz = f'(z_0) \int_{T_n} (z - z_0) dz = 0$$

so if $|\psi(z - z_0)| \leq \varepsilon$ for z lying on T_n ,

$$\begin{aligned} \left| \int_T f(z) dz \right| &\leq 4^n \left| \int_{T_n} f(z) dz \right| = 4^n \left| \int_{T_n} \psi(z - z_0)(z - z_0) dz \right| \\ &\leq 4^n \varepsilon l(T_n) \text{diam}(T_n) = \varepsilon l(T) \text{diam}(T) \end{aligned}$$

and thus as $n \rightarrow \infty$, we can take $\varepsilon \rightarrow 0$ to conclude the integral over the triangle is zero. \square

Remark. Note that the proof of this theorem remains true if we replace the condition that f is holomorphic in D with the weaker condition that f is analytic except at finitely many points z_1, \dots, z_N in the interior of the triangle, with the property that $\lim_{z \rightarrow z_n} (z - z_n)f(z) = 0$ for each n . By splitting our triangles into smaller triangles, we may assume that there is only a single point with this property in the triangle. By splitting our triangle into smaller triangles, and applying the theorem above in all regions but the region containing the triangle, we determine that the integral over the entire region is equal to the integral of f about an equilateral triangle T' centered at z , of arbitrarily small length. But if T' is small enough that $|z - z_n||f(z)| \leq \varepsilon$ for z on T' , then

$$\left| \int_{T'} f(z) dz \right| \leq \varepsilon \left| \int_{T'} \frac{dz}{z - z_n} \right|$$

and one verifies that

$$\left| \int_{T'} \frac{dz}{z - z_n} \right| \lesssim 1$$

since the distance from points on the triangle to z_n is proportional to the perimeter of the triangle. This concludes the argument.

Attaching two triangles together, we can obtain Cauchy's theorem for a rectangle. Given a holomorphic function f on a circle, by defining $F(w)$ to be the line integral from the centre of the circle to w on either half of the rectangle parallel to the axis, we find a function F with $F' = f$, so f has a primitive, and so Cauchy's theorem holds in a circle. More generally, this holds under the weaker condition where f is defined except at finitely many interior points z_1, \dots, z_N with $\lim(z - z_n)f(z) = 0$, because we can with a few edge cases not included in the previous case find a function F with $F' = f$.

Lemma 2.2. *If γ is a closed curve not passing through z_0 , then*

$$\eta(\gamma, z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - z_0}$$

*is an integer, known as the **winding number** of γ around z_0 .*

Proof. If we parameterize γ by z , then consider the continuous function

$$h(t) = \int_a^t \frac{z'(t)}{z(t) - z_0} dt$$

And we have $h'(t) = z'(t)/(z(t) - z_0)$. Thus the derivative of $g(t) = e^{-h(t)}(z(t) - z_0)$ vanishes, so $g(t)$ must be constant, so

$$e^{h(t)} = \frac{z(t) - z_0}{z(a) - z_0}$$

In particular,

$$e^{h(b)} = \frac{z(b) - z_0}{z(a) - z_0} = 1$$

so $h(b)$ must be a multiple of $2\pi i$, completing the proof. \square

Cauchy's theorem implies that if γ is contained in a circle, and z_0 is a point lying outside of the circle, then $\eta(\gamma, z_0) = 0$. Furthermore, if there is a curve between two points z_0 and z_1 not passing through γ , then $\eta(\gamma, z_0) = \eta(\gamma, z_1)$. Since we can replace the curve with a polygon, it suffices to prove this assuming the curve is a line segment. Outside of the segment the function $(z - z_0)(z - z_1)^{-1}$ is never a negative number, so

$\log((z - z_0)/(z - z_1))$ is analytic in the complement of the segment, and its derivative is $(z - z_0)^{-1} - (z - z_1)^{-1}$, hence Cauchy's theorem implies that

$$\int_{\gamma} \frac{dz}{z - z_0} = \int_{\gamma} \frac{dz}{z - z_1}$$

hence $\eta(\gamma, z_0) = \eta(\gamma, z_1)$.

Theorem 2.3. *If f is holomorphic in a disc D containing a point w , and γ is a closed curve in D not passing through w , then*

$$\int_{\gamma} \frac{f(z)}{z - w} = 2\pi i \eta(\gamma, w) f(w)$$

Proof. Since $g(z) = (f(z) - f(w))/(z - w)$ is holomorphic everywhere in D except at w , and $\lim_{z \rightarrow w} (z - w)g(z) = 0$, we can apply our remark to conclude that

$$\int_{\gamma} \frac{f(z)}{z - w} dz = \int_{\gamma} \frac{f(w)}{z - w} dz = 2\pi i \eta(\gamma, w) f(w)$$

Thus the theorem becomes an explicit calculation. □

Remark. Integral representation formulas are very important in analysis, because they enable us to control a function from its behaviour on a very small set, and also use it to show the function is suitably regular.

2.1 Consequences of Cauchy's Formula

A simple corollary of Cauchy's integral formula, proved by induction, shows any holomorphic function is completely smooth.

Theorem 2.4. *If f is holomorphic, then f has derivatives of all orders, with*

$$f^{(n)}(w) = \frac{n!}{2\pi i \eta(\gamma, w)} \int_{\gamma} \frac{f(z)}{(z - w)^{n+1}} dz$$

where γ is some closed curve with nonzero winding number about w .

Proof. To prove this, it would suffice to reduce the question to the theory of differentiation under the integral sign in real variable theory. But there is a more elementary proof. We know this is true for $n = 0$. And now by induction, if it is true for n , then

$$\begin{aligned} \frac{n!}{2\pi i \eta(\gamma, w)} \int_{\gamma} \left(\frac{f(z)}{(z-w)^{n+1}} - \frac{f(z)}{(z-w_0)^{n+1}} \right) dz \\ = \frac{n!}{2\pi i \eta(\gamma, w)} \int_{\gamma} \frac{f(z)}{w_0 - w} \left(\frac{1}{z-w} - \frac{1}{z-w_0} \right) \left(\sum_{k=0}^n \frac{1}{(z-w)^{n-k}(z-w_0)^k} \right) \\ = \frac{n!}{2\pi i \eta(\gamma, w)} \int_{\gamma} \frac{f(z)}{(z-w)(z-w_0)} \left(\sum_{k=0}^n \frac{1}{(z-w)^{n-k}(z-w_0)^k} \right) \end{aligned}$$

and as $w \rightarrow w_0$, we conclude that

$$f^{(n+1)}(w_0) = \frac{(n+1)!}{2\pi i \eta(\gamma, w)} \int_{\gamma} \frac{f(z)}{(z-w_0)^{n+2}} dz$$

which completes the proof. \square

Remark. Because of this, we obtain that for each z upon which f is holomorphic,

$$\|f^{(n)}(z)\| \leq \frac{n!}{2\pi \eta(\gamma, z)} \frac{\|f\|_{L^\infty(\gamma)}}{d(z, \gamma)^n} l(\gamma)$$

a result known as Cauchy's inequality. In the case that γ is just a disk centered at z with radius R , the formula simplifies to

$$\|f^{(n)}(z)\| \leq n! \frac{\|f\|_{L^\infty(\gamma)}}{R^n}$$

Another regularity property enables us to show every holomorphic function is analytic, because

$$\frac{1}{z-w} = \frac{1}{(z-z_0)-(w-z_0)} = \sum_{n=0}^{\infty} \frac{(w-z_0)^n}{(z-z_0)^{n+1}}$$

which converges uniformly on ∂D , hence we can interchange integration and summation in Cauchy's integral formula to obtain a power series representation of f from Cauchy's integral formula. For the next theorem, one needs to know that an **entire** function is one which is holomorphic on the entire complex plane.

Theorem 2.5 (Louville). *Every bounded and entire function is constant.*

Proof. Consider Cauchy's inequality for $n = 1$, which we just proved. As we take the radius of the disc D to ∞ , the term $\|f\|_{L^\infty(\partial D)}$ remains bounded, so the numerator is bounded, whereas the denominator tends to ∞ for every z . Thus we conclude $f' = 0$, so f is constant. \square

Theorem 2.6 (The Fundamental Theorem of Algebra). *Every nonconstant polynomial over the complex numbers has a root.*

Proof. Let f be a polynomial with no root over the complex numbers. If f has degree n , then for large z , $f(z) = (1 + o(1))z^n$. Thus $1/f(z)$ is an entire function, and it is also bounded, since $|z|^n \lesssim |f(z)|$ for large n . Thus $1/f(z)$ is constant, so f is constant. \square

Theorem 2.7 (Morera). *If f is a function with $\int_T f(z) dz = 0$ for all triangles T in a region, then f is holomorphic.*

Proof. Then we know there is a primitive function F with $F' = f$. But F is holomorphic, hence infinitely differentiable, and so f must also be infinitely differentiable. \square

2.2 Uniqueness of Holomorphic Functions

Theorem 2.8. *Let f be a holomorphic function, with $z_n \rightarrow w$ with $f(z_n) = 0$. Then $f = 0$ everywhere.*

Proof. Without loss of generality, assume $w = 0$. Write $f(z) = \sum_{n=0}^{\infty} a_n z^n$ for z close enough to zero. By continuity, $f(0) = a_0 = 0$. If $N > 0$ is the first index with $a_N \neq 0$, then

$$f(z) = a_N z^N (1 + g(z))$$

where g is analytic in a neighbourhood of the origin, and $g(z) \rightarrow 0$ as $z \rightarrow 0$. But this means f cannot have any zeroes in a neighbourhood of the origin, except for the origin itself. So all coefficients of the power series must vanish, so $f = 0$ uniformly. \square

Thus if f and g are two holomorphic functions agreeing on an open set, or even on a countable set with a convergent point, then $f = g$ everywhere.

Example. From Fourier analysis, we know the function $e^{-\pi x^2}$ is its own Fourier transform. Thus we have the integral equation

$$e^{-\pi \xi^2} = \int_{-\infty}^{\infty} e^{-\pi x^2 - 2\pi i \xi x} dx$$

Since both sides can be interpreted as holomorphic functions in the variable ξ (later we will prove that the integral on the right is holomorphic), if ξ is allowed to be complex, we conclude this integral formula holds for any complex ξ , not just the real ξ which were used for the Fourier transform.

2.3 Meromorphic Functions and Residues

s

2.4 Sequences of Holomorphic Functions

Theorem 2.9. If a sequence f_n of holomorphic functions converges locally uniformly to a holomorphic function f , then f is holomorphic.

Proof. Then for any triangle T , f_n converges to f uniformly on T , so

$$\int_T f(z) dz = \lim \int_T f_n(z) dz = 0$$

and so by Morera's theorem, f is holomorphic. \square

A simple consequence is that if $\sum f_n$ converges locally uniformly, then the limit is holomorphic. Taking limits of this proposition leads to the fact that an integral of a continuous family of holomorphic functions is holomorphic.

Theorem 2.10. If $f(z, s)$ is a continuous function holomorphic in z for each s , then

$$F(z) = \int_0^1 f(z, s) ds$$

is holomorphic.

Proof. Using Morera's theorem, if T is a triangle, then we can interchange integration to conclude that

$$\int_T \int_0^1 f(z, s) ds dz = \int_0^1 \int_T f(z, s) dz ds = \int_0^1 0 = 0$$

so the integral is holomorphic. Alternatively, F is the uniform limit of Riemann sums, which are holomorphic. \square

Now we discuss Runge's theorem, which tells us how well we can approximate harmonic functions on a domain by polynomials.

Theorem 2.11 (Runge). *Every holomorphic function defined in a neighbourhood of a compact set K can be uniformly approximated by rational functions with poles in K^c . If K^c is connected, then every such function can be uniformly approximated by polynomials.*

Proof. First we find a curve γ in K^c such that for any $w \in K$ and f holomorphic on K ,

$$f(w) = \int_{\gamma} \frac{f(z)}{z - w}$$

It therefore suffices to approximate f by rational functions on a very small set, the region encompassed by γ . If we parameterize, writing

$$\int_{\gamma} \frac{f(z)}{z - w} = \int_a^b \frac{f(\gamma(t))}{\gamma(t) - w}$$

then this can be uniformly approximated on K by Riemann sums, and each Riemann sum is a rational function.

To show that on a connected K^c we can approximate by polynomials, it suffices to approximate a function of the form $(z - w)^{-1}$, for $z \in K^c$, by a polynomial in w . If all points in K have modulus bounded by R , and if $|z| > R$, then this is easy, for

$$\frac{1}{z - w} = \sum_{k=0}^{\infty} z^k / w^{k+1}$$

and this sum converges uniformly on the ball of radius R . If $|z| < R$, we form a path γ connecting z to a point z_0 with $|z_0| > R$. If $|w_0 - w_1| <$

$d(K, \gamma)/2$, then

$$\frac{1}{z - w_0} = \frac{1}{(z - w_1) - (z - w_0)} = \sum \frac{(z - w_0)^n}{(z - w_1)^{n+1}}$$

Thus if $(z - w_1)^{-1}$ can be approximated by polynomials, so too can $(z - w_0)^{-1}$. But if we now take a sequence of points along γ at a distance $d(K, \gamma)/2$ apart from one another, we can successively approximate each inverse function by polynomials in the inverse function of the next point. Eventually, this gives us the approximation of $(z - w)^{-1}$ as polynomials in $(z_0 - w)^{-1}$, and this can be approximated by polynomials in w . \square

2.5 The Reflection Principle

Let Ω be a region symmetric under conjugation, so $\overline{\Omega} = \Omega$. Split the region into its upper and lower parts Ω^+ and Ω^- . We now discuss if it's possible to patch together two functions defined on either side of the half plane.

Theorem 2.12. *If f^+ and f^- are functions defined on Ω^+ and Ω^- respectively, and they can be extended to continuous functions extended to be defined on $\Omega \cap \mathbf{R}$ in addition to their original domain, then this extension, when f^+ and f^- are combined to form a function f , is holomorphic.*

Proof. Just apply Morera's theorem, splitting triangles passing through the real line into smaller triangles on either side of the line. \square

Theorem 2.13 (Schwartz Reflection Principle). *If f is holomorphic in Ω^+ , and real valued when extended to the real axis, then f can be extended to Ω .*

Proof. If we define $f(z) = \overline{f(\bar{z})}$ on Ω^- , then f remains holomorphic, and we can apply the last theorem to conclude that this gives a holomorphic extension on the entirety of Ω . \square

Chapter 3

Geometric Function Theory

An intuitive way to visualize analytic functions is how they act on the geometry of the complex plane. The Riemann school believes that this should be the primary way of understanding these maps, giving us interesting geometric results. Here we introduce conformal maps, and pave the way to Riemann's classical mapping theorem.

The study of geometry was fundamentally changed by Felix Klein's Erlangen program. To rationalize the emergence of the paradoxical non-euclidean geometries, The fundamental thesis of the program is that a study of the invariants of space under actions from a certain group. Klein's Euclidean geometry studies invariants of the plane invariant under similarity transformations, which preserve angles and proportions, collected into the Euclidean group $E(2)$. Projective geometry studies invariants of the sphere under transformations which preserve the incidence of straight lines, collected into the projective group $P(2)$. Of course, one may perturb these groups to obtain more interesting results. For instance, to study properties of space which remain after oriented similarity, we weaken our actions to the special euclidean group $SE(2)$, a subgroup of $E(2)$.

Geometrically, the set of all **biholomorphic functions** (holomorphic maps with a holomorphic inverse) forms a group.

After studying all these analytic functions, it is of course, of interest to determine the properties of the plane which remain invariant under the actions of bijective analytic functions, which form a group.

Let us pretend we have never seen the definition of an analytic function. In fact, forget we have ever invented complex numbers at all (we consider \mathbf{C} as another name for the real plane \mathbf{R}^2). \mathbf{C} is a two-dimensional

surface, and therefore possesses a tangent space $T\mathbf{C}$, which naturally can be considered as \mathbf{C}^2 – tangent vectors in this space will be denoted as v_p , the vector v at the point p . $T\mathbf{C}$ is also a Riemannian manifold, since we have a natural inner product on the tangent vectors at each point on the plane, defined by

$$\langle v_p, w_p \rangle = v_1 w_1 + v_2 w_2 = \Re[v \overline{w}]$$

One obtains ‘infinitesimal’ equivalents to the structures on an inner product space. For instance, one may talk about the ‘infinitesimal’ length of a tangent vector, defined by

$$\|v_p\| = \langle v_p, v_p \rangle$$

More interesting to us, we are able to talk about ‘infinitesimal angles’ at points. That is, the angle between two tangent vectors v_p and w_p is the angle θ such that

$$\cos(\theta) = \frac{\langle v_p, w_p \rangle}{\|v\| \|w\|}$$

We are now ready to discuss conformal maps, the geometric method of introducing analytic functions.

Definition. A complex-valued differentiable map f defined on an open subset of the plane is **conformal** if it is oriented, and preserves infinitesimal angles. Pointwise conformality is defined similarly

Lemma 3.1. *A function analytic at a point with non-zero derivative is conformal there.*

Proof. Let f be analytic at p , with $f'(p) = z \neq 0$. The matrix representation of $f'(p)$ is

$$\begin{pmatrix} \Re[z] & \Im[z] \\ -\Im[z] & \Re[z] \end{pmatrix}$$

Whose determinant is $\Re[z]^2 + \Im[z]^2 > 0$, hence the map is oriented at p . A calculation verifies that

$$\langle f^*(v_p), f^*(w_p) \rangle = \langle zv, zw \rangle = |z|^2 \langle v, w \rangle$$

So that, since $|zv| = |z||v|$, and $|zw| = |z||w|$,

$$\frac{\langle zv, zw \rangle}{|zv||zw|} = \frac{|z|^2 \langle v, w \rangle}{|z|^2 |v||w|} = \frac{\langle v, w \rangle}{|v||w|}$$

Thus the map is conformal. \square

We shall show that, the differential map at any point in the plane is just a rotation and scaling. For each unit tangent vector $e^{i\theta}$ at a point p , consider the differential map $f_*((e^{i\theta})_p)$.

Let us now use our knowledge of the complex plane to reintroduce an old friend. We may express the differential of a conformal map $f = u + iv$ as

$$f^*(v_p) =$$

Consider the action of a conformal map f on the unit tangent vectors $(e_1)_p$ and $(e_2)_p$

$$f^*((e_1)_p) = Df(p)(e_1) = \left. \frac{\partial f^1}{\partial x} \right|_p \left. \frac{\partial}{\partial x} \right|_{f(p)} + \left. \frac{\partial f^2}{\partial x} \right|_p \left. \frac{\partial}{\partial y} \right|_{f(p)}$$

$$f^*((e_2)_p) = Df(p)(e_2) = \left. \frac{\partial f^1}{\partial y} \right|_p \left. \frac{\partial}{\partial x} \right|_{f(p)} + \left. \frac{\partial f^2}{\partial y} \right|_p \left. \frac{\partial}{\partial y} \right|_{f(p)}$$

Since f is conformal, these vectors must be at right angles, and $(f^*(1_p), f^*(i_p))$ must be an oriented basis. In two dimensions, the only vectors orthogonal to a vector $v = (a, b)$ are scalar multiples of the vector $w = (-b, a)$. Since $(v, \lambda w)$ must be oriented, we must have $\lambda(a^2 + b^2) > 0$, hence $\lambda > 0$. Taken in the case of our conformally mapped tangent vectors, we must have some $\lambda > 0$ for which,

$$f^*((e_2)_p) = \lambda \left(- \left. \frac{\partial f^2}{\partial x} \right|_p \left. \frac{\partial}{\partial x} \right|_{f(p)} + \left. \frac{\partial f^1}{\partial x} \right|_p \left. \frac{\partial}{\partial y} \right|_{f(p)} \right)$$

Equating our two values of $f^*((e_2)_p)$, we see that

$$\left. \frac{\partial f^1}{\partial y} \right|_p = -\lambda \left. \frac{\partial f^2}{\partial x} \right|_p \quad \left. \frac{\partial f^2}{\partial y} \right|_p = \lambda \left. \frac{\partial f^1}{\partial x} \right|_p$$

Chapter 4

Riemann Surfaces

The study of differentiable manifolds generalizes the calculus of real variables in Euclidean space to spaces which locally behave like Euclidean space. In this section we discuss the theory which generalizes the theory of holomorphic functions on domains in the complex plane to spaces which locally behave like the complex plane. These ‘one-dimensional complex manifolds’ are known as **Riemann surfaces**. Topological intuition suggests that complex manifolds should be 2-dimensional real manifolds, with additional structure which enables us to identify a class of holomorphic functions. The standard way to identify differentiable functions on a real manifold is to introduce a C^∞ structure, considering an atlas of local embeddings $x : U \rightarrow \mathbf{R}$, such that if $y : V \rightarrow \mathbf{R}$ is any other embedding, the transition map $x \circ y^{-1}$ is C^∞ where defined. Similarly, we introduce a C^ω **atlas** to be a cover of a two dimensional manifold by charts such that any two chars (x, U) and (y, V) have a holomorphic transition map $x \circ y^{-1}$ where defined. A two dimensional manifold X with a fixed C^ω atlas is known as a Riemann surface.

Every Riemann surface X is certainly a two dimensional manifold, so we can consider the tangent space $T_p X$. What’s more, we can give these two dimensional tangent spaces a one dimensional complex structure. Recall that a complex vector space is just a real vector space V with a fixed ‘90 degree rotation’, a linear map $J : V \rightarrow V$ satisfying $J^2 = -1$. We can then define $(a + bi)v = a + b(Jv)$. We define this linear map by working in coordinates. For each point p with holomorphic coordinate chart (x, U) , we define $J_p : T_p X \rightarrow T_p X$ by letting $J_p(v) = x_*^{-1}(ix_*(v))$. Since the transition maps of any two coordinate charts x and y are holomorphic, $(y_* \circ x_*^{-1})$ is

essentially a complex linear map on the tangent spaces, which means that J_p is well defined, and gives TX a canonical complex structure.

Remark. Fundamentally, the ‘ J ’ map is what defines the holomorphic properties of a 2-manifold. If a given bundle map $J : TM \rightarrow TM$ is given such that $J^2 = -1$, then this map defines a complex linear structure on the tangent spaces, and we define $f : M \rightarrow N$ to be holomorphic if the corresponding map f_* is complex linear at each point. A holomorphic atlas on M may then be defined to be the set of all ‘holomorphic charts’, for if (x, U) and (y, V) are holomorphic charts, then $x \circ y^{-1}$ is holomorphic, for it is certainly differentiable, and the derivative is complex linear. Thus as long as J is ‘consistent’, in the sense that there exist enough holomorphic charts to cover the manifold, then J is a perfectly reasonable way to define a complex structure on a 2-manifold. In fact, J is what we call a **almost complex structure** on the manifold, for it is almost all we need to define a complex manifold structure to M .

Theorem 4.1. *Every complex manifold is orientable.*

Proof. Let $T : \mathbf{R}^{2n} \rightarrow \mathbf{R}^{2n}$ be a linear endomorphism given by complex multiplication, viewing $\mathbf{R}^{2n} = \mathbf{C}^n$, so $Tv = \lambda v$ for some $\lambda = a + ib$. Then if $v = x + iy$, for $x, y \in \mathbf{R}^n$, we have $Tv = (ax - by) + i(ay + bx)$. If we take a basis being the unit coordinate vectors of x and y , the matrix representation of T is n copies of the matrix

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

along the diagonal. Thus $\det(T) = (a^2 + b^2)^n = |\lambda|^{2n} > 0$. In particular, viewing the holomorphic transition maps $(y \circ x^{-1}) : \mathbf{C}^n \rightarrow \mathbf{C}^n$ as maps from \mathbf{R}^{2n} to itself, we conclude that since $(y \circ x^{-1})'$ is non-vanishing, the determinant of $(y \circ x^{-1})'$ is positive. Thus the atlas preserves orientation, and thus the manifold is orientable. \square

A map $f : X \rightarrow Y$ between Riemann surfaces is holomorphic if the map is holomorphic in every coordinate system. Since we have the tangent bundle, there is a more natural coordinate-invariant definition. Provided f is differentiable (which it must be if it is holomorphic), f induces a map $f_* : TX \rightarrow TY$. f is holomorphic at a point if and only if f_* is complex-linear at the point. This makes sense, because Riemann surfaces should

model the local properties of the complex plane, and holomorphic functions on the complex plane are exactly those functions which are locally complex linear maps. Most tangibly, we often focus on the space $\mathcal{O}(X)$ of holomorphic maps from X into \mathbf{C} , which forms a complex vector space.

Example. *The fundamental example of a Riemann surface is the Riemann sphere $\mathbf{C}_\infty = \mathbf{C} \cup \{\infty\}$, whose topology is defined to be the compactification of \mathbf{C} . The subsets \mathbf{C} and $\mathbf{C}_\infty - \{\infty\}$ both have homeomorphisms onto \mathbf{C} , the first by the identity map, and the second by the map $z \mapsto 1/z$. The maps are compatible since $1/z$ is holomorphic for $z \neq 0$ on \mathbf{C} . A function $f : \mathbf{C}_\infty \rightarrow \mathbf{C}$ is holomorphic if and only if $f(z)$, and $f(1/z)$ are entire, which unfortunately implies that f is constant, since \mathbf{C}_∞ is compact, and therefore f is bounded.*

Since the space $\mathcal{O}(X)$ is trivial when X is a compact complex manifold, it gives little information about the complex structure of X . Thus we must generalize the functions we study to a more general class, which enables us to be ‘non-compact’ in the image. If X is a Riemann surface, we say $f : X \rightarrow \mathbf{C}_\infty$ is **meromorphic** if it is holomorphic as a map between Riemann surfaces, and f is not the constant ∞ function. The *singularities* of such a function are the points p with $f(p) = \infty$.

Example. *Let $f : U \rightarrow \mathbf{C}_\infty$ be meromorphic, as in the definition of Riemann surfaces, where U is a connected, open subset of \mathbf{C} . This means that $1/f(z) \neq 0$, and both $f(z)$ and $1/f(z)$ are holomorphic where they are defined. Since $1/f(z) \neq 0$, this means that the set of points z where $f(z) = \infty$ must form a discrete subset of U . What’s more, if $f(z_0) = \infty$, by the continuity of f , $\lim_{z \rightarrow z_0} f(z) = \infty$. Thus f is meromorphic in the classical sense. On the other hand, if f is meromorphic in the usual sense of the word, then $f(z)$ is holomorphic where finite, and to prove that the definitions are equivalent it suffices to prove $1/f(z)$ is holomorphic. But we know this is true because if $f(z_0) = \infty$, there is a positive number n such that $(z - z_0)^n f(z)$ is holomorphic and nonvanishing in a neighbourhood of z_0 , so*

$$\frac{1}{f(z)(z - z_0)^n}$$

is also holomorphic in a neighbourhood of z_0 . But this means that

$$\frac{1}{f(z)} = \frac{1}{f(z)(z - z_0)^n} (z - z_0)^n$$

is holomorphic at z_0 , completing the proof.

Thus a meromorphic function on a Riemann surface is really just locally a meromorphic function in the usual sense. In particular, it follows from this that the singularities of a meromorphic function forms a discrete set, and that each singularity has a well defined order (since a biholomorphic change of variables on \mathbf{C} does not change the order of a singularity). Furthermore, a meromorphic function is locally just a quotient of two holomorphic functions, which in a more modern context means the meromorphic functions are the rational functions corresponding to the scheme of holomorphic functions. The sum of two meromorphic functions is also meromorphic (where $\infty + z = \infty$) is also meromorphic, and if f is a nonzero meromorphic function, then $1/f$ is also meromorphic. In particular, the family $M(X)$ of all meromorphic functions on X forms a field. For each p , the function $f \mapsto \text{ord}_p(f)$ is algebraically an order function, and therefore the functions for which $\text{ord}_p(f) \geq 0$ forms a discrete valuation ring $\mathcal{O}_p(X)$, the ring of meromorphic functions defined at p .

4.1 Derivations

Even if we forget the tangent space $T_p M$, we may still reconstruct it from a holomorphic atlas by considering derivations. Recall that a p -derivation is a complex-valued function Λ such that for any two functions $f, g \in C^\infty(M, \mathbf{C})$, $\Lambda(fg) = f(p)\Lambda g + g(p)\Lambda f$. The differential operators ∂_x and ∂_y are contained in $T_p \mathbf{C}$, and the same argument as in the real case shows these operators span the complex vector space of all derivations. As with a tangent space, we can consider the cotangent space $T_p^* \mathbf{C}$, the dual space of $T_p \mathbf{C}$. A smooth $f : \mathbf{C} \rightarrow \mathbf{C}$ gives rise to a section df of the cotangent space, and $dx(p)$ and $dy(p)$ span the cotangent space at each point. The most intriguing part of this construction occurs when we examine holomorphic functions. In classic notation, define z to be the identity map on \mathbf{C} . Then z is trivially smooth, and so we may consider $dz = dx + i dy$. Define $\bar{z} : w \rightarrow \bar{w}$. Then $d\bar{z} = dx - i dy$. These equations are linearly independent, and thus span the cotangent space at each point. Thus any $w \in T_p^* \mathbf{C}$ may be written $\alpha dz + \beta d\bar{z}$ for two values α and β . In particular, this holds for df . We define the coefficients $\partial_z f$ and $\partial_{\bar{z}} f$ by the equation $df = \partial_z f dz + \partial_{\bar{z}} f d\bar{z}$. Since $\alpha dz + \beta d\bar{z} = (\alpha + \beta)dx + i(\alpha - \beta)dy$. We obtain the equalities

$$\frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}} = \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial z} - \frac{\partial f}{\partial \bar{z}} = -i \frac{\partial f}{\partial y}$$

Reordering these equations, we determine that

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right) \quad \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right)$$

If f is real-differentiable at a point z , then

$$f(z+w) = f(z) + w \left. \frac{\partial f}{\partial z} \right|_z + \bar{w} \left. \frac{\partial f}{\partial \bar{z}} \right|_z + o(w)$$

so these partial derivatives separate the parts of the derivative as the part corresponding to the identity and the parts corresponding to the conjugate. This is possible because the space of real-linear endomorphisms on \mathbb{C} is four dimensional, and each of the partial derivatives in the equation above has two degrees of freedom.

Note that the Cauchy Riemann equations state exactly that $\partial_{\bar{z}} f = 0$, and it then follows that $f' = \partial_z f$. This explains why the operators $\partial_{\bar{z}}$ rarely come up once we discuss holomorphic functions. And indeed, we find that the ‘tangent space’ corresponding to the space of holomorphic functions is one dimensional, spanned by ∂_z . Since the space of holomorphic functions does not contain bump functions, simply taking derivations on $C^\omega(\mathbb{C})$ would not be local enough to correspond to a tangent space. Thus we define the tangent space to be the space of derivations on the ring $\mathcal{O}_p(\mathbb{C})$ of germs of holomorphic functions defined around p .

Theorem 4.2. *The derivations on $\mathcal{O}_p(\mathbb{C})$ are multiples of ∂_z .*

Proof. Without loss of generality, assume the p is the origin. Let l be a derivation. If f is holomorphic and $f(0) = 0$, then $g(z) = (f(z) - f(0))/z$ is holomorphic with $g(0) = f'(0)$, and we calculate

$$l(f) = l(gz) + f(0)l(1) = f'(0)l(z)$$

This means that $l = l(z)\partial_z$. □

This also means that the cotangent space is spanned by dz . Since $dz(\dot{\gamma}(t)) = \gamma'(t)$, the differential forms definition of integration gives

$$\int_{\gamma} f(z) dz = \int_a^b f(\gamma(t)) dz(\dot{\gamma}(t)) dt = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

which is precisely the definition of contour integration we gave before. By linearity, all the real theorems on integration extend by linearity to this complex situation. In particular, Poincare's theorem implies that if f is a smooth holomorphic function on a simply connected domain, then $dg = f dz$ for some smooth function g , which must be holomorphic because then $\partial_{\bar{z}}g = 0$, and so for any curve γ ,

$$\int_{\gamma} f(z) dz = g(\gamma(b)) - g(\gamma(a))$$

Most importantly, if γ is a closed curve in a simply connected domain, then

$$\int_{\gamma} f(z) dz = 0$$

Since all holomorphic functions are smooth, this very easily justifies the proof of Cauchy's theorem for general curves in simply connected regions.

Bibliography

- [1] Michael Spivak, *A Concise Introduction to Differential Geometry: Vol. One*
- [2] Paul Halmos, *Naive Set Theory*
- [3] Wikipedia, *Lie Groups*