
Using fMRI to Diagnose Schizophrenia

Department of Computing Science
University of Alberta

Department of Computing Science
University of Alberta

Department of Computing Science
University of Alberta

Abstract

1 Introduction

Diagnosis of schizophrenia is a challenging task that yet to be addressed[1]. Although, in recent years methods which use Functional Magnetic Resonance Imaging(fMRI) for mental disorder diagnosis has become more popular, but in case of schizophrenia it still needs to become more robust and reliable. In similar studies[2][3] have been shown that fMRI can be used in conjunction with Sparse Gaussian Markov Random Field to produce high accuracy in diagnosis of illness. However having a dataset with homogeneous distribution of illness makes this result less reliable and creates the need for more evidence using heterogeneous dataset in terms of illness.

References

- [1] Philip McGuire, Oliver D Howes, James Stone, and Paolo Fusar-Poli. Functional neuroimaging in schizophrenia: diagnosis and drug discovery. *Trends in Pharmacological Sciences*, 29(2):91 – 98, 2008.
- [2] Irina Rish, Guillermo Cecchi, Benjamin Thyreau, Bertrand Thirion, Marion Plaze, Marie Laure Paillere-Martinot, Catherine Martelli, Jean-Luc Martinot, and Jean-Baptiste Poline. Schizophrenia as a network disease: Disruption of emergent brain function in patients with auditory hallucinations. *PLoS ONE*, 8(1):e50625, jan 2013.
- [3] Maria J. Rosa, Liana Portugal, John Shawe-Taylor, and Janaina Mourao-Miranda. Sparse network-based models for patient classification using fMRI. In *2013 International Workshop on Pattern Recognition in Neuroimaging*. Institute of Electrical & Electronics Engineers (IEEE), jun 2013.