


Divide-and-Conquer




Outline and Reading

- ◆ Divide-and-conquer paradigm (§5.2)
- ◆ Review Merge-sort (§4.1.1)
- ◆ Recurrence Equations (§5.2.1)
 - Iterative substitution
 - Recursion trees
 - Guess-and-test
 - The master method
- ◆ Integer Multiplication (§5.2.2)

Divide-and-Conquer

- ◆ **Divide-and conquer** is a general algorithm design paradigm:
 - **Divide**: divide the input data S in two or more disjoint subsets S_1, S_2, \dots
 - **Recur**: solve the subproblems recursively
 - **Conquer**: combine the solutions for S_1, S_2, \dots , into a solution for S
- ◆ The base case for the recursion are subproblems of constant size
- ◆ Analysis can be done using **recurrence equations**



Merge-Sort Review

- ◆ Merge-sort on an input sequence S with n elements consists of three steps:
 - **Divide**: partition S into two sequences S_1 and S_2 of about $n/2$ elements each
 - **Recur**: recursively sort S_1 and S_2
 - **Conquer**: merge S_1 and S_2 into a unique sorted sequence

Algorithm *mergeSort*(S, C)

Input sequence S with n elements, comparator C

Output sequence S sorted according to C

if $S.size() > 1$

$(S_1, S_2) \leftarrow partition(S, n/2)$

mergeSort(S_1, C)

mergeSort(S_2, C)

$S \leftarrow merge(S_1, S_2)$


Recurrence Equation Analysis



- ◆ The conquer step of merge-sort consists of merging two sorted sequences, each with $n/2$ elements and implemented by means of a doubly linked list, takes at most bn steps, for some constant b .
- ◆ Likewise, the basis case ($n < 2$) will take at b most steps.
- ◆ Therefore, if we let $T(n)$ denote the running time of merge-sort:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn & \text{if } n \geq 2 \end{cases}$$

- ◆ We can therefore analyze the running time of merge-sort by finding a **closed form solution** to the above equation.
 - That is, a solution that has $T(n)$ only on the left-hand side.



Iterative Substitution

- ◆ In the iterative substitution, or “plug-and-chug,” technique, we iteratively apply the recurrence equation to itself and see if we can find a pattern:

$$\begin{aligned}T(n) &= 2T(n/2) + bn \\&= 2(2T(n/2^2) + b(n/2)) + bn \\&= 2^2 T(n/2^2) + 2bn \\&= 2^3 T(n/2^3) + 3bn \\&= 2^4 T(n/2^4) + 4bn \\&= \dots \\&= 2^i T(n/2^i) + ibn\end{aligned}$$





- ◆ Note that base, $T(1)=b$, case occurs when $2^i=n$. That is, $i = \log n$.
- ◆ So, $T(n) = bn + bn \log n$
- ◆ Thus, $T(n)$ is $O(n \log n)$.

The Recursion Tree



- ◆ Draw the recursion tree for the recurrence relation and look for a pattern:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn & \text{if } n \geq 2 \end{cases}$$

depth	T's	size		time
0	1	n		bn
1	2	$n/2$		bn
i	2^i	$n/2^i$		bn
...		...

Total time = $bn + bn \log n$
(last level plus all previous levels)

Guess-and-Test Method

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn & \text{if } n \geq 2 \end{cases}$$

- ◆ In the guess-and-test method, we guess a closed form solution and then try to prove it is true by induction:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn \log n & \text{if } n \geq 2 \end{cases}$$

- ◆ Guess: $T(n) < cn \log n$.

$$\begin{aligned} T(n) &= 2T(n/2) + bn \log n \\ &= 2(c(n/2) \log(n/2)) + bn \log n \\ &= cn(\log n - \log 2) + bn \log n \\ &= cn \log n - cn + bn \log n \end{aligned}$$

- ◆ Wrong: we cannot make this last line be less than $cn \log n$

Guess-and-Test Method, Part 2

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn \log n & \text{if } n \geq 2 \end{cases}$$

- ◆ Recall the recurrence equation:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn \log n & \text{if } n \geq 2 \end{cases}$$

- ◆ Guess #2: $T(n) < cn \log^2 n$.

$$\begin{aligned} T(n) &= 2T(n/2) + bn \log n \\ &= 2(c(n/2) \log^2(n/2)) + bn \log n \\ &= cn(\log n - \log 2)^2 + bn \log n \\ &= cn \log^2 n - 2cn \log n + cn + bn \log n \\ &\leq cn \log^2 n \end{aligned}$$

- if $c > b$.

- ◆ So, $T(n)$ is $O(n \log^2 n)$.

- ◆ In general, to use this method, you need to have a good guess and you need to be good at induction proofs.



Master Method

- Many divide-and-conquer recurrence equations have the form:

$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

- The Master Theorem:

- if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
- if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
- if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

Master Method, Example 1



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:
$$T(n) = 4T(n/2) + n$$

Solution: $\log_b a = 2$, so case 1 says $T(n)$ is $\Theta(n^2)$.

Master Method, Example 2



◆ The form: $T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 2T(n/2) + n \log n$$

Solution: $\log_b a = 1$, so case 2 says $T(n)$ is $\Theta(n \log^2 n)$.

Master Method, Example 3



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = T(n/3) + n \log n$$

Solution: $\log_b a = 0$, so case 3 says $T(n)$ is $\Theta(n \log n)$.

Master Method, Example 4



◆ The form: $T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 8T(n/2) + n^2$$

Solution: $\log_b a = 3$, so case 1 says $T(n)$ is $\Theta(n^3)$.

Master Method, Example 5



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 9T(n/3) + n^3$$

Solution: $\log_b a = 2$, so case 3 says $T(n)$ is $\Theta(n^3)$.

Master Method, Example 6



◆ The form: $T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = T(n/2) + 1 \quad (\text{binary search})$$

Solution: $\log_b a = 0$, so case 2 says $T(n)$ is $\Theta(\log n)$.

Master Method, Example 7



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$
2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$
3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $a f(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 2T(n/2) + \log n \quad (\text{heap construction})$$

Solution: $\log_b a = 1$, so case 1 says $T(n)$ is $\Theta(n)$.

Iterative “Proof” of the Master Theorem



◆ Using iterative substitution, let us see if we can find a pattern:

$$\begin{aligned}T(n) &= aT(n/b) + f(n) \\&= a(aT(n/b^2) + f(n/b)) + f(n) \\&= a^2T(n/b^2) + af(n/b) + f(n) \\&= a^3T(n/b^3) + a^2f(n/b^2) + af(n/b) + f(n) \\&\quad \text{M} \\&= a^kT(n/b^k) + \sum_{i=0}^{k-1} a^i f(n/b^i) \\&\quad \text{M}\end{aligned}$$

Iterative “Proof” of the Master Theorem



$$\begin{aligned} & M \\ &= a^{\log_b n} T(1) + \sum_{i=0}^{(\log_b n)-1} a^i f(n/b^i) \\ &= n^{\log_b a} T(1) + \sum_{i=0}^{(\log_b n)-1} a^i f(n/b^i) \end{aligned}$$

- ◆ We then distinguish the three cases as
 - The first term is dominant
 - Each part of the summation is equally dominant
 - The summation is a geometric series

Integer Multiplication



◆ Algorithm: Multiply two n-bit integers I and J.

- Divide step: Split I and J into high-order and low-order bits

$$I = I_h 2^{n/2} + I_l$$

$$J = J_h 2^{n/2} + J_l$$

- We can then define $I * J$ by multiplying the parts and adding:

$$\begin{aligned} I * J &= (I_h 2^{n/2} + I_l) * (J_h 2^{n/2} + J_l) \\ &= I_h J_h 2^n + I_h J_l 2^{n/2} + I_l J_h 2^{n/2} + I_l J_l \end{aligned}$$

- So, $T(n) = 4T(n/2) + n$, which implies $T(n)$ is $\Theta(n^2)$.
- But that is no better than the algorithm we learned in grade school.

where did this come from?

An Improved Integer Multiplication Algorithm



◆ Algorithm: Multiply two n-bit integers I and J.

- Divide step: Split I and J into high-order and low-order bits

$$I = I_h 2^{n/2} + I_l$$

$$J = J_h 2^{n/2} + J_l$$

- Observe that there is a different way to multiply parts:

$$\begin{aligned} I * J &= I_h J_h 2^n + [(I_h - I_l)(J_l - J_h) + I_h J_h + I_l J_l] 2^{n/2} + I_l J_l \\ &= I_h J_h 2^n + [(I_h J_l - I_l J_l - I_h J_h + I_l J_h) + I_h J_h + I_l J_l] 2^{n/2} + I_l J_l \\ &= I_h J_h 2^n + (I_h J_l + I_l J_h) 2^{n/2} + I_l J_l \end{aligned}$$

requires only 3 mults!

- So, $T(n) = 3T(n/2) + n$, which implies $T(n)$ is $O(n^{\log_2 3})$, by the Master Theorem.
- Thus, $T(n)$ is $O(n^{1.585})$.

Matrix Multiplication

- ◆ Given $n \times n$ matrices X and Y , wish to compute the product $Z=XY$.
- ◆ Formula for doing this is

$$Z_{ij} = \sum_{k=0}^{n-1} X_{ik} Y_{kj}$$

- ◆ This runs in $O(n^3)$ time
 - In fact, multiplying an $n \times m$ by an $m \times q$ takes nmq operations

Matrix Multiplication

$$\begin{bmatrix} I & J \\ K & L \end{bmatrix} = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} E & F \\ G & H \end{bmatrix}$$

$$I = AE + BG$$

$$J = AF + BH$$

$$K = CE + DG$$

$$L = CF + DH$$

Matrix Multiplication

- ◆ Using the decomposition on previous slide, we can compute Z using 8 recursively computed $(n/2) \times (n/2)$ matrices plus four additions that can be done in $O(n^2)$ time
- ◆ Thus $T(n) = 8T(n/2) + bn^2$
- ◆ Still gives $T(n)$ is $\Theta(n^3)$

Strassen's Algorithm

- ◆ If we define the matrices S_1 through S_7 as follows

$$S_1 = A(F - H)$$

$$S_2 = (A + B)H$$

$$S_3 = (C + D)E$$

$$S_4 = D(G - E)$$

$$S_5 = (A + D)(E + H)$$

$$S_6 = (B - D)(G + H)$$

$$S_7 = (A - C)(E + F)$$

Strassen's Algorithm

◆ Then we get the following:

$$I = S_5 + S_6 + S_4 - S_2$$

$$J = S_1 + S_2$$

$$K = S_3 + S_4$$

$$L = S_1 - S_7 - S_3 + S_5$$

◆ So now we can compute $Z=XY$ using only seven recursive multiplications

Strassen's Algorithm

- ◆ This gives the relation $T(n) = 7T(n/2) + bn^2$ for some $b > 0$.
- ◆ By the Master Theorem, we can thus multiply two $n \times n$ matrices in $\Theta(n^{\log 7})$ time, which is approximately $\Theta(n^{2.808})$.
 - May not seem like much, but if you're multiplying two 100×100 matrices:
 - ◆ n^3 is 1,000,000
 - ◆ $n^{2.808}$ is 413,048
- ◆ With added complexity, there are algorithms to multiply matrices in as little as $\Theta(n^{2.376})$ time
 - Reduces figures above to 56,494