

Prior Authorization for Botulinum Toxin	
Contraindications Check	
Does the beneficiary have a condition that is a contraindication to receiving botulinum toxin? *	☐ No Contraindications ☐ An existing medical condition(s) which could affect the neuromuscular function ☐ A hypersensitivity to any botulinum toxin preparation or to any of the components in the formulation of the serotype ☐ A severe clotting disorder ☐ An infection at the proposed injection site
Order Details	
Order date *	MM/DD/YYYY
Requested procedure date of service *	MM/DD/YYYY
The requested procedure is one of the following: Proposed frequency of injections. Provider Note: Administration must not be given more frequently than every 12 weeks, regardless of diagnosis, unless specifically address in the policy. *	☐ Initial injection ☐ Subsequent injection ☐ Last injection ☐ One time dose/trial dose
Order Description: A request for injection of chemical (botulinum toxin) for destruction of nerve muscles (chemodenervation). *	☐ CPT 64612 - Chemodenervation of muscles(s); muscle(s) innervated by facial nerve, unilateral (e.g.m for blepharospasm, hemifacial spasm) ☐ CPT 64615 - Chemodenervation of muscles(s); muscle(s) innervated by facial, trigeminal, cervical, spinal and accessory nerves, bilateral (e.g., for chronic migraine) ☐ CPT 95873 -Muscle stimulation guidance will be used during chemodenervation to avoid the documented structures that are adjacent to or behind the target muscle, reducing the risk of complications. ☐ CPT 95874 - Electromyographic guidance for chemodenervation to avoid the documented structures that are adjacent to or behind the target muscle, reducing the risk of complications ☒ NA-LCD does not require
Allowable Diagnosis	
Covered Diagnosis. Provider education: There must be objective documentation of the clinical features consistent with the covered diagnosis in the beneficiary record. *	☐ G24.1 Genetic torsion dystonia ☐ G24.2 Idiopathic nonfamilial dystonia ☐ G24.3 Spasmodic torticollis ☐ G24.4 Idiopathic orofacial dystonia ☐ G24.5 Blepharospasm ☐ G24.8 Other Dystonia ☐ G51.2 Melkersson's syndrome ☐ G51.31 Clonic hemifacial spasm, right ☐ G51.32 Clonic hemifacial spasm, left ☐ G51.4 Facial myokymia ☐ H02.041 Spastic entropion of right upper eyelid ☐ H02.042 Spastic entropion of right lower eyelid ☐ H02.044 Spastic entropion of left upper eyelid ☐ H02.045 Spastic entropion of left lower eyelid ☐ H02.141 Spastic ectropion of right upper eyelid ☐ H02.142 Spastic ectropion of right lower eyelid ☐ H02.144 Spastic ectropion of left upper eyelid ☐ H02.145 Spastic ectropion of left lower eyelid ☐ G81.11 Spastic hemiplegia affecting right dominant side ☐ G81.12 Spastic hemiplegia affecting left dominant side ☐ G81.13 Spastic hemiplegia affecting right nondominant side ☐ G81.14 Spastic hemiplegia affecting left nondominant side ☐ H02.431 Paralytic ptosis of right eyelid ☐ H02.432 Paralytic ptosis of left eyelid ☐ H02.59 Other disorders affecting eyelid function ☐ R25.8 Other abnormal involuntary movements ☐ Other
Covered Diagnosis. Provider education: There must be objective documentation of the clinical features consistent with the covered diagnosis in the beneficiary record. *	Select one
Specify Other	Type a value
If the selected ICD-10 code is not on the billing article allowable list, submit robust published evidence to support medical neccessity.	Type a value
Provider Education Note: Specific information must be documented about the conservative methods of treatments such as medication, physical therapy, and other appropriate methods that have been tried and proven unsuccessful. The documentation must include an objective assessment of the intolerance or inadequacy of the response(s). (Excludes hemifacial spasm or blepharospasm)	
Non-Migraine Diagnosis Requirements	
Medication(s) that have been attempted. *	☐ Anticholinergics (Muscarinic receptor antagonist (372679000) ☐ Benzotropine ☐ Trihexyphenidyl ☐ Benzodiazepine ☐ Clonazepam ☐ Diazepam ☐ Lorazepam ☐ Skeletal Muscle relaxants ☐ Baclofen ☐ Tizanidine ☐ Cyclobenzaprine ☐ Methocarbamol ☐ Gabapentin ☐ Dantrolene ☐ Other ☐ N/A -Not Required by LCD
Specify Other	Type a value
Conventional treatment methods that have been attempted. (Excludes hemifacial spasm or blepharospasm) *	☐ Physical Therapy ☐ Biofeedback (Electromyographic) ☐ Acupuncture ☐ Lifestyle modifications ☐ N/A -Not Required by LCD
Beneficiary's therapeutic response to treatments attempted *	☐ Complete ☐ Excellent ☐ Good response ☐ Fair response ☐ Poor/minimal ☐ N/A -Not Required by LCD
Pertinent test/procedure results *	Type a value
Objective scale score of the severity of symptoms *	☐ 0 - No impairment ☐ 1 - Mild impairment ☐ 2 - Moderate impairment ☐ 3 - Severe impairment ☐ 4 - Not possible due to the disease ☐ Not applicable
Migraine Diagnosis Requirements	
The use of the selected serotype is not approved by the FDA and falls outside of treatment guidelines. Submit robust published evidence for this off label request AND the reason an approved serotype is not used.	Type a value
History of Chronic Migraines, select all that apply. *	
	☐ Headache disorders occurring greater than 15 days a month, with a duration of four or more hours, for a period of at least three months ☐ Migraine headache documented to be lasting for a duration ≥4 hours on a migraine day ☐ Headache occurs 15 or more days per month, at least 8 of the headache days per month meet the migraine without aura criteria, or respond to migraine specific treatment. ☐ The beneficiary has significant disability due to the headaches and have been refractory to standard and conventional therapy. ☐ For continuing Botulism toxin therapy the beneficiary demonstrates a significant decrease in the number and frequency of headaches and an improvement in function upon receiving Botulinum toxin
Medication(s) that have been attempted for chronic migraines with at least a 2-month trial. Provider Education Note: The beneficiary has had a trial of and inadequate response to a 2-month trial of at least one agent in any two of the following classes or has contraindication to the following medications. *	☐ Tricyclic Antidepressants ☐ Venlafaxine ☐ Nortriptyline ☐ Duloxetine ☐ Beta Blocker ☐ Propranolol ☐ Timolol (oral) ☐ Metoprolol ☐ Atenolol ☐ Nadolol ☐ Nebivolol ☐ Calcium Channel Blocker ☐ Verapamil ☐ Antiepileptic ☐ Valproate sodium ☐ Divalproex sodium ☐ Topiramate ☐ Gabapentin ☐ Other
Calcitonin gene-related peptide agent for chronic migraine prophylaxis is currently used. *	☐ NA ☐ Yes, and the beneficiary has had a reduction in the overall number of migraine days but still has chronic migraines requiring additional therapy for chronic migraine prevention. ☐ Yes, and the beneficiary has had a reduction in number of severe migraine days per month with CGRP use, but still has chronic migraines requiring additional therapy for chronic migraine prevention.
Specific information is summarized about the conservative methods of treatments such as medication, physical therapy, and other appropriate methods that have been tried and proven unsuccessful. Must include an objective assessment of the intolerance or inadequacy of the response(s). *	Type a value
Objective scale score of the severity of pain, for Chronic Migraines e.g. Numeric Pain Scale, Verbal Descriptor. *	☐ 0 - None (NRS 0) ☐ 1 - Mild (NRS 1 - 3) ☐ 2 - Moderate (NRS 4 - 6) ☐ 3 - Severe (NRS 7 - 9) ☐ 4 –Extreme, worst possible (NRS 10) ☐ Not Applicable
Serotype and Dosage	
Which serotype will be used for the procedure and total dose: *	☐ J0585: Injection, onabotulinumtoxinA(Botox), 1 unit ☐ J0586: Injection abobotulinumtoxinA (Dysport), 5 units ☐ J0587: Injection rimabotulinumtoxinB (Myobloc), 100 units ☐ J0588: Injection incobotulinumtoxinA (Xeomin), 1 unit
Total dose (units) *	☐ 50 units per vial, vacuum-dried powder for reconstitution ☐ 100 units per vial, vacuum-dried powder for reconstitution ☐ 200 units per vial, vacuum-dried powder for reconstitution ☐ 300 units per vial, vacuum-dried powder for reconstitution ☐ 500 units per vial, vacuum-dried powder for reconstitution ☐ 2500 units per vial, vacuum-dried powder for reconstitution
Total dose (units) *	
Total dose (units) *	

	5000 units per vial, vacuum-dried powder for reconstitution 10000 units per vial, vacuum-dried powder for reconstitution
--	--